



**UvA-DARE (Digital Academic Repository)**

**Solubility constraints on aquatic ecotoxicity testing of anionic surfactants**

Hammer, J.; Tukker, A.M.; Postma, J.F.; Haftka, J.J.-H.; Hermens, J.L.M.; de Voogt, P.; Kraak, M.H.S.

*Published in:*  
Bulletin of environmental contamination and toxicology

*DOI:*  
[10.1007/s00128-018-2361-1](https://doi.org/10.1007/s00128-018-2361-1)

[Link to publication](#)

*Citation for published version (APA):*  
Hammer, J., Tukker, A. M., Postma, J. F., Haftka, JJ-H., Hermens, J. L. M., de Voogt, P., & Kraak, M. H. S. (2018). Solubility constraints on aquatic ecotoxicity testing of anionic surfactants. Bulletin of environmental contamination and toxicology, 101(1), 99–104. <https://doi.org/10.1007/s00128-018-2361-1>

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.



# Solubility Constraints on Aquatic Ecotoxicity Testing of Anionic Surfactants

J. Hammer<sup>1,2</sup> · A. M. Tukker<sup>1,2</sup> · J. F. Postma<sup>3</sup> · J. J. -H. Haftka<sup>2</sup> · J. L. M. Hermens<sup>2</sup> · P. de Voogt<sup>1,4</sup> · M. H. S. Kraak<sup>1</sup>

Received: 22 January 2018 / Accepted: 14 May 2018 / Published online: 1 June 2018  
© The Author(s) 2018

## Abstract

In order to develop models that can predict the environmental behavior and effects of chemicals, reliable experimental data are needed. However, for anionic surfactants the number of ecotoxicity studies is still limited. The present study therefore aimed to determine the aquatic ecotoxicity of three classes of anionic surfactants. To this purpose we subjected daphnids (*Daphnia magna*) for 48 h to alkyl carboxylates ( $C_xCO_2^-$ ), alkyl sulfonates ( $C_xSO_3^-$ ), and alkyl sulfates ( $C_xSO_4^-$ ) with different carbon chain lengths ( $x$ ). However, all surfactants with  $x > 11$  showed less than 50% immobility at water solubility. Hence,  $EC_{50}$  values for only few surfactants could be gathered:  $C_9CO_2^-$  ( $16 \text{ mg L}^{-1}$ ),  $C_{11}CO_2^-$  ( $0.8 \text{ mg L}^{-1}$ ) and  $C_{11}SO_4^-$  ( $13.5 \text{ mg L}^{-1}$ ). Data from these compounds showed an increase in ecotoxicity with a factor 4.5 per addition of a hydrocarbon unit to the alkyl chain, and a factor 20 when replacing the sulfate head group by a carboxylate head group. Unfortunately, we could not test carboxylates with a broader variety of chain lengths because solubility limited the range of chain length that can be tested.

**Keywords** Aquatic ecotoxicity · *Daphnia* · Solubility · Anionic surfactants · Alkyl sulfonates · Alkyl sulfates · Alkyl carboxylates

Numerous new organic chemicals are produced yearly for application in industry and consumer products (CEFIC 2014). For environmental risk assessment of new and existing chemicals, an understanding of their environmental behavior and effects is required, but for anionic surfactants the number of ecotoxicity studies is still limited. For the development of predictive models such as quantitative structure–activity relationships (QSARs) for surfactants, more experimental data for these group of compounds are therefore needed. Although some toxicity tests on surfactants have been performed thus far (Schüürmann 1990; Roberts 1991; Versteeg et al. 1997; Wong et al. 1997; Dyer et al.

2000; Roberts and Costello 2003; Boeije et al. 2006; Hodges et al. 2006; Qi et al. 2011), the data is still too limited to compare the effect between different surfactant groups (i.e., surfactants with different head group structures), certainly for anionic surfactants. In this study, we therefore focused on generating aquatic ecotoxicity data for anionic surfactants from three different surfactant groups.

Anionic surfactants are high production volume chemicals which are present in many consumer products and consequently also in the environment (Sanderson et al. 2006; CEFIC 2014). Their amphiphilic and electrostatic properties make them very efficient compounds for the detergent industry. At the same time, these properties result in a very different environmental behavior compared to e.g. neutral organic compounds (Jones et al. 2003; Guo and Gaiki 2005). Unlike for common neutral organic pollutants, their accumulation and potential effects can therefore not always be correlated with predicted octanol–water partition constants ( $\log K_{ow}$ ) (Tolls and Sijm 1995).

The ecotoxicity of organic compounds (quantified by the concentration causing a 50% effect;  $EC_{50}$  value) is usually determined in standardized *Daphnia magna* acute ecotoxicity tests according to OECD guideline 202 (OECD 2004). For some surfactants within a specific surfactant group

✉ J. Hammer  
jorhammer@gmail.com

<sup>1</sup> Institute for Biodiversity and Ecosystem Dynamics (IBED), University of Amsterdam, P.O. Box 94248, 1090 GE Amsterdam, The Netherlands

<sup>2</sup> Institute for Risk Assessment Sciences, Toxicology Division, Utrecht University, P.O. Box 80177, 3508 TD Utrecht, The Netherlands

<sup>3</sup> Ecofide, Singel 105, 1381 AT Weesp, The Netherlands

<sup>4</sup> KWR Watercycle Research Institute, P.O. Box 1072, 3430 BB Nieuwegein, The Netherlands

(i.e., homologues sharing the same head group), toxicity is observed to increase with increasing alkyl chain length due to increased hydrophobicity (Roberts 2000; Roberts et al. 2013; Barmantlo et al. 2015). At the same time, hydrophobicity affects the bioavailability of surfactants by decreasing the solubility, but also by increasing sorption to other phases (Pittinger et al. 1989). Bioavailability of anionic surfactants is also influenced by the electrostatic characteristics of the head group, which can result in ion-pairing with divalent inorganic cations (e.g.,  $\text{Ca}^{2+}$  or  $\text{Mg}^{2+}$ ) (Rodriguez et al. 2001; Yan et al. 2010). The standard medium in the *D. magna* toxicity test (OECD 2004) contains a relatively high total ionic strength that includes divalent cations and solubility problems can therefore be expected for some surfactants. The determination of  $\text{EC}_{50}$  values for (ionic) compounds with a low solubility using OECD guideline 202 can therefore be challenging. However, since experimental data for anionic surfactants is still much needed, the aim of the present study was to employ the standardized *D. magna* ecotoxicity test to determine the aquatic ecotoxicity of three classes of anionic surfactants: alkyl carboxylates, alkyl sulfonates, and alkyl sulfates.

## Materials and Methods

All test compounds had a typical surfactant structure containing a hydrophobic alkyl chain and a hydrophilic ionized head group. Sodium salts of linear alkyl sulfates ( $\text{C}_x\text{SO}_4^-$ ; with alkyl chain lengths  $\text{C}_{11}$ ,  $\text{C}_{13}$ ,  $\text{C}_{15}$  and  $\text{C}_{16}$ ) and linear alkyl sulfonates ( $\text{C}_x\text{SO}_3^-$ ;  $\text{C}_{11}$ ,  $\text{C}_{13}$ ,  $\text{C}_{14}$  and  $\text{C}_{15}$ ) were obtained from Research Plus (South Plainfield, NJ). Sodium salts of linear alkyl carboxylates ( $\text{C}_x\text{CO}_2^-$ ;  $\text{C}_9$ ,  $\text{C}_{11}$ ,  $\text{C}_{13}$ ,  $\text{C}_{14}$ , and  $\text{C}_{15}$ ) were obtained from Sigma-Aldrich, (Zwijndrecht, The Netherlands). All organic compounds had purities higher than 98%. Ammonium acetate was purchased from Sigma-Aldrich. Methanol was obtained from Biosolve (Valkenswaard, The Netherlands). Ultrapure water was obtained from a Millipore water purification system (resistivity > 18  $\text{M}\Omega/\text{cm}$ , Merck Chemicals, Amsterdam, The Netherlands).

The daphnid *D. magna* Straus was selected as test organism to determine the aquatic ecotoxicity of surfactants. Juvenile daphnids (clone 4) aged < 24 h were obtained from adults between 2 and 3 weeks old. Continuous cultures were maintained in Elendt M4 medium and fed with the alga *Chlorella vulgaris*. At regular intervals (about every 3 months), acute toxicity tests were performed with the reference toxicant  $\text{K}_2\text{Cr}_2\text{O}_7$  to check whether the sensitivity of the daphnids culture was within the limits ( $\text{EC}_{50}$ , 24 h = 0.6–2.1  $\text{mg L}^{-1}$ ) as set by the guideline (OECD 2004). The medium used in the toxicity experiments consisted of the standard OECD medium that was prepared according to

OECD guideline 202, containing 266  $\text{mg L}^{-1}$   $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ , and 112  $\text{mg L}^{-1}$   $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ . Concentrations of KCl and  $\text{NaHCO}_3$  were 5 and 65  $\text{mg L}^{-1}$  respectively (OECD 2004). The test media was buffered to pH  $7 \pm 0.3$  with NaOH (66  $\text{mg L}^{-1}$ ) and 3-(*N*-morpholino)propanesulfonic acid (MOPS; 1.046  $\text{g L}^{-1}$ ).

The *D. magna* were exposed to the selected compounds in 48 h immobility tests (OECD 2004). Per experiment five test concentrations, a solvent control (0.25% methanol without the test compound) and a control were tested with four replicates per treatment. Each replicate consisted of a glass tube filled with 20 mL of test solution, spiked with 50  $\mu\text{L}$  (0.25% of total volume) methanol containing the test compound. The tubes were randomly distributed in a climate controlled fume hood ( $20 \pm 1^\circ\text{C}$ ), with a light–dark regime of 16:8 h. The experiment was started by introducing five neonates (younger than 24 h) into each tube. After 48 h, the number of animals not responding to stimulation was scored. Hardness, oxygen concentration, temperature and pH were measured at the start and the end of the experiments and were within the range prescribed by OECD guideline 202 (OECD 2004). The concentration of the test compounds was analyzed by extracting a 200  $\mu\text{L}$  water sample from each replicate at the start and the end of the experiment, an injection standard was added and the sample was subsequently diluted with 750  $\mu\text{L}$  of methanol and stored in a freezer ( $-18^\circ\text{C}$ ) until chemical analysis.

All anionic surfactants were detected with a triple quadrupole mass spectrometer (MDS SCIEX API 3000 MS/MS System from Applied Biosystems, Bleiswijk, The Netherlands) with a Turbo Ion spray source operated at  $400^\circ\text{C}$ . A solvent delay switch (Da Vinci, Rotterdam, The Netherlands) was used to prevent introduction of inorganic constituents from water samples into the MS. Chromatograms were integrated with Analyst 1.4.2 (Applied Biosystems). Concentration–response relationships and the corresponding 48 h  $\text{EC}_{50}$  values were calculated according to Haanstra et al. (1985) by fitting a logistic curve (Eq. 1) to the percentage of mobility (100% – immobilization) versus the surfactant concentration in the water phase.

$$y(x) = \frac{C}{1 + e^{b(\log_{10}x - \log_{10}a)}} \quad (1)$$


where  $y(x)$  is the mobility at concentration  $x$  (in %),  $a$  is the  $\text{EC}_{50}$  value (in  $\text{mg L}^{-1}$ ),  $b$  is the slope of the curve,  $c$  is  $y(0)$  which equals the average mobility of the control and  $x$  is the surfactant concentration in water (in  $\text{mg L}^{-1}$ ). Data analyses were performed with SPSS software (IBM Corp 2013) and Graphpad Prism Version 7.0 (GraphPad Software 2017).

## Results and Discussion

A total of 14 surfactants with varying alkyl chain lengths from three surfactant groups (alkyl sulfates, alkyl sulfonates, and alkyl carboxylates) were tested. Due to their hydrophobicity and electrostatic charge, anionic surfactants with long alkyl chains often poorly dissolve in water containing inorganic cations. We therefore decided to first test the effect of saturated water solutions at maximum aqueous solubility ( $S_w$ ) on the daphnids. To this end we stirred an excess of compound for 48 h in standard OECD medium under the standard conditions of the *D. magna* toxicity tests (OECD 2004). For the compounds that caused more than 50% immobility of the daphnids at  $S_w$ , a concentration range was tested in order to obtain concentration–response relationships and to derive  $EC_{50}$  values.

We were unable to dissolve alkyl sulfonates ( $C_xSO_3^-$ ) in the OECD medium at sufficiently high concentrations to cause any effect. This may have been a result of the presence of (divalent) cations in the aqueous phase. Cations are known to affect the hydration of anionic surfactants and often lowers their critical micelles concentration (CMC) (Yan et al. 2010). Divalent cations such as  $Ca^{2+}$  and  $Mg^{2+}$  can furthermore form ion pairs containing two surfactant monomers and one divalent cation, or form bridges between monomers and charged sites on sorbents (Haftka et al. 2015). For the alkyl carboxylates ( $C_xCO_2^-$ ) and the alkyl sulfates ( $C_xSO_4^-$ ), compounds with an alkyl chain longer than  $C_{11}$  were badly soluble in the OECD medium and showed less than 50% immobility at  $S_w$ . Hence, no further ecotoxicity tests were performed for these compounds.

Because of the solubility problems of the tested compounds in the OECD medium,  $EC_{50}$  values for only few anionic surfactants could be generated:  $C_9CO_2^-$ ,  $C_{11}CO_2^-$  and  $C_{11}SO_4^-$ . Because one pair of these surfactants contains equal alkyl chain lengths and different surfactant head groups ( $C_{11}CO_2^-$  and  $C_{11}SO_4^-$ ), and another pair ( $C_9CO_2^-$  and  $C_{11}CO_2^-$ ) differs in chain length with equal head group, we had two single opportunities to evaluate the effect of head group structure and alkyl chain length on the toxicity of the anionic surfactants. However, note that these interpretations are based on only a single pair of surfactants. For  $C_9CO_2^-$ ,  $C_{11}SO_4^-$  and  $C_{11}CO_2^-$  analyzed concentrations were respectively  $\pm 10\%$ ,  $\pm 10\%$  and  $\pm 30\%$  lower compared to nominal concentrations. During the 48 h *D. magna* toxicity experiments 100% control survival was recorded. From the dose–response curve of  $C_{11}SO_4^-$  an  $EC_{50}$  value of  $13.5 \text{ mg L}^{-1}$  was derived (95% CI  $13.2\text{--}13.8 \text{ mg L}^{-1}$ ) (Fig. 1). We were unable to find any  $EC_{50}$  values of  $C_{11}SO_4^-$  in literature as




**Fig. 1** Effect of head group on ecotoxicity of  $C_{11}SO_4^-$  and  $C_{11}CO_2^-$  to *Daphnia magna* after 48 h of exposure. Both dose–response curves were calculated according to Haanstra et al. (1985). The  $EC_{50}$  concentrations are plotted with their 95% confidence intervals as solid black symbols (the 95% confidence interval of  $C_{11}SO_4^-$  is too small to be seen)

most studies focused on  $C_{12}SO_4^-$ . Persoone et al. (1989) reported an  $EC_{50}$  value of  $9.6 \text{ mg L}^{-1}$  for  $C_{12}SO_4^-$  in a *D. magna* 24 h toxicity test and Dyer et al. (1996) found an  $EC_{50}$  value of  $5.5 \text{ mg L}^{-1}$  in a 48 h *Ceriodaphnia dubia* toxicity test (comparable sensitivity to *D. magna* (Versteeg et al. 1997)). Both values are in line with our data for  $C_{11}SO_4^-$ , as toxicity generally increases from 24 to 48 h exposure and an  $EC_{50}$  value of  $5.5 \text{ mg L}^{-1}$  is close to the expected  $EC_{50}$  concentration increase when a hydrocarbon ( $-CH_2-$ ) unit is added to the alkyl chain of  $C_{11}SO_4^-$  (see next paragraph). The dose–response curve of  $C_{11}CO_2^-$  provided an  $EC_{50}$  concentration of  $0.80 \text{ mg L}^{-1}$  (95% CI  $0.7\text{--}0.9 \text{ mg L}^{-1}$ ) (Fig. 1). Toxicity data for *D. magna* are scarce for  $C_{11}CO_2^-$ , a 36x higher  $EC_{50}$  value ( $EC_{50} = 29 \text{ mg L}^{-1}$ ) was reported by Lundahl and Cabridenc (1978) in a 24 h ecotoxicity test, and an  $EC_{50}$  value of  $1.3 \text{ mg L}^{-1}$  was reported by the European Chemical Agency (2014). While, we were unable to acquire the exact experimental details of the toxicity test of Lundahl and Cabridenc (1978), their analysis was performed using the Methylene Blue Active Substance (MBAS) assay which is meanwhile retracted as a standard method by ASTM.

Comparing the dose–response curves and  $EC_{50}$  values for  $C_{11}SO_4^-$  and  $C_{11}CO_2^-$  shows that the head group has a significant effect on ecotoxicity (Fig. 1). The alkyl chains of both compounds are of the same length and the effect of hydrophobicity is subsequently similar (Hammer et al. 2017). Therefore, the difference in  $EC_{50}$  values is likely a result from the different molecular properties of the surfactant head groups ( $SO_4^-$  vs.  $CO_2^-$ ). Besides the head group structure, the most notable distinction between the properties of these two surfactant groups is the difference in  $pK_a$  [4.8 for  $C_xCO_2^-$  (Haynes 2015), and  $-3.6$  for

$C_xSO_4^-$  (COSMOlogic 2015)]. The  $pK_a$  value is partly a result of the charge distribution over a molecule and shows what fraction of the compound is in the ionic form at certain pH. While these compounds are in the OECD medium both for  $>99\%$  present in their ionic (de-protonated) form, the difference charge distribution between both molecules still affects their behavior in the aqueous phase and their interaction with other phases. For example, alkyl carboxylates are much better hydrated than alkyl sulfates (Vlachy et al. 2009), which also affects their electrostatic interaction with sorbents (Rabin and Stillian 1994). Furthermore, the difference in charge distribution may affect the uptake of the anionic surfactants in cell membranes due to their zwitterionic properties (Scherer and Seelig 1989). Badly hydrated compounds are usually more affected by local charges and have more difficulty to partition into membranes than well hydrated compounds (Jing et al. 2009; Roberts et al. 2013). The  $C_{11}CO_2^-$  surfactant may therefore partition more effectively into cell membranes of the daphnids compared to  $C_{11}SO_4^-$  which explains why alkyl carboxylates were approximately 20 times more toxic compared to their sulfated counterparts.

The effect of the alkyl chain length on surfactant toxicity was studied by comparing the  $EC_{50}$  values of  $C_9CO_2^-$  and  $C_{11}CO_2^-$ . The dose–response curve for  $C_9CO_2^-$  showed an  $EC_{50}$  concentration of  $16.0\text{ mg L}^{-1}$  (95% CI  $14.8\text{--}17.3\text{ mg L}^{-1}$ ), see Fig. 2. Just like for the previously discussed surfactants, literature data on the toxicity of  $C_9CO_2^-$  to *D. magna* is inconsistent and details about the experimental setup were difficult to obtain. We were able to find two  $EC_{50}$  concentrations from literature: first, again a very high  $EC_{50}$  concentration of  $135\text{ mg L}^{-1}$  from a 24 h *D. magna* toxicity test by Lundahl and Cabridenc (1978). Second, a reported  $EC_{50}$  value of  $16\text{ mg L}^{-1}$ , which is equal to our experimentally derived  $EC_{50}$  value and originates from a



**Fig. 2** Effect of alkyl chain length on ecotoxicity of  $C_{11}CO_2^-$  and  $C_9CO_2^-$  on *Daphnia magna* after 48 h of exposure. Both dose–response curves were calculated according to Haanstra et al. (1985). The  $EC_{50}$  concentrations are plotted with their 95% confidence intervals as solid black symbols

report of the European Chemical Agency (2013). The results from Lundahl and Cabridenc are questionable (see previous paragraph) and both studies lack experimental details about medium composition and only mention the duration of the tests. Toxicity between  $C_{11}$  and  $C_9$  carboxylate differed with a factor of  $\sim 23$  compared (Fig. 2), which is a factor of  $\sim 4.5$  per hydrocarbon unit added to the alkyl chain. This is somewhat higher than the increments found for other surfactant groups in previous studies [between 2.4 and 3.4 (Lundahl and Cabridenc 1978; Maki and Bishop 1979; Hodges et al. 2006)]. An increase in the alkyl chain length increases the hydrophobicity of the compound and thus increases the sorption to the membrane lipid (Könnecker et al. 2011). At longer alkyl chain lengths ( $>C_{11}$ ) the toxicity is expected to further increase, but this effect is not detectible using the *D. magna* toxicity test due the low solubility of the compounds in the OECD medium. The factor  $\sim 4.5$  increase in toxicity with addition of a carbon atom to the alkyl chain is based on only two chemicals. This data set is limited and could be regarded as a shortcoming of the study. Unfortunately, we could not test more compounds because of the solubility problems (limits) of the longer chain carboxylates in the calcium rich test medium of the *Daphnia* test. Another test organism that requires another medium composition (less calcium) could avoid this shortcoming.

The main reason why ecotoxicity could not be detected for most of the test compounds probably lies in the presence of cations in the aqueous solution of the *D. magna* tests, which can affect the solubility and bioavailability of anionic surfactants. In an attempt to generate more ecotoxicity data, we decided to change the composition of the original OECD medium and study the effect of divalent cation concentration on the ecotoxicity of  $C_9CO_2^-$  and  $C_{11}CO_2^-$ . To this end, four different media were prepared with different concentrations of  $Ca^{2+}$  and  $Mg^{2+}$ , while maintaining original  $Ca^{2+}:Mg^{2+}$  ratio (Naddy et al. 2002). A concentration of  $Ca^{2+}$  of  $10\text{ mg L}^{-1}$  was selected as the lowest concentration, because lower concentrations will affect with *D. magna* survival (Hessen et al. 2000). The highest concentration of  $Ca^{2+}$  tested was  $80\text{ mg L}^{-1}$ , conform with the original OECD guideline 202. The resulting  $EC_{50}$  concentrations varied slightly, but did not differ significantly between medium compositions. Hence, the medium with the lowest ionic strength may already contain enough cations to cause precipitation of anionic surfactants.

The *D. magna* toxicity test is a well-accepted and standardized toxicity test which has generated ecologically relevant toxicity data for many organic compounds. However, the medium proposed in the OECD guideline for *D. magna* is of high ionic strength and this can result in solubility problems for compounds that are already barely soluble in water and for compounds that maintain an electrostatic charge (Waijers et al. 2013). The *D. magna* toxicity test appeared

unable to produce reliable results for most of the surfactants tested in this study. For hazard assessment purposes of anionic surfactants, alternative approaches should therefore be investigated that either exclude the influence of divalent cations present in the test medium or endpoints should be selected that are affected at concentrations below the aqueous solubility of the surfactants. Furthermore, because anionic surfactants are known to have an affinity for soil surfaces and organic matter (Rico–Rico 2009) toxicity tests that include sediment living organisms (e.g. *Lumbriculus variegatus* or *Chironomus riparius*) may be more suitable for the production of toxicological endpoint data. Despite the obstacles that occurred with anionic surfactants during the *D. magna* tests, we were able to determine the effect of surfactant alkyl chain length and head group composition on the aquatic ecotoxicity of a select group of anionic surfactants. However, these interpretations were based on only a single pair of surfactants.

**Acknowledgements** The present study was supported by the Dutch Technology Foundation STW, which is part of The Netherlands Organization for Scientific Research and is partly funded by the Ministry of Economic Affairs (Stichting voor de Technische Wetenschappen). Additional funding was received from Deltares (Utrecht, The Netherlands) and Environmental Risk Assessment and Management (ERASM), which is a partnership of European detergent and surfactant products. We would like to thank BSc students Tessa de Bruin, Remy Mulders, and Linde de Herder from the University of Amsterdam for their support with the experimental work and Rineke Keijzers from Ecofide for supplying the daphnids.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Barmantlo SH, Stel JM, van Doorn M et al (2015) Acute and chronic toxicity of short chained perfluoroalkyl substances to *Daphnia magna*. *Environ Pollut* 198:47–53. <https://doi.org/10.1016/j.envpol.2014.12.025>
- Boeije GMG, Cano ML, Marshall SJ et al (2006) Ecotoxicity quantitative structure–activity relationships for alcohol ethoxylate mixtures based on substance-specific toxicity predictions. *Ecotoxicol Environ Saf* 64:75–84. <https://doi.org/10.1016/j.ecoenv.2005.08.009>
- CEFIC (2014) The European Chemical Industry Council. <http://www.cefic.org/Facts-and-Figures/Chemicals-Industry-Profile/>. Accessed 1 Jan 2018
- COSMOlogic GmbH (2015) COSMOtherm. pp 1–77
- Dyer SD, Lauth JR, Morrall SW et al (1996) Development of a chronic toxicity structure–activity relationship for alkyl sulfates. *Environ Toxicol Water* 295–303
- Dyer SD, Stanton DT, Lauth JR, Cherry DS (2000) Structure–activity relationships for acute and chronic toxicity of alcohol ether sulfates. *Environ Toxicol Chem* 19:608–616. <https://doi.org/10.1002/etc.5620190312>
- European Chemical Agency (2013) Regulation (EU) No. 528/2012 concerning decanoic acid
- European Chemical Agency (2014) Regulation (EU) No. 528/2012 concerning lauric acid
- GraphPad S (2017) GraphPad Prism version 7.00 for Windows
- Guo Y, Gaiki S (2005) Retention behavior of small polar compounds on polar stationary phases in hydrophilic interaction chromatography. *J Chromatogr A* 1074:71–80. <https://doi.org/10.1016/j.chroma.2005.03.058>
- Haanstra L, Doelman P, Voshaar JHO (1985) The use of sigmoidal dose response curves in soil ecotoxicological research. *Plant Soil* 84:293–297. <https://doi.org/10.1007/BF02143194>
- Haftka JJH, Hammer J, Hermens JLM (2015) Mechanisms of neutral and anionic surfactant sorption to solid-phase microextraction fibers. *Environ Sci Technol* 49:11053–11061. <https://doi.org/10.1021/acs.est.5b02901>
- Hammer J, Haftka JJ-H, Scherpenisse P et al (2017) Fragment-based approach to calculate hydrophobicity of anionic and nonionic surfactants derived from chromatographic retention on a C18 stationary phase. *Environ Toxicol Chem* 36:329–336. <https://doi.org/10.1002/etc.3564>
- Haynes WM (2015) CRC handbook of chemistry and physics, 96th edn. CRC Press, Florida
- Hessen DO, Alstad NEW, Skardal L (2000) Calcium limitation in *Daphnia magna*. *J Plankton Res* 22:553–568. <https://doi.org/10.1093/plankt/22.3.553>
- Hodges G, Roberts DW, Marshall SJ, Dearden JC (2006) The aquatic toxicity of anionic surfactants to *Daphnia magna*—a comparative QSAR study of linear alkylbenzene sulphonates and ester sulphonates. *Chemosphere* 63:1443–1450. <https://doi.org/10.1016/j.chemosphere.2005.10.001>
- IBM Corp (2013) IBM SPSS statistics for Windows version 22.0.0. IBM Corp, Armonk
- Jing P, Rodgers PJ, Amemiya S (2009) High lipophilicity of perfluoroalkyl carboxylate and sulfonate: Implications for their membrane permeability. *J Am Chem Soc* 131:2290–2296. <https://doi.org/10.1021/ja807961s>
- Jones PD, Hu W, De Coen W et al (2003) Binding of perfluorinated fatty acids to serum proteins. *Environ Toxicol Chem* 22:2639–2649. <https://doi.org/10.1897/02-553>
- Könnecker G, Regelman J, Belanger S et al (2011) Environmental properties and aquatic hazard assessment of anionic surfactants: physico-chemical, environmental fate and ecotoxicity properties. *Ecotoxicol Environ Saf* 74:1445–1460. <https://doi.org/10.1016/J.ECOENV.2011.04.015>
- Lundahl P, Cabridenc R (1978) Molecular structure–biological properties relationships in anionic surface-active agents. *Water Res* 12:25–30. [https://doi.org/10.1016/0043-1354\(78\)90191-4](https://doi.org/10.1016/0043-1354(78)90191-4)
- Maki AW, Bishop WE (1979) Acute toxicity studies of surfactants to *Daphnia magna* and *Daphnia pulex*. *Arch Environ Contam Toxicol* 8:599–612. <https://doi.org/10.1007/BF01055040>
- Naddy RB, Stubblefield WA, May JR et al (2002) The effect of calcium and magnesium ratios on the toxicity of copper to five aquatic species in freshwater. *Environ Toxicol Chem* 21:347–352
- OECD (2004) Test No. 202: *Daphnia* sp. acute immobilisation test. OECD Publishing, Paris
- Persoone G, Van de Vel A, Van Steertegem M, De Nayer B (1989) Predictive value of laboratory tests with aquatic invertebrates: influence of experimental conditions. *Aquat Toxicol* 14:149–167. [https://doi.org/10.1016/0166-445X\(89\)90025-8](https://doi.org/10.1016/0166-445X(89)90025-8)
- Pittinger CA, Woltering DM, Masters JA (1989) Bioavailability of sediment-sorbed and aqueous surfactants to *Chironomus*

- riparius* (midge). *Environ Toxicol Chem* 8:1023–1033. <https://doi.org/10.1002/etc.5620081108>
- Qi P, Wang Y, Mu J, Wang J (2011) Aquatic predicted no-effect-concentration derivation for perfluorooctane sulfonic acid. *Environ Toxicol Chem* 30:836–842. <https://doi.org/10.1002/etc.460>
- Rabin S, Stillian J (1994) Practical aspects on the use of organic solvents in ion chromatography. *J Chromatogr A* 671:63–71. [https://doi.org/10.1016/0021-9673\(94\)80222-X](https://doi.org/10.1016/0021-9673(94)80222-X)
- Rico-Rico A (2009) Linear alkylbenzene sulfonates in the aquatic environment: study of the analysis, sorption processes and sediment toxicity. Utrecht University, Utrecht
- Roberts DW (1991) QSAR issues in aquatic toxicity of surfactants. *Sci Total Environ* 109–110:557–568. [https://doi.org/10.1016/0048-9697\(91\)90209-W](https://doi.org/10.1016/0048-9697(91)90209-W)
- Roberts DW (2000) Aquatic toxicity—are surfactant properties relevant? *J Surfactants Deterg* 3:309–315. <https://doi.org/10.1007/s11743-000-0134-z>
- Roberts DW, Costello J (2003) QSAR and mechanism of action for aquatic toxicity of cationic surfactants. *QSAR Comb Sci* 22:220–225. <https://doi.org/10.1002/qsar.200390015>
- Roberts DW, Roberts JF, Hodges G et al (2013) Aquatic toxicity of cationic surfactants to *Daphnia magna*. *SAR QSAR Environ Res* 24:417–427. <https://doi.org/10.1080/1062936X.2013.781538>
- Rodriguez CH, Lowery LH, Scamehorn JF, Harwell JH (2001) Kinetics of precipitation of surfactants. I. Anionic surfactants with calcium and with cationic surfactants. *J Surfactants Deterg* 4:1–14. <https://doi.org/10.1007/s11743-001-0155-7>
- Sanderson H, Dyer SD, Price BB et al (2006) Occurrence and weight-of-evidence risk assessment of alkyl sulfates, alkyl ethoxysulfates, and linear alkylbenzene sulfonates (LAS) in river water and sediments. *Sci Total Environ* 368:695–712. <https://doi.org/10.1016/j.scitotenv.2006.04.030>
- Scherer PG, Seelig J (1989) Electric charge effects on phospholipid headgroups. Phosphatidylcholine in mixtures with cationic and anionic amphiphiles. *Biochemistry* 28:7720–7728. <https://doi.org/10.1021/bi00445a030>
- Schüürmann G (1990) QSAR analysis of the acute toxicity of oxyethylated surfactants. *Chemosphere* 21:467–478. [https://doi.org/10.1016/0045-6535\(90\)90017-N](https://doi.org/10.1016/0045-6535(90)90017-N)
- Tolls J, Sijm DTHM (1995) A preliminary evaluation of the relationship between bioconcentration and hydrophobicity for surfactants. *Environ Toxicol Chem* 14:1675–1685. <https://doi.org/10.1002/etc.5620141007>
- Versteeg DJ, Stanton DT, Pence MA, Cowan C (1997) Effects of surfactants on the rotifer *Brachionus calyciflorus* in a chronic toxicity test and in the development of QSARs. *Environ Toxicol Chem* 16:1051–1058. <https://doi.org/10.1002/etc.5620160527>
- Vlachy N, Jagoda-Cwiklik B, Vácha R et al (2009) Hofmeister series and specific interactions of charged headgroups with aqueous ions. *Adv Colloid Interface Sci* 146:42–47. <https://doi.org/10.1016/j.cis.2008.09.010>
- Waijers SL, Hartmann J, Soeter AM et al (2013) Toxicity of new generation flame retardants to *Daphnia magna*. *Sci Total Environ* 463–464:1042–1048. <https://doi.org/10.1016/j.scitotenv.2013.06.110>
- Wong DCL, Dorn PB, Chai EY (1997) Acute toxicity and structure-activity relationships of nine alcohol ethoxylate surfactants to fathead minnow and *Daphnia magna*. *Environ Toxicol Chem* 16:1970–1976. <https://doi.org/10.1002/etc.5620160929>
- Yan H, Yuan S-L, Xu G-Y, Liu C-B (2010) Effect of Ca<sup>2+</sup> and Mg<sup>2+</sup> ions on surfactant solutions investigated by molecular dynamics simulation. *Langmuir* 26:10448–10459. <https://doi.org/10.1021/la100310w>