UvA-DARE (Digital Academic Repository)

Stress, hippocampal neurogenesis and cognition: functional correlations

Lucassen, P.J.; Oomen, C.A.

DOI

[10.1007/s11515-016-1412-4](https://doi.org/10.1007/s11515-016-1412-4)

Publication date

2016

Document Version

Final published version

Published in

Frontiers in Biology

License

Article 25fa Dutch Copyright Act

[Link to publication](#)

Citation for published version (APA):

Lucassen, P. J., & Oomen, C. A. (2016). Stress, hippocampal neurogenesis and cognition: functional correlations. *Frontiers in Biology*, 11(3), 182-192. <https://doi.org/10.1007/s11515-016-1412-4>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Stress, hippocampal neurogenesis and cognition: functional correlations

Paul J. Lucassen (✉)¹, Charlotte A. Oomen^{1,2}

¹ Centre for Neuroscience, Swammerdam Institute for Life Sciences, University of Amsterdam, Science Park 904, 1098 XH, Amsterdam, The Netherlands

² Department of Molecular and Cellular Neurobiology, Center for Neurogenomics and Cognitive Research, Neuroscience Campus Amsterdam, VU University Amsterdam, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands

© Higher Education Press and Springer-Verlag Berlin Heidelberg 2016

Abstract The brain of many species including humans, harbors stem cells that continue to generate new neurons up into adulthood. This form of structural plasticity occurs in a limited number of brain regions, i.e. the subventricular zone and the hippocampal dentate gyrus and is regulated by environmental and hormonal factors. In this minireview, we provide an overview of the effects of stress and glucocorticoid hormones on adult hippocampal neurogenesis and discuss how these effects may be relevant for cognitive function and possibly, brain disease. While its exact functional role remains elusive, adult neurogenesis has been implicated in learning and memory, fear and mood regulation and recently, adult-born neurons were found to be involved in specific cognitive functions such as pattern separation (i.e. the ability to form unique memory representations) and cognitive flexibility. The process of adult neurogenesis is influenced by several factors; whereas e.g. exercise stimulates, exposure to stress and stress hormones generally inhibit neurogenesis. Effects of acute, mild stress are generally short-lasting and recover quickly, but chronic or severe forms of stress can induce lasting reductions in adult neurogenesis. Some of the inhibitory effects of stress can be rescued by exercise, by allowing a period of recovery from stress, by drugs that target the stress system, or by some, but not all, antidepressants. Stress may, partly through its effects on adult neurogenesis, alter structure and plasticity of the hippocampal circuit. This can lead to subsequent changes in stress responsivity and aspects of memory processing, which may be particularly relevant for stress related psychopathology or brain diseases that involve perturbed memory processing.

Keywords adult neurogenesis, stem cells, hippocampus, stress, memory, pattern separation, cognition

Stress and the stress response

In our daily life, exposure to stress is difficult to avoid. Stress is diverse in nature, duration or intensity; it can be psychological in nature, such as during relationship problems (Ursin and Eriksen, 2004) or can involve biological changes, such as the occurrence of an infection. Exposure to any stress generally elicits a stress response in the body and brain that enables the individual to adapt, maintain homeostasis, and ultimately promote survival. The subjective experience of

stress is highly individual, as is one's response to it, and this response depends largely on genetic make up, sex and personality traits. Gene-environment interactions, particularly during sensitive developmental periods, are further important in “programming” our hormonal and cognitive response to stress and also determine individual differences in vulnerability to disease, as reviewed before (Joëls et al., 2006, 2012; Meaney et al., 2007; Binder et al., 2008; Koolhaas et al., 2011; Lucassen et al., 2013b).

The endocrinological response to stress involves first the release of adrenal (nor)epinephrine that among others, increases blood flow to essential organs. Later, the hypothalamic-pituitary-adrenal (HPA) axis is activated, a classic neuroendocrine circuit that regulates the eventual behavioral, neural and hormonal response to stress (Joëls and Baram

Received June 2, 2016; accepted June 12, 2016

Correspondence: Paul J. Lucassen

E-mail: p.j.lucassen@uva.nl

2009; Joëls et al., 2012). Activation of the HPA-axis starts with hypothalamic corticotropin-releasing hormone (CRH) production, which subsequently results in the release of glucocorticoids (GCs) from the adrenal glands (GCs; corticosterone in rodents; cortisol in man). Negative feedback of GC-release occurs when GCs bind to high-affinity mineralocorticoid (MR) and lower affinity glucocorticoid receptors (GR) in key regions of the brain (de Kloet et al., 2005). Through this, GR activation helps to maintain GC levels within physiological limits (Kretz et al., 1999; Erdmann et al., 2008) and consequently, aberrant GR expression or changes in the GR/MR ratio have been implicated in hypercortisolism, hippocampal changes, stress resistance, anxiety and depression (de Kloet et al., 2005; Ridder et al., 2005; Wei et al., 2007; Wang et al., 2012; Vinkers et al., 2014). GC-release is pulsatile and under circadian and ultradian control (Qian et al., 2012; Lucassen et al., 2013b; Fitzsimons et al., 2016), and GCs act as transcription factors that control protein expression in a slow, genomic manner, however, faster non-genomic actions exist as well (Tasker 2006; Karst et al., 2010).

Upon their release in the periphery, GCs affect energy, inflammatory responses and lipid metabolism, among others. Imbalances in GC regulation can have deleterious consequences, particularly for the brain (de Kloet et al., 2005). Specifically the hippocampus, a region important for learning and memory, is sensitive to GC increases as it contains high levels of GRs (de Kloet et al., 2005; Swaab et al., 2005; Wang et al., 2013). Indeed, GCs influence memory, fear and attention in a negative manner, particularly when exposure to stress is chronic and uncontrollable. However, positive effects of stress-such as enhanced memory have also been described and effects of stress depend on timing, type and controllability of a stressor (Joëls et al., 2006; Schwabe et al., 2012).

While functional changes after stress involve reductions in hippocampal excitability, long-term potentiation and hippocampus-dependent memory, morphological consequences of stress include hippocampal volume reductions as well as a number of cellular changes, notably dendritic atrophy and a suppressed rate of adult neurogenesis (see below) (Sapolsky et al., 1990; Czéh and Lucassen, 2007; Lucassen et al., 2014). Given its peak values after 15 min, most GC effects on brain function occur minutes to hours after the stressful event rather than in the critical first few minutes after which a threat is detected. These slow effects, as well as its morphological effects suggest that an important role of GCs in the brain is to affect future behavior and responsivity.

Effects of stress on the hippocampus may help facilitate adaptation after stressful or challenging conditions, whereas a failure to adjust behavior in response to stress could have negative consequences. Ultimately, the stress response helps to (re)direct energy and focus attention to cope with the most urgent elements of a stressor, at the cost of less urgent bodily processes that are temporarily suppressed by stress (Joëls et al., 2012).

Stress modulates adult neurogenesis

Adult neurogenesis (AN) refers to stem cells present in adult brains that continue to produce new neurons up into old age. These stem cells undergo different stages of proliferation, fate specification, migration and neuronal differentiation, before they eventually become new, functional neurons that integrate into the pre-existing, adult network of the hippocampus (Abrous et al., 2005; Toni et al., 2008; Zhao et al., 2008; Kempermann, 2012; Vivar et al., 2012; Jessberger and Gage, 2014). It has also been reported to occur in other brain structures, like the amygdala, striatum, hypothalamus and neocortex, albeit under specific conditions and with considerable differences between species (Gould, 2007).

AN is dynamically regulated by various environmental factors and declines prominently with age in many species (e.g., (Kuhn et al., 1996; Heine et al., 2004b)). Neurogenesis is potently stimulated by exercise and environmental enrichment, which notably occurs parallel to improvements in hippocampal function (Kempermann et al., 2010; Vivar et al., 2013). A strong inhibitor of AN on the other hand, is stress and GC exposure (Balu and Lucki, 2009; Lucassen et al., 2010a) as will be discussed below in more detail.

Exposure to stress during adult life is one of the best known environmental suppressors of AN. Both psychosocial (Gould et al., 1997; Czéh et al., 2002) and physical stressors (Malberg and Duman, 2003; Pham et al., 2003; Vollmayr et al., 2003) can inhibit one or more phases of the neurogenesis process (Mirescu and Gould, 2006; Lucassen et al., 2010a). In classical studies, rodents exposed to the odor of a predator generated a strong stress hormone response that was associated with significant parallel reductions in hippocampal proliferation (Tanapat et al., 1998). Both acute and chronic stressors generally suppress proliferation and many different types of stressors, including physical restraint, social defeat, inescapable foot shock, sleep deprivation, and mixed types of multiple, unpredictable or mild stressors, generally all decrease numbers of new neurons in the dentate gyrus (Gould et al., 1997; Czéh et al., 2002; Pham et al., 2003; Heine et al., 2004a; Heine et al., 2004c; Simon et al., 2005; Dranovsky and Hen, 2006; Jayatissa et al., 2006; Mitra et al., 2006; Jayatissa et al., 2009; Perera et al., 2011; Lehmann et al., 2013; Schoenfeld and Gould, 2013; Wu et al., 2014).

Notably, exceptions exist in stress inhibition of neurogenesis and negative findings have also been reported (Dagyte et al., 2009; Lyons et al., 2010; Hanson et al., 2011a; Parihar et al., 2011; O'Leary et al., 2012). These might depend on the type of stressor applied, or the species, sex or strain used (Westenbroek et al., 2004; Hanson et al., 2011b; Schoenfeld and Gould, 2013; Kanatsou et al., 2015). Inter-individual variation in the behavioral susceptibility to stress is also a relevant factor (Levone et al., 2015). In some instances, increased AN has been reported after stress, but in these studies the stressors were often predictable, controllable and/

or mild, and may actually have enriched standard laboratory housing conditions (Van der Borgh et al., 2005; Parihar et al., 2011).

AN in the hippocampus is further required for the beneficial effects of an enriched environment on recovery from stress-induced changes in behavior (Schloesser et al., 2010), where stress recovery is correlated with newborn cell survival (Schloesser et al., 2010; Tanti et al., 2012). Surprisingly, housing animals in an enriched environment that includes voluntary exercise, increases GCs (Vivinetto et al., 2013), suggesting that this rise in GC levels is essential for increased AN in the hippocampus (Schloesser et al., 2010; Sampedro-Piquero et al., 2014). When rats are adrenalectomized, admittedly a highly artificial condition, environmental enrichment-induced increases in AN are no longer apparent (Lehmann et al., 2013), indicating that GCs can facilitate adult hippocampal neurogenesis under specific conditions. Again somewhat counterintuitively, exercise, which is considered a potent stimulus for AN (van Praag et al., 1999b), also stimulates GC levels, even though exercise *per se* reduces stress (Kannangara et al., 2011). Cessation of voluntary exercise subsequently impairs AN and can increase anxiety-like behavior (Nishijima et al., 2013) consistent with other studies that indicate that changes in AN often correlate with anxiety measures (Pham et al., 2005; Revest et al., 2009; Hill et al., 2015; Seo et al., 2015; Hu et al., 2016; Yun et al., 2016).

When no other transmitter systems are altered and the stressor is unpredictable or uncontrollable and its nature severe, stress generally reduces AN (Pham et al., 2003; Heine et al., 2004c; Simon et al., 2005; Dranovsky and Hen 2006; Jayatissa et al., 2006; Mitra et al., 2006; Jayatissa et al., 2009; Perera et al., 2011; Lehmann et al., 2013; Schoenfeld and Gould, 2013). In fact, unpredictable/uncontrollable stress can reduce multiple stages of the neurogenic process, including the initial phase of proliferation of the neural stem cells and amplifying progenitor cells, as well as subsequent neuronal differentiation phase and dendritic expansion. Exposure to GCs *per se* was even shown to deplete the neural precursor pool (Yu et al., 2010). Stress not only reduces proliferation and AN in many different species, it may also shift neural stem cells away from neuronal differentiation, and instead 'redirect' them toward the generation of oligodendrocytes (Chetty et al., 2014). Although not studied in great detail yet, such stress-induced fate shifts may have important functional consequences: e.g., for the myelination of axons and/or mossy fibers and hence network connectivity, particularly when they occur during early development when cell division is massive.

Although different types of stress trigger different behavioral and functional responses, adrenal GCs are considered instrumental in mediating the suppressing effects of stress on AN (Schoenfeld and Gould, 2013). The basis for this assertion is as follows. First, exogenous GC administra-

tion to animals has effects similar to those of stress on cell proliferation, neuronal differentiation and cell survival as well as on the production of oligodendrocytes and microglia responses (Mayer et al., 2006; Wong and Herbert 2006; Yu et al., 2010; Hu et al., 2012). Second, the reductions in AN after stress and many of the molecular (Datson et al., 2012) and physiological (Krugers et al., 2006) changes, can be prevented by blocking the GR, for a very short period (Mayer et al., 2006; Oomen et al., 2007; Hu et al., 2012) or by CRH antagonists (Alonso et al., 2004). Furthermore, in a transgenic mouse model of AN inhibition a transient increase in the corticosterone response to stress occurs, as well as an attenuated dexamethasone-induced suppression of corticosterone release (Snyder et al., 2011). This is indicative of a role for the newborn cells in regulating HPA axis activity. On the other hand, ablation of AN by irradiation did not impair basal HPA axis activity (Surget et al., 2011), see also (Lucassen et al., 2013a).

Although general blockers of different elements of the stress axis are thus already effective, the precise mechanism(s) by which GCs decrease the numbers of new neurons remains poorly understood. More information has become available on its molecular control (Schouten et al., 2012; Anacker et al., 2013; Miller et al., 2013). NMDA receptors, GRs and MRs are all present on newly born cells, albeit in different ratios over time, and these likely act in concert to mediate effects of stress on AN (Montaron et al., 2003; Garcia et al., 2004; Wong and Herbert, 2004, 2005). Notably, GR knockdown in newborn cells exclusively accelerates their neuronal differentiation and migration and alters their dendritic complexity. This was paralleled by reduced freezing during contextual fear conditioning. Hence, GR expression in the newborn hippocampal cells is important for structural as well as functional integration into the mature hippocampal circuits involved in fear memory (Fitzsimons et al., 2013).

Furthermore, most precursors in the brain are located closely to blood vessels (Palmer et al., 2000). Although often not distinguished in quantitative analysis, this proximity makes this population particularly sensitive to stress hormones (Heine et al., 2005) and many other peripheral factors. Astrocytes are also of relevance as they closely align the vasculature, express GRs, support the survival of developing neurons and are involved in their synaptic integration (Sultan et al., 2015). Notably, astrocytes are affected by some, but not all, types of stress (Vallières et al., 2002; Czéh et al., 2006; Banasr and Duman, 2007; Oomen et al., 2009; Wang et al., 2013).

Stress further slows down neuronal differentiation of the adult-born cells, as evidenced by the upregulation of markers indicating cell cycle arrest (Heine et al., 2004a) that may be induced by specific changes in DNA methylation (Boku et al., 2015). Stress also reduces the survival of new neurons that were born prior to the actual stressful experience. A change in "corticoid environment" (Wong and Herbert, 2006) is thought

to be mediated by stress-induced reductions in neurotrophins and survival promoting factors such as brain-derived neurotrophic factor (BDNF), insulin-like growth factor-1 (IGF-1) and vascular endothelial growth factor (VEGF), (Schmidt and Duman, 2007; Wilson et al., 2014). Reductions in newborn cell survival may involve microglia, which can phagocytose new neurons. Indeed, stress influences microglia, both in terms of numbers and their responsivity. This may modulate their efficiency in clearing debris or dead neurons (Hinwood et al., 2012; Morris et al., 2013; Sierra et al., 2010) or their capacity to release neurotoxic cytokines (Kreisel et al., 2014; Sierra et al., 2014; Guadagno et al., 2015; Llorens-Martín et al., 2016).

An important aspect relevant for studies on temporal aspects of stress and AN is whether GC levels remain elevated or not after the initial exposure to stress has ended. In some psychosocial stress models, GC levels remain elevated, which has stronger suppressive effects on AN than exposure to severe, but predictable, physical stressors like restraint (Wong and Herbert, 2004). Several examples exist of a persistent and lasting inhibition of AN after an initial exposure to stress, despite a later lowering of GC levels (Czéh et al., 2002; Mirescu and Gould, 2006). In contrast, GC levels can remain elevated after a psychosocial stressor, with AN being suppressed long-term. In milder models of stress, GC levels generally normalize, yet AN remains reduced (Van Bokhoven et al., 2011; Schoenfeld and Gould, 2013). This suggests that while GCs are involved in the initial suppression of proliferation, they are not always necessary for maintaining this effect.

When studying effects of stress on AN in laboratory conditions, it is further important to realize that many variables influence the outcome (Bekinschtein et al., 2011). These variables include inter-individual genetic or gender differences in stress coping and resilience (Levine, 2005), prior handling of the animals, time of day at sacrifice and previous exposure to stressful learning tasks like the water maze, or exercise (Droste et al., 2003; Holmes et al., 2004; Ehninger and Kempermann 2006; Marlatt et al., 2012). Anatomical differences exist such as in projections to specific subregions of the hippocampus or in the larger networks, or neuromatrix (Sousa, 2016). Thus, stress effects on AN might differ between the dorsal or the ventral hippocampus depending upon the stimulus (Tanti and Belzung 2013; O'Leary and Cryan, 2014).

Many other factors may contribute to the stress-induced inhibition of AN, such as the stress-induced increase in glutamate release and NMDA receptor activation (Gould et al., 1997; Nacher and McEwen, 2006; Schoenfeld and Gould, 2013) or through stress effects on various neurotransmitter systems implicated in the regulation of AN such as GABA (Ge et al., 2007), serotonin (Djavanmard, 2004), noradrenaline (Joca et al., 2007), acetylcholine (Brüel-Jungferman et al., 2011), dopamine (Domínguez-Escribà et al., 2006; Takamura

et al., 2014), cannabinoids, opioids, nitric oxide, and gonadal steroids (Galea, 2008; Balu and Lucki, 2009; Mahmoud et al., 2016).

Finally, many antidepressant drugs that interfere with stress-related behavior in animals also modulate AN. The relation between stress, AN, antidepressants and psychopathology like major depression has been extensively discussed (Malberg and Duman, 2003; Santarelli et al., 2003; Dranovsky and Hen, 2006; Sahay and Hen, 2007; Lucassen et al., 2010; Perera et al., 2011; Surget et al., 2011; Tanti and Belzung, 2013; Lucassen et al., 2014; Schoenfeld and Cameron, 2015).

Changes in adult neurogenesis: functional relevance

To understand the potential functional impact of stress-modulation of adult neurogenesis, we first need to know which functional role these new neurons play in the hippocampal circuitry. In functional terms, early studies had already found that changes in AN were commonly paralleled by changes in learning and memory (Gould et al., 1999; van Praag et al., 1999a). This gave rise to the idea that AN may be relevant for at least some forms of hippocampal dependent cognition (Saxe et al., 2006; Zhang et al., 2008; Clelland et al., 2009; Aimone et al., 2010; Sahay et al., 2011a; Sahay et al., 2011b; Oomen et al., 2014). Suppression of AN, e.g. by stress or other means, has also been linked to an impairment in the animal's performance on hippocampal tasks or a change in behavior in mood or anxiety-related paradigms (Lemaire et al., 2000; Montaron et al., 2004; Montaron et al., 2006; Veena et al., 2009; Snyder et al., 2011).

Initial studies on selective elimination of newborn granule neurons administered methylazoxymethanol acetate (MAM), (Shors et al., 2001) and showed decreased cue responding in MAM-treated rats that were trained in a trace cue condition, but not in a delay condition. A similar impairment was also found in freezing in a cued fear conditioning task but not in spatial maze learning or contextual fear conditioning, among others (Shors et al., 2002). With trace fear conditioning being the most slowly learned of these tasks, this suggested that task difficulty is an important variable, and that AN may be particularly involved in more challenging memory tests such as those in which the associations are separated in time (i.e. trace conditioning).

Other studies using toxins replicated some of these findings: trace eyeblink conditioning was also impaired after temozolomide (TMZ) treatment and, consistent with earlier findings, MAM treatment did not affect contextual fear conditioning or water maze acquisition. Contrary to Shors et al. (2002) however, retention of platform location in the water maze was found to be impaired by MAM mediated ablation.

Recently, TMZ induced ablation affected water maze acquisition in juvenile but not in older animals.

Decreased contextual fear conditioning has further been demonstrated in a few studies where AN was depleted using irradiation (Winocur et al., 2006; Wojtowicz et al., 2008; Snyder et al., 2009), although exceptions exist too (Shors et al., 2002; Groves et al., 2013). The Morris water maze is one of the most widely used hippocampus-dependent spatial tasks. In contrast to impairments seen after hippocampal lesions, performance on standard spatial water maze training and probe trials is consistently spared in mice and rats that lack adult neurogenesis (Snyder et al., 2005; Wojtowicz et al., 2008; Arruda-Carvalho et al., 2011; Ben Abdallah et al., 2013; see also Snyder et al., 2016). Also here, more difficult tests that use long delays, or more subtle behavioral analyses of search strategies, identify impairments in animals lacking new neurons in some, though not all, experiments (Snyder et al., 2005; Ben Abdallah et al., 2013; Garthe et al., 2014).

A deficit in spatial performance was recently confirmed using optogenetics, that highlighted the importance of the age of the adult-born neurons as well. Four week old, but not of 2 or 8 week-old, neurons were shown to be involved in retention, but not acquisition, of the water maze (Gu et al., 2012). Studies on contextual fear conditioning and object (location) memory show mixed results and both impaired and unaffected memory performance was found. In addition to these learning tasks, animals lacking adult neurogenesis show an altered behavior in spontaneous investigation tasks (Lagace et al., 2011; Mak and Weiss 2010), which may reflect a depletion of olfactory neurogenesis, rather than of hippocampal neurogenesis.

Mice use specific strategies to solve a spatial task. Mice that lack adult neurogenesis use precise spatial strategies less frequently than normal mice, and mainly during the reversal learning trial but not during the initial learning phase (Garthe et al., 2009). In a probe trial without the platform, mice lacking adult neurogenesis search in both the original location and the new location, whereas normal mice focus much more on the newer location (Arruda-Carvalho et al., 2011). Also in an active place-avoidance spatial reversal task, mice lacking adult neurogenesis showed impairment when a reversal element was built in (Burghardt et al., 2012). Importantly, in all these reversal tasks, the novel memory for the initial platform location may interfere with the memory for the additional location. Distinguishing two similar memories may resemble aspects of pattern separation, which will be discussed below.

The specific involvement of AN in classical learning and memory tests further seems to depend on a number of factors, including species tested, the age of neurons, the phase of memory addressed and the type, design and difficulty of the test used. Together, such differences may contribute to some of the inconsistencies reported in the spatial and emotional paradigms.

Following studies that were in part based on computational models, new neurons were predicted to be important for pattern separation (Becker 2005; Aimone et al., 2009), the process by which a neural circuit reduces overlap between similar input patterns, resulting in more dissimilar output signals. Pattern separation is thought to be particularly necessary for the formation of unique, non- (or less-) overlapping representations and thus for successful memory storage.

The first study showing empirical evidence for a role of AN in pattern separation found that mice lacking neurogenesis were impaired on two spatial pattern separation tests in which spatial similarity was manipulated (Clelland et al., 2009). In particular, mice made more errors when locations were spaced closely, but not when spaced further apart (Clelland et al., 2009). Two different techniques for ablating immature neurons, X-irradiation and lentiviral mediated expression of a dominant-negative version of the Wnt protein were used. Further studies found similar results, whereby mice with impaired neurogenesis are slower to show differential freezing to highly similar contexts (Kheirbek et al., 2012b; Tronel et al., 2012).

Conversely, Sahay et al., (2011a) used a genetic manipulation to increase AN artificially. This resulted in improved context discrimination in a fear conditioning paradigm in which animals were trained to discriminate between two similar contexts across repeated sessions. Mice with increased neurogenesis were quicker to demonstrate discrimination of highly similar contexts (Sahay et al., 2011a). Furthermore, Creer and colleagues (2010) demonstrated that wheel running in mice increased AN as well as pattern separation in a touchscreen-based behavioral task, a treatment that was ineffective in aged animals that lacked running-dependent increase in AN, suggesting it was the increase in AN and not other exercise-induced effects, that were responsible for the improvements (Creer et al., 2010). Finally, brain-derived neurotrophic factor (BDNF) in the dentate gyrus was shown to be important specifically during the consolidation (but not retrieval) of memories for similar (but not dissimilar) spatial locations in an open field. BDNF was found to interact with adult-born neurons in the service of pattern separation (Bekinschtein et al., 2013; Bekinschtein et al., 2014).

For stress-related disorders such as PTSD, it has been hypothesized that impaired pattern separation may lead to overgeneralization: i.e. when events are not stored as unique representations, similar cues may be easily confused with those of the original experience (Kheirbek et al., 2012a; Besnard and Sahay 2016). Since PTSD patients experience panic attacks and relive memories in response to cues similar to those of the traumatic event, overgeneralization may be regarded as a core cognitive mechanism of this disorder. This may be particularly interesting in linking the effects of stress on adult neurogenesis to potential mechanisms of brain disease and psychopathology.

Concluding remarks

Stress and GCs commonly interfere with one or more stages of adult neurogenesis. Such inhibitory effects can normalize after a recovery period, voluntary exercise or antidepressant treatment. Adult neurogenesis has been implicated in cognitive functions and in the regulation of mood and anxiety. A reduced rate of AN may be indicative of impaired hippocampal plasticity. Lasting reductions in AN, or in the turnover rate of the DG granule cells will in time alter the overall composition of the DG cell population and can modify stress responsivity and cognition, and thereby influence overall functioning of the adult hippocampal circuit and its involvement in (aspects of) psychopathology.

Acknowledgments

PJL is supported by the HersenStichting Nederland, Alzheimer Nederland, ISAO, NWO and Amsterdam Brain & Cognition (ABC), CAO is supported by the Amsterdam Brain and Mind Project (ABMP).

Compliance with ethics guidelines

Paul J. Lucassen and Charlotte A. Oomen declare that they have no conflict of interest. This manuscript is a review article and does not involve a research protocol requiring approval by the relevant institutional review board or ethics committee.

References

- Abrous D N, Koehl M, Le Moal M (2005). Adult neurogenesis: from precursors to network and physiology. *Physiol Rev*, 85(2): 523–569
- Aimone J B, Deng W, Gage F H (2010). Adult neurogenesis: integrating theories and separating functions. *Trends Cogn Sci*, 14(7): 325–337
- Aimone J B, Wiles J, Gage F H (2009). Computational influence of adult neurogenesis on memory encoding. *Neuron*, 61(2): 187–202
- Alonso R, Griebel G, Pavone G, Stemmelin J, Le Fur G, Soubrié P (2004). Blockade of CRF(1) or V(1b) receptors reverses stress-induced suppression of neurogenesis in a mouse model of depression. *Mol Psychiatry*, 9(3): 278–286, 224
- Anacker C, Cattaneo A, Luoni A, Musaelyan K, Zunszain P A, Milanese E, Rybka J, Berry A, Cirulli F, Thuret S, Price J, Riva M A, Gennarelli M, Pariante C M (2013). Glucocorticoid-related molecular signaling pathways regulating hippocampal neurogenesis. *Neuropsychopharmacology*, 38(5): 872–883
- Arruda-Carvalho M, Sakaguchi M, Akers K G, Josselyn S A, Frankland P W (2011). Posttraining ablation of adult-generated neurons degrades previously acquired memories. *J Neurosci*, 31(42): 15113–15127
- Balu D T, Lucki I (2009). Adult hippocampal neurogenesis: regulation, functional implications, and contribution to disease pathology. *Neurosci Biobehav Rev*, 33(3): 232–252
- Banasr M, Duman R S (2007). Regulation of neurogenesis and gliogenesis by stress and antidepressant treatment. *CNS Neurol Disord Drug Targets*, 6(5): 311–320
- Becker S (2005). A computational principle for hippocampal learning and neurogenesis. *Hippocampus*, 15(6): 722–738
- Bekinschtein P, Kent B A, Oomen C A, Clemenson G D, Gage F H, Saksida L M, Bussey T J (2014). Brain-derived neurotrophic factor interacts with adult-born immature cells in the dentate gyrus during consolidation of overlapping memories. *Hippocampus*, 24(8): 905–911
- Bekinschtein P, Kent B A, Oomen C A, Clemenson G D, Jr., Gage F H, Saksida L M, Bussey T J (2013). BDNF in the Dentate Gyrus Is Required for Consolidation of “Pattern-Separated” Memories. *Cell Reports*, 5: 1–10
- Bekinschtein P, Oomen C A, Saksida L M, Bussey T J (2011). Effects of environmental enrichment and voluntary exercise on neurogenesis, learning and memory, and pattern separation: BDNF as a critical variable? *Semin Cell Dev Biol*, 22(5): 536–542
- Ben Abdallah N M, Filipkowski R K, Pruschy M, Jaholkowski P, Winkler J, Kaczmarek L, Lipp H P (2013). Impaired long-term memory retention: common denominator for acutely or genetically reduced hippocampal neurogenesis in adult mice. *Behav Brain Res*, 252: 275–286
- Besnard A, Sahay A (2016). Adult Hippocampal Neurogenesis, Fear Generalization, and Stress. *Neuropsychopharmacology*, 41(1): 24–44
- Binder E B, Bradley R G, Liu W, Epstein M P, Deveau T C, Mercer K B, Tang Y, Gillespie C F, Heim C M, Nemeroff C B, Schwartz A C, Cubells J F, Ressler K J (2008). Association of FKBP5 polymorphisms and childhood abuse with risk of posttraumatic stress disorder symptoms in adults. *JAMA*, 299(11): 1291–1305
- Boku S, Toda H, Nakagawa S, Kato A, Inoue T, Koyama T, Hiroi N, Kusumi I (2015). Neonatal maternal separation alters the capacity of adult neural precursor cells to differentiate into neurons via methylation of retinoic acid receptor gene promoter. *Biol Psychiatry*, 77(4): 335–344
- Bruel-Jungerman E, Lucassen P J, Francis F (2011). Cholinergic influences on cortical development and adult neurogenesis. *Behav Brain Res*, 221(2): 379–388
- Burghardt N S, Park E H, Hen R, Fenton A A (2012). Adult-born hippocampal neurons promote cognitive flexibility in mice. *Hippocampus*, 22(9): 1795–1808
- Chetty S, Friedman A R, Taravosh-Lahn K, Kirby E D, Mirescu C, Guo F, Krupik D, Nicholas A, Geraghty A C, Krishnamurthy A, Tsai M K, Covarrubias D, Wong A T, Francis D D, Sapolsky R M, Palmer T D, Pleasure D, Kaufer D (2014). Stress and glucocorticoids promote oligodendrogenesis in the adult hippocampus. *Mol Psychiatry*, 19(12): 1275–1283
- Clelland C D, Choi M, Romberg C, Clemenson G D Jr, Fragniere A, Tyers P, Jessberger S, Saksida L M, Barker R A, Gage F H, Bussey T J (2009). A functional role for adult hippocampal neurogenesis in spatial pattern separation. *Science*, 325(5937): 210–213
- Creer D J, Romberg C, Saksida L M, van Praag H, Bussey T J (2010). Running enhances spatial pattern separation in mice. *Proc Natl Acad Sci USA*, 107(5): 2367–2372
- Czeh B, Lucassen P J (2007). What causes the hippocampal volume decrease in depression? Are neurogenesis, glial changes and apoptosis implicated? *Eur Arch Psychiatry Clin Neurosci*, 257(5): 250–260
- Czeh B, Simon M, Schmeling B, Hiemke C, Fuchs E (2006). Astroglial plasticity in the hippocampus is affected by chronic psychosocial

- stress and concomitant fluoxetine treatment. *Neuropsychopharmacology*, 31(8): 1616–1626
- Czéh B, Welt T, Fischer A K, Erhardt A, Schmitt W, Müller M B, Toschi N, Fuchs E, Keck M E (2002). Chronic psychosocial stress and concomitant repetitive transcranial magnetic stimulation: effects on stress hormone levels and adult hippocampal neurogenesis. *Biol Psychiatry*, 52(11): 1057–1065
- Dagyte G, Van der Zee E A, Postema F, Luiten P G, Den Boer J A, Trentani A, Meerlo P (2009). Chronic but not acute foot-shock stress leads to temporary suppression of cell proliferation in rat hippocampus. *Neuroscience*, 162(4): 904–913
- Datson N A, Speksnijder N, Mayer J L, Steenbergen P J, Korobko O, Goeman J, de Kloet E R, Joëls M, Lucassen P J (2012). The transcriptional response to chronic stress and glucocorticoid receptor blockade in the hippocampal dentate gyrus. *Hippocampus*, 22(2): 359–371
- de Kloet E R, Joëls M, Holsboer F (2005). Stress and the brain: from adaptation to disease. *Nat Rev Neurosci*, 6(6): 463–475
- Djavadian R L (2004). Serotonin and neurogenesis in the hippocampal dentate gyrus of adult mammals. *Acta Neurobiol Exp (Wars)*, 64(2): 189–200
- Dominguez-Escribà L, Hernández-Rabaza V, Soriano-Navarro M, Barcia J A, Romero F J, García-Verdugo J M, Canales J J (2006). Chronic cocaine exposure impairs progenitor proliferation but spares survival and maturation of neural precursors in adult rat dentate gyrus. *Eur J Neurosci*, 24(2): 586–594
- Dranovsky A, Hen R (2006). Hippocampal neurogenesis: regulation by stress and antidepressants. *Biol Psychiatry*, 59(12): 1136–1143
- Droste S K, Gesing A, Ulbricht S, Müller M B, Linthorst A C, Reul J M (2003). Effects of long-term voluntary exercise on the mouse hypothalamic-pituitary-adrenocortical axis. *Endocrinology*, 144(7): 3012–3023
- Ehninger D, Kempermann G (2006). Paradoxical effects of learning the Morris water maze on adult hippocampal neurogenesis in mice may be explained by a combination of stress and physical activity. *Genes Brain Behav*, 5(1): 29–39
- Erdmann G, Berger S, Schütz G (2008). Genetic dissection of glucocorticoid receptor function in the mouse brain. *J Neuroendocrinol*, 20(6): 655–659
- Fitzsimons C P, Herbert J, Schouten M, Meijer O C, Lucassen P J, Lightman S (2016). Circadian and Ultradian Glucocorticoid Rhythmicity: Implications for the Effects of Glucocorticoids on Neural Stem Cells and Adult Hippocampal Neurogenesis. *Front Neuroendocrinol*, (In press)
- Fitzsimons C P, van Hooijdonk L W, Schouten M, Zalachoras I, Brinks V, Zheng T, Schouten T G, Saaltink D J, Dijkmans T, Steindler D A, Verhaagen J, Verbeek F J, Lucassen P J, de Kloet E R, Meijer O C, Karst H, Joels M, Oitzl M S, Vreugdenhil E (2013). Knockdown of the glucocorticoid receptor alters functional integration of newborn neurons in the adult hippocampus and impairs fear-motivated behavior. *Mol Psychiatry*, 18(9): 993–1005
- Galea L A (2008). Gonadal hormone modulation of neurogenesis in the dentate gyrus of adult male and female rodents. *Brain Res Brain Res Rev*, 57(2): 332–341
- Garcia A, Steiner B, Kronenberg G, Bick-Sander A, Kempermann G (2004). Age-dependent expression of glucocorticoid- and mineralocorticoid receptors on neural precursor cell populations in the adult murine hippocampus. *Aging Cell*, 3(6): 363–371
- Garthe A, Behr J, Kempermann G (2009). Adult-generated hippocampal neurons allow the flexible use of spatially precise learning strategies. *PLoS ONE*, 4(5): e5464
- Garthe A, Huang Z, Kaczmarek L, Filipkowski R K, Kempermann G (2014). Not all water mazes are created equal: cyclin D2 knockout mice with constitutively suppressed adult hippocampal neurogenesis do show specific spatial learning deficits. *Genes Brain Behav*, 13(4): 357–364
- Ge S, Yang C H, Hsu K S, Ming G L, Song H (2007). A critical period for enhanced synaptic plasticity in newly generated neurons of the adult brain. *Neuron*, 54(4): 559–566
- Gould E (2007). How widespread is adult neurogenesis in mammals? *Nat Rev Neurosci*, 8(6): 481–488
- Gould E, Beylin A, Tanapat P, Reeves A, Shors T J (1999). Learning enhances adult neurogenesis in the hippocampal formation. *Nat Neurosci*, 2(3): 260–265
- Gould E, McEwen B S, Tanapat P, Galea L A, Fuchs E (1997). Neurogenesis in the dentate gyrus of the adult tree shrew is regulated by psychosocial stress and NMDA receptor activation. *J Neurosci*, 17(7): 2492–2498
- Groves J O, Leslie I, Huang G J, McHugh S B, Taylor A, Mott R, Munafò M, Bannerman D M, Flint J (2013). Ablating adult neurogenesis in the rat has no effect on spatial processing: evidence from a novel pharmacogenetic model. *PLoS Genet*, 9(9): e1003718
- Gu Y, Arruda-Carvalho M, Wang J, Janoschka S R, Josselyn S A, Frankland P W, Ge S (2012). Optical controlling reveals time-dependent roles for adult-born dentate granule cells. *Nat Neurosci*, 15: 1700–1706
- Guadagno J, Swan P, Shaikh R, Cregan S P (2015). Microglia-derived IL-1 β triggers p53-mediated cell cycle arrest and apoptosis in neural precursor cells. *Cell death & disease*, 6: e1779
- Hanson N D, Owens M J, Boss-Williams K A, Weiss J M, Nemeroff C B (2011a). Several stressors fail to reduce adult hippocampal neurogenesis. *Psychoneuroendocrinology*, 36(10): 1520–1529
- Hanson N D, Owens M J, Nemeroff C B (2011b). Depression, antidepressants, and neurogenesis: a critical reappraisal. *Neuropsychopharmacology*, 36(13): 2589–2602
- Harris A P, Holmes M C, de Kloet E R, Chapman K E, Seckl J R (2013). Mineralocorticoid and glucocorticoid receptor balance in control of HPA axis and behaviour. *Psychoneuroendocrinology*, 38(5): 648–658
- Heine V M, Maslam S, Joëls M, Lucassen P J (2004a). Increased P27KIP1 protein expression in the dentate gyrus of chronically stressed rats indicates G1 arrest involvement. *Neuroscience*, 129(3): 593–601
- Heine V M, Maslam S, Joëls M, Lucassen P J (2004b). Prominent decline of newborn cell proliferation, differentiation, and apoptosis in the aging dentate gyrus, in absence of an age-related hypothalamus-pituitary-adrenal axis activation. *Neurobiol Aging*, 25(3): 361–375
- Heine V M, Maslam S, Zareno J, Joëls M, Lucassen P J (2004c). Suppressed proliferation and apoptotic changes in the rat dentate gyrus after acute and chronic stress are reversible. *Eur J Neurosci*, 19(1): 131–144
- Heine V M, Zareno J, Maslam S, Joëls M, Lucassen P J (2005). Chronic stress in the adult dentate gyrus reduces cell proliferation near the vasculature and VEGF and Flk-1 protein expression. *Eur J Neurosci*,

- 21(5): 1304–1314
- Hill A S, Sahay A, Hen R (2015). Increasing adult hippocampal neurogenesis is sufficient to reduce anxiety and depression-like behaviors. *Neuropsychopharmacology*, 40(10): 2368–2378
- Hinwood M, Morandini J, Day T A, Walker F R (2012). Evidence that microglia mediate the neurobiological effects of chronic psychological stress on the medial prefrontal cortex. *Cereb Cortex*, 22(6): 1442–1454
- Holmes M M, Galea L A, Mistlberger R E, Kempermann G (2004). Adult hippocampal neurogenesis and voluntary running activity: circadian and dose-dependent effects. *J Neurosci Res*, 76(2): 216–222
- Hu P, Oomen C, van Dam A M, Wester J, Zhou J N, Joëls M, Lucassen P J (2012). A single-day treatment with mifepristone is sufficient to normalize chronic glucocorticoid induced suppression of hippocampal cell proliferation. *PLoS ONE*, 7(9): e46224
- Hu P, Wang Y, Liu J, Meng F T, Qi X R, Chen L, van Dam A M, Joëls M, Lucassen P J, Zhou J N (2016). Chronic retinoic acid treatment suppresses adult hippocampal neurogenesis, in close correlation with depressive-like behavior. *Hippocampus*: doi: 10.1002/hipo.22574
- Jayatissa M N, Bisgaard C, Tingström A, Papp M, Wiborg O (2006). Hippocampal cytogenesis correlates to escitalopram-mediated recovery in a chronic mild stress rat model of depression. *Neuropsychopharmacology*, 31(11): 2395–2404
- Jayatissa M N, Henningsen K, West M J, Wiborg O (2009). Decreased cell proliferation in the dentate gyrus does not associate with development of anhedonic-like symptoms in rats. *Brain Res*, 1290: 133–141
- Jessberger S, Gage F H (2014). Adult neurogenesis: bridging the gap between mice and humans. *Trends Cell Biol*, 24(10): 558–563
- Joca S R, Ferreira F R, Guimarães F S (2007). Modulation of stress consequences by hippocampal monoaminergic, glutamatergic and nitrgic neurotransmitter systems. *Stress*, 10(3): 227–249
- Joëls M, Baram T Z (2009). The neuro-symphony of stress. *Nat Rev Neurosci*, 10(6): 459–466
- Joëls M, Pu Z, Wiegert O, Oitzl M S, Krugers H J (2006). Learning under stress: how does it work? *Trends Cogn Sci*, 10(4): 152–158
- Joëls M, Sarabdjitsingh R A, Karst H (2012). Unraveling the time domains of corticosteroid hormone influences on brain activity: rapid, slow, and chronic modes. *Pharmacol Rev*, 64(4): 901–938
- Kanatso S, Fearey B C, Kuil L E, Lucassen P J, Harris A P, Seckl J R, Krugers H, Joels M (2015). Overexpression of Mineralocorticoid Receptors Partially Prevents Chronic Stress-Induced Reductions in Hippocampal Memory and Structural Plasticity. *PLoS ONE*, 10(11): e0142012
- Kannangara T S, Lucero M J, Gil-Mohapel J, Drapala R J, Simpson J M, Christie B R, van Praag H (2011). Running reduces stress and enhances cell genesis in aged mice. *Neurobiol Aging*, 32(12): 2279–2286
- Karst H, Berger S, Erdmann G, Schütz G, Joëls M (2010). Metaplasticity of amygdalar responses to the stress hormone corticosterone. *Proc Natl Acad Sci USA*, 107(32): 14449–14454
- Kempermann G (2012). New neurons for ‘survival of the fittest’. *Nat Rev Neurosci*, 13(10): 727–736
- Kempermann G, Fabel K, Ehninger D, Babu H, Leal-Galicia P, Garthe A, Wolf S A (2010). Why and how physical activity promotes experience-induced brain plasticity. *Front Neurosci*, 4: 189
- Kheirbek M A, Klemenhagen K C, Sahay A, Hen R (2012a). Neurogenesis and generalization: a new approach to stratify and treat anxiety disorders. *Nat Neurosci*, 15(12): 1613–1620
- Kheirbek M A, Tannenholz L, Hen R (2012b). NR2B-dependent plasticity of adult-born granule cells is necessary for context discrimination. *J Neurosci*, 32(25): 8696–8702
- Koolhaas J M, Bartolomucci A, Buwalda B, de Boer S F, Flügge G, Korte S M, Meerlo P, Murison R, Olivier B, Palanza P, Richter-Levin G, Sgoifo A, Steimer T, Stiedl O, van Dijk G, Wöhr M, Fuchs E (2011). Stress revisited: a critical evaluation of the stress concept. *Neurosci Biobehav Rev*, 35(5): 1291–1301
- Kreisler T, Frank M G, Licht T, Reshef R, Ben-Menachem-Zidon O, Baratta M V, Maier S F, Yirmiya R (2014). Dynamic microglial alterations underlie stress-induced depressive-like behavior and suppressed neurogenesis. *Mol Psychiatry*, 19(6): 699–709
- Kretz O, Reichardt H M, Schütz G, Bock R (1999). Corticotropin-releasing hormone expression is the major target for glucocorticoid feedback-control at the hypothalamic level. *Brain Res*, 818(2): 488–491
- Krugers H J, Goltstein P M, van der Linden S, Joëls M (2006). Blockade of glucocorticoid receptors rapidly restores hippocampal CA1 synaptic plasticity after exposure to chronic stress. *Eur J Neurosci*, 23(11): 3051–3055
- Kuhn H G, Dickinson-Anson H, Gage F H (1996). Neurogenesis in the dentate gyrus of the adult rat: age-related decrease of neuronal progenitor proliferation. *J Neurosci*, 16(6): 2027–2033
- Lagace D C, Donovan M H, DeCarolis N A, Farnbauch L A, Malhotra S, Berton O, Nestler E J, Krishnan V, Eisch A J (2010). Adult hippocampal neurogenesis is functionally important for stress-induced social avoidance. *Proc Natl Acad Sci USA*, 107(9): 4436–4441
- Lehmann M L, Brachman R A, Martinowich K, Schloesser R J, Herkenham M (2013). Glucocorticoids orchestrate divergent effects on mood through adult neurogenesis. *J Neurosci*, 33(7): 2961–2972
- Lemaire V, Koehl M, Le Moal M, Abrous D N (2000). Prenatal stress produces learning deficits associated with an inhibition of neurogenesis in the hippocampus. *Proc Natl Acad Sci USA*, 97(20): 11032–11037
- Levine S (2005). Developmental determinants of sensitivity and resistance to stress. *Psychoneuroendocrinology*, 30(10): 939–946
- Levone B R, Cryan J F, O’Leary O F (2015). Role of adult hippocampal neurogenesis in stress resilience. *Neurobiology of Stress* 1. *Neurobiol Stress*, 1: 147–155
- Liston C, Cichon J M, Jeanneteau F, Jia Z, Chao M V, Gan W B (2013). Circadian glucocorticoid oscillations promote learning-dependent synapse formation and maintenance. *Nat Neurosci*, 16(6): 698–705
- Llorens-Martín M, Jurado-Arjona J, Bolós M, Pallas-Bazarra N, Ávila J (2016). Forced swimming sabotages the morphological and synaptic maturation of newborn granule neurons and triggers a unique pro-inflammatory milieu in the hippocampus. *Brain Behav Immun*, 53: 242–254
- Lucassen P J, Fitzsimons C P, Korosi A, Joels M, Belzung C, Abrous D N (2013a). Stressing new neurons into depression? *Mol Psychiatry*, 18(4): 396–397
- Lucassen P J, Meerlo P, Naylor A S, van Dam A M, Dayer A G, Fuchs E, Oomen C A, Czéh B (2010). Regulation of adult neurogenesis by stress, sleep disruption, exercise and inflammation: Implications for

- depression and antidepressant action. *Eur Neuropsychopharmacol*, 20(1): 1–17
- Lucassen P J, Naninck E F, van Goudoever J B, Fitzsimons C, Joels M, Korosi A (2013b). Perinatal programming of adult hippocampal structure and function; emerging roles of stress, nutrition and epigenetics. *Trends Neurosci*, 36(11): 621–631
- Lucassen P J, Pruessner J, Sousa N, Almeida O F, Van Dam A M, Rajkowska G, Swaab D F, Czéh B (2014). Neuropathology of stress. *Acta Neuropathol*, 127(1): 109–135
- Lyons D M, Buckmaster P S, Lee A G, Wu C, Mitra R, Duffey L M, Buckmaster C L, Her S, Patel P D, Schatzberg A F (2010). Stress coping stimulates hippocampal neurogenesis in adult monkeys. *Proc Natl Acad Sci USA*, 107(33): 14823–14827
- Mahmoud R, Wainwright S R, Galea L A (2016). Sex hormones and adult hippocampal neurogenesis: Regulation, implications, and potential mechanisms. *Front Neuroendocrinol*
- Mak G K, Weiss S (2010). Paternal recognition of adult offspring mediated by newly generated CNS neurons. *Nat Neurosci*, 13: 753–758
- Malberg J E, Duman R S (2003). Cell proliferation in adult hippocampus is decreased by inescapable stress: reversal by fluoxetine treatment. *Neuropsychopharmacology*, 28(9): 1562–1571
- Marlatt M W, Potter M C, Lucassen P J, van Praag H (2012). Running throughout middle-age improves memory function, hippocampal neurogenesis, and BDNF levels in female C57BL/6J mice. *Dev Neurobiol*, 72(6): 943–952
- Mayer J L, Klumpers L, Maslam S, de Kloet E R, Joëls M, Lucassen P J (2006). Brief treatment with the glucocorticoid receptor antagonist mifepristone normalizes the corticosterone-induced reduction of adult hippocampal neurogenesis. *J Neuroendocrinol*, 18(8): 629–631
- Meaney M J, Szyf M, Seckl J R (2007). Epigenetic mechanisms of perinatal programming of hypothalamic-pituitary-adrenal function and health. *Trends Mol Med*, 13(7): 269–277
- Medina A, Seasholtz A F, Sharma V, Burke S, Bunney W Jr, Myers R M, Schatzberg A, Akil H, Watson S J (2013). Glucocorticoid and mineralocorticoid receptor expression in the human hippocampus in major depressive disorder. *J Psychiatr Res*, 47(3): 307–314
- Miller J A, Nathanson J, Franjic D, Shim S, Dalley R A, Shapouri S, Smith K A, Sunkin S M, Bernard A, Bennett J L, Lee C K, Hawrylycz M J, Jones A R, Amaral D G, Šestan N, Gage F H, Lein E S (2013). Conserved molecular signatures of neurogenesis in the hippocampal subgranular zone of rodents and primates. *Development*, 140(22): 4633–4644
- Mirescu C, Gould E (2006). Stress and adult neurogenesis. *Hippocampus*, 16(3): 233–238
- Mitra R, Sundlass K, Parker K J, Schatzberg A F, Lyons D M (2006). Social stress-related behavior affects hippocampal cell proliferation in mice. *Physiol Behav*, 89(2): 123–127
- Montaron M F, Drapeau E, Dupret D, Kitchener P, Aurousseau C, Le Moal M, Piazza P V, Abrous D N (2006). Lifelong corticosterone level determines age-related decline in neurogenesis and memory. *Neurobiol Aging*, 27(4): 645–654
- Montaron M F, Koehl M, Lemaire V, Drapeau E, Abrous D N, Le Moal M (2004). Environmentally induced long-term structural changes: cues for functional orientation and vulnerabilities. *Neurotox Res*, 6(7–8): 571–580
- Montaron M F, Piazza P V, Aurousseau C, Urani A, Le Moal M, Abrous D N (2003). Implication of corticosteroid receptors in the regulation of hippocampal structural plasticity. *Eur J Neurosci*, 18(11): 3105–3111
- Morris G P, Clark I A, Zinn R, Vissel B (2013). Microglia: a new frontier for synaptic plasticity, learning and memory, and neurodegenerative disease research. *Neurobiol Learn Mem*, 105: 40–53
- Nacher J, McEwen B S (2006). The role of N-methyl-D-aspartate receptors in neurogenesis. *Hippocampus*, 16(3): 267–270
- Nishijima T, Llorens-Martín M, Tejeda G S, Inoue K, Yamamura Y, Soya H, Trejo J L, Torres-Alemán I (2013). Cessation of voluntary wheel running increases anxiety-like behavior and impairs adult hippocampal neurogenesis in mice. *Behav Brain Res*, 245: 34–41
- O’Leary O F, Cryan J F (2014). A ventral view on antidepressant action: roles for adult hippocampal neurogenesis along the dorsoventral axis. *Trends Pharmacol Sci*, 35(12): 675–687
- O’Leary O F, O’Connor R M, Cryan J F (2012). Lithium-induced effects on adult hippocampal neurogenesis are topographically segregated along the dorso-ventral axis of stressed mice. *Neuropharmacology*, 62(1): 247–255
- Oomen C A, Bekinschtein P, Kent B A, Saksida L M, Bussey T J (2014). Adult hippocampal neurogenesis and its role in cognition. *Wiley Interdiscip Rev Cogn Sci*, 5(5): 573–587
- Oomen C A, Girardi C E, Cahyadi R, Verbeek E C, Krugers H, Joëls M, Lucassen P J (2009). Opposite effects of early maternal deprivation on neurogenesis in male versus female rats. *PLoS ONE*, 4(1): e3675
- Oomen C A, Mayer J L, de Kloet E R, Joëls M, Lucassen P J (2007). Brief treatment with the glucocorticoid receptor antagonist mifepristone normalizes the reduction in neurogenesis after chronic stress. *Eur J Neurosci*, 26(12): 3395–3401
- Palmer T D, Willhoite A R, Gage F H (2000). Vascular niche for adult hippocampal neurogenesis. *J Comp Neurol*, 425(4): 479–494
- Parihar V K, Hattiangady B, Kuruba R, Shuai B, Shetty A K (2011). Predictable chronic mild stress improves mood, hippocampal neurogenesis and memory. *Mol Psychiatry*, 16(2): 171–183
- Perera T D, Dwork A J, Keegan K A, Thirumangalakudi L, Lipira C M, Joyce N, Lange C, Higley J D, Rosoklija G, Hen R, Sackeim H A, Coplan J D (2011). Necessity of hippocampal neurogenesis for the therapeutic action of antidepressants in adult nonhuman primates. *PLoS ONE*, 6(4): e17600
- Pham K, McEwen B S, Ledoux J E, Nader K (2005). Fear learning transiently impairs hippocampal cell proliferation. *Neuroscience*, 130(1): 17–24
- Pham K, Nacher J, Hof P R, McEwen B S (2003). Repeated restraint stress suppresses neurogenesis and induces biphasic PSA-NCAM expression in the adult rat dentate gyrus. *Eur J Neurosci*, 17(4): 879–886
- Qi X R, Kamphuis W, Wang S, Wang Q, Lucassen P J, Zhou J N, Swaab D F (2013). Aberrant stress hormone receptor balance in the human prefrontal cortex and hypothalamic paraventricular nucleus of depressed patients. *Psychoneuroendocrinology*, 38(6): 863–870
- Qian X, Droste S K, Lightman S L, Reul J M, Linthorst A C (2012). Circadian and ultradian rhythms of free glucocorticoid hormone are highly synchronized between the blood, the subcutaneous tissue, and the brain. *Endocrinology*, 153(9): 4346–4353
- Revest J M, Dupret D, Koehl M, Funk-Reiter C, Grosjean N, Piazza P V, Abrous D N (2009). Adult hippocampal neurogenesis is involved in anxiety-related behaviors. *Mol Psychiatry*, 14(10): 959–967

- Ridder S, Chourbaji S, Hellweg R, Urani A, Zacher C, Schmid W, Zink M, Hörtnagl H, Flor H, Henn F A, Schütz G, Gass P (2005). Mice with genetically altered glucocorticoid receptor expression show altered sensitivity for stress-induced depressive reactions. *J Neurosci*, 25(26): 6243–6250
- Sahay A, Hen R (2007). Adult hippocampal neurogenesis in depression. *Nat Neurosci*, 10(9): 1110–1115
- Sahay A, Scobie K N, Hill A S, O'Carroll C M, Kheirbek M A, Burghardt N S, Fenton A A, Dranovsky A, Hen R (2011a). Increasing adult hippocampal neurogenesis is sufficient to improve pattern separation. *Nature*, 472(7344): 466–470
- Sahay A, Wilson D A, Hen R (2011b). Pattern separation: a common function for new neurons in hippocampus and olfactory bulb. *Neuron*, 70(4): 582–588
- Sampedro-Piquero P, Begega A, Arias J L (2014). Increase of glucocorticoid receptor expression after environmental enrichment: relations to spatial memory, exploration and anxiety-related behaviors. *Physiol Behav*, 129: 118–129
- Santarelli L, Saxe M, Gross C, Surget A, Battaglia F, Dulawa S, Weisstaub N, Lee J, Duman R, Arancio O, Belzung C, Hen R (2003). Requirement of hippocampal neurogenesis for the behavioral effects of antidepressants. *Science*, 301(5634): 805–809
- Sapolsky R M, Uno H, Rebert C S, Finch C E (1990). Hippocampal damage associated with prolonged glucocorticoid exposure in primates. *J Neurosci*, 10(9): 2897–2902
- Sarabdjitsingh R A, Conway-Campbell B L, Leggett J D, Waite E J, Meijer O C, de Kloet E R, Lightman S L (2010). Stress responsiveness varies over the ultradian glucocorticoid cycle in a brain-region-specific manner. *Endocrinology*, 151(11): 5369–5379
- Sarabdjitsingh R A, Jezequel J, Pasricha N, Mikasova L, Kerkhofs A, Karst H, Groc L, Joëls M (2014). Ultradian corticosterone pulses balance glutamatergic transmission and synaptic plasticity. *Proc Natl Acad Sci USA*, 111(39): 14265–14270
- Sarabdjitsingh R A, Joëls M, de Kloet E R (2012). Glucocorticoid pulsatility and rapid corticosteroid actions in the central stress response. *Physiol Behav*, 106(1): 73–80
- Saxe M D, Battaglia F, Wang J W, Malleret G, David D J, Monckton J E, Garcia A D, Sofroniew M V, Kandel E R, Santarelli L, Hen R, Drew M R (2006). Ablation of hippocampal neurogenesis impairs contextual fear conditioning and synaptic plasticity in the dentate gyrus. *Proc Natl Acad Sci USA*, 103(46): 17501–17506
- Schloesser R J, Lehmann M, Martinowich K, Manji H K, Herkenham M (2010). Environmental enrichment requires adult neurogenesis to facilitate the recovery from psychosocial stress. *Mol Psychiatry*, 15(12): 1152–1163
- Schmidt H D, Duman R S (2007). The role of neurotrophic factors in adult hippocampal neurogenesis, antidepressant treatments and animal models of depressive-like behavior. *Behav Pharmacol*, 18(5-6): 391–418
- Schoenfeld T J, Cameron H A (2015). Adult neurogenesis and mental illness. *Neuropsychopharmacology*, 40(1): 113–128
- Schoenfeld T J, Gould E (2013). Differential effects of stress and glucocorticoids on adult neurogenesis. *Curr Top Behav Neurosci*, 15: 139–164
- Schouten M, Buijink M R, Lucassen P J, Fitzsimons C P (2012). New Neurons in Aging Brains: Molecular Control by Small Non-Coding RNAs. *Front Neurosci*, 6: 25
- Schwabe L, Joëls M, Roozendaal B, Wolf O T, Oitzl M S (2012). Stress effects on memory: an update and integration. *Neurosci Biobehav Rev*, 36(7): 1740–1749
- Seo D O, Carillo M A, Chih-Hsiung Lim S, Tanaka K F, Drew M R (2015). Adult Hippocampal Neurogenesis Modulates Fear Learning through Associative and Nonassociative Mechanisms. *J Neurosci*, 35(32): 11330–11345
- Shors T J, Miesegaes G, Beylin A, Zhao M, Rydel T, Gould E (2001). Neurogenesis in the adult is involved in the formation of trace memories. *Nature*, 410(6826): 372–376
- Shors T J, Townsend D A, Zhao M, Kozorovitskiy Y, Gould E (2002). Neurogenesis may relate to some but not all types of hippocampal-dependent learning. *Hippocampus*, 12(5): 578–584
- Sierra A, Beccari S, Diaz-Aparicio I, Encinas J M, Comeau S, Tremblay M E (2014). Surveillance, phagocytosis, and inflammation: how never-resting microglia influence adult hippocampal neurogenesis. *Neural Plast*, 2014: 610343
- Sierra A, Encinas J M, Deudero J J, Chancey J H, Enikolopov G, Overstreet-Wadiche L S, Tsirka S E, Maletic-Savatic M (2010). Microglia shape adult hippocampal neurogenesis through apoptosis-coupled phagocytosis. *Cell Stem Cell*, 7(4): 483–495
- Simon M, Czéh B, Fuchs E (2005). Age-dependent susceptibility of adult hippocampal cell proliferation to chronic psychosocial stress. *Brain Res*, 1049(2): 244–248
- Snyder J S, Glover L R, Sanzone K M, Kamhi J F, Cameron H A (2009). The effects of exercise and stress on the survival and maturation of adult-generated granule cells. *Hippocampus*, 19(10): 898–906
- Snyder J S, Grigereit L, Russo A, Seib D R, Brewer M, Pickel J, Cameron H A (2016). A Transgenic Rat for Specifically Inhibiting Adult Neurogenesis. *eNeuro*, 3(3): e0064-16.2016 1–13
- Snyder J S, Hong N S, McDonald R J, Wojtowicz J M (2005). A role for adult neurogenesis in spatial long-term memory. *Neuroscience*, 130(4): 843–852
- Snyder J S, Soumier A, Brewer M, Pickel J, Cameron H A (2011). Adult hippocampal neurogenesis buffers stress responses and depressive behaviour. *Nature*, 476(7361): 458–461
- Sousa N (2016). The dynamics of the stress neuromatrix. *Mol Psychiatry*, 21(3): 302–312
- Sultan S, Li L, Moss J, Petrelli F, Cassé F, Gebara E, Lopatar J, Pfrieger F W, Bezzi P, Bischofberger J, Toni N (2015). Synaptic Integration of Adult-Born Hippocampal Neurons Is Locally Controlled by Astrocytes. *Neuron*, 88(5): 957–972
- Surget A, Tanti A, Leonardo E D, Laugeray A, Rainer Q, Touma C, Palme R, Griebel G, Ibarguen-Vargas Y, Hen R, Belzung C (2011). Antidepressants recruit new neurons to improve stress response regulation. *Mol Psychiatry*, 16(12): 1177–1188
- Swaab D F, Bao A M, Lucassen P J (2005). The stress system in the human brain in depression and neurodegeneration. *Ageing Res Rev*, 4(2): 141–194
- Takamura N, Nakagawa S, Masuda T, Boku S, Kato A, Song N, An Y, Kitaichi Y, Inoue T, Koyama T, Kusumi I (2014). The effect of dopamine on adult hippocampal neurogenesis. *Prog Neuropsychopharmacol Biol Psychiatry*, 50: 116–124
- Tanapat P, Galea L A, Gould E (1998). Stress inhibits the proliferation of granule cell precursors in the developing dentate gyrus. *Int J Dev Neurosci*, 16(3-4): 235–239
- Tanti A, Belzung C (2013). Neurogenesis along the septo-temporal axis

- of the hippocampus: are depression and the action of antidepressants region-specific? *Neuroscience*, 252: 234–252
- Tanti A, Rainer Q, Minier F, Surget A, Belzung C (2012). Differential environmental regulation of neurogenesis along the septo-temporal axis of the hippocampus. *Neuropharmacology*, 63(3): 374–384
- Tasker J G (2006). Rapid glucocorticoid actions in the hypothalamus as a mechanism of homeostatic integration. *Obesity (Silver Spring)*, 14 (Suppl 5): 259S–265S
- Toni N, Laplagne D A, Zhao C, Lombardi G, Ribak C E, Gage F H, Schinder A F (2008). Neurons born in the adult dentate gyrus form functional synapses with target cells. *Nat Neurosci*, 11(8): 901–907
- Tronel S, Belnoue L, Grosjean N, Revest J M, Piazza P V, Koehl M, Abrous D N (2012). Adult-born neurons are necessary for extended contextual discrimination. *Hippocampus*, 22(2): 292–298
- Ursin H, Eriksen H R (2004). The cognitive activation theory of stress. *Psychoneuroendocrinology*, 29(5): 567–592
- Vallires L, Campbell I L, Gage F H, Sawchenko P E (2002). Reduced hippocampal neurogenesis in adult transgenic mice with chronic astrocytic production of interleukin-6. *J Neurosci*, 22(2): 486–492
- Van Bokhoven P, Oomen C A, Hoogendijk W J, Smit A B, Lucassen P J, Spijker S (2011). Reduction in hippocampal neurogenesis after social defeat is long-lasting and responsive to late antidepressant treatment. *Eur J Neurosci*, 33(10): 1833–1840
- Van der Borgh K, Meerlo P, Luiten P G, Eggen B J, Van der Zee E A (2005). Effects of active shock avoidance learning on hippocampal neurogenesis and plasma levels of corticosterone. *Behav Brain Res*, 157(1): 23–30
- van Praag H, Christie B R, Sejnowski T J, Gage F H (1999a). Running enhances neurogenesis, learning, and long-term potentiation in mice. *Proc Natl Acad Sci USA*, 96(23): 13427–13431
- van Praag H, Kempermann G, Gage F H (1999b). Running increases cell proliferation and neurogenesis in the adult mouse dentate gyrus. *Nat Neurosci*, 2(3): 266–270
- Veena J, Srikumar B N, Mahati K, Bhagya V, Raju T R, Shankaranarayana Rao B S (2009). Enriched environment restores hippocampal cell proliferation and ameliorates cognitive deficits in chronically stressed rats. *J Neurosci Res*, 87(4): 831–843
- Vinkers C H, Jols M, Milaneschi Y, Kahn R S, Penninx B W, Boks M P (2014). Stress exposure across the life span cumulatively increases depression risk and is moderated by neuroticism. *Depress Anxiety*, 31(9): 737–745
- Vivar C, Potter M C, Choi J, Lee J Y, Stringer T P, Callaway E M, Gage F H, Suh H, van Praag H (2012). Monosynaptic inputs to new neurons in the dentate gyrus. *Nat Commun*, 3: 1107
- Vivar C, Potter M C, van Praag H (2013). All about running: synaptic plasticity, growth factors and adult hippocampal neurogenesis. *Curr Top Behav Neurosci*, 15: 189–210
- Vivinetto A L, Surez M M, Rivarola M A (2013). Neurobiological effects of neonatal maternal separation and post-weaning environmental enrichment. *Behav Brain Res*, 240: 110–118
- Vollmayr B, Simonis C, Weber S, Gass P, Henn F (2003). Reduced cell proliferation in the dentate gyrus is not correlated with the development of learned helplessness. *Biol Psychiatry*, 54(10): 1035–1040
- Wang Q, Joels M, Swaab D F, Lucassen P J (2012). Hippocampal GR expression is increased in elderly depressed females. *Neuropharmacology*, 62(1): 527–533
- Wang Q, Van Heerikhuizen J, Aronica E, Kawata M, Seress L, Joels M, Swaab D F, Lucassen P J (2013). Glucocorticoid receptor protein expression in human hippocampus; stability with age. *Neurobiol Aging*, 34(6): 1662–1673
- Wei Q, Hebda-Bauer E K, Pletsch A, Luo J, Hoversten M T, Osetek A J, Evans S J, Watson S J, Seasholtz A F, Akil H (2007). Overexpressing the glucocorticoid receptor in forebrain causes an aging-like neuroendocrine phenotype and mild cognitive dysfunction. *J Neurosci*, 27(33): 8836–8844
- Westenbroek C, Den Boer J A, Veenhuis M, Ter Horst G J (2004). Chronic stress and social housing differentially affect neurogenesis in male and female rats. *Brain Res Bull*, 64(4): 303–308
- Wilson C B, Ebenezer P J, McLaughlin L D, Francis J (2014). Predator exposure/psychosocial stress animal model of post-traumatic stress disorder modulates neurotransmitters in the rat hippocampus and prefrontal cortex. *PLoS ONE*, 9(2): e89104
- Winocur G, Wojtowicz J M, Sekeres M, Snyder J S, Wang S (2006). Inhibition of neurogenesis interferes with hippocampus-dependent memory function. *Hippocampus*, 16(3): 296–304
- Wojtowicz J M, Askew M L, Winocur G (2008). The effects of running and of inhibiting adult neurogenesis on learning and memory in rats. *Eur J Neurosci*, 27(6): 1494–1502
- Wong E Y, Herbert J (2004). The corticoid environment: a determining factor for neural progenitors' survival in the adult hippocampus. *Eur J Neurosci*, 20(10): 2491–2498
- Wong E Y, Herbert J (2005). Roles of mineralocorticoid and glucocorticoid receptors in the regulation of progenitor proliferation in the adult hippocampus. *Eur J Neurosci*, 22(4): 785–792
- Wong E Y, Herbert J (2006). Raised circulating corticosterone inhibits neuronal differentiation of progenitor cells in the adult hippocampus. *Neuroscience*, 137(1): 83–92
- Wu M V, Shamy J L, Bedi G, Choi C W, Wall M M, Arango V, Boldrini M, Foltin R W, Hen R (2014). Impact of social status and antidepressant treatment on neurogenesis in the baboon hippocampus. *Neuropsychopharmacology*, 39(8): 1861–1871
- Yu S, Patchev A V, Wu Y, Lu J, Holsboer F, Zhang J Z, Sousa N, Almeida O F (2010). Depletion of the neural precursor cell pool by glucocorticoids. *Ann Neurol*, 67(1): 21–30
- Yun S, Donovan M H, Ross M N, Richardson D R, Reister R, Farnbauch L A, Fischer S J, Riethmacher D, Gershenfeld H K, Lagace D C, Eisch A J (2016). Stress-Induced Anxiety- and Depressive-Like Phenotype Associated with Transient Reduction in Neurogenesis in Adult Nestin-CreERT2/Diphtheria Toxin Fragment A Transgenic Mice. *PLoS ONE*, 11(1): e0147256
- Zhang C L, Zou Y, He W, Gage F H, Evans R M (2008). A role for adult TLX-positive neural stem cells in learning and behaviour. *Nature*, 451(7181): 1004–1007
- Zhao C, Deng W, Gage F H (2008). Mechanisms and functional implications of adult neurogenesis. *Cell*, 132(4): 645–660