UvA-DARE (Digital Academic Repository)

Detection of Rotational CO Emission From the Red-supergiants in the Large Magellanic Cloud

Matsuura, M.; Sargent, B.; Swinyard, B.; Yates, J.A.; Royer, P.; Barlow, M.J.; Boyer, M.L.; Decin, L.; Khouri, T.; Meixner, M.; van Loon, J.Th.; Woods, P.M.

DOI

[10.1051/eas/1571010](https://doi.org/10.1051/eas/1571010)

Publication date

2015

Document Version

Final published version

Published in

EAS Publications Series

[Link to publication](#)

Citation for published version (APA):

Matsuura, M., Sargent, B., Swinyard, B., Yates, J. A., Royer, P., Barlow, M. J., Boyer, M. L., Decin, L., Khouri, T., Meixner, M., van Loon, J. T., & Woods, P. M. (2015). Detection of Rotational CO Emission From the Red-supergiants in the Large Magellanic Cloud. *EAS Publications Series*, 71-72, 53-56. <https://doi.org/10.1051/eas/1571010>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

DETECTION OF ROTATIONAL CO EMISSION FROM THE RED-SUPERGIANTS IN THE LARGE MAGELLANIC CLOUD

M. Matsuura^{1,2}, B. Sargent³, B. Swinyard^{2,4}, J.A. Yates², P. Royer⁵,
M.J. Barlow², M.L. Boyer⁶, L. Decin^{5,7}, T. Khouri^{7,8}, M. Meixner^{9,10},
J.Th. van Loon¹¹ and P.M. Woods¹²

Abstract. It is yet well understood how mass-loss rates from evolved stars depend on metallicities. With a half of the solar metallicity and the distance of only 50 kpc, the evolved stars of the Large Magellanic Cloud (LMC) are an ideal target for studying mass loss at low metallicity. We have obtained spectra of red-supergiants in the LMC, using the Hershel Space Observatory, detecting CO thermal lines from $J=6-5$ up to 15-14 lines. Modelling CO lines with non-LTE Radiative transfer code suggests that CO lines intensities can be well explained with high gas-to-dust ratio, with no obvious reduction in mass-loss rate at the LMC. We conclude that the luminosities of the stars are dominant factors on mass-loss rates, rather than the metallicity.

¹ School of Physics and Astronomy, Cardiff University, Cardiff CF24 3AA, UK

² University College London, London WC1E 6BT, UK

³ Center for Imaging Science and Laboratory for Multiwavelength Astrophysics, Rochester Institute of Technology, 54 Lomb Memorial Drive, Rochester, NY 14623, USA

⁴ Rutherford Appleton Laboratory, Chilton, Didcot, Oxfordshire OX11 0QX, UK

⁵ Instituut voor Sterrenkunde, KU Leuven, Celestijnenlaan 200D 2401, 3001 Leuven, Belgium

⁶ Observational Cosmology Lab., Code 665, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

⁷ Astronomical Institute Anton Pannekoek, University of Amsterdam, PO Box 94249, 1090 GE Amsterdam, The Netherlands

⁸ Onsala Space Observatory, Dept. of Radio and Space Science, Chalmers University of Technology, 43992 Onsala, Sweden

⁹ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

¹⁰ The Johns Hopkins University, Department of Physics and Astronomy, 366 Bloomberg Center, 3400 N. Charles Street, Baltimore, MD 21218, USA

¹¹ School of Physical & Geographical Sciences, Lennard-Jones Laboratories, Keele University, Staffordshire ST5 5BG, UK

¹² Astrophysics Research Centre, School of Mathematics & Physics, Queen's University, University Road, Belfast BT7 1NN, UK

1 Introduction

During the late phase of the stellar evolution, stars lose a large fraction of their mass from their surfaces. High-mass ($> 8 M_{\odot}$) stars evolve into the red-supergiant (RSG) phase, and lose a large amount of mass through stellar winds. In order to fully understand post-main sequence stellar evolution, it is important to characterise the mass loss processes of RSGs. Due to this dust-driven mechanism, theory predicts that the mass-loss rate from AGB stars, and most likely RSGs, should mainly depend on the luminosity of the central star, and the metallicity of the galaxy (*e.g.* Bowen & Willson 1991).

With the aim of investigating the physics and chemistry of the circumstellar envelopes of evolved stars at low metallicity, we have measured far-infrared (IR) and submillimeter molecular emission lines of two IR RSGs in the LMC, which has approximately half a solar metallicity. We report the first detection of the CO rotational emission lines from RSGs in the LMC.

2 Observations and analysis

We observed two RSGs with the *Herschel* Space Observatory (Pilbratt *et al.* 2010). The primary target, IRAS 05280–6910 was chosen because it is the brightest LMC RSG at far-infrared wavelengths (Boyer *et al.* 2010), with the $250 \mu\text{m}$ flux of 205 mJy. The submillimeter spectrum of IRAS 05280–6910 was obtained, using the SPIRE Fourier Transform spectrometer (Griffin *et al.* 2010; Swinyard *et al.* 2010) on board the *Herschel* Space Observatory. Figure 1 shows the SPIRE spectrum of IRAS 05280–6910 in comparison to the spectrum of the Galactic red-supergiant VY CMa (Royer *et al.* 2010; Matsuura *et al.* 2014). CO rotation lines and some strong H_2O lines are detected. The secondary target is WOH G64, which is one of the most luminous RSGs.

Fig. 1. The SPIRE FTS spectra of the LMC red-supergiant, IRAS 05280–6910, compared with the SPIRE FTS spectra of the Galactic red-supergiant, VY CMa. CO and H_2O are detected in the spectrum of IRAS 05280–6910.

Fig. 2. The spectral line energy distribution rotation diagram of IRAS 05280–6910, VY CMa (Matsuura *et al.* 2014) and W Hya (Khouri *et al.* 2014), with a single CO measurement of WOH G64. The non-LTE model fits are displayed with dotted lines.

Fig. 3. The mass-loss rate ($M_{\odot} \text{ yr}^{-1}$) *vs.* luminosity ($L_{\odot}$) for oxygen-rich AGB stars and red-supergiants in the Galaxy and the LMC. The Galactic data are taken from DeBeck *et al.* (2010), and the LMC data are from Groenewegen *et al.* (2009). The solid line shows the mass-loss *vs.* luminosity relation fitted to Galactic sample, and the dashed line shows the mass-loss *vs.* luminosity relation for 3500 K stars in the LMC (van Loon *et al.* 2005).

A non-local thermodynamic equilibrium (non-LTE) line radiative transfer code was used to analyse the CO thermal emission. The non-LTE modelling of CO lines shows the gas-mass loss rate of $3 \times 10^{-4} M_{\odot} \text{ yr}^{-1}$. For this analysis, we adopted the gas-to-dust ratio of 500, and CO/H₂ of 2.7×10^{-4} , considering the lower abundance of refractory elements and carbon elements in the LMC. Details of analysis and modelling are described by Matsuura *et al.* (2015).

Reduced mass-loss rates are predicted at low metallicity for oxygen-rich AGB stars and RSGs (Bowen & Willson 1991). However, no metallicity effects on mass-loss rate was found for RSGs at least in the range of half of the solar metallicity.

Instead, we found that CO lines and mass-loss rate largely depends on luminosity/luminosity class. Figure 3 shows the mass-loss rate *vs.* luminosity relation for LMC and Galactic evolved stars. The Galactic evolved stars tend to follow an increasing trend of mass-loss rate with higher luminosity.

Dr. Olivier Chesneau helped us interpreting VLTI high-angular resolution observations, of post-AGB star, OH 231.8+4.2 (Matsuura *et al.* 2006). He introduced his then PhD student, Eric Lagadec, who became our long-standing collaborator, and one of the organisers of this nice conference. We will miss Olivier. M.M. is saddened by the death of Prof. Bruce Swinyard, who led the SPIRE FTS construction and development. He helped this work by leading the development of data reduction procedure for faint objects.

References

- Boyer, M.L., Sargent, B.A., van Loon, J.T., *et al.*, 2010, A&A, 518, L142
 Bowen, G.H., & Willson, L.A., 1991, ApJ, 375, L53
 De Beck, E., Decin, L., de Koter, A., *et al.*, 2010, A&A, 523, A18
 Griffin, M.J., *et al.*, A.A., 2010, A&A, 518, L3
 Groenewegen, M.A.T., Wood, P.R., Sloan, G.C., *et al.*, 2007, MNRAS, 376, 313
 Groenewegen, M.A.T., Sloan, G.C., Soszynski, I., & Petersen, E.A., 2009, A&A, 506, 1277
 Khouri, T., De Koter, A., Decin, L., *et al.*, 2014, A&A, 570, A67
 Meynet, G., Chomienne, V., Ekstrom, S., *et al.*, 2015, A&A, 575, A60
 Knapp, G.R., & Morris, M., 1985, ApJ, 292, 640
 Matsuura, M., Chesneau, O., Zijlstra, A.A., *et al.*, 2006, ApJ, 646, L123
 Matsuura, M., Yates, J.A., Barlow, M.J., *et al.*, 2014, MNRAS, 437, 532
 Matsuura, M., Sargent, B., Swinyard, B., *et al.*, MNRAS, submitted
 Pilbratt, G.L., Riedinger, J.R., Passvogel, T., *et al.*, 2010, A&A, 518, L1
 Ramstedt, S., Schöier, F.L., Olofsson, H., & Lundgren, A.A., 2008, A&A, 487, 645
 Royer, P., Decin, L., Wesson, R., *et al.*, 2010, A&A, 518, L145
 Swinyard, B.M., Ade, P., Baluteau, J.-P., *et al.*, 2010, A&A, 518, L4
 van Loon, J.T., Cioni, M.-R.L., Zijlstra, A.A., & Loup, C., 2005, A&A, 438, 273