UvA-DARE (Digital Academic Repository)

Hydrological Connectivity Does Change Over 70 Years of Abandonment and Afforestation in the Spanish Pyrenees

López-Vicente, M.; Nadal-Romero, E.; Cammeraat, E.L.H.

DOI

[10.1002/ldr.2531](https://doi.org/10.1002/ldr.2531)

Publication date

2017

Document Version

Final published version

Published in

Land Degradation & Development

License

Article 25fa Dutch Copyright Act

[Link to publication](#)

Citation for published version (APA):

López-Vicente, M., Nadal-Romero, E., & Cammeraat, E. L. H. (2017). Hydrological Connectivity Does Change Over 70 Years of Abandonment and Afforestation in the Spanish Pyrenees. *Land Degradation & Development*, 28(4), 1298–1310. <https://doi.org/10.1002/ldr.2531>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

HYDROLOGICAL CONNECTIVITY DOES CHANGE OVER 70 YEARS OF ABANDONMENT AND AFFORESTATION IN THE SPANISH PYRENEES

Manuel López-Vicente^{1*}, Estela Nadal-Romero^{2,3}, Erik L. H. Cammeraat²¹Department of Soil and Water, EEAD-CSIC, Avda. Montañana 1005, Zaragoza 50059, Spain²Institute for Biodiversity and Ecosystem Dynamics (IBED), University of Amsterdam, PO Box 94240, Amsterdam 1090 GE, The Netherlands³Department of Geography, Environmental Sciences Institute, University of Zaragoza, Pedro Cerbuna 12, Zaragoza 50009, Spain

Received: 17 February 2016; Revised: 11 April 2016; Accepted: 11 April 2016

ABSTRACT

Runoff connectivity depends on topography, rainfall, man-made elements (terraces, trails, roads and drainage systems) and vegetation. In this study, we quantified the effects of 70 years of human activities on runoff connectivity in the mountainous Aragón afforested sub-catchment (17.2 ha; Central Spanish Pyrenees). The IC index of hydrological connectivity was chosen to perform this metric over six land use scenarios at high spatial resolution (1 × 1 m of cell size). The current scenario (year 2012) was simulated with three flow accumulation algorithms (MD, MD8 and D8). MD8 was linked with the most frequent hydrological response of the sub-catchment (rainfall intensity and stream flow during 7 years) and generated the most representative pattern of connectivity, especially in the linear landscape elements (LLE). This algorithm was chosen to simulate five past scenarios (1945, 1956, 1973, 1980 and 2006). In all scenarios, the highest connectivity appeared related to trails and roads, as well as to streams and gullies, whereas the lowest appeared related to stonewalls in 1945 and 1956 to hillslopes in 1973, and the following afforestation. Changes in connectivity mainly depended on the changes in the vegetation factor and in a minor way in the total length, spatial location and type of LLE. Afforestation promoted lower and more stable connectivity at both local and catchment scales. Copyright © 2016 John Wiley & Sons, Ltd.

KEY WORDS: runoff connectivity; stream flow; land abandonment; afforestation; IC model

INTRODUCTION

The connectivity concept allows investigating the effect of heterogeneities on the general behaviour of a system that contains those heterogeneities, and it is an essential factor when modelling dissimilar systems (Antoine *et al.*, 2009). Reaney *et al.* (2014) related hydrological connectivity to the time required for upslope-generated runoff and sediment fluxes to reach an efficient flow channel. According to these authors, including Bracken & Croke (2007), runoff connectivity depends on climate, hillslope runoff potential, landscape position, delivery pathway and lateral connectivity. In the last decade, the attention for runoff connectivity increased to serve as a geostatistical tool in hydrology and geomorphology (Parsons *et al.*, 2015) and became a key factor to understand the redistribution dynamics of the soil components in the different compartments of the landscape (Chartin *et al.*, 2013). In this study, we quantified the effects of 70 years of human activities (land abandonment, new infrastructures and afforestation) on runoff connectivity throughout a mountainous sub-catchment in the Central Spanish Pyrenees.

During low-intensity rainfall events, the kinematic dispersion of runoff is dominant, whereas geomorphologic dispersion becomes more dominant for larger intensities (Rossel

et al., 2014). Extreme events activate the entire catchment (Lana-Renault *et al.*, 2014), whereas some parts do not significantly contribute to runoff during low-intensity and medium-intensity events (López-Vicente *et al.*, 2013a). In Mediterranean and other dryland environments that are characterized by patchy vegetation cover (Foerster *et al.*, 2014), the size, length, spatial distribution and connectivity between the different patches determine the magnitude of the runoff processes (Cammaraat, 2004).

Human organization of landscape elements has a significant control on runoff and soil redistribution processes, so that similar events can produce very different responses (and vice versa) (Lesschen *et al.*, 2009). In the second half of the past century, many European and Mediterranean mountainous agricultural landscapes suffered from land abandonment and depopulation but also were subject to significant afforestation (García-Ruiz & Lana-Renault, 2011). In the Spanish mountains, afforestation with conifers was used for land reclamation during many years, causing landscape disturbances. The effects at catchment scale were expressed by a clear decrease of connectivity, lower peak flows and runoff and sediment supply to streams (Ortigosa & García-Ruiz, 1995; Sanjuán *et al.*, 2014). Construction of paths, forest roads and firewalls influences runoff connectivity in forested catchments (Croke & Mockler, 2001). Stonewalls and agricultural terraces form an important feature in the Mediterranean landscapes and have an important and clear effect, reducing soil loss, runoff coefficients and connectivity (Lesschen *et al.*, 2009; Arnáez *et al.*, 2015).

*Correspondence to: M. López-Vicente, Department of Soil and Water, EEAD-CSIC, Avda. Montañana 1005, Zaragoza 50059, Spain.
E-mail: mvlcente@eead.csic.es; mvlcente@eead.csic.es

Land abandonment promoted the mismanagement and collapse of these terraces increasing runoff connectivity and soil erosion (López-Vicente *et al.*, 2013b). However, slope terracing without any supporting structure is an aggressive and common man-made technique in afforestation practices (Linares *et al.*, 2002).

In the last decade, several hydrological connectivity indices appeared in the literature, most of them based on geographic information system and stream-power approaches, such as the 'volume to breakthrough' of Bracken & Croke (2007) and the Network Index of Lane *et al.* (2009). Recently, Reaney *et al.* (2014) used the Connectivity of Runoff Model, 2D version, in semi-arid environments. A few years ago, Borselli *et al.* (2008) developed the index of runoff and sediment connectivity (IC) testing successfully this approach in a large catchment in Central Italy against field observations. Afterwards, Cavalli *et al.* (2013) proposed some improvements on the original equation related to very steep slopes. Gay *et al.* (2015) executed the index in Central France and improved it for lowlands considering infiltration properties. The ability of the IC model has been proven in different fields and catchments around the world, such as the studies of Heckmann & Schwanghart (2013) in Austria, D'Haen *et al.* (2013) in Turkey, Vigiak *et al.* (2012) in Australia and Chartin *et al.* (2013) in Japan. In north-eastern Spain (López-Vicente *et al.*, 2013b; Foerster *et al.*, 2014), several studies proved its capacity to map runoff and sediment connectivity.

There are no studies dealing with the long-term effects of combined land use changes on runoff connectivity, especially after afforestation practices. Therefore, we did the following: (i) evaluated the current conditions (year 2012: no crops, tall pines, three trails, one unpaved forest road and many broken stonewalls) of runoff connectivity using three flow accumulation algorithms and the IC model; (ii) analysed the relationship between each pattern of connectivity and the hydrological response at the sub-catchment outlet (stream flow and rainfall depth) during 7 years (October 2007–September 2014); (iii) simulated the runoff connectivity during five past scenarios (1945, 1956, 1973, 1980 and 2006) with the most appropriate algorithm and the IC model, and the land use and linear landscape elements (LLE) of each scenario; (iv) evaluated the role of the combined and purely human-induced changes on runoff connectivity for six scenarios; and (v) proposed potential management practices to reduce the negative impacts of high runoff connectivity.

MATERIALS AND METHODS

Study Area

The study area corresponds to the afforested sub-catchment of the Araguás catchment (Rebulella stream) that is located in the Central Spanish Pyrenees near Jaca town (Figure 1A) (42°36'15.18"N, 0°37'27.10"W). The elevation ranges between 899 and 1,159 m above sea-level, and the total extension is 17.2 ha. The landscape is hilly with a minimum and average slope steepness of 4% and 43%, respectively. There

is a lack of flat areas, and apart from the main gully that works as the stream, there are several small gullies directly connected to it. In the lower part and near the gauging station, badlands are present (Nadal-Romero *et al.*, 2010). The area represented an intensely cultivated landscape until the 1950s, which was afforested with coniferous during the 1960s and 1970s (*Pinus sylvestris* and *Pinus nigra*) (Figure 1B). Modified soils show truncation signs in the surface horizons, low organic matter content and poor structure (Nadal-Romero *et al.*, 2016). The climate is sub-Mediterranean with Oceanic and Continental influences. Annual rainfall varies between 500 and 1,000 mm (average annual rainfall approximately 800 mm), with two major rainfall periods (autumn and spring); convective storms usually occur in summer and occasional snowfalls in winter (Nadal-Romero *et al.*, 2008).

Land Use Change and Scenarios

A total of six aerial photos were used from the digital aerial photo repository of the Spanish National Geographic Institute (<http://fototeca.cnig.es/>). These photos covered a period of almost 70 years: 1945, 1956, 1973, 1980, 2006 and 2012 (Figure 1C). The four oldest photos had no geometric correction, and 15 geographic control points were used to identify the boundary of the sub-catchment. The two more recent images were orthophotos, and the complete boundary of the sub-catchment was derived from the digital elevation model (DEM). In the oldest scenario (1945), almost the entire sub-catchment was cultivated with cereals (wheat and barley), and hardly any trees were present and mainly located along the stream. Many stonewalls (~1.2-m height) and terraces controlled the size and shape of the fields, and two narrow paths (~1.5 m wide) crossed the hillslope (Table I). In 1956, almost all fields were continuously cultivated, but mismanagement practices of the LLE triggered the breakdown of some of these structures and thus an increment in the number and length of ephemeral gullies. The third scenario (1973) described the early stage (short pines) of the massive coniferous plantation (59% of total surface). Most LLE were still preserved although walls and terraces showed many breaking points. A new narrow trail appeared in the upper part related to the afforestation works. In 1980, the trees were taller, and a new unpaved forest road (~3 m wide) appeared. In 2006, the canopy of the pines was well developed and many shrubs covered the abandoned fields and open areas. In the current scenario (2012; Figure 2A), more sections of terraces and stonewalls were disintegrated because of the long duration of abandonment. The current height of the *P. sylvestris* plantation is ~6 m whereas *P. nigra* pines are slightly taller, ~6.5-m height (Figure 2B). Some small areas underwent natural plant colonization (secondary succession) with plant species like *Genista scorpius*, *Juniperus communis*, *Rosa gr. canina* and *Buxus sempervirens* (~1 m tall), as well as some stands of young *P. sylvestris*. The different vegetation types appear as heterogeneous patches with many spots of open scrubland and bare soil, with a density of 81.9 patches per hectare.

Figure 1. Location of the Araguás catchment within the Aragón catchment in NE Spain (Central Pyrenees) (A). Images of the pine forest plantation on terraces and a forest road (B). Old aerial photos and recent orthophotos of the study area over six different years in a 70-year period (C). [Colour figure can be viewed at wileyonlinelibrary.com]

Field Data and Monitoring

The sub-catchment is equipped with two tipping buckets that collect rainfall every 5 min. At the outlet of the sub-catchment, stream flow is measured at a gauging station (V-notch weir) using a water pressure sensor (Keller DCX-22 AA, KELLER, Winterthur, Switzerland; Figure 2C). A 7-year record (1 October 2007–30 September 2014) of rainfall depth and stream flow was measured and analysed. We derived a high-spatial-resolution DEM at 1×1 m of cell size from the lidar-derived DEM at 5×5 m of cell size, generated by the Spanish National Geographic Institute (IGN; <http://centrodedescargas.cnig.es/CentroDescargas/>). We selected this spatial resolution to guarantee that the narrow and in some places short LLE were included in the simulation.

The Index of Runoff and Sediment Connectivity Model

We used part of the modifications made by Cavalli *et al.* (2013) on the IC proposed by Borselli *et al.* (2008). As the study site does not present flat areas, we did not consider the modifications made by Gay *et al.* (2015) for lowlands. The IC model accounts the characteristics of the drainage area (D_{up} , upslope module) and the flow path length that a particle has to travel to arrive at the nearest sink (D_{dn} , downslope module):

$$IC_K = \log_{10} \left(\frac{D_{up,K}}{D_{dn,K}} \right) = \log_{10} \left(\frac{\overline{W_K} \cdot \overline{S_K} \cdot \sqrt{A_K}}{\sum_{i=K,n_K} \frac{d_i}{W_i \cdot S_i}} \right) \quad (1)$$

Where: $\overline{W}$ is the average weighing factor of the upslope

Table I. Land use characteristics and average values of the C-RUSLE factor for each scenario and map, and description and length of the linear landscape elements

Scenario	Land use and land cover			Linear landscape elements					
	Description	C-RUSLE			All elements	Total length (m)		Change relative to 2012 values (%)	Description
		M	SD	Change relative to 2012 values (%)		All stonewall and terraces	All trails and forest road		
1945	85% surface as cereal fields. Trees only appeared near the stream.	0.2752	0.0710	227.9	5,216.9	4,550.6	666.3	69.8%	2 trails. Most stonewalls and terraces well preserved: few breakpoints.
1956	84% surface as cereal fields. First abandoned fields near the divide. Trees only appeared near the stream.	0.2747	0.0711	228.7	4,302.7	3,636.4	666.3	40.0	2 trails. Many small breakpoints in the stonewalls and terraces.
1973	Afforestation (59% of total surface): early stage, short pines (~1-m height). Low dense scrublands.	0.1418	0.0987	69.6	3,090.9	2,225.9	865.0	0.6	3 trails: 2 old plus 1 new linked to the afforestation. More breakpoints in the stonewalls and terraces.
1980	Afforestation: short–medium pines (~2-m height). Low–medium dense scrublands. Very few areas with natural forest.	0.1118	0.1209	33.8	3,193.0	1,847.4	1,345.6	3.9	1 new forest road and 3 trails. More breakpoints in the stonewalls and terraces.
2006	Afforestation: tall pines (~6.1-m height). 3 recent abandoned fields. Medium-dense scrublands. Small areas with natural forested areas.	0.0839	0.1388	0.4	3,121.5	1,606.6	1,514.9	1.6	1 forest road and 3 trails (2 old, 1 new and 1 abandoned). Many breakpoints in the stonewalls and terraces.
2012	Afforestation: tall pines (~6.3-m height on average). 3 recent abandoned fields. Dense scrublands.	0.0836	0.1388		3,073.0	1,558.1	1,514.9		1 forest road and 3 trails. Many breakpoints in the stonewalls and terraces.

Figure 2. Map of current land use in the afforested Araguás sub-catchment (A). Images of stonewalls and trails in the study area (B) and of the gauging station at the outlet (C). [Colour figure can be viewed at wileyonlinelibrary.com]

contributing area (dimensionless and equal to the C-RUSLE factor), $\bar{S}$ is the average slope gradient of the upslope contributing area (m m^{-1}), A is the upslope contributing area (m^2), d_i is the length of the i th cell along the downslope path (m), W_i is the weight of the i th cell (dimensionless) and S_i is the slope gradient of the i th cell (m m^{-1}). Values of slope steepness lower than 0.005 must be replaced by the value $S_i = 0.005$, and those higher than 1 must be set to a maximum value of 1. The subscript K indicates that each cell i has its own IC value. This index is defined in the range of $[-\infty, +\infty]$, and connectivity increases when IC grows towards $+\infty$. The role of the LLE (agricultural terraces, stonewalls, trails and roads) is added by modifying the original map of flow direction with a mask with two values, 0 for the LLE and 1 for the remaining area.

Connectivity changes because of changes in runoff magnitude; thus, we computed different IC maps with three flow accumulation algorithms for the current scenario. We tested algorithms with single direction (D8), multiple directions (MD) and multiple directions with threshold value for linear flow (MDD8) (more details in López-Vicente & Navas, 2010; López-Vicente *et al.*, 2014). $\bar{W}$, $\bar{S}$ and A were also calculated with these three algorithms. The 'stream mask'

was associated with the area where the gully showed a permanent stream (see Appendix A in Borselli *et al.*, 2008). We used the values of the C-RUSLE factor obtained by López-Vicente & Navas (2009) and López-Vicente *et al.* (2011) in two catchments of the Pyrenees with similar vegetation as that of the study area. For badlands and gullies, we used the value proposed by Capolongo *et al.* (2008) in a semi-arid area in Southern Italy. For the low-elevation (early stage of plantation) and middle-elevation and high-elevation (current scenario) pine forest, we used the values proposed by De Tar *et al.* (1980) and Miller *et al.* (2003).

RESULTS

Current Scenario of Stream Flow and Runoff Connectivity

The average annual rainfall (R) and stream flow (Q) during the seven hydrological years were 783 and 244 mm y^{-1} (Figure 3A), obtaining an average runoff coefficient of 31%. A total of 1,339 rainfall events (e) were recorded, and only 13% of these events ($n = 168$) were considered as erosive events (ee) (Figure 3B). According to Renard *et al.* (1997), an erosive event had a total rainfall depth equal to

Figure 3. Daily rainfall (R) and stream flow (Q) depth during the study period (A). Monthly number of rainfall events (B) and of the maximum rainfall intensity during all the rainfall events (I_{30-e}) and the erosive events (I_{30-ee}) (C). Average monthly values of R , Q , I_{30-e} , I_{30-ee} and percentage of erosive events (ee) (D). [Colour figure can be viewed at wileyonlinelibrary.com]

or higher than 12.7 mm or lower than 12.7 mm but with a minimum maximum rainfall intensity of 6.35 mm in 15 min. Values of R , Q and maximum rainfall intensity during all events (I_{30-e}) including the erosive events (I_{30-ee}) showed a marked variability (Figure 3C). The mean values of R , Q , I_{30-e} and I_{30-ee} were 61.8 and 19.8 mm month⁻¹ and 3.1 and 10.2 mm h⁻¹, respectively, and their coefficients of variation were 0.7, 1.3, 1.0 and 1.1. Values of daily rainfall did not correlate well with those of stream flow ($r=0.237$). Thus, we selected the monthly values of Q and I_{30-ee} to describe and identify three different hydrological periods (HyP). HyP1 is from January to March when the highest values of Q (~41.2 mm month⁻¹) were registered and in August when the highest values of I_{30-ee} (~29.2 mm h⁻¹) appeared. This period represented the most intense hydrological response of the sub-catchment. HyP2 had the longest period (6 months), from April to July and from September to October, when moderate values of Q and I_{30-ee} were registered, and it represented the most predominant conditions. The short HyP3, from November to December, represented the period with the lowest activity, when the lowest values of Q (~8.4 mm month⁻¹) and I_{30-ee} (~4.4 mm h⁻¹) were observed (Figure 3D).

In the current scenario, the total length of the LLE was 3,073 m, and the C-RUSLE factor had a mean value of 0.0836 (Figure 4 and Table I). The IC map obtained with the MD algorithm represented dispersed overland flow patterns without the presence or limited occurrence of concentrated runoff (Figure 5A). The IC map with the MD8

algorithm represented the occurrence of concentrated runoff from a threshold value downwards and disperse flow near the divides (Figure 5B). The threshold value was associated with the initiation of the gullies and the permanent rills of the ruined terraces. The IC map with the D8 algorithm represented very concentrated overland flow from divides to the stream (Figure 5C). Each map showed a different mean value and range because of the specific characteristics of each algorithm. Because the different values of rainfall intensity and depth trigger different patterns of runoff, we linked the IC-MD map with the HyP3 period, the IC-MD8 map with HyP2 and the IC-D8 map with HyP1. The frequency histograms of the three maps of connectivity showed a similar pattern although the histogram of the MD8 approach was the closest to a normal distribution (Figure 5D). Finally, the MD8 algorithm was the most appropriate to describe the average spatial pattern and variability of runoff connectivity, and thus, we ran the IC model with this algorithm in five past scenarios.

The highest values of connectivity appeared on the road and trails, followed by stonewalls and terraces, and in the stream and gullies (Table II). Lower values of connectivity appeared upslope the forest road as compared with downslope, highlighting the key role of this infrastructure on the magnitude of connectivity throughout the hillslopes. The afforested areas presented the lowest values of connectivity owing to the protective effect of the pine plantation, as reflected in the low values of the C-RUSLE factor of both pine species.

Figure 4. Maps of linear landscape elements (LLE) for the six simulated scenarios. [Colour figure can be viewed at wileyonlinelibrary.com]

Runoff Connectivity During the Past Scenarios

Values and spatial patterns of connectivity clearly changed in the five past scenarios in comparison with the current situation (Figure 6 and Table III). The analysis was performed from the most recent past scenario to the oldest. In 2006, the total length of the LLE was slightly longer, and differences in vegetation were small (Figure 4 and Table I). The mean connectivity in 2006 was the lowest of the six scenarios although it was very similar to that in 2012 (Figure 6E). In 1980, the total length of the LLE was 4% longer than the total current length although the mean connectivity was 19% higher owing to the small/medium size of the pine plantation (Figure 6D). The abandonment of an old trail and the creation of a new one located upslope modified the spatial patterns and values of connectivity in 2006 in comparison with those in 1980. In 1973, the total length of the LLE was quite similar to that in 1980, although in 1973 the forest road that controlled the spatial patterns of connectivity did not yet exist in 1980 and in later scenarios. Hence, the changes in connectivity in 1980 in comparison with 1973 were explained by both the changes in the C-RUSLE factor and the spatial location of the LLE (Figures 6C and 7A).

In 1956 and 1945, the lengths of the LLE were, respectively, 40% and 70% longer than in 2012 (Figure 4), and

the land use maps were clearly different, with many cereal fields. Hence, the maps of connectivity in 1945 and 1956 showed different spatial patterns and the highest values of connectivity of the six scenarios (Figure 6A and B). The collapse of some sections of stonewalls in 1956 and the main land use (crop) resulted in the highest values of connectivity in that year. The mean connectivity in 1945 was 42% higher than in 2012, and conversely connectivity in 2012 was 82% lower than in 1945 (Figure 6A and F). Surfaces that currently present higher values of connectivity only affected 6% of the total surface and were associated with a forest road, a new trail, some sections of stonewalls, abandoned fields and small areas near the stream and bare lands. Areas showing a decrease in connectivity were predominant and affected 94% of the surface. During the six scenarios, the highest values of connectivity appeared on the trails and road, followed by stream and gullies, whereas the lowest values appeared in stonewalls and terraces in 1945 and 1956, and for the hillslopes in 1973 and the following scenarios due to the afforestation (Table III).

The density of LLE also modified the mean and maximum lengths of the flow path lines (FPL) in each scenario (Table III). The 1945 presented the shortest mean value of the FPL due to the highest density of LLE in the study area.

Figure 5. Maps of runoff connectivity calculated with the multiple direction (MD) (A), multiple direction with threshold value for linear flow (MD8) (B) and single direction (D8) (C) algorithms. Frequency histograms of the values of runoff connectivity for each map (D). [Colour figure can be viewed at wileyonlinelibrary.com]

This value increased in 1956 and reached the highest value in 1973 because of the tumble down of many sections of the stonewalls. The construction of the forest road reduced the mean and maximum lengths of the FPL in 1980 and next scenarios. The beginning of afforestation in 1973 slightly reduced the maximum length of FPL. The variability of the IC values at short distance, estimated by analysing the values of the eight surrounding pixels of each pixel, also changed during the six scenarios (Table III). The highest local variability appeared in 1945 and 1956, whereas the most stable local

patterns appeared in 2012. Hence, afforestation promoted lower and more stable values of runoff connectivity.

DISCUSSION

The different patterns and values of connectivity that we found for the Araguás sub-catchment throughout the year were also revealed by Zimmermann *et al.* (2014) at monthly resolution, and in the forests of the Panama Canal Watershed, supporting the importance of selecting an adequate

Table II. Values of runoff connectivity for the current scenario (year 2012) in the main physiographic and land use units, calculated with the three flow accumulation algorithms

Algorithm	IC value	Total area	Hillslopes ^a	LLEs ^b	Stream and gullies	Upslope forest road ^c	Road and trails	Downslope forest road ^c	Afforested area
MD	Mean	-4.0878	-4.2086	-2.5305	-3.0340	-4.3163	-2.1004	-3.8535	-4.4902
	SD	0.9816	0.8497	1.4182	1.1023	0.8733	1.2867	0.9705	0.6601
MD8	Mean	-4.0858	-4.2073	-2.4965	-3.0524	-4.3080	-2.0535	-3.8627	-4.4910
	SD	1.0150	0.8844	1.4620	1.1362	0.9115	1.3356	0.9996	0.7101
D8	Mean	-4.5097	-4.6639	-2.5180	-3.1666	-4.7416	-2.0338	-4.2819	-5.0798
	SD	1.0873	0.9150	1.4508	1.1563	0.9461	1.2596	1.1069	0.6056

LLE, linear landscape elements.

^aExcluding all LLEs.

^bAll types of LLEs.

^cIncluding afforested areas.

Figure 6. Maps of runoff connectivity in the five oldest scenarios (A, B, C, D and E) and of the differences in runoff connectivity between the most recent scenario (2012) and the oldest scenario (1945) (F). [Colour figure can be viewed at wileyonlinelibrary.com]

spatial approach to study overland flow patterns. Foerster *et al.* (2014) also found different connectivity maps for contrasting seasons in a 70-km² catchment in the Spanish Pyrenees. They obtained the highest connectivity values in August comparable with what we described in the IC-D8 map for HyP1. Few evaluations of the hydrological effects of afforestation in humid Mediterranean areas have been carried out at catchment scale, stressing the lack of available information. Nadal-Romero *et al.* (2016) showed that seasonal differences in hydrological response and connectivity exist in the same afforested sub-catchment we were studying, and they suggested different runoff generation processes: an alternation between a wet period, when the catchment was hydrologically responsive, and a dry summer period when the catchment was not or rarely responsive and two transitional periods. This pattern is partially coherent with the analyses presented here. The HyP1 from January to March coincides with the wet period and the most intense hydrological response, including August resulting from high rainfall intensities. The HyP2 (from April to July and from September to October) could correspond to the two

transitional periods. According to Nadal-Romero *et al.* (2016), the HyP3 would not correspond with summer. This time gap could be due to the frequent occurrence of intense rainstorms during the summer period, demonstrating a lower efficiency to reduce flooding risk after afforestation programmes. In a neighbouring catchment, the Arnás river catchment (abandoned fields with natural revegetation), three periods were identified by Lana-Renault *et al.* (2007): (i) the wet period, from the beginning of December to the middle of May (linked with HyP1); (ii) the drying-up period, from May to June to the end of the summer season (similar to our HyP2); and (iii) the wetting-up period, which started with the first autumn rainfall events and lasted until the end of November (or later) and which could be associated to HyP3.

The high values of connectivity in the two oldest scenarios and the marked decrease in connectivity after land abandonment and afforestation agreed with the results of connectivity, runoff and sediment yield dynamics reported by Sanjuán *et al.* (2014) and Buendía *et al.* (2016) in the Spanish Pyrenees and Pre-Pyrenees, which were linked to

Table III. Values of runoff connectivity in the five past scenarios and in the main landscape units

Scenario (year)	Length FPL (m) ^a	Local change IC	IC value				
			Total area	Hillslopes ^b	LLE wall ^c	LLE T&R ^d	Stream and gullies
1945	Mean	0.0285	-2.3602	-2.370	-2.614	-1.517	-2.018
	SD		0.6193	0.594	0.829	0.904	0.674
	Maximum						
	Change compared with 2012 scenario (%)	11.2	42.2	43.7	22.1	26.1	33.9
1956	Mean	0.0280	-2.3495	-2.364	-2.592	-1.441	-1.949
	SD		0.6253	0.599	0.838	0.936	0.671
	Maximum						
	Change compared with 2012 scenario (%)	9.5	42.5	43.8	22.7	29.8	36.1
1973	Change compared with previous scenario (%)	-1.5	0.5	0.3	0.8	5.0	3.4
	Mean	0.0262	-2.9835	-3.021	-2.857	-1.861	-2.447
	SD		0.6447	0.601	0.922	1.049	0.804
	Maximum						
1980	Change compared with 2012 scenario (%)	2.2	27.0	28.2	14.8	9.4	19.8
	Change compared with previous scenario (%)	-6.7	-27.0	-27.8	-10.2	-29.1	-25.6
	Mean	0.0257	-3.2980	-3.367	-3.054	-1.979	-2.556
	SD		0.7319	0.657	0.980	1.062	0.857
2006	Maximum						
	Change compared with 2012 scenario (%)	0.3	19.3	20.0	9.0	3.6	16.3
	Change compared with previous scenario (%)	-1.9	-10.5	-11.4	-6.9	-6.3	-4.4
	Mean	0.0264	-4.1431	-4.257	-3.553	-2.267	-3.131
	SD		0.9816	0.859	1.260	1.363	1.105
	Maximum						
	Change compared with 2012 scenario (%)	3.2	-1.4	-1.2	-5.9	-10.4	-2.6
	Change compared with previous scenario (%)	2.9	-25.6	-26.4	-16.3	-14.5	-22.5

Length of the flow path lines and values of the local changes (eight surrounding pixels) in connectivity.

^aLength of the flow path lines (FPL) excluding the main stream.^bExcluding all LLEs, the stream and the gullies.^cOnly stonewalls and terraces.^dOnly trails and forest road.

Figure 7. Relative changes in the values of overland flow connectivity (IC), total length of the landscape linear elements (LLE) and the C-RUSLE factor during the five past scenarios in comparison with the current scenario (year 2012) (A). Specific contribution played by the alone changes in the LLE and the C-RUSLE factor on the IC value for each past scenario in comparison with the current scenario (B). [Colour figure can be viewed at wileyonlinelibrary.com]

land cover changes and afforestation, respectively. Hawtree *et al.* (2015) also found in a large and wet catchment in north-central Portugal that pine afforestation promoted slower flow pathways, especially during the wet season, by increasing the amount of water entering the soil matrix via infiltration due to the ground preparation and planting operations used. However, these authors also described a faster catchment response with young eucalypt and older pine plantations, possibly due to soil water repellency. Baartman *et al.* (2013) found with both virtual and real catchments that connectivity decreased with increasing landscape morphological complexity. We also observed this association in the Araguás sub-catchment in the following: (i) between the complex patterns of patched vegetation in 2006 and 2012 and the low values of connectivity and (ii) between the simple land uses in 1945 and 1956 and the highest values of connectivity.

Changes in the total length and position of the LLE and those of the vegetation (C-RUSLE factor) played a different role in the evolution of the simulated values of runoff connectivity in the six scenarios (Table III). These results agreed with those obtained by Marchamalo *et al.* (2015) in a semi-arid environment where man-made lines (terraces and tracks) and the presence of vegetation influenced differently the local and catchment values of sediment connectivity. In 2012, when vegetation hardly changed, as was also the case in 2006 (C-RUSLE 0.4% higher), the small increment of the total length of the LLE (1.6%) explained the slight decrease in connectivity (−1.4%). However, when both factors changed, such as in 1980, the role played by vegetation changes (C-RUSLE 33.8% higher) was more important than the increment of the total length of the LLE (3.9%), and thus, the runoff connectivity clearly increased (19.3%) (Figure 7A). In order to estimate the specific contribution of the changes in the LLE and the vegetation cover to the total change of connectivity in each scenario, we ran the IC model in the five past scenarios considering only the changes in the LLE (fixed map of C-RUSLE). Then, we ran the IC model considering that changes only happen in the vegetation cover (fixed map of LLE). When only LLE changed, the values of runoff connectivity varied between

−1.7% and 5.7%, whereas the range of changes triggered by vegetation changes was between 0.05% and 41.3% (Figure 7B).

In spite of the positive effects of afforestation reducing runoff connectivity, the mean length of the FPL increased because of the loss of stonewalls. This fact could explain the similar and very high values of connectivity as reported by Lana-Renault *et al.* (2014) in other comparable areas in the Pyrenees during high-intensity summer rainstorms, when entire catchments were activated. Monitoring of the changes in LLE, vegetation growth and stream flow is included in current and future research planning. We also plan to run the spatially distributed hydrological DR2 model (López-Vicente *et al.*, 2014) to make a stronger analysis of the hydrological response of the soils. In order to reduce the negative impacts of high runoff connectivity of gullies and degraded (bare soil and open scrubland) areas on the Araguás sub-catchment, we propose to rebuild some of the destructed stonewalls. This practice should focus on those areas where the length of the FPL reaches higher values. Twenty years ago, Gallart *et al.* (1994) highlighted in a small catchment in the Eastern Pyrenees the environmental hazard of the spontaneous reorganization of the unmaintained artificial drainage network of abandoned terraces. Zimmermann *et al.* (2014) showed that the length of the FPL and its location are crucial to understand the magnitude of runoff connectivity. The recovery of stonewalls can be also considered as a structural measure to prevent off-site effects of soil erosion and corresponds well with the suggestions recently proposed by Mekonnen *et al.* (2015) on soil conservation through sediment trapping. Knowledge of the spatial pattern of connectivity and its dynamics over time is essential for land and water resource management and for understanding the potential environmental effects of land use changes. Research on hydrological connectivity can also help to increase the understanding of the effects of afforestation, land abandonment, timberland, skid trails and other forest practices in mountain landscapes and thus can provide crucial knowledge to land managers, policy makers and stakeholders.

CONCLUSIONS

In the current scenario, the combined analysis of the spatially distributed values of runoff connectivity and the temporal changes in the hydrological response of the sub-catchment improved our ability to understand the complexity of water dynamics. The multiple-flow algorithm with threshold value for linear flow (MD8) generated the most representative map in the current scenario, and thus, it was selected to run the index of connectivity for the past scenarios. The observed changes in connectivity mainly depended on the changes in the vegetation factor (C-RUSLE) and in a minor way on the total length, spatial location and type of the LLE. However, when small vegetation differences occurred, the role of the changes in the LLE became more important than the changes in the vegetation. Afforestation promoted lower and more stable values of runoff connectivity, reducing the local variability of the values of runoff connectivity. However, the collapse of many sections of stonewalls and terraces favoured the increment of the mean length of the FPL. We propose to rebuild the ruined stonewalls in those areas where the length of the FPL reached a high value to avoid significant increases of runoff connectivity especially during intense summer storm events. These results and the conservation proposal can be of interest for hydrological and forestry projects for both the public and private sectors.

ACKNOWLEDGEMENT(S)

This research was supported by a Marie Curie Intra-European Fellowship in the project 'MED-AFFOREST' (PIEF-GA-2013-624974). Dr López-Vicente acknowledges the financial support of his postdoctoral contract to the project 'EroCostModel' (CGL2014-54877-JIN; Programme Jóvenes Investigadores), and Dr Nadal-Romero was a recipient of a *Ramón y Cajal* research contract. Both contracts are of the Spanish Ministry of Economy and Competitiveness.

REFERENCES

- Antoine M, Javaux M, Bielders C. 2009. What indicators can capture runoff-relevant connectivity properties of the micro-topography at the plot scale? *Advances in Water Resources* **32**: 1297–1310. DOI:10.1016/j.advwatres.2009.05.006.
- Amáez J, Lana-Renault N, Lasanta T, Ruiz-Flaño P, Castroviejo J. 2015. Effects of farming terraces on hydrological and geomorphological processes. A review. *Catena* **128**: 122–134. DOI:10.1016/j.catena.2015.01.021.
- Baartman JEM, Masselink R, Keesstra SD, Temme AJAM. 2013. Linking landscape morphological complexity and sediment connectivity. *Earth Surface Processes and Landforms* **38**: 1457–1471. DOI:10.1002/esp.3434.
- Borselli L, Cassi P, Torri D. 2008. Prolegomena to sediment and flow connectivity in the landscape: a GIS and field numerical assessment. *Catena* **75**: 268–277. DOI:10.1016/j.catena.2008.07.006.
- Bracken LJ, Croke J. 2007. The concept of hydrological connectivity and its contribution to understanding runoff-dominated geomorphic systems. *Hydrological Processes* **21**: 1749–1763. DOI:10.1002/hyp.6313.
- Buendia C, Bussi G, Tuset J, Vericat D, Sabater S, Palau A, Batalla RJ. 2016. Effects of afforestation on runoff and sediment load in an upland Mediterranean catchment. *Science of the Total Environment* **540**: 144–157. DOI:10.1016/j.scitotenv.2015.07.005.
- Cammeraat ELH. 2004. Scale dependent thresholds in hydrological and erosion response of a semi-arid catchment in southeast Spain. *Agriculture, Ecosystems and Environment* **104**: 317–332. DOI:10.1016/j.agee.2004.01.032.
- Capolongo D, Pennetta L, Piccarreta M, Fallacara G, Boenzi F. 2008. Spatial and temporal variations in soil erosion and deposition due to land-levelling in a semi-arid area of Basilicata (Southern Italy). *Earth Surface Processes and Landforms* **33**: 364–379. DOI:10.1002/esp.1560.
- Cavalli M, Trevisani S, Comiti F, Marchi L. 2013. Geomorphometric assessment of spatial sediment connectivity in small Alpine catchments. *Geomorphology* **188**: 31–41. DOI:10.1016/j.geomorph.2012.05.007.
- Chartin C, Evrard O, Onda Y, Patin J, Lefèvre I, Ottlé C, Ayrault S, Lepage H, Bonte P. 2013. Tracking the early dispersion of contaminated sediment along rivers draining the Fukushima radioactive pollution plume. *Anthropocene* **1**: 23–34. DOI:10.1016/j.ancene.2013.07.001.
- Croke J, Mockler S. 2001. Gully initiation and road-to-stream linkage in a forested catchment, southeastern Australia. *Earth Surface Processes and Landforms* **26**: 205–217. DOI:10.1002/1096-9837(200102)26:2<205::AID-ESP168>3.0.CO;2-G.
- De Tar WR, Ross JJ, Cunningham RL. 1980. Estimating the C factor in the universal soil loss equation for landscape slopes. *Journal of Soil and Water Conservation* **35**: 40–41.
- D'Haen K, Duser B, Verstraeten G, Degryse P, De Brue H. 2013. A sediment fingerprinting approach to understand the geomorphic coupling in an eastern Mediterranean mountainous river catchment. *Geomorphology* **197**: 64–75. DOI:10.1016/j.geomorph.2013.04.038.
- Foerster S, Wilczok C, Brosinsky A, Segl K. 2014. Assessment of sediment connectivity from vegetation cover and topography using remotely sensed data in a dryland catchment in the Spanish Pyrenees. *Journal of Soils and Sediments* **14**: 1982–2000. DOI:10.1007/s11368-014-0992-3.
- Gallart F, Llorens P, Latron J. 1994. Studying the role of old agricultural terraces on runoff generation in a small Mediterranean mountainous basin. *Journal of Hydrology* **159**: 291–303. DOI:10.1016/0022-1694(94)90262-3.
- García-Ruiz JM, Lana-Renault N. 2011. Hydrological and erosive consequences of farmland abandonment in Europe, with special reference to the Mediterranean region – a review. *Agriculture, Ecosystems and Environment* **140**: 317–338. DOI:10.1016/j.agee.2011.01.003.
- Gay A, Cerdan O, Mardhel V, Desmet M. 2015. Application of an index of sediment connectivity in a lowland area. *Journal of Soils and Sediments*. DOI:10.1007/s11368-015-1235-y.
- Hawtree D, Nunes JP, Keizer JJ, Jacinto R, Santos J, Rial-Rivas ME, Boulet A-K, Tavares-Wahren F, Feger K-H. 2015. Time series analysis of the long-term hydrologic impacts of afforestation in the Águeda watershed of north-central Portugal. *Hydrology and Earth System Sciences* **19**: 3033–3045. DOI:10.5194/hess-19-3033-2015.
- Heckmann T, Schwanghart W. 2013. Geomorphic coupling and sediment connectivity in an alpine catchment – exploring sediment cascades using graph theory. *Geomorphology* **182**: 89–103. DOI:10.1016/j.geomorph.2012.10.033.
- Lana-Renault N, Latron J, Regüés D. 2007. Streamflow response and water-table dynamics in a sub-Mediterranean research catchment (Central Pyrenees). *Journal of Hydrology* **347**: 497–507. DOI:10.1016/j.jhydrol.2007.09.037.
- Lana-Renault N, Nadal-Romero E, Serrano-Muela MP, Alvera B, Sánchez-Navarrete P, Sanjuan Y, García-Ruiz JM. 2014. Comparative analysis of the response of various land covers to an exceptional rainfall event in the central Spanish Pyrenees, October 2012. *Earth Surface Processes and Landforms* **39**: 581–592. DOI:10.1002/esp.3465.
- Lane SN, Reaney SM, Heathwaite AL. 2009. Representation of landscape hydrological connectivity using a topographically driven surface flow index. *Water Resources Research* **45**: W08423. DOI:10.1029/2008WR007336.
- Lesschen JP, Schoorl JM, Cammeraat ELH. 2009. Modelling runoff and erosion for a semi-arid catchment using a multi-scale approach based on hydrological connectivity. *Geomorphology* **109**: 174–183. DOI:10.1016/j.geomorph.2009.02.030.
- Linares R, Rosell J, Palli L, Roque C. 2002. Afforestation by slope terracing accelerates erosion. A case study in the Barranco de Barcedana (Conca de Tremp, NE Spain). *Environmental Geology* **42**: 11–18. DOI:10.1007/s00254-002-0527-x.

- López-Vicente M, Lana-Renault N, García-Ruiz JM, Navas A. 2011. Assessing the potential effect of different land cover management practices on sediment yield from an abandoned farmland catchment in the Spanish Pyrenees. *Journal of Soils and Sediments* **11**: 1440–1455. DOI:10.1007/s11368-011-0428-2.
- López-Vicente M, Navas A. 2009. Predicting soil erosion with RUSLE in Mediterranean agricultural systems at catchment scale. *Soil Science* **174**: 272–282. DOI:10.1097/SS.0b013e3181a4bf50.
- López-Vicente M, Navas A. 2010. Routing runoff and soil particles in a distributed model with GIS: implications for soil protection in mountain agricultural landscapes. *Land Degradation & Development* **21**: 100–109. DOI:10.1002/ldr.901.
- López-Vicente M, Navas A, Gaspar L, Machín J. 2013a. Advanced modeling of runoff and soil redistribution for agricultural systems: the SERT model. *Agricultural Water Management* **125**: 1–12. DOI:10.1016/j.agwat.2013.04.002.
- López-Vicente M, Pérez-Bielsa C, López-Montero T, Lambán LJ, Navas A. 2014. Runoff simulation with eight different flow accumulation algorithms: recommendations using a spatially distributed and open-source model. *Environmental Modelling & Software* **62**: 11–21. DOI:10.1016/j.envsoft.2014.08.025.
- López-Vicente M, Poesen J, Navas A, Gaspar L. 2013b. Predicting runoff and sediment connectivity and soil erosion by water for different land use scenarios in the Spanish Pre-Pyrenees. *Catena* **102**: 62–73. DOI:10.1016/j.catena.2011.01.001.
- Marchamalo M, Hooke JM, Sandercock PJ. 2015. Flow and sediment connectivity in semi-arid landscapes in SE Spain: patterns and controls. *Land Degradation & Development*. DOI:10.1002/ldr.2352.
- Mekonnen M, Keesstra SD, Stroosnijder L, Baartman JEM, Maroulis J. 2015. Soil conservation through sediment trapping: a review. *Land Degradation & Development* **26**: 544–556. DOI:10.1002/ldr.2308.
- Miller JD, Nyhan JW, Yool SR. 2003. Modeling potential erosion due to the Cerro Grande fire with a GIS-based implementation of the revised universal soil loss equation. *International Journal of Wildland Fire* **12**: 85–100. DOI:10.1071/WF02017.
- Nadal-Romero E, Cammeraat E, Serrano P, Lana-Renault N, Regüés D. 2016. Hydrological response of an afforested catchment in a Mediterranean humid mountain area: a comparative study with a natural forest. *Hydrological Processes*. DOI:10.1002/hyp.10820.
- Nadal-Romero E, Regüés D, Latron J. 2008. Relationships among rainfall, runoff, and suspended sediment in a small catchment with badlands. *Catena* **74**: 127–136. DOI:10.1016/j.catena.2008.03.014.
- Nadal-Romero E, Regüés D, Serrano-Muela P. 2010. Respuesta hidrológica en una pequeña cuenca experimental pirenaica con dos ambientes extremos: cárcavas y bosque de repoblación. *Pirineos* **165**: 135–155. DOI:10.3989/Pirineos.2010.165007.
- Ortigosa L, García-Ruiz JM. 1995. Geomorphological consequences of afforestation at a basin scale, an example from the Central Pyrenees. *Physics and Chemistry of the Earth* **20**: 345–349. DOI:10.1016/0079-1946(95)00047-X.
- Parsons AJ, Bracken L, Poepl RE, Wainwright J, Keesstra SD. 2015. Introduction to special issue on connectivity in water and sediment dynamics. *Earth Surface Processes and Landforms* **40**: 1275–1277. DOI:10.1002/esp.3714.
- Reaney SM, Bracken LJ, Kirkby MJ. 2014. The importance of surface controls on overland flow connectivity in semi-arid environments: results from a numerical experimental approach. *Hydrological Processes* **28**: 2116–2128. DOI:10.1002/hyp.9769.
- Renard KG, Foster GR, Weesies GA, McCool DK, Yoder DC. 1997. Predicting soil erosion by water: a guide to conservation planning with the Revised Universal Soil Loss Equation (RUSLE). Handbook #703. US Department of Agriculture: Washington, DC.
- Rossel F, Gironás J, Mejía A, Rinaldo A, Rodríguez F. 2014. Spatial characterization of catchment dispersion mechanisms in an urban context. *Advances in Water Resources* **74**: 290–301. DOI:10.1016/j.advwatres.2014.09.005.
- Sanjuán Y, Gómez-Villar A, Nadal-Romero E, Álvarez-Martínez J, Arnáez J, Serrano-Muela MP, Rubiales JM, González-Sampériz P, García-Ruiz JM. 2014. Linking land cover changes in the sub-alpine and montane belts to changes in a torrential river. *Land Degradation & Development*. DOI:10.1002/ldr.2294.
- Vigiak O, Borselli L, Newham LTH, McInnes J, Roberts AM. 2012. Comparison of conceptual landscape metrics to define hillslope-scale sediment delivery ratio. *Geomorphology* **138**: 74–88. DOI:10.1016/j.geomorph.2011.08.026.
- Zimmermann B, Zimmermann A, Turner BL, Francke T, Elsenbeer H. 2014. Connectivity of overland flow by drainage network expansion in a rain forest catchment. *Water Resources Research* **50**: 1457–1473. DOI:10.1002/2012WR012660.