



UvA-DARE (Digital Academic Repository)

Path Sampling Simulations of the Mechanisms and Rates of Transitions between Watson-Crick and Hoogsteen Base Pairing in DNA

Vreede, J.; Bolhuis, P.G.; Swenson, D.W.H.

DOI

[10.1016/j.bpj.2016.11.1181](https://doi.org/10.1016/j.bpj.2016.11.1181)

Publication date

2017

Document Version

Final published version

Published in

Biophysical Journal

License

Other

[Link to publication](#)

Citation for published version (APA):

Vreede, J., Bolhuis, P. G., & Swenson, D. W. H. (2017). Path Sampling Simulations of the Mechanisms and Rates of Transitions between Watson-Crick and Hoogsteen Base Pairing in DNA. *Biophysical Journal*, 112(3, Supplement 1), 214a. <https://doi.org/10.1016/j.bpj.2016.11.1181>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

1058-Pos Board B126**Path Sampling Simulations of the Mechanisms and Rates of Transitions between Watson-Crick and Hoogsteen Base Pairing in DNA****Jocelyne Vreede**, Peter G. Bolhuis, David W.H. Swenson.

van't Hoff Institute for Molecular Sciences, University of Amsterdam, Amsterdam, Netherlands.

DNA duplexes predominantly contain Watson-Crick (WC) base pairs. However, at any time, a non-negligible fraction of base pairs exhibit a different interaction pattern, called Hoogsteen (HG) where the purine base is rotated approximately 180° relative to its orientation in the WC pairing. The conversion from WC to HG alters the conformation of DNA, and may play a role in several processes, including recognition and replication. The transient nature of these transitions hamper thorough experimental investigation. Molecular dynamics simulations complement experimental work by providing insights at very high spatial and temporal resolution. Path sampling methods focus the molecular dynamics effort on the dynamics during a transition, thus avoiding the long waiting times in stable states. We apply two different path sampling techniques to study the transitions between the WC and HG base pairing motifs. By using transition path sampling we observed that the WC to HG conversion can proceed along several mechanistic routes with varying degrees of exposure of the purine. We computed the rate for each process by employing transition interface sampling.