UvA-DARE (Digital Academic Repository)

10 Micron detection of the hard x-ray transient GRO J0422+23: Free-free emission from an x-ray-driven accretion disk wind?

van Paradijs, J.A.; Telesco, C.M.; Kouveliotou, C.; Fishman, G.J.

DOI

[10.1086/187403](https://doi.org/10.1086/187403)

Publication date

1994

Published in

Astrophysical Journal

[Link to publication](#)

Citation for published version (APA):

van Paradijs, J. A., Telesco, C. M., Kouveliotou, C., & Fishman, G. J. (1994). 10 Micron detection of the hard x-ray transient GRO J0422+23: Free-free emission from an x-ray-driven accretion disk wind? *Astrophysical Journal*, 429, L19-L23. <https://doi.org/10.1086/187403>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

10 MICRON DETECTION OF THE HARD X-RAY TRANSIENT GRO J0422+32: FREE-FREE EMISSION FROM AN X-RAY-DRIVEN ACCRETION DISK WIND?

J. VAN PARADIJS

Astronomical Institute “Anton Pannekoek,” University of Amsterdam and Center for High Energy Astrophysics, The Netherlands; and
 Department of Physics, University of Alabama Huntsville, AL 35899

C. M. TELESKO¹

Space Science Laboratory, NASA/Marshall Space Flight Center, ES-84, Huntsville, AL 35812

C. KOUVELIOTOU²

Universities Space Research Association

AND

G. J. FISHMAN

Space Science Laboratory, NASA Marshall Space Flight Center ES-81, Huntsville, AL 35812

Received 1993 August 19; accepted 1994 April 12

ABSTRACT

We report the detection of 10 μm emission from the transient low-mass X-ray binary (LMXB) and optical nova GRO J0422+32 near the maximum of its outburst. We discuss this result in terms of (1) a “standard” model according to which low-energy radiation of LMXB is caused by reprocessing of X-rays in an accretion disk; (2) emission from a cool secondary star; (3) emission from dust grains heated by the transient X-rays, and (4) free-free emission from an X-ray-driven wind from the accretion disk. Only the fourth alternative provides a viable explanation for the observed 10 μm emission, with a mass-loss rate in the disk wind that may be substantially higher than the rate of accretion onto the compact star. The presence of such a wind may have a profound effect on the evolution of the binary, and contribute to the solution of the “birthrate problem” of millisecond radio pulsars.

Subject headings: accretion, accretion disks — binaries: close — stars: individual (GRO J0422+32)

1. INTRODUCTION

On 1992 August 9, the bright transient X-ray source GRO J0422+32 was discovered with the Burst and Transient Source Experiment (BATSE) on board the *Compton Gamma Ray Observatory* (Paciesas et al. 1992). In 5 days the (25–300 keV) flux rose to $\sim 4 \times 10^{-8} \text{ ergs cm}^{-2} \text{ s}^{-1}$, or ~ 3 times that of the Crab Nebula. The flux remained at this level for ~ 10 days, and then decreased smoothly, with an e -folding decay time of 40 days (Paciesas et al. 1994). About 110 days after the peak of the outburst, the X-ray flux showed a temporary increase by a factor of about 2, after which the X-ray source continued to decrease at approximately the initial decay rate (Paciesas et al. 1994).

GRO J0422+32 has been identified with an optical nova (Castro-Tirado et al. 1992, 1993; Shrader et al. 1993) that reached a peak visual magnitude $V = 13.2$ from a quiescent value ≥ 17 (Chevalier & Ilovaisky 1993); this indicates that the source is a low-mass X-ray binary (LMXB; for a recent review see van Paradijs & McClintock 1994). The source has also been detected as a radio and ultraviolet transient (Han & Hjellming 1992; Shrader et al. 1993). The optical flux decayed smoothly with an e -folding time of ~ 90 days (Chevalier & Ilovaisky 1993). During the decay of the outburst, Chevalier & Ilovaisky (1993) found a periodic (5.1 hr) brightness variation. This period (or double its value) is likely to be the orbital

period (see also Chevalier & Ilovaisky 1994; cf. Bailyn 1992 and Remillard et al. 1992, on Nova Muscae 1991).

We report here details of observations of 10 μm emission from GRO J0422+32, the detection of which was announced previously (Telesko et al. 1992). In § 2 we describe the observations. In § 3 we discuss the results of our observations in terms of (1) reprocessing of X-rays in an accretion disk; (2) emission by dust near the X-ray source, and (3) free-free emission from an X-ray driven wind from the disk. In § 4 we summarize our conclusions. To our knowledge this is the first detection of an LMXB at such long infrared wavelengths.

2. OBSERVATIONS

GRO J0422+32 was observed at the NASA Infrared Telescope Facility (IRTF) on 1992 UT September 9.52–9.58 (near the peak of the outburst) and on 1993 April 3.21–3.25 (then the source had become undetectable with BATSE, i.e., its flux had decreased by at least a factor of 30 below its peak value). For the 1992 observations we used the Marshall Space Flight Center 20 pixel bolometric array (see, e.g., Telesko, Joy, & Sisk 1990) at 10.8 μm ($\Delta\lambda = 5.3 \mu\text{m}$). The FWHM pixel size was $4''.3 \times 4''.3$, and the separation between the source and the sky-reference positions was $15''$ in declination. The total integration time was 1.17 hr. The photometric standard for both dates was α Tau, with a 10.8 μm magnitude of -3.03 , corresponding to 570 Jy, based on values in the IRTF list of standard stars. We offset to GRO J0422+32 from the *HST* guide star SAO 057182 (1950: $04^{\text{h}}17^{\text{m}}58^{\text{s}}.4$, $+32^{\circ}42'43''$) using coordinates kindly provided by Dr. C. Shrader of the *GRO* Science Support Center. We determined that the coordinates of the optical counterpart of GRO J0422+32 are (1950) $04^{\text{h}}18^{\text{m}}29^{\text{s}}.67$,

¹ Visiting Astronomer at the Infrared Telescope Facility which is operated by the University of Hawaii under contract to the National Aeronautics and Space Administration.

² Postal address: NASA/Marshall Space Flight Center, ES-84 Huntsville, AL 35812.

+32°47'22".4, with an uncertainty of about $\pm 1''$ in each coordinate, which agrees within the uncertainties with those communicated to us by Dr. Shrader (see also McCroskey 1992). Only the pixel centered on the optical counterpart of GRO J0422+32 showed a statistically significant detection, corresponding to a flux density of 51 ± 9 mJy at $10.8 \mu\text{m}$; the error does not include the $\pm 7\%$ uncertainty in the absolute calibration. (The system sensitivity was relatively poor but stable on that date, due to significant degradation on the previous night of the hygroscopic pressure window.)

At the time of this observation the V magnitude of GRO J0422+32 was ~ 13.4 (Chevalier & Ilovaisky 1993); with $E_{B-V} = 0.4$ (Shrader et al. 1993) this corresponds to a reddening-corrected V band flux of 50 mJy.

The observations on 1993 April 3 were made at $10.2 \mu\text{m}$ ($\Delta\lambda = 5.0 \mu\text{m}$) using the IRTF single-channel bolometer. The total integration time was 0.67 hr. The FWHM size of the circular beam was $4''.7$, and the separation between the source and sky-reference positions was $10''$ in declination. Again we offset to GRO J0422+32 from the *HST* guide star, but this time the optical counterpart was barely visible using the IRTF optical acquisition camera; we estimate its V magnitude to be 15–16, based on a comparison with offset star B for *IRAS* 10214+4724, which was observed during the same run and is of comparable magnitude (see Telesco 1993). This estimate is in good agreement with the long-term light curve of GRO J0422+32 presented by Chevalier & Ilovaisky (1993). The $10.2 \mu\text{m}$ flux density of J0422+32 was 19 ± 12 mJy, which corresponds to a 3σ upper limit of 36 mJy.

3. DISCUSSION

During the second observation the $10 \mu\text{m}$ flux had decreased, but not by a sufficient amount (just over 2σ) to exclude that the flux had remained constant. We have investigated the possibility that the $10 \mu\text{m}$ source is not related to the X-ray source, but is rather a chance superposition. Using the *IRAS* point source catalog ($12 \mu\text{m}$) we have, for a circular field (3° radius) around the X-ray source determined the (cumulative) distribution of sources as a function of their flux, down to 250 mJy (we have ignored the 10% difference in wavelength between the *IRAS* and IRTF passbands). From a conservative (i.e., on the high side) extrapolation we find that there are 30 such sources per square degree with fluxes of at least 50 mJy. The corresponding probability of finding, by chance coincidence, a source in the pixel that contains the X-ray source is 4×10^{-5} . We conclude that the $10 \mu\text{m}$ emission we detected is related to the X-ray source.

We have considered three possible explanations for this $10 \mu\text{m}$ emission from GRO J0422+32. (1) The $10 \mu\text{m}$ flux is the long-wavelength extension of the optical/UV emission of the LMXB, which originates from reprocessing of X-rays in the accretion disk; we have also investigated the possibility of a contribution from the nonirradiated parts of a cool companion star. (2) The $10 \mu\text{m}$ emission comes from dust grains near the X-ray source that are heated by X-rays. (3) Free-free emission from an X-ray-driven wind from the accretion disk.

3.1. Thermal Emission from the Accretion Disk

To estimate the $10 \mu\text{m}$ (and optical) flux from the disk in GRO J0422+32 we made the following simplifying assumptions. (1) Everywhere in the disk the optical and $10 \mu\text{m}$ emission is Planckian. For $\lambda > 0.5 \mu\text{m}$, disk spectra for this simple model differ from those calculated for detailed models of

X-ray-irradiated disks by an amount that is small enough for our purpose (see, e.g., Vrtilek et al. 1990); (2) For a given X-ray luminosity L_X , the temperature T in the disk varies with radial distance r from the X-ray source as $r^{-3/7}$ (Vrtilek et al. 1990); (3) At a given value of r , T is proportional to $L_X^{1/4}$. We ignore the possible dependence of the $T-L_X$ relation on the shape of the X-ray spectrum.

A crude measure of the average disk temperature of GRO J0422+32 can be obtained from the relative rates of decay of the X-ray and optical light curves (Endal et al. 1976); it is $\sim 20,000$ K.

To relate $T(r)$ and L_X we use the properties of the Z-type (Hasinger & van der Klis 1989) LMXB Sco X-1, for which $V = 12.2$ at the transition from the normal to the flaring branch (Augusteijn et al. 1992). At this transition, L_X is close to the Eddington limit L_{Edd} (van der Klis & Lamb 1994), which equals $\sim 2 \times 10^{38}$ ergs s^{-1} for matter of cosmic abundances accreting onto a neutron star ($1.4 M_\odot$). The average X-ray flux of Sco X-1 (Warwick et al. 1981) then corresponds to a distance d to Sco X-1 of 2.3 kpc. For an assumed outer disk radius in Sco X-1 ($R_{\text{Sco X-1}}$) of 85% of the average radius of the Roche lobe of the X-ray source, the system parameters as given by Crampton et al. (1976) imply $R_{\text{Sco X-1}} \sim 1.2 \times 10^{11}$ cm. From a numerical integration [$F_V \sim (2\pi \cos i/d^2) \sum_i B_V(r_i) r_i \Delta r_i$, where $B_V(r_i)$ is the Planck function at radius r_i] we then find that, for $L_X = 2 \times 10^{38}$ ergs s^{-1} the temperature T_0 at the outer edge of the disk of Sco X-1 equals 22,500 K. (In the following, we will indicate with T_0 the disk temperature in any disk, at $r = R_{\text{Sco X-1}}$.)

We calculated, for a range in disk sizes, the ratio of the V band and $10 \mu\text{m}$ fluxes by numerical integration of the disk brightness (as above) in the two passbands from the center to the outer disk radius. For the 10.2 hr orbital period of GRO J0422+32 the disk radius R_{disk} (in units of $R_{\text{Sco X-1}}$) is between 0.6 (if the X-ray source is a neutron star) and 1.6 (if it is a $10 M_\odot$ black hole). We used values of T_0 between 10,000 K and 30,000 K (corresponding to L_X between 8×10^{36} and 6×10^{38} ergs s^{-1}). In all cases the expected $10 \mu\text{m}$ flux is less than 1% of the V -band flux. This result strongly disagrees with their observed approximate equality.

The basic reason why the observed $10 \mu\text{m}$ flux is not reproduced by the standard accretion disk reprocessing model is that the disk is much too small to contain the large cool outer regions required for efficient IR emission. We have estimated the disk size that, for a given value of T_0 , would reproduce the observed ratio $F_{10\mu}/F_V$ by continuing the numerical integration of the disk brightness following the $r^{-3/7}$ temperature dependence. We find that for T_0 between 10,000 K and 30,000 K the disk radius is between ~ 70 and $\sim 10^3$ times that of Sco X-1. The corresponding range in orbital period is 1.3 to 55 yr for a neutron star. For a $10 M_\odot$ black hole, these boundaries are about a factor of 3 lower. It should be noted that the outer disk regions may become optically thin for very large disk radii, which would make the required disk radii (and orbital periods) even larger.

We conclude that the standard accretion disk reprocessing does not explain the observed $10 \mu\text{m}$ flux of GRO J0422+32, unless its orbital period is very long.

3.2. Contribution from the Companion Star

Can the $10 \mu\text{m}$ flux come from the cool, nonirradiated parts of the companion star? Since the heated part of the companion is generally much less bright than the disk (van Paradijs &

McClintock 1994), we have approximated the observed emission as the sum of the disk emission (see § 3.1) and a blackbody with trial temperatures in the range 2000–4000 K.

For specific values of T_0 and R_{disk} , the radius R_{bb} of the blackbody emitter is given by $\Delta F_{10.8\mu} = \pi R_{\text{bb}}^2 d^{-2} B_{10.8\mu}(T_{\text{bb}})$, where $\Delta F_{10.8\mu}$ is the amount of 10.8 μm flux not accounted for by the disk (we know the value of this amount quantity because the observed ratio $F_{10.8\mu}/F_V$ has to be reproduced). As R_{disk} increases, $\Delta F_{10.8\mu}$ and R_{bb} decrease. We have taken as acceptable solutions those for which the ratio $R_{\text{bb}}/R_{\text{disk}}$ is below a value dictated by the fact that GRO J0422+32 is an LMXB (for a companion mass of $0.5 M_\odot$ and a $1.4 M_\odot$ neutron star this value is ~ 0.7 ; with a $10 M_\odot$ black hole it is ~ 0.3). Conversely, this requirement leads to solutions which, for chosen values of T_0 and T_{bb} , provide a lower limit to R_{disk} .

The next question then is, do these solutions satisfy the constraint that in quiescence the V magnitude can be as high as 19 (Zhao et al. 1994)? This puts a strong constraint on T_{bb} , since the quiescent flux (i.e., no contribution from the disk) should be at most 0.6% of that of the disk (this corresponds to the difference between the observed V magnitudes, 13.4 and >19 , respectively). It turns out that, virtually independent of the assumed value of T_0 , this requirement implies very low values of T_{bb} : below 2000 K if the compact star is a neutron star, and below 2200 K if it is a $10 M_\odot$ black hole.

Once T_0 has been selected the size is determined for systems that obey these constraints. For $T_0 = 10,000$ – $30,000$ K, the corresponding orbital periods (with neutron star accretors) are between 0.9 and 40 yr, i.e., $\sim 75\%$ of the periods for the corresponding disk-only models in the previous section. For $10 M_\odot$ black holes the periods are a factor ~ 3 shorter.

Therefore, the conclusion reached in § 3.1 remains unaffected, even if we include a possible contribution from a cool companion star.

3.3. 10 Micron Emission from X-Ray-Heated Dust Grains

We assume that X-ray heating of dust grains takes place inside a “heating sphere” whose radius R_h equals the distance traveled by X-rays in the time interval since the start of the outburst till our first 10 μm observation ($R_h \sim 9 \times 10^{16}$ cm). We used the theoretical IR spectral energy distributions emitted by an X-ray-irradiated “average dust grain surface” (Voit 1991), for incident X-ray fluxes f_X between 10^2 and 10^8 ergs $\text{cm}^{-2} \text{s}^{-1}$. This dust grain surface is an average over a distribution of grain sizes a given by the power law $f(a)da \propto a^{-3.5} da$ proposed by Mathis, Rumpl, & Nordsieck (1977). Voit presents his results in terms of νf_ν , where f_ν is the total flux density leaving the “average” dust grain per unit area; the specific intensity $I_\nu = f_\nu/\pi$.

For a given L_X (the source distance d is fixed by the observed X-ray flux F_X), we have taken $f_X = L_X/(4\pi R_h^2)$ and used the results of Voit (his Figs. 13, 15, and 16) to interpolate to the corresponding value of f_ν at 10 μm .

The BATSE X-ray flux refers to the 20–300 keV range. To estimate the total X-ray flux F_X we extrapolated the X-ray spectrum near the time of our first observation to lower energies using the best-fit exponential spectrum ($kT \sim 115$ keV; Paciesas et al. 1994). We find $F_X = 2.4 \times 10^{-8}$ ergs $\text{cm}^{-2} \text{s}^{-1}$. Since GRO J0422+32 may harbor a black hole (Paciesas et al. 1994), this estimate may be too low because it does not include a possible ultrasoft spectral component, as is often seen in the spectra of accreting black holes (Tanaka & Lewin 1994). However, *Granat* observations in 1992 September show no evidence for such a component (Sunyaev et al. 1993).

We can estimate an upper limit to the expected 10 μm flux according to $F_{10\mu\text{m}} = I_{10\mu\text{m}} \Omega$, where $\Omega = \pi R_h^2/d^2$ is the solid angle of the X-ray-heated dust sphere. In making this estimate we assume that dust grains fill the whole solid angle Ω . We find that for assumed values $L_X = 10^{37}$, 10^{38} , and 10^{39} ergs s^{-1} , the expected 10 μm fluxes are 110, 230, and 420 μJy , respectively, i.e., about two orders of magnitude below the observed value.

Dust grains are also heated by UV photons produced by reprocessing of X-rays in the accretion disk. UV photons are much more effective (by up to a factor 30; see Voit 1991) than X-rays in heating dust particles. However, since the luminosity of this reprocessed emission is at most a few percent of the X-ray luminosity (van Paradijs & McClintock 1994; Vrtilek et al. 1991) this contribution cannot alleviate the large discrepancy noted above.

Unless the density of dust around the X-ray source is extremely high, this inability of X-ray-heated dust grains to account for the observed 10 μm flux of GRO J0422+32 is exacerbated by the fact that, the filling factor of dust inside the solid angle Ω is much smaller than unity. For the above dust grain size distribution (with minimum and maximum grain sizes of 0.0005 and 0.25 μm) we find that the surface area (A) to volume (V) ratio of the dust grains equals $A/V = 2 \times 10^5 \text{ cm}^{-1}$. For a typical interstellar gas density of 1 atom cm^{-3} , a dust to gas mass ratio of 0.012, and a dust specific density of 3 g cm^{-3} , we find that the filling factor is $\sim 10^{-4}$. Thus, for a normal interstellar medium around GRO J0422+32 the gap between the predicted and observed 10 μm fluxes is six orders of magnitude.

We conclude that X-ray heating of dust cannot explain the observed 10 μm emission of GRO J0422+32 unless it is surrounded by a highly nonstandard interstellar medium.

3.4. Free-Free Emission from an X-Ray-driven Disk Wind

We have considered the possibility that the 10 μm emission originates from (optically thin) free-free emission in a wind flowing out of the disk (see, e.g., Wagner et al. 1991). Such a wind may be formed by X-ray heating of the disk for X-ray luminosities in excess of a few percent of the Eddington limit (Begelman, McKee, & Shields 1983). This possibility is suggested by the relative ease with which Oke & Greenstein (1977) were able to explain the near-IR ($<3 \mu\text{m}$) excess they observed from the black-hole transient A0620–00 by optically thin thermal bremsstrahlung (OTTB) from a hot ($\sim 10^6$ K) plasma.

In the case of GRO J0422+32 such a high-temperature OTTB interpretation faces the problem that the expected low-frequency spectrum is flat. That is, if the 10.8 μm flux (~ 50 mJy) is bremsstrahlung, one expects a flux density of 50 mJy in the V band as well, which leaves little room for reprocessing of X-rays into optical emission in the (optically thick) accretion disk. This would contradict a large body of observational evidence that the optical emission of LMXB is dominated by such reprocessing (see van Paradijs & McClintock 1994). This problem does not arise if the temperature of the plasma is sufficiently low that the OTTB dominates at 10.8 μm but not in the V band; this “solution” requires that the temperature be less than $\sim 30,000$ K.

To test roughly the viability of this model, we have estimated the expected free-free emission from a constant-velocity wind composed of ionized hydrogen at a temperature of 20,000 K. The electron density in this wind is given by

$$n_e = \dot{M}_w / 4\pi v_w r^2 m_p,$$

where $\dot{M}_w$ is the mass-loss rate in the wind, v_w its outflow

velocity, and m_p the proton mass. For the emissivity in this wind we use the expression (Rybicki & Lightman 1979, eq. [6.14b]; we have set the Gaunt factor equal to unity)

$$j_\nu = 6.8 \times 10^{-38} T^{-1/2} \exp(-h\nu/kT) n_e^2 \text{ ergs cm}^{-2} \text{ s}^{-1}.$$

The expected 10.8 μm luminosity in the wind is then given by

$$L_{10.8 \mu\text{m}} = 4\pi \int_{R_0}^{\infty} j_{10.8 \mu\text{m}}(r) r^2 dr = 1.5 \times 10^7 \dot{M}_w / (v_w^2 R_0).$$

The observed luminosity equals $6 \times 10^{19} d_{\text{kpc}}^2$. From this we infer $\dot{M}_w / (v_w^2 R_0) = 4 \times 10^{12} d_{\text{kpc}}^2$; this result is not sensitive to the assumed temperature of the free-free emitting wind.

Following Begelman et al. (1983) we have taken for R_0 the radial distance at which the radiation temperature T_{rad} corresponds to the escape velocity. This escape velocity is not expected to be very different from the wind velocity v_w , so that $v_w^2 R_0 \sim 2GM_X$. Therefore, we infer $\dot{M}_w \sim 8 \times 10^{12} GM_X d_{\text{kpc}}^2$, which leads to a mass-loss rate in the wind of $\sim 3 \times 10^{19} (M_X/M_\odot)^{1/2} d_{\text{kpc}} \text{ g s}^{-1}$. This mass-loss rate is very high, but not so high that this model for the 10.8 μm emission is ruled out: according to the calculations by Begelman et al. (1983) the mass-loss rate in the disk wind may substantially exceed the rate of inflow of matter that eventually makes it to the compact star.

At the time of our first observation the (5 GHz) radio flux was ~ 5 mJy, and this flux was nonthermal (Shrader et al. 1993; Hjellming 1994). This nondetection of the expected free-free radio emission can be understood as an optical-depth effect (i.e., the luminosity of a uniform source with optical depth $\tau \gg 1$ is smaller than that calculated on the assumption that it is optically thin by roughly a factor τ). The optical depth $\tau_\nu(\text{ff})$ in the wind above described is given by

$$\begin{aligned} \tau_\nu(\text{ff}) &= \int_{R_0}^{\infty} \kappa_\nu(\text{ff}) dr \\ &= 3.7 \times 10^8 v^{-3} T^{-1/2} (1 - e^{-h\nu/kT}) \int_{R_0}^{\infty} n_e^2 dr. \end{aligned}$$

Using the above expressions for $\dot{M}_w$ and v_w we find

$$\tau_\nu(\text{ff}) = 4.9 \times 10^{20} v^{-3} T^{-1/2} (1 - e^{-h\nu/kT}) (4\pi m_p)^{-2} R_0^{-2} d_{\text{kpc}}^2,$$

which for the 10 μm and radio (5 GHz) bands gives: $\tau_{10 \mu\text{m}}(\text{ff}) \sim 25 (R_0/3 \times 10^{10} \text{ cm})^{-2} d_{\text{kpc}}^2$, and $\tau_{5 \text{ GHz}}(\text{ff}) \sim 10^9 (R_0/3 \times 10^{10} \text{ cm})^{-2} d_{\text{kpc}}^2$. In view of the uncertainties in the system parameters and the crudeness of our wind model these results serve to show that at 10 μm the optical depth of the disk wind can be small, while in the radio it is very large. The implied turnover of the IR spectrum is a specific prediction of the free-free model, which can be tested by more detailed observations of the IR spectral energy distribution of LMXB (see, e.g., Wynn-Williams & Becklin 1979).

A similar argument shows that the observed rather small equivalent width ($\sim 5 \text{ \AA}$) of the H α line (Shrader et al. 1993) may be reconciled with the large value expected for an optically thin emitter ($\sim 400 \text{ \AA}$ using the emissivity coefficients given by Osterbrock 1974). In the center of the H α line the optical depth is given by

$$\tau_\alpha = \sigma_\alpha N_2 = \sigma_\alpha \int_{R_0}^{\infty} n_2(r) dr.$$

Here $n_2(r)$ indicates the number density of hydrogen atoms in the lower level of the H α transition. Using Saha's equation (which confirms that hydrogen is almost fully ionized), and assuming that in the line center thermal broadening dominates, we find for our assumed wind parameters $\tau_\alpha \sim 4 \times 10^3 (R_0/3 \times 10^{10} \text{ cm})^{-2}$, which shows that optical depth effects may play a major role.

4. CONCLUSIONS

We have detected 10 μm emission from the X-ray transient GRO J0422+32. We have considered several models to explain this emission: reprocessing of X-rays in an accretion disk, X-ray heating of nearby interstellar dust grains, and free-free emission from an X-ray induced wind from the accretion disk. The first model requires that the orbital period of GRO J0422+32 be very long, which seems inconsistent with the currently available data. The second model requires highly nonstandard interstellar-medium properties. Only the last model appears to account reasonably for the observed 10.8 μm emission.

Through its effect on the angular-momentum balance a high mass-loss rate in the form of a disk wind can have a major influence on the evolution of the system. Also, the usual estimates of the lifetimes of LMXB (roughly the ratio of the mass stored in the secondary envelope to the mass accretion rate) may be substantial overestimates. The presence of a strong disk wind may therefore contribute to the solution of the "birthrate problem" of millisecond radio pulsars (see Lorimer 1994 for a recent assessment of this problem). Infrared observations may be an effective way of studying accretion disk winds in LMXB, as is the case, e.g., for stellar winds of Be stars (see, e.g., Waters & Marlborough 1994).

We thank the referee, R. M. Wagner, for remarks which led us to realize the possible importance of accretion disk winds. We thank X. Tielens for useful correspondence on interstellar dust properties, M. Groenewegen for providing us with a list of IRAS sources near the position of the X-ray source, and J. Cassinelli for a useful discussion, and R. Piña and S. Fajando-Acousta for assistance with the observations.

REFERENCES

- Augusteijn, T., et al. 1992, *A&A*, 265, 177
 Bailyn, C. D. 1992, *ApJ*, 391, 298
 Begelman, M. C., McKee, C. F., & Shields, G. A. 1983, *ApJ*, 271, 70
 Castro-Tirado, A., et al. 1992, *IAU Circ.*, No. 5588
 ———, 1993, *A&A*, 276, L37
 Chevalier, C., & Ilovaisky, S. A. 1993, *La Lettre de l'OHP*, No. 11
 ———, 1994, *IAU Circ.*, No. 5928
 Crampton, D., Cowley, A. P., Hutchings, J. B., & Kaat, C. 1976, *ApJ*, 207, 907
 Endal, A. S., et al. 1976, *Astrophys. Lett.*, 17, 131
 Han, X., & Hjellming, R. M. 1992, *IAU Circ.*, 5593
 Hasinger, G., & van der Klis, M. 1989, *A&A*, 225, 79
 Hjellming, R. M. 1994, private communication
 Lorimer, D. 1994, in *The Lives of Neutron Stars*, ed. A. Alpar, U. Kiziloglu, & J. van Paradijs (Dordrecht: Kluwer) in press
 Mathis, J. S., Ruml, W., & Nordsieck, K. H. 1977, *ApJ*, 217, 425
 McCroskey, R. E. 1992, *IAU Circ.*, 5597
 Oke, J. B., & Greenstein, J. L. 1977, *ApJ*, 211, 872
 Osterbrock, D. E. 1974, *Astrophysics of Gaseous Nebulae* (San Francisco: Freeman)
 Paciasas, W. S., et al. 1992, *IAU Circ.*, No. 5580
 ———, et al. 1994, in preparation
 Remillard, R. E., McClintock, J. E., & Bailyn, C. D. 1992, *ApJ*, 399, L145
 Rybicki, G. B., & Lightman, A. P. 1979, *Radiative Processes in Astrophysics* (New York: Wiley)

- Shrader, C. R., Wagner, R. M., Starrfield, S. G., Hjellming, R. M., & Han, X. H. 1993, AIP Conf. Proc. 280, *Compton Gamma-Ray Observatory* ed. M. Friedlander, N. Gehrels, & D. J. Macomb (New York: AIP), 324
- Sunyaev, R. A., et al. 1993, A&A, 280, L1
- Tanaka, Y., & Lewin, W. H. G. 1994, in X-Ray Binaries, ed. W. H. G. Lewin, J. van Paradijs, & E. P. J. van den Heuvel (Cambridge Univ. Press), in press
- Telesco, C. M. 1993, MNRAS, 263, L37
- Telesco, C. M., Joy, M., & Sisk, C. 1990, ApJ, 358, L17
- Telesco, C. M., Pina, R., Fajardo, S., Fishman, G. J., Kouveliotou, C., & Van Paradijs, J. 1992, IAU Circ., No. 5613
- van der Klis, M., & Lamb, F. K. 1994, ARA&A, in preparation
- van Paradijs, J., & McClintock, J. E. 1994, in X-ray Binaries, ed. W. H. G. Lewin, J. van Paradijs, & E. P. J. van den Heuvel (Cambridge Univ. Press), in press
- Voit, G. M. 1991, ApJ, 379, 122
- Vrtilek, S. D., et al. 1990, A&A, 235, 162
- . 1991, ApJ, 376, 278
- Wagner, R. M., et al. 1991, ApJ, 378, 293
- Warwick, R. S., et al. 1981, MNRAS, 197, 865
- Waters, L. B. F. M., & Marlborough, J. M. 1994, in IAU Symp. 162, Pulsation, Rotation and Mass Loss in Early Type Stars, ed. L. A. Balona, H. F. Henrichs, & J.-M. LeContre (Dordrecht: Kluwer), in press
- Wynn-Williams, C. G., & Becklin, E. 1979, Nature, 282, 810
- Zhao, P., et al. 1994, IAU Circ., No. 5923