UvA-DARE (Digital Academic Repository)

Bisimulation is two-way simulation

Bergstra, J.A.; Stefanescu, G.

DOI

[10.1016/0020-0190\(94\)00165-0](https://doi.org/10.1016/0020-0190(94)00165-0)

Publication date

1994

Published in

Information Processing Letters

[Link to publication](#)

Citation for published version (APA):

Bergstra, J. A., & Stefanescu, G. (1994). Bisimulation is two-way simulation. *Information Processing Letters*, 52, 285-287. [https://doi.org/10.1016/0020-0190\(94\)00165-0](https://doi.org/10.1016/0020-0190(94)00165-0)

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Bisimulation is two-way simulation

J.A. Bergstra^a, Gh. Ștefănescu^{b,*}

^a *Programming Research Group, University of Amsterdam, Kruislaan 403, 1098 SJ Amsterdam, The Netherlands*

^b *Institute of Mathematics, Romanian Academy, P.O. Box 1-764, RO-70700 Bucharest, Romania*

Communicated by H. Ganzinger; received 24 February 1994; revised 9 September 1994

Abstract

We give here a simple proof of the fact that on transition systems bisimulation is the equivalence relation generated by simulation via functions. The proof entirely rests on simple rules of the calculus of relations.

Keywords: Theory of computation; Concurrency; Transition systems; Equivalence

Simulation is a standard notion of graph homomorphism that has been used in the study of flow diagram programs (see, e.g. [6,7,11]). Bisimulation is an equivalence on transition systems introduced by Park [9] in connection with Milner's work on concurrency [8]. In [3] we have shown that bisimulation is the equivalence relation generated by simulation via functions by using a translation between flowchart schemes and process graphs. We give here a simple proof of this fact using transition systems and simple rules of the calculus of relations, only.

To this end we show that (1) bisimulation via functions coincides with simulation via functions and (2) for two bisimilar processes one may find a common refinement which is bisimilar via functions with each of them.

In a categorical formulation this result also appears in [10]. A similar decomposition of bisimulation in terms of simpler relations was given by Castellani in [5]. She has shown that two nondeterministic processes (i.e., certain acyclic graphs that occur by unfolding transition systems) are bisimilar iff they may

be reduced to a common one. The reductions used by Castellani are given by certain abstract homomorphisms, which roughly correspond to the relation on the unfolded processes induced by our simulation via functions defined on the level of (possible cyclic) transition systems. (See also the comments in the final part of the present paper.) This result was further extended in [1] to the logical equivalence of transition systems.

The result of this paper is used in [4] as a key point in order to extend the axiomatisation of process graphs modulo isomorphism to the case of bisimilar ones.

Remarks on notation. We use “ $\circ$ ” to denote relational composition of two binary relations (i.e., $R \circ T = \{(x, z) \mid \exists y: x R y \text{ and } y T z\}$) and Id_A to denote the identity relation on a set A (i.e., $\text{Id}_A = \{(x, x) \mid x \in A\}$). Moreover, if R is a binary relation and T a ternary one, then $R \circ T = \{(x, z, w) \mid \exists y: (x, y) \in R \text{ and } (y, z, w) \in T\}$ and $T \circ R = \{(x, y, w) \mid \exists z: (x, y, z) \in T \text{ and } (z, w) \in R\}$.

Definitions. Let A be a set of atomic actions. A *transition system over A with multiple entries and multiple*

* Corresponding author.

exists $P : I \xrightarrow{S} O$ consists of three sets of vertices: I , S and O with $S \cap (I \cup O) = \emptyset$ (I specifies the start vertices, S the internal ones and O the end ones) and a transition relation $T \subseteq (I \cup S) \times A \times (O \cup S)$. Two transition systems $P : I \xrightarrow{S} O$ and $P' : I \xrightarrow{S'} O$ are similar via a function $\phi : S \rightarrow S'$ – written $P \rightarrow_\phi P'$ – if the corresponding transition relations T and T' fulfill

- $(\text{Id}_I \cup \phi) \circ T' = T \circ (\text{Id}_O \cup \phi)$.

P and P' are bisimilar via a relation $\rho \subseteq S \times S'$ – written $P \leftrightarrow_\rho P'$ – if the corresponding transition relations T and T' fulfill

- $(\text{Id}_I \cup \rho) \circ T' \subseteq T \circ (\text{Id}_O \cup \rho)$,
- $(\text{Id}_I \cup \rho^{-1}) \circ T \subseteq T' \circ (\text{Id}_O \cup \rho^{-1})$.

Note that simulation $\rightarrow$ is a transitive, but not a symmetric relation, whilst bisimulation $\leftrightarrow$ is an equivalence relation. We denote by $\leftarrow$ the converse of $\rightarrow$.

Lemma 1. For two transition systems $P : I \xrightarrow{S} O$ and $P' : I \xrightarrow{S'} O$ (with the corresponding transition relations T and T') and a function $\phi : S \rightarrow S'$ the following equivalence holds:

$$\begin{aligned} \text{(i)} \quad & (\text{Id}_I \cup \phi^{-1}) \circ T \subseteq T' \circ (\text{Id}_O \cup \phi^{-1}) \\ \iff \\ \text{(ii)} \quad & T \circ (\text{Id}_O \cup \phi) \subseteq (\text{Id}_I \cup \phi) \circ T'. \end{aligned}$$

Proof. Since ϕ is a function, $\phi \circ \phi^{-1} \supseteq \text{Id}_S$ (i.e., ϕ is a total relation) and $\phi^{-1} \circ \phi \subseteq \text{Id}_{S'}$ (i.e., ϕ is a univocal relation).

($\Rightarrow$) Compose (i) on the left with $\text{Id}_I \cup \phi$ and on the right with $\text{Id}_O \cup \phi$. Then

$$\begin{aligned} & (\text{Id}_I \cup \phi \circ \phi^{-1}) \circ T \circ (\text{Id}_O \cup \phi) \\ & \subseteq (\text{Id}_I \cup \phi) \circ T' \circ (\text{Id}_O \cup \phi^{-1} \circ \phi) \end{aligned}$$

and this implies (ii) by the above properties of ϕ .

($\Leftarrow$) Similar. $\square$

Corollary. Simulation via a function coincides with bisimulation via a function. $\square$

In order to pass from bisimilar transition systems to chains of similar ones we need a characterization of bisimulation in terms of bisimulation via functions. The result given here is based on a construction of

a common refinement of two bisimilar transition systems.

Lemma 2 (Interpolation property). Suppose two transition systems $P' : I \xrightarrow{S'} O$ and $P'' : I \xrightarrow{S''} O$ and a relation $\rho \subseteq S' \times S''$ are given such that $P' \leftrightarrow_\rho P''$. Then there exist a transition system $P : I \xrightarrow{S} O$ and two functions $\phi : S \rightarrow S'$ and $\psi : S \rightarrow S''$ such that $P \leftrightarrow_\phi P'$ and $P \leftrightarrow_\psi P''$.

Proof. Take a decomposition of ρ as $\phi^{-1} \circ \psi$ for two functions $\phi : S \rightarrow S'$ and $\psi : S \rightarrow S''$. Bisimulation $P' \leftrightarrow_\rho P''$ means

$$\begin{aligned} \text{(i)} \quad & (\text{Id}_I \cup \phi^{-1} \circ \psi) \circ T'' \\ & \subseteq T' \circ (\text{Id}_O \cup \phi^{-1} \circ \psi) \end{aligned}$$

and

$$\begin{aligned} \text{(ii)} \quad & (\text{Id}_I \cup \psi^{-1} \circ \phi) \circ T' \\ & \subseteq T'' \circ (\text{Id}_O \cup \psi^{-1} \circ \phi) \end{aligned}$$

Take the transition system $P : I \xrightarrow{S} O$ given by the transition relation

$$\begin{aligned} T = & [(\text{Id}_I \cup \phi) \circ T' \circ (\text{Id}_O \cup \phi^{-1})] \\ & \cap [(\text{Id}_I \cup \psi) \circ T'' \circ (\text{Id}_O \cup \psi^{-1})]. \end{aligned}$$

Then $P \leftrightarrow_\phi P'$. Indeed, one inclusion is straightforward:

$$\begin{aligned} & (\text{Id}_I \cup \phi^{-1}) \circ T \\ & \subseteq (\text{Id}_I \cup \phi^{-1} \circ \phi) \circ T' \circ (\text{Id}_O \cup \phi^{-1}) \\ & \subseteq T' \circ (\text{Id}_O \cup \phi^{-1}) \end{aligned}$$

where in the last step we have used the fact that ϕ is a univocal relation, hence $\phi^{-1} \circ \phi \subseteq \text{Id}_{S'}$.

For the second one we use (ii). By a left composition of (ii) with $\text{Id}_I \cup \psi$ we get

$$\begin{aligned} & (\text{Id}_I \cup \psi \circ \psi^{-1} \circ \phi) \circ T' \\ & \subseteq (\text{Id}_I \cup \psi) \circ T'' \circ (\text{Id}_O \cup \psi^{-1} \circ \phi) \end{aligned}$$

By using $\text{Id}_S \subseteq \psi \circ \psi^{-1}$ (i.e., ψ is a total relation) it follows that the left-hand side of this inclusion contains $(\text{Id}_I \cup \phi) \circ T'$. On the other hand, the right-hand side of this inclusion is equal to the second part of the

definition of T composed on right with $\text{Id}_O \cup \phi$, hence is included in $T \circ (\text{Id}_O \cup \phi)$. Consequently,

$$(\text{Id}_I \cup \phi) \circ T' \subseteq T \circ (\text{Id}_O \cup \phi)$$

In a similar way one may prove that $P \leftrightarrow_\psi P''$. $\square$

Theorem. *Bisimulation is the equivalence relation generated by simulation via functions. More precisely,*

$$\leftrightarrow = \leftarrow \circ \rightarrow.$$

Proof. From the lemmas above. $\square$

Simulation is an useful tool to speak about minimization. Simulation via surjective functions models identification of the states with the same behaviour and the *converse* of the simulation via injective functions models the deletion of the nonaccessible parts. We may give an alternative proof of the Theorem by replacing Lemma 2 with another one using minimization (as in [11] or as in Theorem 2.7.13 in [2] regarding normal forms), but the characterization is less direct, i.e., instead of the present characterization

$$\leftrightarrow = \leftarrow \circ \rightarrow,$$

we get

$$\leftrightarrow = \rightarrow_{\text{sur}} \circ \text{inj} \leftarrow \circ \rightarrow_{\text{inj}} \circ \text{sur} \leftarrow,$$

where $\rightarrow_{\text{sur}}$ and $\rightarrow_{\text{inj}}$ are the restrictions of simulation to simulation via surjective and injective functions, respectively. The latter characterization says that two transition systems are bisimilar iff by the identification of the states with the same behaviour and the deletion of the nonaccessible ones both systems may be reduced to the same minimal one.

References

- [1] A. Arnold and A. Dicky, An algebraic characterization of transition systems equivalences, *Inform. and Comput.* **82** (1989) 198–229.
- [2] J.C.M. Baeten and W.P. Weijland, *Process Algebra* (Cambridge University Press, Cambridge, 1990).
- [3] J.A. Bergstra and Gh. Ştefănescu, Translations between flowchart schemes and process graphs, in: *Proc. FCT'93*, Lecture Notes in Computer Science **710** (Springer, Berlin, 1993) 153–162.
- [4] J.A. Bergstra and Gh. Ştefănescu, Processes with multiple entries and exits modulo isomorphism and modulo bisimulation, Tech. Rept. PRG-9403, Programming Research Group, University of Amsterdam, 1994.
- [5] I. Castellani, Bisimulation and abstraction homomorphisms, *J. Comput. System Sci.* **34** (1987) 210–235.
- [6] C.C. Elgot, Some “geometrical” categories associated with flowchart schemes, in: *Proc. FCT'77*, Lecture Notes in Computer Science **56** (Springer, Berlin, 1977) 256–259.
- [7] J.A. Goguen, On homomorphism, correctness, termination, unfoldments and equivalence of flow diagram programs, *J. Comput. System Sci.* **8** (1974) 333–365.
- [8] R. Milner, *A Calculus of Communicating Systems*, Lecture Notes in Computer Science **92** (Springer, Berlin, 1980).
- [9] D. Park, Concurrency and automata on infinite sequences, in: *Proc. 5th GI Conf.*, Lecture Notes in Computer Science **104** (Springer, Berlin, 1981) 167–183.
- [10] J. Rutten and D. Turi, Initial algebra and final coalgebra semantics for concurrency, in: *A Decade of Concurrency*, Lecture Notes in Computer Science **803** (Springer, Berlin, 1994) 530–582.
- [11] Gh. Ştefănescu, On flowchart theories. Part I: The deterministic case, *J. Comput. System Sci.* **35** (1987) 163–191.