



UvA-DARE (Digital Academic Repository)

The golden years

Pippias, M.

[Link to publication](#)

Citation for published version (APA):

Pippias, M. (2018). The golden years: Kidney transplant outcomes in the 21st century

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Chapter 5

The European Renal Association- European Dialysis and Transplant Association Registry Annual Report 2012: a summary

Maria Pippias, Vianda S. Stel, José Maria Abad Diez, Nikolaos Afentakis, Jose Antonio Herrero-Calvo, Manuel Arias, Boris Bikbov, Encarnación Bouzas Caamaño, Jadranka Buturovic-Ponikvar, Svjetlana Čala, Fergus J. Caskey, Pablo Castro de la Nuez, Harijs Cernevskis, Frederic Collart, Ramón Alonso de la Torre, Maria de los Ángeles García Bazaga, Johan De Meester, Joan Manuel Díaz, Ljubica Djukanovic, Manuel Ferrer Alamar, Patrik Finne, Liliana Garneata, Eliezer Golan, Raquel González Fernández, Gonzalo Gutiérrez Avila, James Heaf, Andries Hoitsma, Nino Kantaria, Mykola Kolesnyk, Reinhard Kramar, Anneke Kramer, Mathilde Lasalle, Torbjørn Leivestad, Frantisek Lopot, Fernando Macario, Angela Magaz, Eduardo Martín-Escobar, Wendy Metcalfe, Marlies Noordzij, Runolfur Palsson, Ülle Pechter, Karl G. Prutz, Marina Ratkovic, Halima Resić, Boleslaw Rutkowski, Carmen Santiuste de Pablos, Viera Spustová, Gültekin Süleymanlar, Karlijn van Stralen, Nestor Thereska, Christoph Wanner and Kitty J. Jager

Clin Kidney J. 2015 Jun; 8(3): 248–261

Abstract

Background: This article summarises the 2012 ERA-EDTA Registry Annual Report (available at www.era-edta-reg.org) with a specific focus on older patients (defined as ≥ 65 years).

Methods: Data provided by 45 national or regional renal registries in 30 countries in Europe and bordering the Mediterranean Sea were used. Individual patient level data were received from 31 renal registries, whereas 14 renal registries contributed data in an aggregated form. The incidence, prevalence and survival probabilities of patients with ESRD requiring RRT and renal transplantation rates for 2012 are presented.

Results: In 2012, the overall unadjusted incidence rate of patients with ESRD receiving RRT was 109.6 per million population (pmp) (N=69,035), ranging from 219.9 pmp in Portugal to 24.2 pmp in Montenegro. The proportion of incident patients ≥ 75 years varied from 15% to 44% between countries. The overall unadjusted prevalence on 31st December 2012 was 716.7 pmp (N=451,270), ranging from 1670.2 pmp in Portugal to 146.7 pmp in the Ukraine. The proportion of prevalent patients ≥ 75 years varied from 11% to 32% between countries. The overall renal transplantation rate in 2012 was 28.3 pmp (N=15,673), with the highest rate seen in the Spanish region of Catalonia. The proportion of patients ≥ 65 years receiving a transplant ranged from 0% to 35%. Five-year adjusted survival for all RRT patients was 59.7% (95% confidence interval, CI: 59.3; 60.0) which fell to 39.3% (95%CI: 38.7; 39.9) in patients 65-74 years and 21.3% (95%CI: 20.8; 21.9) in patients ≥ 75 years.

Introduction

The ERA-EDTA Registry annually collects data on RRT for ESRD from national and regional renal registries in Europe and those bordering the Mediterranean sea. In this paper we provide a summary of the 2012 ERA-EDTA Registry Annual Report (available at www.era-edta-reg.org) on the incidence and prevalence of ESRD patients receiving RRT, on kidney transplant rates and on patient and graft survival. As the older patient (defined in this paper as 65 years and over and presented in two age groups i.e. 65-74 years and 75 years and over) now represents 42% of the RRT population within Europe, this article will pay particular attention to this patient group.

Methods

In 2014, the ERA-EDTA Registry received the 2012 data from the national and regional renal registries in 30 countries in Europe and those bordering the Mediterranean Sea (Figure 5.1a). Data sets with individual patient data were received from 31 national or regional renal registries in 17 countries, whereas 14 national renal registries contributed data in an aggregated form. The percentage coverage of the general population by the renal registries was above 98% for all participating countries with the exception of Romania (96%), Latvia (85%) and Georgia (63.5%). The total coverage of the European population by the ERA-EDTA Registry in 2012 was 71.3%. A list of the European countries with available data and those without is provided in Appendix 5.1.

The RRT incidence and prevalence data were presented as unadjusted rates and standardized for the age and sex distribution of the EU27 population in 2010 (104). Patient survival on RRT and dialysis were based on incident RRT patients. Patient and graft survival post-transplantation only included patients receiving their first renal transplant between 2003 and 2007 or 2006 and 2010. The survival analyses were based solely on data from renal registries providing individual patient records. A detailed description of the methods on patient and graft survival is given in Appendix 5.1. More detailed data than those presented in this current paper were published in the full 2012 ERA-EDTA Registry Annual Report (28) which is also available online at www.era-edta-reg.org.

The incidence of RRT for ESRD

In 2012 the total number of patients starting RRT for ESRD among all the registries reporting to the ERA-EDTA Registry was 69,035. These registries covered a total population of 629.7 million, resulting in an unadjusted incidence rate of 109.6 per million

population (pmp). Figure 5.1a and Figure 5.2 show the incidence of ESRD patients receiving RRT per pmp on day 1, among all national and regional renal registries in 30 countries reporting to the ERA-EDTA Registry in 2012. The highest unadjusted incidence rates were reported in Portugal (219.9 pmp), Greece (209.7 pmp) and Georgia (199.6 pmp), whereas the lowest unadjusted incidence rates were reported in Montenegro (24.2 pmp), Ukraine (27.6 pmp) and Russia (48.1 pmp). The incidence rate standardised for the EU27 population was highest in Israel (280.5 pmp), Georgia (227.5 pmp) and French-speaking Belgium (201.1 pmp). The mean age at the start of RRT across Europe was 62.0 years in the renal registries providing individual level patient data and 58.2 years in the renal registries providing aggregated data (Figure 5.2). Figure 5.1b and Figure 5.3 provide information on the incidence of RRT, per million age-related population (pmarp), on day 1 for older patients. For the countries providing individual level data the incidence of RRT for patients aged 65-74 years or 75 years and over varied considerably between the renal registries (Figure 5.3). The lowest incidence rate of patients aged 65-74 and 75 years and over was reported in Finland (199.8 pmarp and 220.8 pmarp respectively). A more than three times as high incidence rate was reported in French-speaking Belgium (619.4 pmarp) for patients aged 65-74 years and Dutch-speaking Belgium (893.6 pmarp) for patients aged over 75 years (Figure 5.3).

The lowest incidence rate of patients aged 65-74 and 75 years and over was reported in Finland (199.8 pmarp and 220.8 pmarp respectively). A more than three times as high incidence rate was reported in French-speaking Belgium (619.4 pmarp) for patients aged 65-74 years and Dutch-speaking Belgium (893.6 pmarp) for patients aged over 75 years (Figure 5.3). Within each country or region, there was a wide variation in the proportion of ESRD patients aged 75 years and over commencing RRT with the highest proportion reported in Dutch-speaking Belgium (44%) and Greece (43%) and the lowest proportion in Romania (15%) and Serbia (15%). Within the countries supplying individual level data, the four national or regional registries with the highest adjusted incidence of RRT on day 1 (Greece, French- and Dutch-speaking Belgium and France) also had the highest incidence of RRT on day 1 for patients 75 years and over (Figure 5.1a, 5.1b and 5.3).

Table 5.1 presents the unadjusted incidence of RRT pmp on day 91, by treatment modality (haemodialysis, peritoneal dialysis and transplantation) for the renal registries providing individual and aggregated data. The incidence rate of haemodialysis was highest in Portugal (184.7 pmp), Greece (171.0 pmp) and Israel (153.7 pmp), whereas the incidence rate of peritoneal dialysis was highest in the Spanish regions of Asturias (38.2 pmp) and Galicia (32.9 pmp) and Sweden (32.6 pmp). The incidence rate of renal transplants on day 91 was highest in Norway (16.5 pmp), the Netherlands (14.9 pmp) and Iceland (12.5 pmp).

Table 5.1. Incident counts and rates, per million population (pmp) of renal replacement therapy at day 91 (unadjusted) in 2012 by established therapy for countries/regions providing individual patient data and for countries/regions providing aggregated data.

Country / Region	All		HD	PD	Tx	Unknown / Missing
	N	pmp	pmp	pmp	pmp	pmp
Country / Region providing individual patient data						
Austria	1054	124.8	104.1	15.0	5.7	0
Belgium: Dutch-speaking*	1089	171.1	150.0	16.8	4.2	0
Belgium: French-speaking*	800	170.2	147.4	17.9	4.0	0.9
Bosnia and Herzegovina	407	116.0	112.3	3.4	0	0.3
Croatia †	595	139.4	123.4	13.7	2.0	0.2
Denmark	654	115.8	75.6	31.9	8.3	0
Finland	433	80.0	57.8	21.6	0.6	0
France	9305	142.2	117.0	14.9	6.2	4.0
Greece	2059	185.6	171.0	13.3	1.3	0
Iceland	19	59.2	28.1	18.7	12.5	0
Norway	499	99.4	63.4	19.5	16.5	0
Romania	2518	130.8	122.0	7.4	1.4	0
Serbia	837	116.3	102.2	12.4	1.7	0
Slovenia	240	116.7	110.8	5.3	0.5	0
Spain: Andalusia	947	112.9	89.3	17.9	5.7	0
Spain: Aragon	171	127.5	105.1	17.9	4.5	0
Spain: Asturias	131	121.9	81.0	38.2	2.8	0
Spain: Basque Country	251	115.1	84.4	24.3	6.4	0
Spain: Cantabria*	55	93.0	67.7	20.3	5.1	0
Spain: Castile and León*	287	113.5	87.4	22.9	3.2	0
Spain: Castile-La Mancha*	198	94.3	76.2	17.1	1.0	0
Spain: Catalonia†	854	112.8	82.3	19.0	11.5	0
Spain: Community of Madrid	659	102.6	82.1	14.5	6.1	0
Spain: Extremadura	127	114.6	96.6	17.1	0.9	0
Spain: Galicia	365	131.9	94.3	32.9	4.7	0
Spain: Region of Murcia†	159	107.6	93.1	12.4	2.0	0
Spain: Valencia	711	138.6	109.8	24.6	4.3	0
Sweden	969	101.8	60.8	32.6	8.4	0
the Netherlands	1892	112.9	79.6	18.5	14.9	0
UK: All countries*†	6372	100.0	71.0	20.0	9.0	0
UK: England*†	5371	100.4	70.6	20.7	9.1	0
UK: Northern Ireland*†	168	92.3	65.8	15.0	11.5	0
UK: Scotland	508	95.6	74.5	13.7	7.3	0
UK: Wales*†	332	108.0	75.4	23.2	9.4	0
Country / Region providing aggregated patient data						
Albania	177	63.3	57.9	4.6	0.7	0
Czech Republic§	1396	134.1	126.0	8.1		0
Estonia	105	79.4	61.2	16.6	1.5	0
Georgia	475	166.3	143.2	21.0	1.1	1.1
Israel	1356	171.4	153.7	12.3	5.4	0
Latvia	138	80.2	64.5	15.7	0.0	0
Montenegro	13	21.0	21.0	0.0	0.0	0
Portugal#	2190	207.3	184.7	20.8	1.8	0
Slovakia§	772	142.7	136.8	5.9		0

Footnotes: Abbreviations used: HD: haemodialysis; PD: peritoneal dialysis; TX: transplant; pmp: per million population. *: Patients younger than 20 years of age are not reported. †: The incident counts at day 91 are estimated. §: Data include dialysis patients only. #: Only pre-emptive transplantations (at day 1) are included. When cells are left empty, the data are unavailable.

Figure 5.1a. The unadjusted incidence of ESRD patients on renal replacement therapy, per million population (pmp) on day 1, among all national and regional renal registries in 30 countries reporting to the ERA-EDTA Registry in 2012.





Figure 5.1b. The unadjusted incidence of ESRD patients aged 75 years and older on renal replacement therapy, per million age-related population (pmarp) at day 1, among all national and regional renal registries in 17 countries providing individual level data to the ERA-EDTA Registry in 2012.



Footnotes for Figures 5.1a & 5.2: The incidence data for the Czech Republic and Slovakia only include dialysis patients. Mean and median age for Slovakia is based on dialysis patients and for Turkey only on haemodialysis patients. Dutch- and French-speaking Belgium, the Spanish regions of Cantabria, Castile & León & Castile-La Mancha and the UK do not report on patients <20 years.




Figure 5.2. Incidence of renal replacement therapy (RRT), per million population (pmp) at day 1 in 2012, unadjusted (black bars) and standardised to the age and gender distribution of the EU27 population in 2010 (grey bars) and the mean and median age (years) at the start of RRT.

Table 5.2 presents the percentage of patients by established treatment modality (haemodialysis, peritoneal dialysis and transplantation) on day 91 after the start of RRT, by age group. For patients 75 years and over, the vast majority were treated with haemodialysis, and almost no patients had a kidney transplant. Notably within this age group the Scandinavian countries (Norway, Sweden, Finland and Denmark) had the highest percentage (about a quarter) of patients treated with peritoneal dialysis.

For children (aged 0 to 19 years), we performed a separate analysis of the incidence rate by age group and initial treatment modality from a selected number of renal registries which are presented in Table 5.3. A more detailed overview of the paediatric RRT data in Europe can be found at www.espn-reg.org.

Table 5.2. Percentages of established renal replacement therapy modality at day 91 (unadjusted) by age group for the countries/regions providing individual patient data.

Country / Region	All			0-19			20-44			45-64			65-74			≥75								
	HD	PD	TX	Oth	HD	PD	TX	Oth	HD	PD	TX	Oth	HD	PD	TX	Oth	HD	PD	TX					
	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%					
Austria	75	12	5	8	39	17	44	0	55	14	18	13	67	16	6	11	82	13	0	5	88	5	0	7
BE: Dutch-speaking*	59	10	3	29					38	26	11	26	57	15	4	24	61	10	3	26	63	4	0	33
BE: French-speaking*	65	11	2	22					48	22	11	19	66	12	3	20	70	6	2	22	65	9	0	26
Bosnia and Herzegovina	94	3	0	3	75	25	0	0	88	7	0	5	93	2	0	5	96	2	0	3	98	2	0	0
Croatia †	88	10	2	1	13	50	38	0	81	10	6	3	85	14	1	1	92	7	0	1	95	5	1	0
Denmark	65	28	7	0	50	25	25	0	64	23	14	0	58	30	12	0	63	31	5	0	78	22	0	0
Finland	69	27	1	3	29	57	14	0	51	44	4	2	70	26	0	4	75	22	0	4	77	22	0	1
France	72	11	4	14	35	21	26	19	64	13	12	11	71	9	7	13	76	9	3	13	73	12	0	15
Greece	89	7	1	3	61	28	11	0	85	9	4	2	87	9	1	3	89	7	0	4	93	6	0	2
Iceland	47	32	21	0	0	0	0	0	0	0	0	0	44	22	33	0	67	0	33	0	43	57	0	0
Norway	64	20	17	0	20	20	60	0	62	17	22	0	56	20	24	0	68	16	16	0	73	24	3	0
Romania	93	6	1	0	68	32	0	0	89	6	5	0	94	6	1	0	94	6	0	0	96	4	0	1
Serbia	86	11	1	2	86	0	14	0	68	20	9	4	90	8	0	2	90	9	0	1	87	13	0	1
Slovenia	95	5	0	0	100	0	0	0	88	12	0	0	90	9	1	0	97	3	0	0	100	0	0	0
ES: Andalusia	79	16	5	0	42	25	33	0	67	19	14	0	75	18	7	0	82	16	2	0	90	10	0	0
ES: Aragon	81	14	4	2	0	0	100	0	58	21	21	0	77	23	0	0	81	15	0	4	93	5	0	2
ES: Asturias	66	31	2	0	0	100	0	0	38	63	0	0	52	44	4	0	77	19	3	0	80	20	0	0
ES: Basque Country	73	21	6	0	0	0	100	0	54	26	20	0	65	31	5	0	82	17	2	0	88	12	0	0
ES: Cantabria*	73	22	5	0					43	14	43	0	64	36	0	0	93	7	0	0	83	17	0	0
ES: Castile and León*	77	20	3	0					52	28	20	0	63	35	2	0	83	16	2	0	91	9	0	0
ES: Castile-La Mancha*	81	18	1	0					56	38	6	0	75	24	1	0	75	25	0	0	98	2	0	0
ES: Catalonia†	53	17	10	20	36	27	38	0	40	16	29	15	42	24	17	17	59	16	4	20	67	9	0	25
ES: Community of Madrid	84	15	1	0	0	0	0	0	61	39	0	0	76	21	2	0	100	0	0	0	90	10	0	0
ES: Extremadura	72	25	4	0	67	33	0	0	43	46	11	0	55	38	7	0	85	15	0	0	87	13	0	0
ES: Galicia	80	14	6	0	57	14	29	0	55	31	13	0	73	20	7	0	87	9	4	0	93	5	2	0
ES: Region of Murcia†	87	12	2	0	50	50	0	0	60	30	11	0	92	8	0	0	93	7	0	0	94	6	0	0
ES: Valencia	79	18	3	0	44	33	22	0	59	32	9	0	69	26	5	0	87	12	1	0	92	8	0	0
Sweden	60	32	8	0	7	41	52	0	51	30	19	0	57	34	9	0	59	37	5	0	74	25	1	0
the Netherlands	71	16	13	0	31	10	59	0	50	17	32	0	61	18	21	0	74	19	7	0	87	13	0	0
UK: All countries*†	69	20	9	2					51	26	21	2	63	23	12	2	75	18	5	2	83	14	0	3
UK: England*†	68	21	9	3					49	27	21	2	62	23	12	2	74	19	5	3	82	14	0	4
UK: Northern Ireland*†	70	16	13	1					43	22	36	0	65	18	17	0	69	26	3	3	95	4	0	1
UK: Scotland	78	14	8	0	20	40	40	0	67	17	17	0	73	15	11	0	88	12	1	0	86	13	1	0
UK: Wales†	70	21	9	0					52	29	19	0	60	29	11	0	70	21	9	0	91	9	0	0

Categories may not add up because of missing values or rounding off; percentages are row percentages. When cells are left empty (complete) data are unavailable. *:Patients younger than 20 years of age are not reported. †: The incident counts at day 91, on which the data presented in this table are based, are estimated.

Table 5.3. Incident rates per million age-related population of renal replacement therapy at day 1 over the time period 2011-2012 for patients aged 0-19 by age and treatment modality.

	0-19			0-4			5-9			10-14			15-19							
	HD	PD	TX	Oth	HD	PD	TX	Oth	HD	PD	TX	Oth	HD	PD	TX					
2011-2012	7.6	3.8	2.0	1.7	7.2	2.2	4.2	0.7	3.7	1.2	0.8	1.5	6.9	3.1	1.3	2.3	12.6	8.5	1.6	2.2

Data from France includes a coverage of 99.37% in 2011 and 100% in 2012. The Spanish region of Murcia is only included in 2012. Abbreviations used: HD: haemodialysis; PD: peritoneal dialysis; TX: transplant; oth: other (includes haemofiltration, haemodiafiltration and unknown); pmarp: per million age-related population.




Figure 5.3. Incidence of ESRD patients on renal replacement therapy, per million age related population (pmarp), unadjusted at day 1, in 2012 for patients aged 65 to 74 years (black bars) and 75 years and over (grey bars), and with the percentage of patients by age group.

The prevalence of RRT for ESRD

On the 31st December 2012 a total of 451,270 patients with ESRD were receiving RRT among the renal registries reporting to the ERA-EDTA Registry, corresponding to an unadjusted prevalence of 716.7 pmp. There was a substantial difference in the prevalence pmp throughout Europe, both in the unadjusted prevalence and the prevalence standardized for the age and sex distribution of the EU27 population (Figure 5.4). The unadjusted

prevalence was highest in Portugal (1670.2 pmp), French- and Dutch-speaking Belgium (1251.3 and 1206.2 pmp, respectively), whereas the unadjusted prevalence was lowest in Ukraine (146.7 pmp), Russia (213.9 pmp) and Montenegro (309.7 pmp). The prevalence pmp standardised to the EU27 was highest in French-speaking Belgium (1324.1 pmp) and the Spanish regions of Murcia (1303.1 pmp) and Catalonia (1221.3 pmp). In 2012, the mean age of the prevalent RRT patients was 60.3 years within the renal registries providing individual level patient data and 56.6 years within the renal registries providing aggregated data, with considerable variation across renal registries (Figure 5.4).

The unadjusted prevalence of patients aged 65-74 years and 75 years and over was 2400.2 pmarp and 2388.9 pmarp respectively within the renal registries providing individual level patient data (Figure 5.5). Prevalent patients aged 65-74 years formed 22% of the total prevalent patient population from these registries and patients aged 75 years and over formed 20%. The prevalence pmarp for patients aged 65-74 years was highest in French-speaking Belgium (3654.2 pmarp) and the Spanish regions of Murcia (3501.5 pmarp) and Catalonia (3382.4 pmarp). The prevalence pmarp for patients aged 75 years and above was highest in French- and Dutch-speaking Belgium (4262.6 and 4093.8 pmarp respectively) and the Spanish region of Murcia (3701.5 pmarp). Table 5.4 presents the unadjusted prevalence of ESRD patients receiving RRT pmp on 31st December 2012, by treatment modality (haemodialysis, peritoneal dialysis and transplantation) for the renal registries providing individual and aggregated data. The prevalence of haemodialysis was highest in Portugal (997.6 pmp), Greece (844.4 pmp) and Israel (691.0 pmp), whereas the prevalence of peritoneal dialysis was highest in the Spanish regions of Galicia (97.6 pmp), Asturias (96.8 pmp) and Basque Country (92.6 pmp). The prevalence of patients living with a functioning renal transplant was highest in the Spanish regions of Basque Country (670.8 pmp) and Catalonia (659.1 pmp) and Norway (639.2 pmp). Table 5.5 presents the percentage of patients by established treatment modality (haemodialysis, peritoneal dialysis and transplantation) on 31st December 2012, by age group. For prevalent patients 75 years and over, the percentage of patients receiving haemodialysis varied from 54% in Finland to 98% in Slovenia. In this age group, the percentage of patients receiving peritoneal dialysis was highest in Denmark (16%) and Sweden (15%), whereas Norway reported the highest number of patients living with a functioning kidney allograft (38%). These results were based only on countries with less than 10% missing values for treatment modality.

The prevalence of children with ESRD by age group and treatment modality is presented in Table 5.6. Overall on the 31st December 2012 the prevalence of children with ESRD was 54.7 pmarp: the majority (41.5 pmarp) had a functioning renal transplant whereas 8.1 pmarp were receiving haemodialysis and 4.6 pmarp peritoneal dialysis.

Table 5.4. Prevalence of renal replacement therapy on 31st December 2012, by established therapy as per million population (unadjusted) for individual and aggregated data.

Country / Region	All		HD	PD	Tx	Unknown / Missing
	N	pmp	pmp	pmp	pmp	pmp
Country / Region providing individual patient data						
Austria	8635	1022.7	460.5	46.0	516.3	0
Belgium: Dutch-speaking*	7679	1206.2	641.0	57.3	507.8	0
Belgium: French-speaking*	5883	1251.3	665.1	59.8	523.0	3.5
Bosnia and Herzegovina	2519	718.1	642.8	23.4	51.6	0.3
Croatia †	4411	1033.2	606.5	42.9	383.9	0
Denmark	4927	872.3	367.5	90.5	410.7	3.5
Finland	4373	807.7	266.0	59.3	482.5	0
France	74521	1138.7	580.3	41.9	507.4	9.2
Greece	12598	1135.7	844.4	59.9	231.3	0
Iceland	219	682.8	177.7	65.5	439.6	0
Norway	4453	887.3	208.8	39.3	639.2	0
Romania	14747	765.9	628.9	78.9	57.9	0.2
Serbia	5414	752.0	586.9	58.6	106.4	0.1
Slovenia	2048	995.5	662.6	22.8	310.1	0
Spain: Andalusia	8905	1062.0	478.0	46.6	537.2	0.1
Spain: Aragon	1452	1082.7	424.3	46.2	612.2	0
Spain: Asturias	1152	1072.3	382.6	96.8	592.9	0
Spain: Basque Country	2488	1140.8	377.4	92.6	670.8	0
Spain: Cantabria*	575	972.6	340.0	66.0	566.6	0
Spain: Castile and León*	2689	1063.6	432.3	69.2	552.2	9.9
Spain: Castile-La Mancha*	2114	1006.6	397.1	57.1	547.1	5.2
Spain: Catalonia†	9086	1200.1	492.4	48.6	659.1	0
Spain: Community of Madrid	6338	987.2	379.6	54.7	552.9	0
Spain: Extremadura	1163	1049.5	475.6	66.8	465.6	41.5
Spain: Galicia	3271	1182.2	526.6	97.6	558.0	0
Spain: Region of Murcia†	1703	1155.0	619.2	48.8	486.3	0.7
Spain: Valencia	6111	1191.4	647.7	64.9	478.4	0.4
Sweden	8882	933.0	320.2	82.9	530.0	0
the Netherlands	15472	923.4	325.8	58.4	539.1	0.1
UK: All countries*†	55238	867.1	372.0	60.4	434.6	0.1
UK: England*†	46456	868.4	373.3	62.1	433.0	0.1
UK: Northern Ireland*†	1557	853.8	391.0	47.2	415.7	0
UK: Scotland	4557	857.6	362.1	45.7	449.8	0
UK: Wales*†	2743	892.3	358.8	67.3	465.8	0.3
Country / Region providing aggregated patient data						
Albania	909	325.1	234.6	17.9	72.6	0
Czech Republic§	10371	996.2	554.4	47.0	394.8	0
Estonia	733	554.0	194.3	31.7	328.0	0
Georgia	1559	545.8	479.0	14.7	48.7	3.5
Israel§	5795	732.6	691.0	41.6		0
Latvia	926	538.2	195.3	52.9	290.0	0
Montenegro	192	309.7	164.5	9.7	135.5	0
Poland	28226	747.5	463.9	29.4	254.2	0
Portugal	17641	1670.2	997.6	70.3	602.3	0
Russia	30349	213.9	156.8	13.2	43.9	0
Slovakia§	3220	595.1	577.5	17.6		0
Spain (18 of 19 regions)	50909	1092.1	478.3	59.2	554.6	0
Turkey		815.6				
Ukraine	6669	146.7	109.0	19.3	18.5	0

Footnotes for Table 5.4 (overleaf): Abbreviations used: HD: haemodialysis; PD: peritoneal dialysis; TX: transplant; pmp: per million population; *: Patients younger than 20 years of age are not reported; ||: The overall prevalence of RRT is underestimated by approximately 3% due to an estimated 30% under-reporting of patients living with a functioning graft; §: Data include dialysis patients only; When cells are left empty, the data are unavailable.

Renal transplants in 2012

During 2012 there were a total of 15,673 renal transplants performed within the countries and regions reporting to the ERA-EDTA Registry, corresponding to a renal transplantation rate of 28.3 pmp (Figure 5.6). Of these 12,252 were from a deceased donor source (22.1 pmp, 78%) and 3,322 renal transplants were from a living donor source (6.0 pmp, 21%). The kidney transplantation rate varied considerably between the countries and regions reporting to the ERA-EDTA Registry, with the highest rates pmp seen in the Spanish regions of Catalonia and Cantabria (74.0 and 60.9 pmp, respectively) and in Northern Ireland (64.2 pmp). The proportion of living donor to deceased donor transplants also varied considerably within each country and region (Figure 5.6). Notably, for the countries with a transplantation rate above 20.0 pmp, Israel and the Netherlands were the only two countries with a higher proportion of living to deceased donor renal transplants (1.54 and 1.03 respectively).

The mean age at renal transplantation (available from the renal registries supplying individual level data and limited to adult patients) was 51.9 years (Figure 5.6). The mean recipient age at the time of renal transplantation was greatest in the Spanish regions of Aragon (57.0 years) and Asturias (55.7 years), and Norway (55.1 years). The youngest mean recipient ages at the point of renal transplantation were reported by Romania (38.7 years), Bosnia and Herzegovina (39.5 years) and Serbia (40.7 years). The proportion of patients aged 65 years and over at the time of renal transplantation also varied across the registries from 35% in the Spanish region of Aragon to 0% in Iceland, Slovenia and Bosnia and Herzegovina.

Patient and graft survival

Survival analyses included data from 22 registries in 11 countries which provided individual patient data for the period of 2003 to 2010. The 1-, 2- and 5-year adjusted survival probabilities for patients with ESRD commencing RRT between 2003 and 2007, are presented for the total cohort and for the older age groups in Table 5.7. Adjustments were made for age, gender and the distribution of primary renal disease (see Appendix 5.1). Five-year survival for all RRT patients was 59.7% (95%CI: 59.3; 60.0) which fell to 39.3% (95%CI: 38.7; 39.9) in patients 65 to 74 years and 21.3% (95%CI: 20.8; 21.9) in patients over 75 years.

Table 5.5. Percentages of established renal replacement therapy modality (unadjusted) of prevalent patients on 31st December 2012, by age group for the countries/regions providing individual patient data.

Country / Region	All			0-19			20-44			45-64			65-74			≥75								
	HD	PD	TX	Oth	HD	PD	TX	Oth	HD	PD	TX	Oth	HD	PD	TX	Oth	HD	PD	TX					
	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%					
Austria	34	4	50	11	9	7	79	5	21	4	65	10	24	5	60	11	39	5	46	11	64	4	19	14
BE: Dutch-speaking*	32	5	42	21					18	8	64	10	19	5	63	13	31	5	44	20	52	3	11	34
BE: French-speaking*	41	5	42	13					23	6	62	9	30	4	57	9	39	5	41	15	64	5	13	18
Bosnia and Herzegovina	81	3	7	9	65	8	27	0	60	6	20	13	79	3	7	11	91	2	1	5	95	1	0	3
Croatia †	58	4	37	1	4	27	69	0	28	6	66	1	46	4	49	1	72	3	23	2	93	3	3	1
Denmark	42	10	47	0	11	7	82	0	25	5	69	1	33	9	58	0	51	14	34	1	78	16	6	0
Finland	26	7	60	7	3	7	90	1	16	8	71	5	19	7	68	6	31	7	53	9	54	9	27	10
France	41	4	45	10	11	5	79	6	26	2	67	5	32	2	58	8	44	4	41	11	66	7	10	18
Greece	58	5	20	16	31	21	45	3	32	5	45	18	44	5	30	20	66	5	13	16	84	5	1	10
Iceland	26	10	64	0	0	20	80	0	10	4	87	0	23	2	75	0	29	7	64	0	58	36	6	0
Norway	24	4	72	0	5	2	93	0	14	2	84	0	16	3	80	0	27	4	69	0	50	12	38	0
Romania	82	10	8	0	53	28	19	0	69	9	21	1	83	9	7	0	88	11	1	0	87	13	0	0
Serbia	66	8	14	12	32	13	48	6	40	6	39	15	66	7	12	14	80	10	1	8	83	9	0	8
Slovenia	67	2	31	0	29	14	57	0	40	4	56	0	56	3	41	0	74	2	24	0	98	1	1	0
ES: Andalusia	45	4	51	0	9	3	88	0	27	4	69	0	33	4	63	0	52	4	44	0	83	5	12	0
ES: Aragon	39	4	57	0	13	0	88	0	19	8	73	0	27	3	69	0	42	6	51	0	68	2	29	0
ES: Asturias	35	9	55	0	13	25	63	0	21	11	67	0	23	10	67	0	35	6	59	0	64	9	27	0
ES: Basque Country	33	8	59	0	3	6	90	0	21	7	72	0	22	10	68	0	34	7	59	0	65	7	28	0
ES: Cantabria*	35	7	58						25	2	73	0	22	8	70	0	36	6	58	0	71	10	19	0
ES: Castile and León*	41	7	52	1					23	5	71	1	25	6	69	0	38	7	54	0	72	8	18	2
ES: Castile-La Mancha*	39	6	54	1					17	10	73	0	30	6	64	0	40	5	55	1	71	3	26	1
ES: Catalonia†	25	4	55	16	6	5	88	2	13	3	76	8	17	4	66	12	25	5	53	17	48	3	22	27
ES: Community of Madrid	45	6	44	4	0	0	100	0	28	10	56	6	30	7	57	5	52	4	40	4	84	4	12	0
ES: Extremadura	45	8	47	0	11	28	61	0	24	8	68	0	31	10	60	0	49	6	45	0	81	8	12	0
ES: Galicia	38	6	56	0	11	8	81	0	20	8	72	0	27	6	66	0	41	5	54	0	71	3	26	0
ES: Region of Murcia†	54	4	42	0	10	20	70	0	33	7	61	0	36	4	60	0	62	4	34	0	92	3	6	0
ES: Valencia	54	5	40	0	17	11	72	0	33	7	61	0	41	6	52	0	56	5	39	0	88	4	8	0
Sweden	34	9	57	0	8	9	83	0	20	5	75	0	25	7	68	0	38	11	52	0	69	15	15	0
the Netherlands	35	6	58	0	6	5	89	0	20	4	77	0	24	6	70	0	38	7	54	0	74	9	17	0
UK: All countries*†	41	7	50	2					24	5	69	1	32	6	60	2	50	9	39	2	76	9	12	3
UK: England*†	41	7	50	2					24	5	69	1	32	6	60	2	50	9	39	3	75	10	11	4
UK: Northern Ireland*†	44	6	49	2					22	7	70	1	33	4	61	2	60	7	30	3	78	7	12	3
UK: Scotland	42	5	52	0	8	15	77	0	27	3	70	0	34	5	61	0	56	8	36	0	80	7	14	0
UK: Wales*†	40	8	52	0					22	6	73	0	30	6	64	0	48	10	42	0	77	10	13	0

Abbreviations used: HD: haemodialysis; PD: peritoneal dialysis; TX: transplant; oth: other (includes haemofiltration, haemodiafiltration and unknown); *: Patients younger than 20 years of age are not reported; †: The overall prevalence of RRT is underestimated by approximately 3% due to an estimated 30% under-reporting of patients living with a functioning graft.

Categories may not add up because of missing values or rounding off. Percentages are row percentages. When cells are left empty (complete) data are unavailable.

Figure 5.6. Renal transplants performed per million population (pmp) in 2012, unadjusted, with mean and median age at transplantation. Data are from renal registries providing individual patient level data and aggregated data. For the registries providing individual patient data the percentage of patients aged over 65 years and percentage of patients aged over 75 years at the point of transplantation are presented. For some countries transplantation data were obtained from alternate sources: UK: NHS Blood and Transplant, Slovakia: Slovak Centre of Organ Transplantation, Spain (18 of 19 regions): Organización Nacional de Trasplantes. For Romania the transplantation activity reflects 70% of the total transplantation activity in the country, due to an under-reporting of pre-emptive transplantations. Mean/median age and percentage >65 or >75 years is limited to the adult patient cohort only.




Table 5.6. Prevalent rates per million age-related population of renal replacement therapy on 31st December in 2012 for patients aged 0-19 by age and treatment modality.

	0-19				0-4				5-9				10-14				15-19			
	HD	PD	TX	Oth	HD	PD	TX	Oth	HD	PD	TX	Oth	HD	PD	TX	Oth	HD	PD	TX	Oth
2012	54.7	8.1	4.6	41.5	19.4	3.9	7.3	8.1	35.9	3.5	2.9	29.3	58.4	5.9	3.4	48.8	104.0	19.0	4.8	79.1

Abbreviations used: HD: haemodialysis; PD: peritoneal dialysis; TX: transplant; pmarp: per million age-related population. Data from France includes a coverage of 99.37% in 2011 and 100% in 2012. The Spanish region of Murcia is only included in 2012.

Table 5.7. 1-, 2- and 5-year patient and graft survival probabilities (95% confidence interval), for all patients and for patients above 65 years of age, adjusted for age, gender and primary diagnosis.

	2006-2010 cohort		2003-2007 cohort		
	1 year	2 year	1 year	2 year	5 year
Incident RRT patients					
All patients	89.7 (89.5; 89.8)	82.2 (82.0; 82.5)	88.7 (88.5; 88.9)	80.6 (80.3; 80.9)	59.7 (59.3; 60.0)
65-74	82.3 (81.8; 82.8)	70.5 (70.0; 71.1)	80.3 (79.8; 80.7)	67.6 (67.1; 68.2)	39.3 (38.7; 39.9)
75+	72.6 (72.1; 73.2)	56.7 (56.1; 57.3)	70.4 (69.8; 70.9)	53.2 (52.6; 53.8)	21.3 (20.8; 21.9)
Patient survival on dialysis					
All patients	88.3 (88.1; 88.5)	79.7 (79.4; 80.0)	86.9 (86.7; 87.1)	77.5 (77.2; 77.8)	52.5 (52.0; 52.9)
65-74	81.9 (81.5; 82.4)	69.7 (69.1; 70.2)	79.9 (79.4; 80.4)	66.9 (66.3; 67.4)	36.7 (36.1; 37.3)
75+	72.4 (71.9; 73.0)	56.4 (55.8; 57.0)	70.1 (69.5; 70.7)	52.9 (52.2; 53.5)	20.9 (20.4; 21.5)
Patient survival after first renal transplantation (deceased donor)					
All patients	97.4 (97.1; 97.6)	96.1 (95.8; 96.4)	97.2 (97.0; 97.4)	96.0 (95.7; 96.2)	91.5 (91.0; 91.9)
65+	91.6 (90.4; 92.8)	87.8 (86.4; 89.2)	89.7 (88.3; 91.1)	85.6 (84.0; 87.3)	72.6 (70.5; 74.8)
Patient survival after first renal transplantation (living donor)					
All patients	98.9 (98.7; 99.2)	98.3 (98.0; 98.6)	98.5 (98.1; 98.8)	97.6 (97.1; 98.0)	95.1 (94.4; 95.8)
65+	95.4 (93.5; 97.3)	92.9 (90.5; 95.3)	95.4 (92.8; 98.1)	92.9 (89.7; 96.3)	84.3 (79.7; 89.3)
Graft survival after first renal transplantation (deceased donor)					
All patients	91.7 (91.3; 92.2)	89.3 (88.8; 89.7)	91.0 (90.6; 91.4)	88.3 (87.9; 88.8)	79.9 (79.2; 80.5)
65+	85.8 (84.2; 87.3)	81.5 (79.8; 83.2)	82.7 (80.9; 84.6)	78.4 (76.4; 80.4)	64.1 (61.8; 66.5)
Graft survival after first renal transplantation (living donor)					
All patients	95.7 (95.2; 96.2)	93.8 (93.2; 94.5)	95.0 (94.4; 95.7)	93.1 (92.3; 93.8)	86.8 (85.8; 87.8)
65+	93.0 (90.6; 95.4)	90.2 (87.4; 93.1)	92.5 (89.1; 96.0)	89.4 (85.4; 93.5)	81.3 (76.2; 86.6)

Based on data from Austria, Dutch- and French-speaking Belgium, Denmark, Finland, Greece, Iceland, Norway, the Spanish renal registries of Andalusia, Aragon, Asturias, Basque Country, Cantabria, Castile and León, Catalonia, Extremadura and Valencia, Sweden, the Netherlands and the United Kingdom (all countries). Methods are described in Appendix 5.1 on page 93.

Appendix 5.1

Coverage of the European population by the ERA-EDTA Registry

European countries supplying data to the ERA-EDTA Registry in 2012: Albania, Austria, Belgium, Bosnia and Herzegovina, Croatia, Czech Republic, Denmark, Estonia, Finland, France, Georgia, Greece, Iceland, Latvia, Montenegro, the Netherlands, Norway, Poland, Portugal, Romania, Russia, Serbia, Slovakia, Slovenia, Spain, Sweden, Turkey, Ukraine and the United Kingdom. The total coverage of the European population by the ERA-EDTA Registry in 2012 was 71.3%. For this calculation only the proportion of the population residing in the European part of Russia and Turkey were considered.

Statistical methods used to determine survival outcomes as presented in Table 5.7

The survival probabilities of all incident ESRD patients receiving RRT, dialysis patients and of patients receiving a first renal transplant between 2003 and 2007 or between 2006 and 2010 are presented in Table 5.7. The Cox regression model was used for the adjusted survival analyses for both the total group and for the older subgroup of patients (65-74 and 75 years and over). The patients were followed until 31st December 2012.

For the analysis of patient survival on RRT, the first day of RRT was taken as the starting point and the event studied was death. Censored observations were recovery of renal function, loss to follow-up and end of follow-up time. Adjustments were made for fixed values for age (60 years), gender (60% men) and primary renal disease (20% diabetes mellitus, 17% hypertension/renal vascular disease, 15% glomerulonephritis and 48% other cause). In all the analyses of the older subgroups the corresponding fixed values were used with the exception of age. For the analysis of patient survival on dialysis the first day on dialysis was taken as the starting point, the event studied was death and censored observations were recovery of renal function, loss to follow-up, end of follow-up time and renal transplantation. Adjustments were made for fixed values of age (60 years), gender (60% men) and primary renal disease (20% diabetes mellitus, 17% hypertension/renal vascular disease, 15% glomerulonephritis and 48% other cause). For the analysis of patient and graft survival after transplantation the date of the first renal transplantation was defined as the first day of follow-up. The event studied for the patient survival after transplantation was death, while for the graft survival the events were graft failure and death. Censored observations were loss to follow-up and end of the follow-up time. Adjustments were made for fixed values of age (45 years), gender (60% men) and primary renal disease (10% diabetes mellitus, 8% hypertension/renal vascular disease, 28% glomerulonephritis and 54% other cause). Patients for whom age, gender, or primary renal disease was missing were excluded from the analysis.