UvA-DARE (Digital Academic Repository)

In vitro splicing of pre-mRNA containing bromouridine.

Wansink, D.G.; Nelissen, R.L.H.; de Jong, L.

DOI

[10.1007/BF00997156](https://doi.org/10.1007/BF00997156)

Publication date

1994

Published in

Mol. Biol. Rep.

[Link to publication](#)

Citation for published version (APA):

Wansink, D. G., Nelissen, R. L. H., & de Jong, L. (1994). In vitro splicing of pre-mRNA containing bromouridine. *Mol. Biol. Rep.*, 19, 109-113. <https://doi.org/10.1007/BF00997156>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

***In vitro* splicing of pre-mRNA containing bromouridine**

Derick G. Wansink¹, Rob L. H. Nelissen² & Luitzen de Jong¹

¹E. C. Slater Institute, University of Amsterdam, Plantage Muidergracht 12, 1018 TV Amsterdam, The Netherlands

²Department of Biochemistry, University of Nijmegen, PO Box 9101, 6500 HB Nijmegen, The Netherlands

Received 21 January 1994; accepted 31 January 1994

Key words: bromouridine, nucleotide analogue, pre-mrna, polypyrimidine tract, splicing

Abstract

The artificial UTP-analogue 5-bromouridine 5'-triphosphate (BrUTP) has been used to label pre-mRNA *in vitro* and *in vivo* [1, 2]. We have investigated the effect of bromouridine (BrU) in pre-mRNA on the efficiency of splicing. An adenovirus major late II construct was used to prepare four different transcripts, each containing a different amount of BrU. These four transcripts were tested in an *in vitro* splicing assay. We found that splicing is strongly inhibited if all uridines (U) in the transcript were substituted for BrU. Splicing was restored to some extent if 50% of the Us were replaced by BrU. The splicing efficiency returned to an almost normal level if only 1 out of every 10 Us was substituted for BrU. This demonstrates that only a pre-mRNA containing a small amount of BrU can be spliced normally *in vitro*. Furthermore, these results strongly suggest that some Us in the adenoviral transcript, probably those at the splice sites, cannot be replaced by BrU and are therefore critical in the splicing reaction.

Abbreviations: BrU = bromouridine; BrUTP = 5-bromouridine 5'-triphosphate; snRNP = small nuclear ribonucleoprotein particle; U = uridine.

Introduction

Splicing includes the removal of introns and the subsequent ligation of exons of pre-mRNAs. Splicing takes place in so-called spliceosomes, which are complexes composed of small nuclear ribonucleoprotein particles (snRNPs) and non-snRNP proteins [reviewed in 3, 4, 5]. The splicing process consists of two transesterification steps and involves the successive binding of several snRNPs and non-snRNP proteins to specific sites of the pre-mRNA. Several conserved sequences, which are essential for splicing, have been characterized in pre-mRNAs [3, 5]. These conserved sequences include the 5'-splice site, the branch point, the polypyrimidine tract, and the 3' splice site (see consensus sequences in Fig. 1).

Recently, we and others have reported an immunofluorescence method to visualize active sites of transcription in the interphase cell nucleus *in vitro* and *in vivo* [1, 2]. This method is based on the incorporation of the UTP-analogue 5-bromouridine 5'-triphosphate

(BrUTP) into nascent RNA and the subsequent detection of the nucleotide with a specific monoclonal antibody. We have shown that BrUTP is efficiently incorporated by RNA polymerase II and does hardly affect the kinetics of nuclear run-on transcription when it replaces UTP [2]. Likewise, labeling pre-mRNA with BrU may also be useful in studying RNA processing in the nucleus. A prerequisite would be that BrU-labeled pre-mRNA is properly spliced. Therefore, in this study we have investigated the effect of BrU on the efficiency of splicing *in vitro*. Transcripts containing BrU were synthesized and splicing was tested in an *in vitro* splicing assay. We find that splicing is strongly inhibited if all Us in the pre-mRNA are replaced by BrU. However, substitution of only a limited fraction of the Us in a transcript hardly affects the efficiency of *in vitro* splicing.

Materials and methods

BrUTP

BrUTP was obtained from Sigma Chemical Co. (St. Louis, MO, USA). A sample was tested for contaminating nucleotides by HPLC on an ion exchange column as described earlier [6]. Two contaminants were found, probably corresponding to BrUDP (12%) and BrUMP (1%). No UTP could be detected.

In vitro transcription

BrUTP is an artificial nucleotide that substitutes for UTP during RNA synthesis [2, 7]. An adenovirus major late II DNA construct, linearized with Sca I [8] (see Fig. 2), was used to synthesize BrU-labeled RNA *in vitro* using the SP6 RNA polymerase system. Four transcription reactions were done with SP6 RNA polymerase in a final volume of 50 μ l to obtain pre-mRNAs containing different amounts of BrU. The transcription reaction was performed in the presence of 0.6 mM GpppG, 1 mM CTP, 1 mM GTP, 24 μ M [α - 32 P]ATP, and either 1 mM BrUTP, 0.5 mM BrUTP + 0.5 mM UTP, 0.1 mM BrUTP + 0.9 mM UTP, or 1 mM UTP. It is expected that SP6 RNA polymerase has, like eukaryotic RNA polymerases, about the same K_m for BrUTP as for UTP [2, 7]. As a result four different types of transcripts were obtained, containing 100%, ~50%, ~10% and 0% BrUTP (percentage of the total UTP content), respectively.

The RNA transcription products were separated on a 3.5% preparative polyacrylamide gel (0.5 $\times$ TBE, 7 M urea). A piece of the gel containing the full length pre-mRNA was cut out and the pre-mRNA was eluted (16 h, 4 $^{\circ}$ C) in 1 ml elution buffer (0.5 M ammonium acetate, pH 7.5; 10 mM MgCl₂; 0.1% SDS). After phenol/chloroform extraction and ethanol precipitation the pre-mRNA pellet was dissolved in 25 μ l H₂O.

In vitro splicing

HeLa S100 splicing extracts were prepared from HeLa cells as described previously [9]. Our standard *in vitro* splicing reaction (12.5 μ l) contained 6 μ l HeLa splicing extract, 2 μ l adenovirus major late II pre-mRNA, 10% glycerol, 10 mM Hepes, pH 7.9, 2.5 mM MgCl₂, 50 mM KCl, 2 mM ATP, 10 mM creatine phosphate, 0.25 mM DTE, and 0.1 mM EDTA. Reaction mixtures were incubated at 30 $^{\circ}$ C for 45 or 120 min. The reactions were terminated by adding 90

μ l ice-cold stop mix (0.3 M sodium acetate, pH 5.2; 100 μ g/ml *E. coli* tRNA; 0.1% SDS), immediately followed by phenol/chloroform extraction and ethanol precipitation. 32 P-labeled RNAs were separated on a 10% polyacrylamide gel (0.5 $\times$ TBE, 7 M urea) and visualized by autoradiography (Kodak Xomat).

Results and discussion

To investigate the effect of BrU in pre-mRNA on the efficiency of splicing, we have tested the splicing of BrU-labeled transcripts in an *in vitro* splicing assay. As substrate a 32 P-labeled adenovirus major late II transcript (253 nt), containing two exons and one intron, was used (Fig. 2). Four different types of this transcript were synthesized, characterized by the substitution of 100%, ~50%, ~10%, or 0% of all Us for BrU, respectively (see Materials and methods). These transcripts were incubated in a splicing assay with HeLa cell splicing extract for 45 or 120 min. Subsequently, RNA splicing products and intermediates were separated by gel electrophoresis and visualized by autoradiography (Fig. 3).

Adenoviral transcripts containing no BrU served as a control in the splicing reaction (i.e. 0% BrU; Fig. 3, lanes 4). After 45 min of incubation, splicing had taken place as is illustrated by the presence of the spliced product and all reaction intermediates. After 2 h more spliced product was formed, but still a lot of pre-mRNA was present. Obviously, pre-mRNA was present in an excess amount and splicing had not completely finished after 2 h of incubation.

Almost no spliced product was detected when pre-mRNA, in which all Us were replaced by BrU, was incubated in the splicing assay (i.e. 100% BrU; lanes 1). The free 5'-exon was the only splicing intermediate that was observed suggesting that the intron-3'-exon lariat and the intron lariat were degraded. To exclude the possibility that the small amount of spliced product was the result of a small UTP contamination in the BrUTP starting material, we have tested the purity of this reagent by HPLC (see Materials and methods). No UTP could be detected, which implies that pre-mRNA used in lanes 1 does not contain any U. In conclusion, BrUTP incorporation strongly inhibited splicing of the adenoviral major late II transcript. Still, adenoviral major late II transcripts in which all Us were replaced by BrU were spliced *in vitro*, albeit at an extremely low rate.

Fig. 1. Consensus splice site sequences in a mammalian intron [3, 5]. N is any nucleotide, R is any purine, Y is any pyrimidine.

Adenovirus major late II construct (253 nt)

Fig. 2. Sequence of the adenovirus major late II transcript that was used in the *in vitro* splicing assay [8]. The transcript contains one intron and two exons as depicted. The pyrimidine tract is underlined. The adenosine residue at the branch site is in *italic* and underlined. The 5'- and 3'-splice sites of the intron are doubly underlined. Uridine residues at consensus sites are depicted in bold. Note that uridines may be replaced by bromouridines as is described in Materials and methods.

If at random 50% or 10% of the Us were replaced by BrU the splicing efficiency increased significantly (Fig. 3, lanes 2 and lanes 3). The rate of splicing was still lower than in the control reaction (lanes 4), but all splicing intermediates and the spliced product were formed. Comparing the relative amounts of the splicing intermediates and spliced product in lanes 4 with those in lanes 2 and 3 may indicate the mode of inhibition of BrU. Since [free 5'-exon] + [spliced

product] is proportional to the rate of the first step of splicing, and [spliced product]/[free 5'-exon] is proportional to the rate of the second step of the splicing reaction, we conclude that both transesterification steps are inhibited by the incorporation of BrU. In the case of 50% BrU RNA (lanes 2) the intron-3'-exon lariat was present in somewhat smaller amounts than the free 5'-exon. This again indicates that the intron-3'-exon lariat is relatively unstable or actively degraded.

Fig. 3. *In vitro* splicing of RNA. Adenovirus major late II pre-mRNAs containing a certain amount of BrU were incubated in a HeLa S100 splicing extract as described in Materials and methods. Lanes 1, RNA containing 100% BrU (percentage of total U amount); lanes 2, 50% BrU/50% U; lanes 3, 10% BrU/90% U; lanes 4, 100% U (control). The different pre-mRNAs were incubated in the splicing assay for 45 or 120 min as indicated at the top of the figure. A schematic representation of the adenovirus major late II pre-mRNA (see Fig. 2), splicing intermediates, and spliced product is indicated on the right of the autoradiograph.

Although BrU-labeled pre-mRNA does not seem to be unstable (Fig. 3), we cannot exclude the possibility that instability of BrU-containing splicing intermediates plays a role in the overall splicing efficiency of BrU-containing pre-mRNA.

The consensus splice site sequences of a mammalian intron contain a few Us, which are probably important in the splicing reaction (Fig. 1). The corresponding Us in the sequence of the adenovirus major late II transcript are depicted bold in Figure 2. We cannot conclude that any of these Us, when replaced by BrU, is responsible for the inhibition of splicing. The incorporation of BrU may modify or disturb specific RNA-protein interactions or base pairs, but may also result in a general change in RNA conformation. Especially prominent is the polypyrimidine tract, which contains eight successive Us. This sequence may be

responsible for the strong inhibition of the second step of the splicing reaction [see also 10].

Summarizing, we have shown that the presence of BrU in pre-mRNA inhibits splicing *in vitro*. The degree of inhibition depends strongly on the number of BrUs that are present in the pre-mRNA. Probably, some Us in the pre-mRNA are important for splicing and can therefore not be replaced by BrU. Pre-mRNA labeled with a limited amount of BrU is apparently spliced properly. This pre-mRNA may be used to study the dynamics of pre-mRNA metabolism in the interphase cell nucleus in pulse chase experiments. BrU-labeled RNA can also be useful in immunoprecipitation experiments, and in crosslinking experiments to study RNA-protein interactions, as has been described earlier [11, 12].

Acknowledgements

We thank Mr. Louis Hartog (E.C. Slater Institute) for determining the purity of BrUTP by HPLC. We thank Professor Roel van Driel (E.C. Slater Institute) for critically reading the manuscript.

References

1. Jackson DA, Hassan AB, Errington RJ & Cook PR (1993) *EMBO J.* 12: 1059–1065
2. Wansink DG, Schul W, van der Kraan I, van Steensel B, van Driel R & de Jong L (1993) *J. Cell Biol.* 122: 283–293
3. Lührmann R, Kastner B & Bach M (1990) *Biochim. Biophys. Acta* 1087: 265–292
4. Green MR (1991) *Annu. Rev. Cell Biol.* 7: 559–599
5. Lamond AI (1993) *Bioessays* 15: 595–603
6. Wise JG, Hicke BJ & Boyer PD (1987) *FEBS Letters* 223: 395–401
7. Nakayama C & Saneyoshi M (1984) *J. Biochem.* 96: 1501–1509
8. Frendewey D & Keller W (1985) *Cell* 42: 355–367
9. Dignam JD, Lebovitz RM & Roeder RG (1983) *Nucl. Acids Res.* 11: 1475–1489
10. Roscigno RF, Weiner M & Garcia-Blanco MA (1993) *J. Biol. Chem.* 268: 11222–11229
11. Murray MT, Krohne G & Franke WF (1991) *J. Cell Biol.* 112: 1–11
12. Willis MC, Hicke BJ, Uhlenbeck OC, Cech TR & Koch TH (1993) *Science* 262: 1255–1257