UvA-DARE (Digital Academic Repository)

From Complex I to hydrogenase and back

Albracht, S.P.J.; de Jong, A.M.P.; Kotlyar, A.B.

Publication date
1994

Published in
Journal of inorganic biochemistry

[Link to publication](#)

Citation for published version (APA):

Albracht, S. P. J., de Jong, A. M. P., & Kotlyar, A. B. (1994). From Complex I to hydrogenase and back. *Journal of inorganic biochemistry*, 56, 36.

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

L01 FROM COMPLEX I TO HYDROGENASE AND BACK

S.P.J. Albracht^a, A.M.Ph. de Jong^a and A.B. Kotlyar^b

^a*E.C. Slater Institute, BioCentrum Amsterdam, Plantage Muidergracht 12, NL-1018 TV Amsterdam, The Netherlands* and ^b*Laser Laboratory for Fast Reactions in Biology, Department of Biochemistry, George S. Wise Faculty of Life Sciences, Ramat Aviv, 69978, Tel Aviv, Israel*

Since the discovery of iron-sulphur clusters in mitochondrial Complex I by Beinert and Sands in 1960 [1] quite some research groups have been studying this most complicated enzyme. At present at least four different Fe-S clusters have been detected with EPR, but their precise function in the energy-linked electron transfer catalyzed by the enzyme is not really understood. The analysis of Weidner et al. [2] of the operon encoding Complex I in *Escherichia coli* indicates that only 14 of the 41 polypeptides of the bovine-mitochondrial enzyme [3] are essential for coupled electron transfer. Five polypeptides show conservative Cys patterns that might accommodate Fe-S clusters. Four of these are quite likely inherited from hydrogenases [see e.g. 4]. The remaining polypeptide, the TYKY subunit [3], contains a Cys pattern typical for two classical cubane clusters. With this information a monomeric model, rather than a dimeric one [5] can be constructed, explaining most physico-chemical and kinetic properties of the enzyme (see figure). All Fe-S

clusters of Complex I are fully reduced within 5 ms, when SMP are mixed with NADH. Within 40 ms the g_z line of the EPR signal of the clusters 2, but not the g_{xy} line, disappears in coupled particles. This effect is sensitive to uncouplers. It is also reversed upon anaerobiosis. It is concluded that we have detected an

'energized' form of Complex I in which the protein structure around the clusters 2 has changed. It is proposed here that the TYKY subunit holds the Fe-S clusters 2 and renders the enzyme the ability to perform coupled electron transfer.

1. H. Beinert and R.H. Sands, *Biochem. Biophys. Res. Commun.* **3**, 41 (1960).
2. U. Weidner, S. Geier, A. Ptock, T. Friedrich, H. Leif and H. Weiss, *J. Mol. Biol.* **233**, 109 (1993).
3. J.E. Walker, *Quart. Rev. Biophys.* **25**, 253 (1992).
4. S.P.J. Albracht, *Biochim. Biophys. Acta* **1144**, 221 (1993)
5. R.van Belzen, M.C.M. van Gaalen, P.A. Cuypers and S.P.J. Albracht, *Biochim. Biophys. Acta* **1017**, 152 (1990)