



## UvA-DARE (Digital Academic Repository)

### Initial NICER observations of a broadened iron line and QPOs in MAXI J1535a571

Gendreau, K.; Arzoumanian, Z.; Markwardt, C.; Okajima, T.; Strohmayer, T.; Remillard, R.; Chakrabarty, D.; Prigozhin, G.; LaMarr, B.; Pasham, D.; Steiner, J.; Homan, J.; Miller, J.; Bult, P.; Cackett, E.; Iwakiri, W.; Enoto, T.; Uttley, P.

**Publication date**

2017

**Document Version**

Final published version

**Published in**

The astronomer's telegram

**License**

Unspecified

[Link to publication](#)

**Citation for published version (APA):**

Gendreau, K., Arzoumanian, Z., Markwardt, C., Okajima, T., Strohmayer, T., Remillard, R., Chakrabarty, D., Prigozhin, G., LaMarr, B., Pasham, D., Steiner, J., Homan, J., Miller, J., Bult, P., Cackett, E., Iwakiri, W., Enoto, T., & Uttley, P. (2017). Initial NICER observations of a broadened iron line and QPOs in MAXI J1535a571. *The astronomer's telegram*, 10768. <http://www.astronomerstelegam.org/?read=10768>

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

**Outside**  
 GCN  
 IAUCs

**Other**  
 ATel on Twitter and Facebook  
 ATELstream  
 ATel Community Site



[ [Previous](#) | [Next](#) | [ADS](#) ]

## Initial NICER observations of a broadened iron line and QPOs in MAXI J1535â 571

ATel #10768; *K. Gendreau, Z. Arzoumanian, C. Markwardt, T. Okajima, T. Strohmayer (GSFC), R. Remillard, D. Chakrabarty, G. Prigozhin, B. LaMarr, D. Pasham, J. Steiner, J. Homan (MIT), J. Miller (University of Michigan), P. Bult (GSFC), E. Cackett (Wayne State), W. Iwakiri (RIKEN), T. Enoto (Kyoto University), P. Uttley (University of Amsterdam) for the NICER Team*

on 22 Sep 2017; 04:07 UT

Credential Certification: Keith Gendreau ([keith.c.gendreau@nasa.gov](mailto:keith.c.gendreau@nasa.gov))

Subjects: X-ray, Binary, Black Hole

Referred to by ATel #: [10816](#), [11020](#)

[Tweet](#)


NICER has observed the new X-ray transient MAXI J1535â 571 (GCN #21788, ATels #10699, #10700, #10702, #10704, #10708, #10711, #10714, #10716, #10734, #10745) several times from 2017 September 9 through September 20. Over this time the flux has grown from 3E-8 ergs/cm<sup>2</sup>/s to 1.2E-7 ergs/cm<sup>2</sup>/s (2-10 keV). The heavily absorbed source had a NICER count rate grew from 3200 to 17,000 counts per second over this period. We fit the time-averaged 1-9.5 keV spectrum of 5.4 ksec of MAXI J1535â 571 data taken on September 13 with a model consisting of a disk blackbody and relativistic reflection including an intrinsic power-law (relxill), modified by interstellar absorption (tbabs). A number of Gaussian lines were included in the model to account for instrument-related residuals that will be corrected in later calibrations. We measured a column density of  $N_H = (4.89 \pm 0.06)E+22 \text{ cm}^{-2}$ ; this is about twice the value derived from Swift data (ATel #10731). The disk parameters were  $kT = 0.58 \pm 0.03 \text{ keV}$ , with a normalization of  $K = 1.6 (+0.1, -0.4) E+4$ . The reflection fraction was  $1.3 \pm 0.2$ , the iron abundance was  $A_{Fe} = 2.0 (+0.1, -0.4)$ , and the ionization parameter of the disk was constrained to be  $\log(\xi) = 3.7 (+0.1, -0.2)$ . Parameters related to the inner disk and the black hole are of special interest. The black hole spin parameter was  $a = 0.88 (+0.1, -0.2)$ , and the inner disk inclination was  $i = 27 (+1, -5) \text{ degrees}$ . These fits assumed a radial emissivity profile with a broken power-law form, with power-law indices  $q_1 = 6 (+1, -4)$  and  $q_2 = 2.2 \pm 0.1$ , and a break radius  $r_{\text{break}} = 5.5 (+0.9, -0.3) \text{ GM}/c^2$ . We note that preliminary fits to subsequent spectra yielded consistent values for the black hole spin and the inner disk inclination. With this particular model, we infer an absorbed source flux of  $6.1 E-8 \text{ erg}/\text{cm}^2/\text{s}$ , and an unabsorbed flux of  $1.7 E-7 \text{ erg}/\text{cm}^2/\text{s}$  in the 0.5-10.0 keV band. Significant variations on time scales ranging from a second to several hundred seconds are clearly visible in the 0.2-12 keV NICER light curves. To quantify this variability further, we extracted an average Leahy normalized (mean noise level of 2) power density spectrum from two observation periods

### Related

- 11884 INTEGRAL detects re-brightening of the black hole candidate MAXI J1535-571
- 11682 MAXI/GSC detection of an undergoing soft-to-hard state transition of MAXI J1535-571
- 11652 Re-brightening of the black hole candidate MAXI J1535-571 as it transitioned back to the soft state
- 11611 X-ray spectral hardening and radio non-detection of MAXI J1535-571
- 11568 Rapidly fading the black hole X-ray nova MAXI J1535-571 from MAXI/GSC observations
- 11020 MAXI J1535-571 currently shows softest spectrum
- 10899 Radio re-brightening of MAXI J1535-571 as it transitions back towards the hard state
- 10816 Brackett-Gamma emission in MAXI J1535-571
- 10768 Initial NICER observations of a broadened iron line and QPOs in MAXI J1535â 571
- 10761 MAXI/GSC observations indicate further softening of MAXI J1535-571 toward the high/soft state
- 10745 Extremely Bright Radio and (Sub-)Millimetre Detections of MAXI J1535-571
- 10734 Detection of the low-frequency QPOs in MAXI J1535-571
- 10733 Swift/BAT sees MAXI J1535-571 declining in 15-50 keV
- 10731 Swift/XRT sees MAXI J1535-571 brightening and softening
- 10729 MAXI/GSC detects an X-ray spectral softening of the X-ray transient MAXI J1535-571
- 10716 The near-infrared counterpart of X-ray transient MAXI J1535-571
- 10714 MAXI J1535-571 (Candidate Black Hole X-ray Binary): Coordination
- 10711 ATCA radio detection of MAXI J1535-571 indicates it is a strong black hole X-ray binary candidate

between September 12, 10:53:39 and September 13, 22:40:40, totaling about 15 ksec in duration. The overall shape of both power spectra can be characterized as a flat-top at the lowest frequencies ( $< 0.2$  Hz) breaking into a power-law above 0.2 Hz and becoming constant (noise) above roughly 30 Hz. We see a clear set of low-frequency QPOs superimposed on the power-law portion of the power spectrum. The centroid of the lower frequency QPO drifts between 1.9 and 2.8 Hz (Coherence,  $Q = 6$ ) and the centroid of the second drifts between 3.8 and 5.6 Hz ( $Q = 3$ ). The centroid frequencies of the two QPOs are always in a 1:2 ratio. The fractional root-mean-squared amplitude of the two QPOs is 6% and 5%, respectively. We stress that the instrument calibration is preliminary, that backgrounds have been neglected given the extremely high source flux, and that these fits should be regarded as an initial characterization of the data.

- 10708** Further brightening of the X-ray transient MAXI J1535-571, suggesting the presence of a black hole
- 10704** Erratum to ATel #10702: MAXI J1535-571 optical counterpart coordinates
- 10702** Discovery of the optical counterpart of MAXI J1535-571
- 10700** MAXI J1535-571: Swift detection and localization
- 10699** MAXI/GSC discovery of a new hard X-ray transient MAXI J1535-571



Reach **550 MILLION** professionals with LinkedIn ads

See how

in

[ [Telegram Index](#) ]

R. E. Rutledge, Editor-in-Chief

Derek Fox, Editor

Mansi M. Kasliwal, Co-Editor

[rrutledge@astronomerstelegam.org](mailto:rrutledge@astronomerstelegam.org)

[dfox@astronomerstelegam.org](mailto:dfox@astronomerstelegam.org)

[mansi@astronomerstelegam.org](mailto:mansi@astronomerstelegam.org)