



UvA-DARE (Digital Academic Repository)

Swift J181723.1-164300 is likely a new bursting neutron star low-mass X-ray binary

Parikh, A.; Wijnands, R.; Degenaar, N.; Altamirano, D.

Publication date

2017

Document Version

Final published version

Published in

The astronomer's telegram

License

Unspecified

[Link to publication](#)

Citation for published version (APA):

Parikh, A., Wijnands, R., Degenaar, N., & Altamirano, D. (2017). Swift J181723.1-164300 is likely a new bursting neutron star low-mass X-ray binary. *The astronomer's telegram*, 10612. <http://www.astronomerstelegam.org/?read=10612>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

The Astronomer's Telegram

[Post](#) | [Search](#) | [Policies](#)
[Credential](#) | [Feeds](#) | [Email](#)

7 Nov 2018; 13:40 UT

This space for free for your conference.

Outside

GCN
IAUCs

Other

ATel on [Twitter](#) and [Facebook](#)
[ATELstream](#)
[ATel Community Site](#)[[Previous](#) | [Next](#) | [ADS](#)]

Swift J181723.1-164300 is likely a new bursting neutron star low-mass X-ray binary

ATel #10612; *Aastha Parikh, Rudy Wijnands, Nathalie Degenaar, Diego Altamirano*
on 2 Aug 2017; 13:58 UTCredential Certification: *Rudy Wijnands (radwijnands@gmail.com)*

Subjects: X-ray, Binary, Neutron Star, Transient

Referred to by ATel #: [10624](#), [10665](#)

On 28 July 2017 Swift/BAT triggered (#00765081) on an event corresponding to a previously unknown source (Barthelmy et al. 2017, GCN #[21369](#), #[21385](#)). Its properties suggested it was likely a Galactic source and not a gamma-ray burst. The source was therefore named Swift J181723.1-164300.

The BAT light curve of the event shows a peak lasting ~ 10 s. We extracted the spectrum of the central 8 s of the burst. When fitting a black-body model to the spectrum, we find a temperature of ~ 3 keV with an emitting region having a radius of ~ 8 km, based on our distance estimate of ~ 7 kpc (see below). The peak of the burst has a flux (15-150 keV) of $\sim 1E-8$ erg/cm²/s and a bolometric flux of $\sim 6E-8$ erg/cm²/s. The temperature and the peak flux are very similar to what is seen for thermonuclear type-I bursts observed from accreting neutron stars in Galactic low-mass X-ray binaries (LMXBs). Therefore, we suggest that this source is a neutron star LMXB as well. Assuming, the BAT burst was a thermonuclear burst occurring at Eddington luminosity ($L_{\text{Edd}} = 3.8E38$ erg/s; Kuulkers et al. 2003, A&A, 399, 663), we calculate the source to be at a distance of $7(+/- 1)$ kpc. On 30 July 2017, another burst was seen from this source (#00765422; Palmer et al. 2017, GCN #[21400](#)). Likely this trigger was also caused by a thermonuclear burst from the source, ~ 2.2 days after the first trigger. This was supported by the fact that the BAT spectrum and the BAT flux during this burst were very similar as observed during the first burst.

Swift/XRT began monitoring the source ~ 90 s after the first BAT trigger. The XRT light curve shows a decaying count rate (0.5-10 keV) from ~ 4 c/s at the start to ~ 1.4 c/s ~ 200 s after the BAT trigger, after which it remained approximately constant. We fit an exponential decay curve to the initial ~ 100 s of the light curve and found a e-folding time of ~ 50 s. We carry out time resolved spectroscopy of the data separating it into two intervals that correspond to the peak in the light curve and average accretion level. The average accretion level was well fit by an absorbed power-law model with an N_{H} of $\sim 8E22$ cm⁻² and with with an index of ~ 1.5 . The unabsorbed (0.5-10 keV) flux was $\sim 5E-10$ erg/s/cm², resulting in a 0.5-10 keV luminosity of $\sim 3E36$ erg/s. Both this luminosity as well as the photon index are typical of a neutron star in its hard state. The spectrum of the decay part of the XRT light curve was fit using a two component model: a black-body

Related

- [10665](#) [Fermi Gamma Ray Burst Monitor detects bursts from Swift J181723.1-164300](#)
- [10624](#) [Swift J181723.1-164300 has entered quiescence](#)
- [10612](#) [Swift J181723.1-164300 is likely a new bursting neutron star low-mass X-ray binary](#)

model representing emission due to the type-I burst, and a power-law model representing the persistent accretion emission. For the power-law model we used the same photon index and normalization as obtained for the persistent flux after the burst. The black-body component indicated a temperature of ~ 1 keV, showing a decrease since the BAT peak. This is indeed expected for a thermonuclear burst, further confirming that the BAT trigger was likely indeed due to such an event. Since this first observation, the source has been monitored using the XRT for a total of ~ 13 ks (until 2 August 2017). The count rates were between 0.8-1.4 c/s and the spectral properties of the source were very similar during these additional observations as those seen right after the end of the first type-I burst.

The source has previously been in the Swift/XRT field of view twice - on 10 February 2011 and 8 October 2012. Both observations were very short (with a combined exposure time of ~ 500 s) and our source was not detected. We determined upper limits on the count rate using the prescription by Gehrels (1986, APJ, 303, 336). We calculated the count rate upper limits to be $< 5E-3$ c/s. We simulated a spectrum in Xspec to determine the flux upper limit (0.5-10 keV), which was found to be $< 8E-13$ erg/cm²/s. Based on our distance estimate, this corresponds to a luminosity upper limit of $< 5E33$ erg/s (0.5-10 keV), a factor ~ 600 lower than its current activity level.

[**Telegram Index**]

R. E. Rutledge, Editor-in-Chief

rrutledge@astronomerstelegam.org

Derek Fox, Editor

dfox@astronomerstelegam.org

Mansi M. Kasliwal, Co-Editor

mansi@astronomerstelegam.org