UvA-DARE (Digital Academic Repository)

What features of psychopathy might be central? A network analysis of the psychopathy checklist-revised (PCL-R) in three large samples

Verschuere, B.; van Ghesel Grothe, S.; Waldorp, L.; Watts, A.L.; Lilienfeld, S.O.; Edens, J.F.; Skeem, J.L.; Noordhof, A.

DOI

[10.1037/abn0000315](https://doi.org/10.1037/abn0000315)

Publication date

2018

Document Version

Final published version

Published in

Journal of Abnormal Psychology

License

Article 25fa Dutch Copyright Act

[Link to publication](#)

Citation for published version (APA):

Verschuere, B., van Ghesel Grothe, S., Waldorp, L., Watts, A. L., Lilienfeld, S. O., Edens, J. F., Skeem, J. L., & Noordhof, A. (2018). What features of psychopathy might be central? A network analysis of the psychopathy checklist-revised (PCL-R) in three large samples. *Journal of Abnormal Psychology*, 127(1), 51-65. <https://doi.org/10.1037/abn0000315>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

What Features of Psychopathy Might Be Central? A Network Analysis of the Psychopathy Checklist-Revised (PCL-R) in Three Large Samples

Bruno Verschuere, Sophia van Ghesel Grothe,
and Lourens Waldorp
University of Amsterdam

Ashley L. Watts and Scott O. Lilienfeld
Emory University

John F. Edens
Texas A&M University

Jennifer L. Skeem
University of California, Berkeley

Arjen Noordhof
University of Amsterdam

Despite a wealth of research, the core features of psychopathy remain hotly debated. Using network analysis, an innovative and increasingly popular statistical tool, the authors mapped the network structure of psychopathy, as operationalized by the Psychopathy Checklist—Revised (PCL-R; Hare, 2003) in two large U.S. offender samples ($n_{NIMH} = 1559$; $n_{Wisconsin} = 3954$), and 1 large Dutch forensic psychiatric sample ($n_{TBS} = 1937$). Centrality indices were highly stable within each sample, and indicated that callousness/lack of empathy was the most central PCL-R item in the 2 U.S. samples, which aligns with classic clinical descriptions and prototypicality studies of psychopathy. The similarities across the U.S. samples offer some support regarding generalizability, but there were also striking differences between the U.S. samples and the Dutch sample, wherein the latter callousness/lack of empathy was also fairly central but irresponsibility and parasitic lifestyle were even more central. The findings raise the important possibility that network-structures do not only reflect the structure of the constructs under study, but also the sample from which the data derive. The results further raise the possibility of cross-cultural differences in the phenotypic structure of psychopathy, PCL-R measurement variance, or both. Network analyses may help elucidate the core characteristics of psychopathological constructs, including psychopathy, as well as provide a new tool for assessing measurement invariance across cultures.

General Scientific Summary

What is psychopathy? Network analyses in three large samples (totaling 7,450 offenders) indicate that affective-interpersonal features, most notably callousness and lack of empathy, may be central to psychopathy.

Keywords: psychopathy, Psychopathy Checklist—Revised (PCL-R), network analysis, empathy, cross-cultural

Research on psychopathy is thriving. Despite the accumulation of knowledge on psychopathy, there remains active debate on its origins, assessment, and clinical and legal implications (DeMatteo et al., 2014; Skeem, Polaschek, Patrick, & Lilienfeld, 2011). At a more basic level, much disagreement surrounds the mere definition of psychopathy, such as what fea-

tures are integral, irrelevant, or peripheral to psychopathy (e.g., Miller & Lynam, 2012; Lilienfeld et al., 2012; Skeem & Cooke, 2010). Indeed, an essential question remains: What is psychopathy?

Psychopathy has often been conceptualized in terms of a constellation of traits. Following the structure of the widely used

This article was published Online First November 27, 2017.

Bruno Verschuere and Sophia van Ghesel Grothe, Department of Clinical Psychology, University of Amsterdam; Lourens Waldorp, Department of Psychological Methods, University of Amsterdam; Ashley L. Watts and Scott O. Lilienfeld, Department of Psychology, Emory University; John F. Edens, Department of Psychology, Texas A&M University; Jennifer L. Skeem, School of Social Welfare, Goldman School of Public Policy, University of California, Berkeley; Arjen Noordhof, Department of Clinical Psychology, University of Amsterdam.

We are very grateful to Joseph Newman for sharing the data of the Wisconsin sample. We thank the Expertisecentrum Forensische Psychiatrie for making the Dutch data available. We thank Lena Christ and Miguel Beira for the initial analyses.

Correspondence concerning this article should be addressed to Bruno Verschuere, Department of Clinical Psychology, University of Amsterdam, Postbus 15933, 1001 NK Amsterdam, the Netherlands. E-mail: B.J.Verschuere@uva.nl

Psychopathy Checklist—Revised (PCL-R; Hare, 2003), these traits can be situated within affective (e.g., fearlessness, shallow affect, callousness/lack of empathy), interpersonal (e.g., detached, manipulative, pathological lying), lifestyle (e.g., irresponsible, poor behavioral control), and antisocial domains (e.g., early behavioral problems, criminal versatility). Nevertheless, there is no consensus on which features should be included in the definition of psychopathy, let alone how important they are to this condition (see Patrick, Fowles, & Krueger, 2009, for a review). To illustrate, we highlight three points of discussion.¹ First, the question of whether overt criminal behavior or antisocial behavior more broadly is intrinsic to psychopathy is controversial; this possibility is suggested by the inclusion of explicit criminality among the PCL-R items. Some authors contend that criminality and antisocial behavior, rather than being intrinsic to psychopathy, are merely downstream consequences of psychopathy as conceptualized in terms of affective and interpersonal traits (e.g., Hare & Neumann, 2010; Skeem & Cooke, 2010).

Second, although many psychopathy researchers consider affective features to lie at the heart of psychopathy, there is no clear consensus on what constitutes this deficient affective experience. Some have argued for a broad, pervasive emotional deficit: “Like the color-blind person, the psychopath lacks an important element of experience—in this case, emotional experience” (Hare, 1993; pp. 129; see also Cleckley, 1941/1976). Others have offered alternatives to this supposition. For instance, Lykken (1995) posited a narrower fearlessness deficit and argued that psychopathic individuals would not engage in problematic behaviors if they were incapable of experiencing positive affect, whereas McCord and McCord (1964) contended that the lack of anxiety and emotional coldness in psychopathy arises from profound lovelessness, assigning a prominent role to positive emotions. Blair (2005) and others have argued that psychopathy stems from a broader social-emotional deficit, namely the capacity to experience empathy. Research pinpointing specific emotional deficits among psychopathic individuals has been decidedly mixed. An early startle eyeblink study (Patrick, Bradley, & Lang, 1993) found that psychopathic individuals showed deficient emotional responding to negative (e.g., fear-eliciting) stimuli, but normal emotional responding to positive (e.g., erotic) stimuli. Physiological hyporesponsivity in psychopathy is also most pronounced for aversive stimuli (Lorber, 2004). In contrast, a meta-analysis of emotion recognition tasks found that psychopathy was associated with poorer recognition not only of fear and sadness, but also of happiness and surprise (Dawel, O’Kearney, McKone, & Palermo, 2012). Owing to inconsistent research findings, a recent review contrasting a general emotional blunting with specific emotional deficits concluded that the “overall pattern of findings is not clearly consistent with any of the dominant theoretical perspectives of emotion processing in psychopathy” (Brook, Brieman, & Kosson, 2013, p. 979).

Third, although prominent authors view impulsivity as “one of the hallmarks of psychopathy” (Hare, 2003, p. 139) and “a cardinal feature of the [psychopathy] construct” (Hart & Dempster, 1997, p. 212), Cleckley (1941/1976) argued that psychopathic individuals are not driven by powerful impulses. Furthermore, the review by Poythress and Hall (2011)

questions the value of impulsivity—at least when viewed or measured as a unitary construct—as a marker for the assessment of psychopathy. The evidence reviewed here suggests that widely held and long-standing belief that <psychopaths are impulsive> must be reconsidered. (p. 132)

In sum, there is little consensus on the core characteristics of psychopathy (Lilienfeld, Watts, Francis Smith, Berg, & Latzman, 2015).

Network analyses, a relatively novel set of statistical techniques, may help to identify the core characteristics of psychological constructs (Borsboom & Cramer, 2013). The network approach was introduced as an alternative view on covariance between phenomena that departs from the largely untested and often faulty or overly stringent assumptions that underlie factor analytic models. The first, ontological, assumption is that if certain phenomena covary there necessarily exists an underlying latent factor (or class, or disorder) that explains the covariance. Although a latent factor may exist, the a priori assumption that there must be one is logically and empirically unfounded. The second, methodological, assumption in all factor analytic models (as well as in item response theory) is local independence, meaning that phenomena (i.e., symptoms or items) are not directly related. Hence, their covariance ostensibly approaches zero after taking into account their common latent factor(s). This assumption is implausible for many psychopathological syndromes. For instance, depressive symptoms may involve a negative feedback loop in which symptoms like lack of sleep contribute to other depressive symptoms, such as trouble concentrating and anhedonia. This may well also apply to psychopathy, for instance, early behavior problems may well explain juvenile delinquency. And it is also very conceivable that a lack of remorse or guilt explains a failure to accept responsibility. The third assumption is that correlations between scales or syndromes (or with, e.g., genetic markers or brain-structures) result from a direct relation between latent variables. These correlations may result as well or instead from more fine-grained bidirectional interactions at the level of signs and symptoms. Thus, the disadvantage of adhering to factor analytic models is that these possibilities are not taken into account.

With network analyses, one plots the features (e.g., symptoms) of the construct as nodes and their associations as edges within a network. The shortest path length between two nodes is the minimum number of edges required to travel from one node to another (if B can only be reached from A through C thus $A > C > B$, then the shortest path length between A and B is 2). The stronger the association between two nodes, the larger their edge weight (displayed in the plot by wider, more saturated edges). The strength of the associations as well as the position of the features in the network provide information about the importance of the feature to the network. A central, well-connected feature is likely to be particularly important. Importantly, network-analyses go beyond eye-balling the interitem correlation matrix (or factor loadings) by providing a growing set of indices that can be used to assess item

¹ In the present study, we can only assess the importance of symptoms tapped by the PCL-R. Therefore, we do not go into the discussion on features not directly covered by the PCL-R such as boldness or fearless dominance (Patrick et al., 2009; Vize et al., 2016). Also, we focus on features at the descriptive level (symptoms) rather than cognitive, physiological, or neural features.

covariance. The centrality of a feature to the network can be assessed in several ways (Costantini et al., 2015).² *Strength* reflects the overall connectivity with other features by summing the (absolute values of) weights of the feature's associations to other features. *Closeness* reflects the distance of a feature to other features by computing the mean weighted shortest path lengths to all other features.

Network analyses recently have been applied to several mental disorders, including major depressive disorder (Bringmann, Lemmens, Huibers, Borsboom, & Tuerlinckx, 2015), posttraumatic stress disorder (McNally, Robinaugh, Wu, Wang, Deserno, & Borsboom, 2015), and substance use disorder (Rhemtulla, Fried, Aggen, Tuerlinckx, Kendler, & Borsboom, 2016). Applying network analysis to *DSM-IV* symptoms of substance abuse in adult twins, for instance, has indicated that using a drug more than planned was among the most central substance abuse indicators (Rhemtulla et al., 2016). Network analyses similarly seem to be a promising tool for identifying and clarifying the core features of psychopathy and for helping to inform longstanding debates in the field.

At the same time, it is still unclear to what extent findings from network analyses are stable (within a sample, design, or instrument) and replicable (across samples, designs, or instruments). Network analyses on mental disorders have been based largely on single samples (Fried & Cramer, 2017). Moreover, network analyses on different instruments, designs or samples have not always converged on the same outcome. For example, cross-sectional analyses of Inventory of Depressive Symptomatology items in 3,463 currently depressed individuals indicated that energy loss and sadness were among the most central depression symptoms (Fried, Epskamp, Nesse, Tuerlinckx, & Borsboom, 2016). In contrast, network analyses on repeated administrations of the Beck Depression Inventory in 182 currently depressed individuals during the course of treatment indicated that past failure and loss of pleasure were the most central depression symptoms (Bringmann et al., 2015). Thus, the stability and replicability of network analyses of mental disorders is unclear and also a point of current controversy (see Forbes, Wright, Markon, & Krueger, 2017; but see reply by Borsboom et al., in press).

In the present study, we apply network analyses to the PCL-R, which is the most extensively researched measure of psychopathy (Hare, 2003). Our aim is to begin to identify the core characteristics of PCL-R operationalized psychopathy. The construct of psychopathy, like any construct (Cronbach & Meehl, 1955), cannot be reduced to a single instrument or ostensible "gold standard" (Skeem & Cooke, 2010)—because that instrument will almost inevitably omit some essential features of the construct (construct underrepresentation) and include others that are peripheral (construct overrepresentation). Nevertheless, the PCL-R is a well-validated and reasonably comprehensive measure that represents an excellent starting point for network analyses. To ascertain the stability of our PCL-R network analyses within the sample, we obtained large samples so we could determine whether indicators of centrality were robust across random subsets of persons within the same sample (Epskamp, Borsboom, & Fried, 2016). Further, to ascertain the replicability of PCL-R network results, we examined three samples: a North American sample of offenders ordered to either prison or substance abuse treatment ($n = 1,661$; hereafter called the NIMH sample; Poythress et al., 2010); a Dutch forensic

psychiatric sample of violent mentally disordered offenders ($n = 1,962$; hereafter called the 'TBS sample'); and a North American sample of offenders ordered to correctional institutes in Wisconsin ($n = 3,963$; hereafter called the Wisconsin sample), and we assessed the similarities by calculating the Spearman rho correlation between the rank orders of the centrality indices between the samples.

Method

This study was approved by the ethical committee of the Psychology Department of the University of Amsterdam (2017-CP-7957).

Sample Characteristics

Exclusion criteria. We excluded data from participants with substantial missing data on the PCL-R (i.e., more than two items missing on either of the two original two factors or more than five items missing in total); see the *Samples* section. For the included participants, the proportion of missing values for the PCL-R items averaged 0.5% (range: 0% to 4%) for the NIMH sample, 1.50% (range: 0% to 6%) for the TBS sample, and 1.66% (range: 0% to 13%) for the Wisconsin sample. Therefore, although the proportion of missing data was overall very low, there were substantial missing data for some items in some samples. For the calculation of the PCL-R total score, data from participants with one to four missing items were prorated following the guidelines of the PCL-R manual (Table 1; Hare, 2003). For the network analyses, correlations were calculated for all included participants based upon pairwise complete observations. Analyses on the three subsamples with only participants without any missing values produced similar findings to those with missing values (see Appendix I; all appendices are on <https://osf.io/4habq/>). In case of multiple PCL-R assessments for each participant, only the most recent assessment available was included.

Samples. The National Institute of Mental Health (NIMH) sample included offenders ($n = 1,661$; 82.5% male; $M_{\text{age}} = 31.0$ years, $SD = 6.6$) ordered to either prison or substance abuse treatment in Florida, Nevada, Oregon, Utah, and Texas. The PCL-R was administered by trained research assistants for research purposes. The recruitment strategy favored men (80%). Exclusion criteria included (a) estimated IQ below 70, (b) non-English-speaking, and (c) residing in a prison mental health unit or receiving medication for active symptoms of psychosis (see Poythress et al., 2010, for further details regarding participant recruitment). We excluded 102 participants because of excessive missing PCL-R items (see the *Exclusion Criteria* section). The final NIMH sample consisted of 1559 offenders (83.6% male; $M_{\text{age}} = 31.0$ years, $SD = 6.5$).

The TBS sample ($n = 1,962$; 73.0% male; $M_{\text{age}} = 38.8$ years, $SD = 10.0$) consisted of violent mentally disordered offenders under mandatory inpatient treatment in the Netherlands (TBS or

² Another often used index is betweenness, which indicates how often a feature functions as a hub in the network, by counting how often the feature lies on the shortest connection of two other features. Because we found this index to be of less relevance to the PCL-R network, and there were indications that this index was less stable, we focus on strength and closeness.

Table 1
PCL-R Items With Their Abbreviation as Used in the Network Plots

Item number	Abbreviation	Item label	PCL-R facet (Hare, 2003)
1	GLI	Glibness/superficial charm	Interpersonal
2	SEL	Grandiose sense of self-worth	Interpersonal
3	STI	Need for stimulation	Lifestyle
4	LIE	Pathological lying	Interpersonal
5	CON	Conning/manipulative	Interpersonal
6	GUI	Lack of remorse or guilt	Affect
7	AFF	Shallow affect	Affect
8	EMP	Callous/lack of empathy	Affect
9	PAR	Parasitic lifestyle	Lifestyle
10	BEV	Poor behavioral controls	Antisocial
11	SEX	Promiscuous sexual behavior	—
12	PRO	Early behavioral problems	Antisocial
13	GOA	Lack of realistic, long-term goals	Lifestyle
14	IMP	Impulsivity	Lifestyle
15	IRR	Irresponsibility	Lifestyle
16	RES	Failure to accept responsibility	Affect
17	SHO	Many short-term marital relations	—
18	DEL	Juvenile delinquency	Antisocial
19	REL	Revocation of conditional release	Antisocial
20	CRI	Criminal versatility	Antisocial

ter beschikking stelling, which can be translated to disposal to be treated on behalf of the state; Philipse, 2005). TBS is imposed by court on high-risk offenders who have committed violent crimes that were determined to result from psychopathology, leading to judgments of diminished responsibility. The duration of the TBS-order is indeterminate—lasting as long as the offender is considered to be high-risk—but averages about 8 years. The PCL-R was administered by trained clinicians as part of a mandatory risk assessment test battery. We obtained PCL-R data from 12 of the 14 treatment facilities that collect mandatory PCL-R data (<http://www.efp.nl/projecten/ldr-tbs>). We excluded 25 participants because of excessive missing PCL-R items (see “Exclusion criteria”). The final TBS sample consisted of $n = 1,937$ offenders (73.4% male; $M_{\text{age}} = 38.8$ years, $SD = 10.0$).

The Wisconsin sample ($n = 3,963$; 63.3% male; $M_{\text{age}} = 30.3$ years, $SD = 7.0$) consisted of offenders from state prisons in the U.S. state of Wisconsin. The sample has been collected over many years, with the PCL-R being scored by trained research assistants (e.g., Baskin-Sommers & Newman, 2014; Newman, MacCoon, Vaughan, & Sadeh, 2005). During recruitment, participants with low IQ scores (<70) and with significant mental illness (particularly psychosis) were excluded. We excluded nine participants because of excessive missing PCL-R items (see the *Exclusion Criteria* section). The final Wisconsin sample consisted of $n = 3,954$ offenders (63.3% male; $M_{\text{age}} = 30.3$ years, $SD = 7.0$).

Measures

The PCL-R (Hare, 2003) consists of a checklist of 20 items (see Table 1). In the TBS sample, the authorized Dutch translation was used (Vertommen, Verheul, de Ruiter, & Hildebrand, 2002). Based upon an extensive interview and collateral file information, trained raters score each item as 0 = absent, 1 = maybe or partly present, or 2 = definitely present. Thus, the PCL-R sum score lies on a continuum ranging from 0 to 40. The mean PCL-R score was

22.54 ($SD = 7.47$) for the NIMH sample, 23.3 ($SD = 7.0$) for the Wisconsin sample, and 20.9 ($SD = 7.3$) for the TBS sample. A cutoff score of 26 (Europe) or 30 (United States) has been used for a clinical psychopathy diagnosis. The proportion of the sample scoring above the country-specific cutoff (Hare, Clark, Grann, & Thornton, 2000) was 28.03% (TBS), 21.93% (Wisconsin), and 19.69% (NIMH). Cronbach’s alpha was high for the total score in all three samples: $\alpha = .82$ for the NIMH sample, $\alpha = .81$ for Wisconsin sample, and $\alpha = .83$ for the TBS sample. In the NIMH sample, the interrater reliability (ICC) for the total PCL-R score based upon $n = 51$ was .88 (Poythress et al., 2010). The ICC of PCL-R total scores by diagnostic staff in Dutch TBS clinics based upon $n = 16$ was found to be .76 (Nentjes, Bernstein, Meijer, Arntz, & Wiers, 2016). The ICC in the Wisconsin sample was very high: .99 based upon $n = 16$ (Baskin-Sommers & Newman, 2014), and .96 based upon $n = 101$ (Newman et al., 2005).

Table 1 shows the facet labels corresponding to the PCL-R 4-facet structure (Hare, 2003). There is still disagreement regarding the optimal factor structure of the PCL-R (see Footnote 3). Early factor analytic work on the measure revealed a two-factor structure, with Factor 1 encompassing affective-interpersonal features (selfish, callous, remorseless use of others: Items 1, 2, 4, 5, 6, 7, 8, and 16) and Factor 2 encompassing chronic antisocial lifestyle (chronically unstable and antisocial lifestyle: 3, 9, 10, 12, 13, 14, 15, 18, and 19; Hare, 1991). Cooke and Michie (2001) later proposed a three-factor structure that essentially (a) splits Factor 1 into an affective factor (deficient affective experience; Items 6, 7, 8, and 16) and an interpersonal factor (deceitful interpersonal style; Items 1, 2, 4, and 5), and (b) excludes overt criminal items from Factor 2 (now labeled the lifestyle factor; impulsive and irresponsible behavioral style; Items 3, 9, 13, 14, and 15). Finally, like the three-factor structure, the four-factor structure (Hare, 2003), also has identical affective, interpersonal, and lifestyle factors, but reintroduces

duces the items primarily focused on prior criminal activities into an antisocial factor (Items 10, 12, 18, 19, and 20).

Statistical Analyses

Our network analyses were based upon polychoric correlations, which allow for estimation of two observed ordinal variables that have presumed theoretical normal distributions, as to account for the limited range in PCL-R item responses (response options: 0, 1, and 2). We elected not to examine the regularized partial correlation network (glasso). The glasso method downsizes indirect relations among PCL-R items (e.g., the correlation between two PCL-R items disappears if a third PCL-R item that strongly connects to those PCL-R items is taken into account), often with the aim of finding unique possible casual relations. Our focus was on centrality, irrespective of whether centrality would be a result of direct or of indirect relations. Moreover, given the covariance between PCL-R items, the meaning of each PCL-R item typically becomes unclear once the variance shared with all other PCL-R items has been controlled (Lynam, Hoyle, & Newman, 2006). Although partialing can rule out spurious connections, it adapts an extreme form of control that may in fact create more inferential problems than the one it attempts to solve. Indeed, Appendix II shows that partialing led to several, implausible and unexpected, negative associations between partialled PCL-R items, indicating that partialing created spurious connections due to common effects.

Network plots. The PCL-R network was constructed with the qgraph package (Epskamp, Cramer, Waldorp, Schmittmann, & Borsboom, 2012) using the statistical application R (version 3.2.4; R Core Team, 2016). In visualization of the network, nodes represent the PCL-R items and the thickness and the color of edges represents their association strength and valence (red: negative association, black: positive association), respectively. The layout of the network was based on Fruchterman and Reingold (1991)'s algorithm, which places more influential nodes central to the network and stronger connected nodes closer together in the network.

Centrality indices. In addition to network visualization, we used the qgraph package to provide two indices of centrality: The overall connectivity with other nodes (strength) as well as the average shortest path lengths to all other nodes (closeness) (Opsahl, Agneessens, & Skvoretz, 2010; Epskamp et al., 2016).³

Within-sample stability. We assessed the stability of the centrality indices. Stability in the network perspective refers to resistance to change: Is the network robust to the dropping of selected participants from the analyses? Stability across participants indicates that a somewhat different sample would not change the rank order of the centrality of the nodes. The stability of the centrality order can be determined with subset bootstrapping using the R-package entitled *bootnet* (Epskamp, 2015), providing the centrality of items over a wide range of sampled participants. Moreover, stability can be quantified by the correlation stability coefficient or conditional stimulus (CS)-coefficient. Specifically, CS ($\text{cor} = 0.70$) reflects the proportion of participants that can be dropped to retain with 95% probability that the correlation between the centrality based on the entire sample and that of the bootstrapped subsamples is at least 0.70 (representing a very large effect). Based upon simulation studies, Epskamp et al.

(2016) recommended that CS ($\text{cor} = 0.70$) should be at least 0.25 and preferably greater than 0.50 to interpret centrality differences.

Centrality differences. The network plots and centrality indices indicate the centrality of the PCL-R items. We further used *bootnet* to conduct bootstrapped difference tests on the centrality indices to examine whether there are significant differences in centrality. Although (a) this approach yields numerous comparisons and (b) there is presently no definitive solution for multiple testing in network analyses (Epskamp et al., 2016), these difference tests provide additional information on differences in centrality of the PCL-R items.

Replicability across samples. Rather than examining the correspondence of the absolute centrality positions of the PCL-R items across the samples (Forbes et al., 2017), we examined whether there was correspondence in their relative centrality positioning (Borsboom et al., *in press*), because absolute positioning is a too strict measure of replicability. Whether a particular PCL-R item is ranked 17th, 18th, or 19th may not matter that much, but if this would be the observed centrality rank across three samples, it would be clear that the PCL-R item is replicably low in centrality. To assess replicability across samples, we calculated the Spearman rho correlation of rank-ordered centrality between samples.

Results

Network Plots

Figure 1 displays the correlational structure of the PCL-R items in the TBS sample (left, Figure 1a), the NIMH sample (middle, Figure 1b), and the Wisconsin sample (right, Figure 1c). The strength of the relations between PCL-R items translates into the thickness of the edges between them and the distance that they are plotted from each other. Glibness/superficial charm (GLI) and grandiose sense of self-worth (SEL), for instance, show strong interrelations in all three samples. Groups of PCL-R items cluster together in the densely connected network, except for many short-term marital relationships (SHO), which appears peripheral to the PCL-R psychopathy network in all three samples.

Centrality Indices

Figure 2A–C depicts the closeness and strength of the 20 PCL-R items for each of the three samples. In the NIMH sample, callous/lack of empathy (EMP) was the most central item, followed by shallow affect (AFF). SHO and revocation of conditional release (REL) were

³ Our network analyses do not serve nor allow to test the best fit of the two-, three-, or four-factor model. Descriptively though, a number of observations are of interest to the factor analytic work. First, the network plots show that the clustering of items generally fit with the two-, three-, or four-factor models. Second, a noteworthy exception is *Lack of realistic long term goals*, which does not cluster with the other items of the factor it is expected to load on (which holds for both the two-, three-, or four-factor model) and appeared rather peripheral to the PCL-R network in all three samples. Third, while *Criminal versatility* was not included in the original Factor2, it clusters with the other antisocial items in the Wisconsin sample and with the antisocial and lifestyle items in the Dutch sample. Fourth, the clustering of items according to two-, three-, or four-factor analytic solution was least clear in the NIMH sample (where there appeared a clear affective-interpersonal clustering of items, but such clustering was less apparent for the behavioral-lifestyle items).

Figure 1. a–c: Network of the Psychopathy Checklist—Revised (PCL-R; Hare, 2003) based upon polychoric correlations between PCL-R items for the three samples (1a: Dutch sample; 1b: NIMH sample; 1c: Wisconsin sample). Nodes represent the PCL-R items: GLI = glibness/superficial charm; SEL = grandiose sense of self-worth; STI = need for stimulation; LIE = pathological lying; CON = conning/manipulative; GUI = lack of remorse or guilt; AFF = shallow affect; EMP = callous/lack of empathy; PAR = parasitic lifestyle; BEV = poor behavioral controls; SEX = promiscuous sexual behavior; PRO = early behavioral problems; GOA = lack of realistic, long-term goals; IMP = impulsivity; IRR = irresponsibility; RES = failure to accept responsibility; SHO = many short-term marital relations; DEL = juvenile delinquency; REL = revocation of conditional release; CRI = criminal versatility. Positive correlations plotted in green (online figure) / grey (print figure). See the online article for the color version of this figure.

the least central items in the NIMH sample. In the Wisconsin sample, EMP was clearly the most central item. SHO, REL, and GOA were the least central in the Wisconsin sample. In contrast, in the Dutch sample, parasitic lifestyle (PAR) and irresponsibility (IRR) were the most central items, promiscuous sexual behavior (SEX), SHO, and—perhaps surprisingly—AFF, were the least central items in the Dutch sample.

Within-Sample Stability

Figure 3A–C depict the stability of the centrality indices, by plotting centrality over increasingly smaller bootstrapped sub-samples for each of the three samples. Closeness and strength display high stability in all three samples for a wide range of

Figure 2. Centrality closeness and strengths of the Psychopathy Checklist—Revised items in the three samples (2a: Dutch sample; 2b: NIMH sample; 2c: Wisconsin sample).

Figure 3. A–C: Bootstrapped differences in centrality strength and closeness in the three samples (3A: Dutch sample; 3B: NIMH sample; 3C: Wisconsin sample). See the online article for the color version of this figure.

increasingly smaller bootstrapped subsamples. These impressions were confirmed by the correlation stability coefficients. CS ($\text{cor} = 0.70$) for strength was .89 for the Dutch sample, .91 for the NIMH sample, and .92 for the Wisconsin sample. CS ($\text{cor} = 0.70$) for closeness was .87 for the Dutch sample, .89 for the NIMH sample, and .92 for the Wisconsin sample. These findings suggest that the centrality indices were highly stable and well over the recommended value of .50, which allows for interpreting differences in centrality.

Centrality Differences

Figure 4A–C displays the results of the bootstrapped difference tests in centrality for all three samples. Gray boxes indicate that the PCL-R items do not significantly differ in centrality; black boxes indicate that the PCL-R items differ significantly in centrality. The values in the white box on the diagonal depict the centrality values. We highlight notable significant differences in centrality. For the NIMH sample, the difference tests confirm that EMP and AFF were significantly more central than most of the other PCL-R items, and that REL and SHO were significantly less central than most of the other PCL-R items. For the Wisconsin sample, the difference tests show that CAL was significantly more central than all other PCL-R items. They also confirm that SHO, REL, and GOA were significantly less central than most other PCL-R items in the Wisconsin sample. For the Dutch sample, the difference tests confirm the high centrality of IRR, as it was significantly more central than nearly all other PCL-R items. They also confirm the low centrality of SHO, AFF, and SEX in that sample, as they were significantly less central than many other PCL-R items.

Replicability Across Samples

The centrality rankings of the PCL-R items of the NIMH and the Wisconsin sample show important similarities. EMP was the most central item in both samples, for both closeness and for strength, and largely irrespective of the number of persons taken into the analyses. In both samples, SHO and REL were peripheral to

the PCL-R network. These observations are also apparent from the strong Spearman rho correlation between the rank orders of the NIMH and the Wisconsin sample for closeness, $\rho = .64$, and strength, $\rho = .60$.

The centrality ranking of the PCL-R items of the U.S. samples differs in important respects from that of the TBS sample. Most notably, PAR and particularly IRR were most central in the TBS sample whereas they showed modest to low centrality in the U.S. samples. The Spearman rho correlation between the rank orders of the TBS and the Wisconsin sample were moderate (for closeness, $\rho = .38$, and strength, $\rho = .42$), and weak for the TBS and the NIMH sample (for closeness, $\rho = .11$, and strength, $\rho = .15$).

Discussion

This study was an effort to shed light on an actively debated and theoretically important question: What are the most central features of psychopathy, at least as operationalized by the most widely used measure of this construct? Using network analyses on three large samples, we examined the centrality of PCL-R features of psychopathy, as well as their stability within each sample, and the replicability across samples. We found a densely connected network, with EMP being stably most central to the PCL-R network in the U.S. offender samples; SHO was stably peripheral to the PCL-R network in all three samples. The network results did not generalize to the Dutch forensic psychiatric sample, wherein EMP was also fairly central but IRR and PAR were even more central.

The Centrality of Callous/Lack of Empathy to PCL-R Psychopathy

EMP was moderately central in the Dutch forensic psychiatric sample, and exhibited the highest centrality in the two U.S. offenders samples. The importance of a lack of empathy fits with early clinical descriptions of psychopathy. For instance, Gough (1948) regarded a deficiency in the ability to mentally “take the place” of other individuals as the core feature of psychopathy, and

A. Dutch Sample: Closeness

Dutch Sample: Strength

Figure 4. A–C: Bootstrapped differences in centrality closeness and strength for the three samples (4A: Dutch sample; 4B: NIMH sample; 4C: Wisconsin sample). Gray boxes indicate nonsignificant differences, black boxes indicate significant differences. Centrality closeness and strength values are plotted on the diagonal.

B. NIMH Sample: Closeness

NIMH Sample: Strength

Figure 4 (continued)

C. Wisconsin Sample: Closeness

Wisconsin Sample: Strength

Figure 4 (continued)

he premised his influential Socialization scale of the California Psychological Inventory on this model. Later, McCord and McCord (1964), focusing more on affective than on cognitive empathy, viewed a lack of social emotions—specifically lovelessness (inability to form deep attachments) and guiltlessness (absence of remorse)—as the essence of psychopathy.

Our finding also dovetails with the work by Frick and colleagues (e.g., Frick, Ray, Thornton, & Kahn, 2014), who view callous-unemotional (CU) traits, including lack of empathy and guilt, to be of critical importance in delineating youth psychopathy, as well with proponents of the triarchic model of psychopathy, who accord a central role to meanness, coldheartedness, or cognate constructs in the conceptualization of psychopathy (Berg, Hecht, Latzman, & Lilienfeld, 2015; Patrick et al., 2009). Likewise, DSM-5 (American Psychiatric Association, 2013) included a specifier to define a subgroup of youth with conduct disorder who display limited prosocial emotions, including weak empathy and guilt. Furthermore, the central role of EMP accords with Blair's (1995) Violence Inhibition Model, which proposes that a reduced emotional reaction to distress in others (i.e., less victim empathy) predisposes to psychopathy-like traits.

Interestingly, the results of these network analyses converge with other methods of evaluating the centrality of various features of psychopathy. A number of recent studies have asked mental health experts as well as laypersons in North America and Europe to judge items from the Comprehensive Assessment of Psychopathic Personality (CAPP; Cooke, Hart, Logan, & Michie, 2012) on prototypicality for psychopathy (Flórez et al., 2015; Hoff, Rypdal, Mykletun, & Cooke, 2012; Kreis, Cooke, Michie, Hoff, & Logan, 2012; Sörman et al., 2014). The CAPP assesses 33 features, including features that closely correspond with PCL-R items (e.g., lacks emotional depth and lacks remorse), but also features that were not (fully) captured by the PCL-R (e.g., aggressive or lacks anxiety). Kreis et al. (2012) found that 30 out of the 33 CAPP items were rated at least moderately prototypical, and that 25 out of the 33 CAPP items were rated as highly prototypical. So although most CAPP items were deemed prototypical for psychopathy and their average prototypicality rating were very close to one another, the most prototypical items were lacks remorse, unempathic, self-centered, manipulative, and lacks emotional depth. These items—particularly lacks remorse and unempathic—broadly corroborate the high centrality of EMP in the NIMH and Wisconsin sample, and its close connection with lack of remorse or guilt.

Moreover, the network findings also align with the findings obtained with the elemental view on psychopathy from a basic trait perspective (Miller & Lynam, 2015). Studies drawing on the five factor model of personality, using both expert ratings and the relations between psychopathy instruments and five factor personality measures, have shown that psychopathy can be characterized by facets of low agreeableness (e.g., low altruism), low conscientiousness (e.g., low self-discipline), high neuroticism (e.g., high hostility), low (e.g., low warmth) and high (e.g., high excitement seeking) extraversion (e.g., Miller, Lynam, Widiger, & Leukefeld, 2001; Lynam & Widiger, 2001; for a recent review, see Lynam & Miller, 2015). Interestingly, the five-factor model of psychopathy suggests that low agreeableness (encompassing EMP) is a central element of psychopathy. Such findings, which derive from very different research designs, support the generalizability of affective-

interpersonal features, and callousness and deficient empathy, in particular, as central components of psychopathy.

Affect, Criminality, and Impulsivity: Core Characteristics of Psychopathy?

Whatever their theoretical differences, virtually all psychopathy researchers agree that psychopathy is characterized by abnormal emotional processing, but there is no consensus on the nature of that affective deficiency (Brook et al., 2013). Our findings speak to the issue of whether the affective deficit is general or largely specific. The PCL-R contains several items that relate to abnormal affect, some of which suggest specific deficits (lack of remorse or guilt, EMP) and others of which reflect a more global deficit (AFF). AFF was marked by high centrality only in the NIMH sample, and it showed low centrality in the two other samples. In contrast, lack of remorse or guilt showed modest centrality in all three samples, and EMP showed modest to high centrality in all three samples. Awaiting a more comprehensive analysis across the full spectrum of positive and negative emotions, our findings provisionally speak to a specific rather than a global affective deficit in psychopathy, broadly corroborating psychometric findings that psychopathy is characterized not by poverty in affect more broadly (cf. Cleckley, 1941/1976) but by deficits in social detachment more specifically (Lilienfeld & Andrews, 1996; Patrick et al., 2009).

The question of whether antisocial and criminal behavior are central to psychopathy remains heavily contested (see Skeem & Cooke, 2010; Hare & Neumann, 2010, for diverging viewpoints). Although limited to the PCL-R, our network analyses sheds some light on this issue. We found that juvenile delinquency and revocation of conditional release (both indices of criminal behavior) were relatively peripheral to the network. In contrast, criminal versatility, early behavioral problems and poor behavioral controls (the latter two of which mix indices of criminal and antisocial behavior) were moderately central. None of the items, however, was highly central to PCL-R psychopathy, calling into question the assertion that nonspecific antisocial behavior is pivotal to the conceptualization and operationalization of psychopathy (see also Lykken, 1995).

Our findings may shed light on the importance of impulsivity for conceptualizing PCL-R defined psychopathy. Across samples, impulsivity did not appear among the most central symptoms. This finding fits with the conclusion of Poythress and Hall (2011) that the "blunt assertion that 'psychopaths are impulsive' is no longer defensible" (p. 120). At the same time, these authors argued that there may be links between psychopathy and impulsivity if both concepts are further refined. Impulsivity is a very heterogeneous concept, leading Whiteside and Lynam (2001, p. 677) to distinguish among (lack of) premeditation ("the tendency to delay action in favor of careful thinking and planning"), urgency ("the tendency to commit rash or regrettable actions as a result of intense negative affect"), sensation seeking ("the tendency to seek excitement and adventure"), and (lack of) perseverance ("one's ability to remain with a task until completion and avoid boredom"). Notably, these differing subdimensions or pathways of impulsivity bear substantially different implications for different forms of psychopathology (Berg, Latzman, Bliwise, & Lilienfeld, 2015). Future research should examine whether these more refined subcomponents of

impulsivity play a more prominent role in psychopathy, or in some forms of (e.g., secondary) psychopathy (Karpman, 1948). Anestis, Anestis, and Joiner (2009), for instance, found preliminary evidence that secondary psychopathy was positively related primarily to the tendency to act rashly when distressed (negative urgency).

Network Analyses: Stability Within Samples and Replicability Across Samples

Although our findings were highly stable within each sample, our study highlights the importance of examining replicability of network analyses across samples (Fried & Cramer, 2017), as we found sizable and potentially important differences between the U.S. samples and the Dutch sample. EMP was the most central item in the two U.S. samples, whereas IRR and PAR were most central in the Dutch sample. We can consider at least three possible reasons for this divergence.

First, in the U.S. samples, the PCL-R was assessed for research purposes by well-trained research assistants, whereas the PCL-R in the TBS sample was assessed by clinicians for the purpose of risk assessment. Although the reliability of the PCL-R, particularly Factor 1 items, in North America and Europe tends to be lower in field than in research settings (Jeandarme et al., 2017; Miller, Kimonis, Otto, Kline, & Wasserman, 2012), there were no marked differences in the reported interrater reliability statistics across samples for the total PCL-R score in the present study—with the caveat that interrater reliability estimates were based on small subsamples. Furthermore, self-selection in the U.S. sample is not evident from a comparison of the average PCL-R score of our U.S. samples that is very similar to that obtained in previous U.S. samples (range of average score was 20–24 across seven U.S. prison and forensic psychiatric samples; Hare et al., 2000). Finally, although the potential consequences of the PCL-R assessment differed greatly between the U.S. samples (zero) and the Dutch sample (impact on the nature and duration of the mandatory treatment, providing incentives to downplay psychopathic tendencies), it is important to bear in mind that the PCL-R assessment is based not only on an interview, but also on extensive file review.

Second, the U.S. samples consisted mostly of non-mentally ill prisoners, whereas the Dutch sample consisted of forensic psychiatric patients. To illustrate, the prevalence of psychotic disorders in the Dutch sample has been estimated to be as high as 39% (Nieuwenhuizen et al., 2011). In the U.S. prison samples, a prevalence of 3–7% psychotic disorders can be expected (Fazel & Danesh, 2002), and given the applied exclusion criteria, that prevalence is expected to be even lower in the U.S. samples of the current study. To examine whether major psychopathology could explain the U.S.-Dutch differences, we extracted a subsample of the Dutch sample, excluding those participants with indications of current or past severe psychopathology. PAR and IRR were still most central in this subsample ($n = 357$; see Appendix III), rendering it less likely that severe psychopathology can explain the differences between the samples.

Third, there are geographic differences between the U.S. (NIMH and Wisconsin) and the Dutch (TBS) samples. The question of whether PCL-R psychopathy is similar in the Netherlands and the U.S. entails two possibilities (Skeem, Edens, Camp, & Colwell, 2004). The psychopathy construct itself could differ, such that genetic and sociocultural factors give rise to EMP being central to

the phenotypic expression of psychopathy in the U.S., and IRR and PAR being central to the phenotypic expression of psychopathy in the Netherlands. This possibility, however, would be at odds with the fact that also Dutch clinicians view affective-interpersonal traits, most particularly a callous lack of empathy, as being prototypical to psychopathy (Verschuere & te Kaat, 2017). Alternatively, the psychopathy measurement could differ between the countries. At least some of the PCL-R items seem to have country-specific content. For instance, it is quite uncommon in the Netherlands to engage in marriage or registered partnership more than twice before the age of 30 (required for a positive score on SHO) and consequently the Dutch sample had a moderately lower mean score for SHO compared with the U.S. samples, with the vast majority (78%) scoring 0 (see Appendix IV). To the extent that the cross-country differences are replicable, they might require us to reconsider differences in the scoring or thresholds of these items across countries. Moreover, replicable differences would reopen the debate of whether the PCL-R as currently implemented in Dutch forensic psychiatric settings captures core psychopathic traits, or rather a deviant, antisocial lifestyle characterized by IRR and PAR (see, e.g., Skeem & Mulvey, 2001; Spreen, Ter Horst, Lutjehuis, & Brand, 2008).

Limitations

Our study was marked by several limitations. First, although our choice of the PCL-R can be justified given that it is the most extensively validated measure of psychopathy, arguably the most important limitation of the present study is that it was restricted to a single instrument. This limitation raises the risk of mono-measurement bias (see, e.g., Skeem & Cooke, 2010). It is therefore important not to commit the “error of reification” by assuming that our findings on the PCL-R necessarily bear on all operationalizations of the construct of psychopathy. As noted later, for example, the PCL-R does not explicitly assess fearlessness and other features that some (e.g., Lykken, 1995) regard as central to psychopathy. Also, the typical PCL-R assessment may bring about local dependencies between PCL-R features, creating artificial correlations. For instance, the same (e.g., unempathic) behavior, such as an especially callous crime, may bear on the ratings of several PCL-R items (Cooke & Michie, 2001). Likewise, the centrality of EMP could be a result of a negative halo-effect (Thorndike, 1920)—that is, a cognitive bias in which the rating of one (undesirable) feature influences the judgment of other features (e.g., one believes that an unattractive person is also unintelligent. To the extent that clinicians consider EMP to be crucial to psychopathy (as indicated by prototypicality ratings), the rating of some other PCL-R items may be influenced by their rating of this feature. More broadly, if clinicians form a global negative impression of an interviewee largely on the basis of his or her callousness and lack of empathy (“This interviewee does not seem to be a nice person”), this impression could inadvertently shape their ratings of other PCL-R items. It will be crucial to extend our observations to other well-validated psychopathy instruments (e.g., self-report measures; Lilienfeld & Fowler, 2006), and to the combinations of instruments (e.g., combine PCL-R items with CAPP items; Cooke et al., 2012).

Second, the use of other measures will help to clarify the importance of psychopathy features that are poorly represented in

the PCL-R. Although some authors have placed great emphasis on fearlessness and/or boldness in the conceptualization of psychopathy (Lykken, 1995; Patrick et al., 2009), others have argued that these features should be de-emphasized or even “be dropped from psychopathy” altogether (Vize, Lynam, Lamkin, Miller, & Pardini, 2016; p. 1). Although there is some evidence that aspects of boldness are viewed by clinicians and researchers as prototypical features of psychopathy (Berg, Lilienfeld, & Sellbom, 2017; Sörman et al., 2016), network analysis on psychopathy instruments that measure boldness, such as the Psychopathic Personality Inventory—Revised (Lilienfeld & Widows, 2005), may shed further light on the relevance of this trait to the psychopathy construct.

Third, a deeper conceptual question is whether it is appropriate to apply network analyses to the concept of psychopathy and to personality disorders more broadly. Borsboom (2017) argued that network theory may be more likely to serve as an explanatory model for some disorders (e.g., posttraumatic stress disorder [PTSD]) than for other, particularly slowly developing, disorders; the latter conditions may well include personality disorders, such as psychopathy. Note, however, that the concern applies to network theory, rather than to network analyses. According to the network theory of mental disorders (Borsboom, 2017) no common cause explains covariance of psychiatric signs and symptoms; signs and symptoms instead cause each other (for hybrid models that combine network and latent variables models, see Epskamp, Rhemtulla, & Borsboom, 2017). For PTSD, for instance, anger and hypervigilance might lead to poor sleep, which in turn makes it difficult to concentrate at work. Network analyses help to elucidate such possible causal effects. In the current study, we used network analyses to reveal which features are most central to PCL-R defined psychopathy. Whether network theory, which typically speaks to rapidly developing feature-feature relations (e.g., poor sleep causing concentration difficulties), also applies to feature-feature relations that are likely to unfold more slowly over time (e.g., EMP causing early behavioral problems) remains to be investigated. Longitudinal designs with repeated measurements of psychopathy symptoms allow for testing this possibility.

Conclusion

Applying network analysis to the PCL-R revealed several important findings, including the identification of callous/lack of empathy as a central feature of psychopathy in all three samples, especially the American sample. At the same time, our findings are restricted to one instrument—the PCL-R—and point to noteworthy differences between the U.S. and the Dutch samples, especially in the importance of irresponsibility and PAR in the latter. Extending network analyses to different measures, samples, and cultures should shed further light on the core characteristics of psychopathy and perhaps ultimately on the unresolved question of what psychopathy is.

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders, 5th ed. (DSM-5)*. Washington, DC: Author.
- Anestis, M. D., Anestis, J. C., & Joiner, T. E. (2009). Affective considerations in antisocial behavior: An examination of negative urgency in primary and secondary psychopathy. *Personality and Individual Differences, 47*, 668–670. <http://dx.doi.org/10.1016/j.paid.2009.05.013>
- Baskin-Sommers, A. R., & Newman, J. P. (2014). Psychopathic and externalizing offenders display dissociable dysfunctions when responding to facial affect. *Personality Disorders, 5*, 369–379. <http://dx.doi.org/10.1037/per0000077>
- Berg, J. M., Hecht, L. K., Latzman, R. D., & Lilienfeld, S. O. (2015). Examining the correlates of the coldheartedness factor of the Psychopathic Personality Inventory—Revised. *Psychological Assessment, 27*, 1494–1499. <http://dx.doi.org/10.1037/pas0000129>
- Berg, J. M., Latzman, R. D., Bliwise, N. G., & Lilienfeld, S. O. (2015). Parsing the heterogeneity of impulsivity: A meta-analytic review of the behavioral implications of the UPPS for psychopathology. *Psychological Assessment, 27*, 1129–1146. <http://dx.doi.org/10.1037/pas0000111>
- Berg, J. M., Lilienfeld, S. O., & Sellbom, M. (2017). The role of boldness in psychopathy: A study of academic and clinical perceptions. *Personality Disorders*. Advance online publication. <http://dx.doi.org/10.1037/per0000247>
- Blair, R. J. (1995). A cognitive developmental approach to mortality: Investigating the psychopath. *Cognition, 57*, 1–29. [http://dx.doi.org/10.1016/0010-0277\(95\)00676-P](http://dx.doi.org/10.1016/0010-0277(95)00676-P)
- Blair, R. J. R. (2005). Responding to the emotions of others: Dissociating forms of empathy through the study of typical and psychiatric populations. *Consciousness and Cognition: An International Journal, 14*, 698–718. <http://dx.doi.org/10.1016/j.concog.2005.06.004>
- Borsboom, D. (2017). A network theory of mental disorders. *World Psychiatry, 16*, 5–13. <http://dx.doi.org/10.1002/wps.20375>
- Borsboom, D., & Cramer, A. O. (2013). Network analysis: An integrative approach to the structure of psychopathology. *Annual Review of Clinical Psychology, 9*, 91–121. <http://dx.doi.org/10.1146/annurev-clinpsy-050212-185608>
- Borsboom, D., Fried, E., Epskamp, E., Waldorp, L. J., van Borkulo, C. D., Van der Maas, H. L. J., & Cramer, A. O. J. (in press). Replicability of psychopathology networks: The right question but the wrong answer. A comment on “Evidence that psychopathology symptom networks have limited replicability” by Forbes, Wright, Markon, and Krueger. *Journal of Abnormal Psychology*.
- Bringmann, L. F., Lemmens, L. H., Huibers, M. J., Borsboom, D., & Tuerlinckx, F. (2015). Revealing the dynamic network structure of the Beck Depression Inventory-II. *Psychological Medicine, 45*, 747–757. <http://dx.doi.org/10.1017/S0033291714001809>
- Brook, M., Brieman, C. L., & Kosson, D. S. (2013). Emotion processing in Psychopathy Checklist-assessed psychopathy: A review of the literature. *Clinical Psychology Review, 33*, 979–995. <http://dx.doi.org/10.1016/j.cpr.2013.07.008>
- Cleckley, H. (1941/1976). *The mask of sanity*. St. Louis, MO: Mosby.
- Cooke, D. J., Hart, S. D., Logan, C., & Michie, C. (2012). Explicating the construct of psychopathy: Development and validation of a conceptual model, the Comprehensive Assessment of Psychopathic Personality (CAPP). *The International Journal of Forensic Mental Health, 11*, 242–252. <http://dx.doi.org/10.1080/14999013.2012.746759>
- Cooke, D. J., & Michie, C. (2001). Refining the construct of psychopathy: Towards a hierarchical model. *Psychological Assessment, 13*, 171–188. <http://dx.doi.org/10.1037/1040-3590.13.2.171>
- Costantini, G., Epskamp, S., Borsboom, D., Perugini, M., Möttus, R., Waldorp, L. J., & Cramer, A. (2015). State of the aRt personality research: A tutorial on network analysis of personality data in R. *Journal of Research in Personality, 54*, 13–29. <http://dx.doi.org/10.1016/j.jrp.2014.07.003>
- Cronbach, L. J., & Meehl, P. E. (1955). Construct validity in psychological tests. *Psychological Bulletin, 52*, 281–302. <http://dx.doi.org/10.1037/h0040957>
- Dawel, A., O’Kearney, R., McKone, E., & Palermo, R. (2012). Not just fear and sadness: Meta-analytic evidence of pervasive emotion recognition deficits for facial and vocal expressions in psychopathy. *Neuro-*

- science and Biobehavioral Reviews*, 36, 2288–2304. <http://dx.doi.org/10.1016/j.neubiorev.2012.08.006>
- DeMatteo, D., Edens, J. F., Galloway, M., Cox, J., Smith, S. T., & Formon, D. (2014). The role and reliability of the Psychopathy Checklist-Revised in U.S. sexually violent predator evaluations: A case law survey. *Law and Human Behavior*, 38, 248–255. <http://dx.doi.org/10.1037/lhb0000059>
- Epskamp, S. (2015). *bootnet: Bootstrap methods for various network estimation routines*. Retrieved from <https://cran.r-project.org/web/packages/bootnet/bootnet.pdf>
- Epskamp, S., Borsboom, D., & Fried, E. (2016). Estimating psychological networks and their stability: A tutorial paper. *arXiv preprint arXiv:1604.08462*.
- Epskamp, S., Cramer, A., Waldorp, L., Schmittmann, V., & Borsboom, D. (2012). qgraph: Network visualizations of relationships in psychometric data. *Journal of Statistical Software*, 48, 1–18. <http://dx.doi.org/10.18637/jss.v048.i04>
- Epskamp, S., Rhemtulla, M., & Borsboom, D. (2017). Generalized network psychometrics: Combining network and latent variable models. *Psychometrika*. Advance online publication. <http://dx.doi.org/10.1007/s11336-017-9557-x>
- Fazel, S., & Danesh, J. (2002). Serious mental disorder in 23000 prisoners: A systematic review of 62 surveys. *The Lancet*, 359, 545–550. [http://dx.doi.org/10.1016/S0140-6736\(02\)07740-1](http://dx.doi.org/10.1016/S0140-6736(02)07740-1)
- Flórez, G., Casas, A., Kreis, M. K., Forti, L., Martínez, J., Fernández, J., . . . Cooke, D. J. (2015). A prototypicality validation of the Comprehensive Assessment of Psychopathic Personality (CAPP) model Spanish version. *Journal of Personality Disorders*, 29, 707–718. <http://dx.doi.org/10.1521/pedi.2014.28.167>
- Forbes, M. K., Wright, A. G. C., Markon, K. E., & Krueger, R. F. (2017). Evidence that psychopathology symptom networks do not replicate. *Journal of Abnormal Psychology*.
- Frick, P. J., Ray, J. V., Thornton, L. C., & Kahn, R. E. (2014). Can callous-unemotional traits enhance the understanding, diagnosis, and treatment of serious conduct problems in children and adolescents? A comprehensive review. *Psychological Bulletin*, 140, 1–57. <http://dx.doi.org/10.1037/a0033076>
- Fried, E. I., & Cramer, A. O. J. (2017). Moving forward: Challenges and directions for psychopathological network theory and methodology. *Perspectives on Psychological Science*. Advance online publication. <http://dx.doi.org/10.1177/1745691617705892>
- Fried, E. I., Epskamp, S., Nesse, R. M., Tuerlinckx, F., & Borsboom, D. (2016). What are “good” depression symptoms? Comparing the centrality of DSM and non-DSM symptoms of depression in a network analysis. *Journal of Affective Disorders*, 189, 314–320. <http://dx.doi.org/10.1016/j.jad.2015.09.005>
- Fruchterman, T. M., & Reingold, E. M. (1991). Graph drawing by force-directed placement. *Software, Practice & Experience*, 21, 1129–1164. <http://dx.doi.org/10.1002/spe.4380211102>
- Gough, H. G. (1948). A sociological theory of psychopathy. *American Journal of Sociology*, 53, 359–366. <http://dx.doi.org/10.1086/220203>
- Hare, R. D. (1991). *The Hare Psychopathy Checklist—Revised*. Toronto, Canada: Multi-Health Systems.
- Hare, R. D. (1993). *Without conscience: The disturbing world of the psychopaths among us*. New York, NY: Simon & Schuster.
- Hare, R. D. (2003). *The Hare Psychopathy Checklist—Revised* (2nd ed.). Toronto, Canada: Multi-Health Systems.
- Hare, R. D., Clark, D., Grann, M., & Thornton, D. (2000). Psychopathy and the predictive validity of the PCL-R: An international perspective. *Behavioral Sciences & the Law*, 18, 623–645. [http://dx.doi.org/10.1002/1099-0798\(200010\)18:5<623::AID-BSL409>3.0.CO;2-W](http://dx.doi.org/10.1002/1099-0798(200010)18:5<623::AID-BSL409>3.0.CO;2-W)
- Hare, R. D., & Neumann, C. S. (2010). The role of antisociality in the psychopathy construct: Comment on Skeem and Cooke (2010). *Psychological Assessment*, 22, 446–454. <http://dx.doi.org/10.1037/a0013635>
- Hart, S. D., & Dempster, R. J. (1997). Impulsivity and psychopathy. In C. D. Webster & M. A. Jackson (Eds.), *Impulsivity: Theory, assessment, and treatment* (pp. 212–232). New York, NY: Guilford Press.
- Hoff, H. A., Rypdal, K., Mykletun, A., & Cooke, D. J. (2012). A prototypicality validation of the Comprehensive Assessment of Psychopathic Personality model (CAPP). *Journal of Personality Disorders*, 26, 414–427. <http://dx.doi.org/10.1521/pedi.2012.26.3.414>
- Jeandarme, I., Edens, J. F., Habets, P., Bruckers, L., Oei, K., & Bogaerts, S. (2017). PCL-R field validity in prison and hospital settings. *Law and Human Behavior*, 41, 29–43. <http://dx.doi.org/10.1037/lhb0000222>
- Karpman, B. (1948). The myth of the psychopathic personality. *The American Journal of Psychiatry*, 104, 523–534. <http://dx.doi.org/10.1176/ajp.104.9.523>
- Kreis, M. K., Cooke, D. J., Michie, C., Hoff, H. A., & Logan, C. (2012). The Comprehensive Assessment of Psychopathic Personality (CAPP): Content validation using prototypical analysis. *Journal of Personality Disorders*, 26, 402–413. <http://dx.doi.org/10.1521/pedi.2012.26.3.402>
- Lilienfeld, S. O., & Andrews, B. P. (1996). Development and preliminary validation of a self-report measure of psychopathic personality traits in noncriminal populations. *Journal of Personality Assessment*, 66, 488–524. http://dx.doi.org/10.1207/s15327752jpa6603_3
- Lilienfeld, S. O., & Fowler, K. A. (2006). The Self-Report Assessment of Psychopathy: Problems, pitfalls, and promises. In C. J. Patrick (Ed.), *Handbook of psychopathy* (pp. 107–133). New York, NY: Guilford Press.
- Lilienfeld, S. O., Patrick, C. J., Benning, S. D., Berg, J., Sellbom, M., & Edens, J. F. (2012). The role of fearless dominance in psychopathy: Confusions, controversies, and clarifications. *Personality Disorders: Theory, Research, and Treatment*, 3, 327–340. <http://dx.doi.org/10.1037/a0026987>
- Lilienfeld, S. O., Watts, A. L., Francis Smith, S., Berg, J. M., & Latzman, R. D. (2015). Psychopathy deconstructed and reconstructed: Identifying and assembling the personality building blocks of Cleckley’s chimera. *Journal of Personality*, 83, 593–610. <http://dx.doi.org/10.1111/jopy.12118>
- Lilienfeld, S. O., & Widows, M. R. (2005). *PPI-R: Psychopathic Personality Inventory Revised: Professional manual*. Lutz, FL: Psychological Assessment Resources, Incorporated.
- Lorber, M. F. (2004). Psychophysiology of aggression, psychopathy, and conduct problems: A meta-analysis. *Psychological Bulletin*, 130, 531–552. <http://dx.doi.org/10.1037/0033-2909.130.4.531>
- Lykken, D. T. (1995). *The antisocial personalities*. Hillsdale, NJ: Erlbaum.
- Lynam, D. R., Hoyle, R. H., & Newman, J. P. (2006). The perils of partialling: Cautionary tales from aggression and psychopathy. *Assessment*, 13, 328–341. <http://dx.doi.org/10.1177/1073191106290562>
- Lynam, D. R., & Miller, J. D. (2015). Psychopathy from a Basic Trait Perspective: The Utility of a Five-Factor Model Approach. *Journal of Personality*, 83, 611–626. <http://dx.doi.org/10.1111/jopy.12132>
- Lynam, D. R., & Widiger, T. A. (2001). Using the five-factor model to represent the DSM-IV personality disorders: An expert consensus approach. *Journal of Abnormal Psychology*, 110, 401–412. <http://dx.doi.org/10.1037/0021-843X.110.3.401>
- McCord, W., & McCord, J. (1964). *The psychopath: An essay on the criminal mind*. Princeton, NJ: Van Nostrand.
- McNally, R. J., Robinaugh, D. J., Wu, G. W., Wang, L., Deserno, M., & Borsboom, D. (2015). Mental disorders as causal systems: A network approach to posttraumatic stress disorder. *Clinical Psychological Science*, 3, 1–14. <http://dx.doi.org/10.1177/2167702614553230>
- Miller, C. S., Kimonis, E. R., Otto, R. K., Kline, S. M., & Wasserman, A. L. (2012). Reliability of risk assessment measures used in sexually violent predator proceedings. *Psychological Assessment*, 24, 944–953. <http://dx.doi.org/10.1037/a0028411>
- Miller, J. D., & Lynam, D. R. (2012). An examination of the Psychopathic Personality Inventory’s nomological network: A meta-analytic review.

- Personality Disorders: Theory, Research, and Treatment*, 3, 305–326. <http://dx.doi.org/10.1037/a0024567>
- Miller, J. D., & Lynam, D. R. (2015). Understanding psychopathy using the basic elements of personality. *Social and Personality Psychology Compass*, 9, 223–237. <http://dx.doi.org/10.1111/spc3.12170>
- Miller, J. D., Lynam, D. R., Widiger, T. A., & Leukefeld, C. (2001). Personality disorders as extreme variants of common personality dimensions: Can the five-factor model adequately represent psychopathy? *Journal of Personality*, 69, 253–276. <http://dx.doi.org/10.1111/1467-6494.00144>
- Nentjes, L., Bernstein, D. P., Meijer, E., Arntz, A., & Wiers, R. W. (2016). The Mask of Sanity: Facial Expressive, Self-Reported, and Physiological Consequences of Emotion Regulation in Psychopathic Offenders. *Journal of Personality Disorders*, 30, 828–838. http://dx.doi.org/10.1521/pedi_2016_30_235
- Newman, J. P., MacCoun, D. G., Vaughan, L. J., & Sadeh, N. (2005). Validating a distinction between primary and secondary psychopathy with measures of Gray's BIS and BAS constructs. *Journal of Abnormal Psychology*, 114, 319–323. <http://dx.doi.org/10.1037/0021-843X.114.2.319>
- Nieuwenhuizen, C., van Bogaerts, S., Ruijter, E. A. W., Bongers, I. L., Coppens, M., & Meijers, R. A. A. C. (2011). *TBS-behandeling geprofileerd. Een gestructureerde casussenanalyse*. Den Haag, the Netherlands: WODC.
- Opsahl, T., Agneessens, F., & Skvoretz, J. (2010). Node centrality in weighted networks: Generalizing degree and shortest paths. *Social Networks*, 32, 245–251. <http://dx.doi.org/10.1016/j.socnet.2010.03.006>
- Patrick, C. J., Bradley, M. M., & Lang, P. J. (1993). Emotion in the criminal psychopath: Startle reflex modulation. *Journal of Abnormal Psychology*, 102, 82–92. <http://dx.doi.org/10.1037/0021-843X.102.1.82>
- Patrick, C. J., Fowles, D. C., & Krueger, R. F. (2009). Triarchic conceptualization of psychopathy: Developmental origins of disinhibition, boldness, and meanness. *Development and Psychopathology*, 21, 913–938. <http://dx.doi.org/10.1017/S0954579409000492>
- Philipse, M. G. W. (2005). *Predicting criminal recidivism, empirical studies and clinical practice in forensic psychiatry*. Enschede, the Netherlands: Febodruk.
- Poythress, N. G., & Hall, J. R. (2011). Psychopathy and impulsivity reconsidered. *Aggression and Violent Behavior*, 16, 120–134. <http://dx.doi.org/10.1016/j.avb.2011.02.003>
- Poythress, N. G., Lilienfeld, S. O., Skeem, J. L., Douglas, K. S., Edens, J. F., Epstein, M., & Patrick, C. J. (2010). Using the PCL-R to help estimate the validity of two self-report measures of psychopathy with offenders. *Assessment*, 17, 206–219. <http://dx.doi.org/10.1177/1073191109351715>
- R Core Team. (2016). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>
- Rhemtulla, M., Fried, E. I., Aggen, S. H., Tuerlinckx, F., Kendler, K. S., & Borsboom, D. (2016). Network analysis of substance abuse and dependence symptoms. *Drug and Alcohol Dependence*, 161, 230–237. <http://dx.doi.org/10.1016/j.drugalcdep.2016.02.005>
- Skeem, J. L., & Cooke, D. J. (2010). Is criminal behavior a central component of psychopathy? Conceptual directions for resolving the debate. *Psychological Assessment*, 22, 433–445. <http://dx.doi.org/10.1037/a0008512>
- Skeem, J. L., Edens, J. F., Camp, J., & Colwell, L. H. (2004). Are there ethnic differences in levels of psychopathy? A meta-analysis. *Law & Human Behavior*, 28, 505–527. <http://dx.doi.org/10.1023/B:LAHU.0000046431.93095.d8>
- Skeem, J. L., & Mulvey, E. P. (2001). Psychopathy and community violence among civil psychiatric patients: Results from the MacArthur Violence Risk Assessment Study. *Journal of Consulting and Clinical Psychology*, 69, 358–374. <http://dx.doi.org/10.1037/0022-006X.69.3.358>
- Skeem, J. L., Polaschek, D. L., Patrick, C. J., & Lilienfeld, S. O. (2011). Psychopathic personality: Bridging the gap between scientific evidence and public policy. *Psychological Science in the Public Interest*, 12, 95–162. <http://dx.doi.org/10.1177/1529100611426706>
- Sörman, K., Edens, J. F., Smith, S. T., Clark, J. W., Kristiansson, M., & Svensson, O. (2016). Boldness and its relation to psychopathic personality: Prototypicality analyses among forensic mental health, criminal justice, and layperson raters. *Law and Human Behavior*, 40, 337–349. <http://dx.doi.org/10.1037/lhb0000176>
- Sörman, K., Edens, J. F., Smith, S. T., Svensson, O., Howner, K., Kristiansson, M., & Fischer, H. (2014). Forensic mental health professionals' perceptions of psychopathy: A prototypicality analysis of the Comprehensive Assessment of Psychopathic Personality in Sweden. *Law and Human Behavior*, 38, 405–417. <http://dx.doi.org/10.1037/lhb0000072>
- Spren, M., Ter Horst, P., Lutjehuis, B., & Brand, E. (2008). De kwaliteit en de rol van de PCL-R in het Nederlands tbs-systeem [The quality and the role of the PCL-R in the Dutch TBS-system]. *Developmental Psychology*, 43, 78–98.
- Thorndike, E. L. (1920). A constant error in psychological ratings. *Journal of Applied Psychology*, 4, 25–29. <http://dx.doi.org/10.1037/h0071663>
- Verschuere, B., & te Kaat, L. (2017, May). *What are the core features of psychopathy? A prototypical analysis using the Psychopathy Checklist-Revised (PCL-R)*. Paper presented at the Annual Conference of the European Association of Psychology and Law, Mechelen, Belgium.
- Vertommen, H., Verheul, R., De Ruiter, C., & Hildebrand, M. (2002). *Hare's Psychopathie checklist: Handleiding* [Dutch version of the Hare Psychopathy Checklist-Revised-Manual]. Lisse, the Netherlands: Swets Test Publisher.
- Vize, C. E., Lynam, D. R., Lamkin, J., Miller, J. D., & Pardini, D. (2016). Identifying essential features of juvenile psychopathy in the prediction of later antisocial behavior: Is there an additive, synergistic, or curvilinear role for fearless dominance? *Clinical Psychological Science*, 4, 572–590. <http://dx.doi.org/10.1177/2167702615622384>
- Whiteside, S. P., & Lynam, D. R. (2001). The five factor model and impulsivity: Using a structural model of personality to understand impulsivity. *Personality and Individual Differences*, 30, 669–689. [http://dx.doi.org/10.1016/S0191-8869\(00\)00064-7](http://dx.doi.org/10.1016/S0191-8869(00)00064-7)

Received April 11, 2017

Revision received July 28, 2017

Accepted August 4, 2017 ■