

CORRIGENDUM

Widespread distribution of proteorhodopsins
in freshwater and brackish ecosystems

Nof Atamna-Ismaeel, Gazalah Sabehi, Itai Sharon, Karl-Paul Witzel, Matthias Labrenz,
Klaus Jürgens, Tamar Barkay, Maayke Stomp, Jef Huisman and Oded Beja

The ISME Journal (2010) 4, 462; doi:10.1038/ismej.2009.130

Correction to: *The ISME Journal* (2008) 2, 656–662;
doi:10.1038/ismej.2008.27

The authors would like to apologize for any
inconvenience this may have caused.

Since the publication of this paper, the authors have
noticed an error in Table 2. The correct table is
shown below.

Table 2 The proteorhodopsin forward and reverse primers used
in this study, designed based on known PR sequences found in
the NCBI and Global Ocean Sampling datasets

<i>Forward</i>	<i>Reverse</i>
RYIDW (5'-MGNTAYATHGA YTGG-3')	GWAIYP (5'-GGRTADATNGCCC ANCC-3') GWLIIYP (5'-GGRTADATNARCC ANCC-3') GWVIYP (5'-GGRTADATNACCC ANCC-3')
RYADW (5'-MGNTAYGCNGA YTGG-3')	GWIIYP (5'-GGRTADATDATCC ANCC-3') GWSIYP (5'-GGRTADATNSWC CANCC-3') GWGTYP (5'-GGRTANGTNCCC CANCC-3')
RYFDW (5'-MGNTAYTTYGA YTGG-3')	GWAVYP (5'-GGRTANACNGCC CANCC-3') GWLVIYP (5'-GGRTANACNARC CANCC-3') GWVVIYP (5'-GGRTANACNACC CANCC-3')
RYVDW (5'-MGNTAYGTNGA YTGG-3')	GWGVYP (5'-GGRTANACNCCC CANCC-3') GWSVIYP (5'-GGRTANACNSWC CANCC-3') GWIVIYP (5'-GGRTANACDATCC ANCC-3')

Variable amino acids in the translated primer regions are marked in
bold letters.