UvA-DARE (Digital Academic Repository)

Amiodarone decreases gene expression of low-density lipoprotein receptor at both the mRNA and the protein level

Hudig, F.; Bakker, O.; Wiersinga, W.M.

DOI

[10.1016/S0026-0495\(98\)90277-3](https://doi.org/10.1016/S0026-0495(98)90277-3)

Publication date

1998

Published in

Metabolism

[Link to publication](#)

Citation for published version (APA):

Hudig, F., Bakker, O., & Wiersinga, W. M. (1998). Amiodarone decreases gene expression of low-density lipoprotein receptor at both the mRNA and the protein level. *Metabolism*, 47, 1052-1057. [https://doi.org/10.1016/S0026-0495\(98\)90277-3](https://doi.org/10.1016/S0026-0495(98)90277-3)

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Amiodarone Decreases Gene Expression of Low-Density Lipoprotein Receptor at Both the mRNA and the Protein Level

Francisca Hudig, Onno Bakker, and Wilmar M. Wiersinga

Amiodarone, a potent antiarrhythmic drug, decreases plasma and tissue triiodothyronine (T_3) and increases plasma cholesterol levels, resembling changes seen during hypothyroidism. The increase of serum cholesterol during amiodarone medication is associated with a decreased expression of the hepatic low-density lipoprotein (LDL) receptor mRNA. To further elucidate the mechanism of amiodarone-induced hypercholesterolemia, we investigated whether the decreased mRNA levels are the result of decreased transcription or increased degradation or both, and whether protein expression is decreased accordingly. Relative to pair-fed controls, amiodarone treatment increased plasma cholesterol by 69% and decreased expression of the mRNA encoding for the hepatic LDL receptor by 45%. To study this decrease in mRNA, we performed a run-on assay, from which it appears that amiodarone acts by decreasing LDL receptor mRNA expression 2.5-fold at the transcriptional level. The decay rate of liver LDL receptor mRNA, measured at different time points after injecting actinomycin D, was not different between amiodarone-treated and control animals (116 ± 32 minutes and 84 ± 10 minutes, $P = .44$). Hepatocytes in primary culture isolated from amiodarone-treated and control animals were used to determine specific binding of [125 I]-LDL to hepatic LDL receptors. Amiodarone decreased specific LDL binding and Scatchard analysis demonstrated that amiodarone treatment reduced the number of LDL receptors by 69%, without affecting the dissociation constant (K_d). In conclusion, amiodarone-induced hypercholesterolemia can be explained by decreased transcription of the LDL receptor gene, resulting in lower mRNA and protein levels.

Copyright © 1998 by W.B. Saunders Company

AMIODARONE IS AN IODINATED benzofuran derivative that is used as an antiarrhythmic drug. It contains 39.4% iodine on weight basis, and pharmacological doses of iodine are released during the biotransformation of the drug. This may cause hyperthyroidism or hypothyroidism in susceptible subjects.¹ Apart from these effects on the thyroid gland itself, amiodarone profoundly affects extrathyroidal metabolism of thyroid hormones. It inhibits the transport of thyroxine (T_4) across the plasma membrane into hepatocytes. The subsequent decreased availability of the substrate T_4 in the liver cells results in a diminished 5'-deiodination of T_4 , which results in a decreased production of triiodothyronine (T_3).² Amiodarone thus appears to induce a hypothyroid-like condition in extrathyroidal tissues. The antiarrhythmic effect of amiodarone is related to a lengthening of the cardiac action potential, quite similar to what is observed in hypothyroid patients. It has been hypothesized that the mechanism of action of amiodarone is partly due to the induction of a local hypothyroid-like condition by inhibition of the extrathyroidal conversion of T_4 into T_3 .

Thyroid hormone deficiency results in hypercholesterolemia, and a dose-dependent increase of plasma cholesterol has indeed been observed in patients on long-term amiodarone treatment.³ The amiodarone-induced increase of plasma cholesterol is largely caused by an increase in low-density lipoprotein (LDL) cholesterol, and has been reproduced in rabbits⁴ and rats.⁵ We hypothesized that the increase of plasma cholesterol by amiodarone is related to the induction of a hypothyroid-like condition in the liver by the drug. The hepatic LDL receptor plays a key role in maintaining cholesterol homeostasis, with lower levels

being associated with a higher plasma cholesterol concentration due to diminished uptake of LDL cholesterol, partly in the liver. We have recently demonstrated that amiodarone decreases the LDL receptor mRNA in the liver, whereas mRNA's encoding for hepatic hydroxymethyl glutaryl coenzyme A (HMG-CoA) reductase and cholesterol 7 α -hydroxylase (the rate-limiting enzymes in cholesterol synthesis and degradation, respectively) remained unchanged.⁶ The present studies were designed to answer the following questions: (1) Are the decreased LDL receptor mRNA's caused by decreased gene transcription or by increased mRNA degradation? (2) Does amiodarone treatment also reduce the expression of the LDL receptor protein?

MATERIALS AND METHODS

Materials

Amiodarone was a kind gift of Sanofi B.V. (Maassluis, The Netherlands). Formaldehyde, formamide, β -mercaptoethanol, and guanidine isothiocyanate were from Merck, Darmstadt, Germany. Collagenase (type H); actinomycin D, rat-tail collagen, and β -actin riboprobe (digoxigenin-labeled) were purchased from Boehringer Mannheim Biochemica (Mannheim, Germany), and 125 I (carrier-free) in NaOH from Amersham (Buckinghamshire, UK). Ham's F10, Hanks balanced salt solution (HBSS) and fetal calf serum (FCS) were obtained from Biowithacker (Brussels, Belgium). Fatty acid-free bovine serum albumin (BSA) was from Sigma (St Louis, MO). The cDNA probe for human LDL receptor (pLDLR3) was a generous gift from Dr D.W. Russell (Dallas, TX).⁷ All other reagents were of the highest grade possible.

Animals

Male Wistar rats (weight, 270 to 325 g) were housed under normal conditions with free access to tap water. All animal experiments were approved by the local animal welfare committee. Rats were divided into two groups. Group A was fed ad libitum on standard laboratory chow, and treated daily with 10 mg/100 g body weight (BW) amiodarone by gastric tube for 2 weeks. Group B was treated daily with water by gastric tube, and served as pair-fed controls of group A.

Experiment I was designed to evaluate the stability of the mRNA encoding for the LDL receptor. Twenty-four hours after the last gastric

From the Department of Endocrinology, Academic Medical Centre, Amsterdam, The Netherlands.

Submitted June 21, 1997; accepted March 13, 1998.

Address reprint requests to Onno Bakker, PhD, Department of Endocrinology, F5-171, Academic Medical Centre, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands.

Copyright © 1998 by W.B. Saunders Company
0026-0495/98/4709-0005\$03.00/0

tubing, rats were intraperitoneally injected with actinomycin D (0.2 mg/100 g BW; reconstituted in water 0.4 mg/mL at 4°C) at 30, 60, or 120 minutes before they were killed (three rats per time point). Blood (EDTA) was collected by cardiac puncture and plasma stored at -20°C for determination of plasma lipids and thyroid hormones. The liver was removed from each rat and immediately frozen and stored in liquid nitrogen for further analysis (described later).

Experiment II was designed to evaluate the effect of amiodarone on LDL receptor protein. Rats (four per group) were treated with amiodarone (group A) or water (pair-fed controls, group B) as described earlier. Twenty-four hours after the last gastric tubing, rats were anesthetized by intraperitoneal injection of Nembutal (Sanofi, France), and rat livers were isolated and perfused in a recirculating system at 37°C to isolate the parenchymal cells.

LDL Receptor mRNA Determination

RNA was isolated from liver samples using a modified Chomczynski single-step RNA isolation method.^{8,9} mRNA was isolated from total RNA using the polyAT tract mRNA isolation system from Promega (Madison, WI). A 5-μg aliquot of polyA⁺ RNA was denatured and electrophoresed in 1% agarose/formaldehyde vertical gels. The separated RNAs were transferred to nylon membranes (Boehringer Mannheim) by pressure blotting and membranes were fixed using UV crosslinking (1 J/cm²). The filters were hybridized at 55°C with a digoxigenin-labeled LDL receptor RNA probe. Hybridization and detection were performed as described previously.⁶ The blots were rehybridized at 68°C with a β-actin probe, labeled with digoxigenin, to standardize the amount of mRNA in each lane. The hybridization pattern of three identical experiments were analyzed. Several exposures of each blot were done on x-ray film (Fuji RX, Japan) and the relative amount of the mRNA was calculated using Eagle Eye II video-imaging system and OneD-scan software (Stratagene, La Jolla, CA). The exposure time was plotted against the intensity of each band. The optimal exposure time chosen was the one with the highest band intensity within the linear range, which was determined for each blot separately. The ratio between LDL receptor and β-actin was calculated. The nuclear run-on assay was performed as described using digoxigenin-labeled uridine triphosphate (UTP) on nuclei isolated from livers of amiodarone-treated and control rats,^{10,11} and signals were quantified using a Lumi-Imager (Boehringer Mannheim, Germany).

LDL-Binding Assay

Isolation of rat hepatocytes. Rat parenchymal cells were isolated as described previously.¹² Briefly, the liver was perfused with a carbogen saturated buffer containing 142 mmol/L NaCl, 6.7 mmol/L KCl, and 6.7 mmol/L Hepes, pH 7.4. The liver was removed and transferred to a perfusion apparatus, which maintains the perfusate temperature at 37°C while constantly oxygenating it. In the second step, collagenase (0.042 U/mL) and 5 mmol/L CaCl₂ were added to the perfusion medium and perfused (recirculating via backflow through the caval vein) for ±10 minutes. The cell suspension was filtered through four sheets sterile gauze and washed four times in HBSS with 2% (wt/vol) BSA. The cells were resuspended in Ham's F10 culture medium containing 10% (vol/vol) FCS, and 3 × 10⁶ cells per well were seeded in collagen-coated six-well culture dishes (2.5 mL/cm² of a 2-mg/mL collagen-coating solution). After incubation at 37°C with 5% CO₂ for 4 hours, the medium and nonattached cells were removed and replaced by serum-free medium containing 2% BSA.

Isolation and labeling of LDL. Human LDL was obtained from the blood of healthy volunteers who had fasted overnight. Isolation of LDL (1.019 < *d* < 1.063) was performed by density-gradient centrifugation according to Redgrave et al.¹³ LDL was dialyzed extensively at 4°C against phosphate-buffered saline containing 1 mmol/L EDTA. Human LDL was iodinated with ¹²⁵I by the ICI method of McFarlane.¹⁴ Labeled

LDL was separated from unbound ¹²⁵I by passage through a Sephadex G-25 column (Pharmacia, Sweden). The specific activity was approximately 125 cpm/ng LDL.

Lipoprotein binding. The binding of [¹²⁵I]-LDL to rat hepatocytes was determined by a modification of the procedure of van Berkel et al.¹⁵ After 24 hours, the dishes with primary hepatocytes were placed on ice. The cells were washed with ice-cold serum-free Ham's F10 medium containing 2% BSA. Fresh albumin-containing medium was then added containing 5 to 100 μg/ml [¹²⁵I]-LDL in the presence or absence of 500 mg/mL unlabeled LDL. Each incubation was performed in triplicate. The cells were incubated for 2 hours at 4°C. After incubation, the cells were washed five times with 0.15 mmol/L NaCl buffered with 0.05 mmol/L Tris/HCl pH 7.4 containing 0.2% BSA, and twice with buffer without BSA. Finally, the cells were lysed with 1 mL 0.1-mmol/L NaOH and the radioactivity counted. The protein content of the samples was determined using a Biorad (München, Germany) protein assay.

Assays

Plasma cholesterol concentrations were determined by a fully enzymatic, kinetic UV method (Cobas analyzer; Boehringer Mannheim). Thyroid hormones were measured with in-house radioimmunoassays (RIAs),¹⁶ and thyroid-stimulating hormone (TSH) by a specific rat TSH RIA. Reagents used in the TSH-RIA were r-TSH-RP-2 (AFP-5153B) as reference preparation, anti r-TSH-S-5 (C 21381) as antibody, and r-TSH-I-8 (AFP-8334B) as antigen for iodination (courtesy of National Institute of Arthritis, Diabetes, Digestive & Kidney Diseases, Bethesda, MD).

Calculations and Statistical Analysis

Student's *t* test was used to evaluate difference in serum cholesterol and thyroid hormones between amiodarone-treated and control rats. Data from experiment I and II were analyzed by ANOVA. The Excel 5 statistical package (Microsoft, Redmond, WA) was used.

Calculation of the binding of LDL to LDL receptor protein in experiment II was performed as follows. Specific binding was calculated by subtracting nonspecific binding from the total binding. The radioligand binding analysis program LIGAND (Biosoft, Cambridge, UK) was used to estimate the affinity constant and maximal-binding capacity (MBC). We obtained nonlinear Scatchard plots, most probably caused by an inequality of the affinity constant of the labeled and unlabeled ligand.¹⁷ The ratio (*a*) between the affinity constant of the unlabeled ligand and the labeled ligand, and the ratio (*n*) between the concentration of unlabeled and labeled ligand were calculated from the binding data according to Hollemans and Toubert¹⁸ and Hollemans and Bertina,¹⁷ and subsequently applied to calculate the MBC and the affinity constant for the unlabeled ligand.

RESULTS

Amiodarone treatment, as compared with pair-fed controls, resulted in a significant increase of plasma cholesterol and T₄ and a decrease of plasma T₃ (Table 1). Plasma TSH did not increase significantly.

Table 1. Effects of Amiodarone on Plasma Cholesterol and Thyroid Hormones

Group	n	T ₄ (nmol/L)	T ₃ (nmol/L)	TSH (ng/dL)	Cholesterol (mmol/L)
Control	16	60 ± 3	1.26 ± 0.05	1.04 ± 0.14	1.6 ± 0.1
Amiodarone	16	130 ± 5*	1.13 ± 0.04†	2.93 ± 1.04	2.7 ± 0.1*

NOTE. Values are means ± SE.

**P* < .0001, *v* controls.

†*P* < .01 *v* controls.

The rate of LDL receptor mRNA disappearance after administration of the transcription inhibitor actinomycin D was determined by Northern blot analysis, using β -actin as internal standard (Fig 1A). The hybridization patterns of three identical experiments were analyzed. The relative mRNA expression at time zero of the pair-fed controls was taken as one. The amount of LDL receptor mRNA (log values) was plotted against time (Fig 1B). Treatment with amiodarone reduced the level of LDL receptor mRNA by approximately 45% relative to pair-fed controls, as evident from time-zero levels in Fig 2. The mRNA half-lives ($t_{1/2}$) were calculated assuming first-order kinetics. The $t_{1/2}$ of LDL-R mRNA in controls was not different from that in amiodarone-treated rats (84 ± 10 v 116 ± 32 minutes, respectively; $P = .44$, $n = 3$). We also performed a run-on experiment on nuclei isolated from livers of control rats and those treated with amiodarone. As can be seen in Fig 2, the transcription from the LDL receptor gene is reduced 2.5-fold in the amiodarone-treated animals compared with controls.

Figure 3 shows the specific binding of LDL to isolated hepatocytes. The binding of LDL to LDL receptors increased with increasing amounts of added LDL. The binding of LDL to hepatocytes isolated from amiodarone-treated rats was approximately two times lower than the binding to control hepatocytes. Data obtained from this experiment were also evaluated by Scatchard analysis. A representative plot is depicted in the insert of Fig 3. The slope and the intercept of the fitted line indicate the dissociation constant (K_d) of the binding of LDL to LDL receptors and the MBC or total number of LDL receptors, respectively. The K_d was not different between amiodarone-treated animals and pair-fed controls (Table 2), but the number of LDL receptors was decreased by 69% (Table 2).

DISCUSSION

In agreement with our previous study, amiodarone treatment resulted in an increase of plasma T_4 and a decrease of plasma T_3 , explained by inhibition of T_4 5'-deiodination in the liver. The

Fig 1. (A) Effect of amiodarone on degradation of hepatic LDL receptor mRNA. Northern blot analysis of liver LDL receptor mRNA in amiodarone-treated animals (lanes 1 to 4) and pair-fed controls (lanes 5 to 8). Actinomycin D was given at 30, 60, or 120 minutes before death. (B) Effect of amiodarone on decay rate of liver LDL receptor mRNA. Degradation of LDL receptor mRNA in amiodarone-treated rats and pair-fed controls (values are mean $\pm$ SE of 3 independent experiments). LDL receptor mRNA values are relative to β -actin mRNA; the level of LDL receptor mRNA of control rats not treated with actinomycin D is set at 1.0. The decay curves are described by the following equations: $\log(y) = -0.0033 \times -0.04$ for control and $\log(y) = -0.0029 \times -0.25$ for amiodarone-treated rats.

Fig 2. Effect of amiodarone on LDL receptor gene transcription. Dots represent the result of a run-on experiment performed on nuclei purified from livers of control or amiodarone-treated rats. β -actin was used as a control to normalize the LDL receptor expression data.

reduction in plasma T_3 in this study is comparable to that previously published.^{5,19} The plasma and tissue concentrations of rats treated with 10 mg/100 g BW closely resemble those in patients on long-term amiodarone treatment.²⁰ Plasma cholesterol increased by 69%; this cannot be explained by the mild increase of TSH, because levels up to 7.6 ng/dL in experimen-

Table 2. Effects of Amiodarone on Liver LDL Receptors

Group	n	MBC (ng/mg protein)	K_d (mg/mL)
Control	4	64 ± 7.5	77 ± 15
Amiodarone	4	$20 \pm 8.6^*$	41 ± 21

NOTE. Values are means $\pm$ SE.

* $P < .01$ v controls.

tally induced hypothyroidism do not increase plasma cholesterol as observed previously.⁵ In a previous study, we demonstrated that the increase in plasma cholesterol during amiodarone treatment is due to an increase in both LDL and HDL cholesterol, whereas triglyceride levels remained unchanged.⁶ In line with our previous study, the liver mRNA encoding the LDL receptor was decreased by approximately 45% after 2 weeks of amiodarone treatment. We evaluated the degradation of the LDL receptor mRNA by injecting actinomycin D at various time intervals before the animals were killed. The dose used was twice the dose shown to decrease total RNA transcription by more than 80% in rat liver.²¹ The rate of LDL receptor mRNA degradation was not different between the amiodarone-treated animals and the pair-fed controls, as evident by the similar slope of the regression lines (Fig 2). We calculated the $t_{1/2}$ of the LDL receptor mRNA as 84 minutes in control rats. This figure is comparable to that obtained by other investigators using human cells.²²⁻²⁴ The only other report in the literature on the $t_{1/2}$ of the LDL receptor mRNA in relation to thyroid hormone status is that by Ness et al.²⁵ They observed a $t_{1/2}$ of approximately 30 minutes in young hypophysectomized rats, which did not change after treatment with T_3 . The discrepancy with our results might be due to difference in age (adult v

Fig 3. Effect of amiodarone treatment on expression of LDL receptor. Specific binding of [125 I]-LDL to LDL receptors in isolated hepatocytes from amiodarone-treated (I) and control rats (II) (values are mean $\pm$ SE of 4 independent experiments). Insert: Scatchard analysis of binding of LDL to LDL receptors in isolated hepatocytes from amiodarone-treated (I) and control animals (II). Representative plot shown.

young) or hormonal status (intact *v* hypophysectomized rats). The latter seems more plausible, because rats easily become hypercholesterolemic after hypophysectomy, by which they lose their natural resistance to develop high cholesterol levels when the cholesterol content of the diet is increased.²⁶ Because the stability of the LDL receptor mRNA is not affected, whereas its gene expression is, we conclude that the amiodarone-induced decrease in LDL receptor mRNA is caused by decreased transcription.

Our second set of experiments demonstrates that amiodarone also decreases the expression of the LDL receptor at the protein level: the number of LDL receptors in the liver of amiodarone-treated animals were on average 69% lower than that of pair-fed controls. This is evident from Scatchard analyses of the binding of LDL to isolated hepatocytes in primary culture. The experiments were performed at 4°C to prevent internalization of LDL receptors.²⁷ Although the number of LDL receptors is in the same order of magnitude as reported by others,^{28,29} the observed K_d is higher than reported. This could (partly) be the result of the collagenase treatment during the preparation of the primary hepatocytes. It is interesting to note that amiodarone did not decrease the affinity of the LDL receptor for LDL binding, although the variation in the K_d between the experiments was rather large. Given the inherent variability of the assays, we believe that amiodarone decreases the LDL receptor mRNA and protein levels about equally. An additional mechanism by which the number of cell surface receptors is decreased could be via a amiodarone-induced decrease in growth hormone concentration. It has been shown that growth hormone can restore LDL receptor expression in livers of hypophysectomized rats.^{30,31} On the other hand, we have recently obtained evidence using transfection experiments that points to a thyroid hormone-responsive element in the promoter of the LDL receptor gene (F. Hudig, unpublished observations, 1997) and we therefore suggest that at least a large part of the amiodarone effect is direct via the LDL receptor promoter. Another way in which amiodarone could influence the number of LDL receptors on the cell surface could be a, nongenomic, effect on LDL receptor recycling, as has been shown for the β -adrenergic receptor.³²

The present results strengthen the hypothesis that a local hypothyroid-like condition in the liver is connected to the amiodarone-induced hypercholesterolemia. Hypothyroidism in experimental animals is, in addition to the increase of plasma

cholesterol, associated with a decrease in hepatic LDL receptor mRNA, which is not caused by changes in mRNA stability.²⁵ The expression of the LDL receptor protein in hepatocytes in culture from hypothyroid rats is also decreased relative to euthyroid controls.³³ Amiodarone mimics these effects of thyroid hormone deficiency. The mechanism by which amiodarone exerts these hypothyroid-like effects remains to be elucidated. Thyroid hormones increase the expression of LDL receptors in rat liver both at mRNA and at the protein level.³³ Putative thyroid hormone-responsive elements are present in the promoter region of the gene encoding for the LDL receptor.³⁴ It might thus be that the potent inhibition of extrathyroidal T_3 production induced by amiodarone, which causes a very low T_3 content of several tissues, including the liver,³⁵ which results in a low T_3 receptor occupancy and thereby decreases transcription of the LDL receptor gene. The recently observed inhibition of the binding of T_3 to its nuclear α_1 and β_1 receptor by desethylamiodarone, the major metabolite of amiodarone, could reduce the occupancy of T_3 receptors even further.^{36,37} We have recently shown that T_3 counteracts the amiodarone-induced increase in plasma cholesterol and decrease in liver LDL receptor gene expression, which suggests that the inhibitory effect of amiodarone on LDL receptor gene expression is mediated via T_3 -dependent pathways.³⁸ However, other mechanisms are also possible. It has been shown that amiodarone, a lipophilic drug that accumulates to a great extent in liver and adipose tissue, stiffens membranes.³⁹ This could interfere with the internalization of membrane-bound receptors and affect the number of membrane receptors, like the LDL receptor.⁴⁰

We conclude that the amiodarone-induced increase of plasma cholesterol levels can be explained by a reduced transcription of the LDL receptor gene, which results in a decrease of the LDL receptor protein on hepatocytes. It is likely that the reduced transcription is related to an amiodarone-induced hypothyroid-like condition in the liver.

ACKNOWLEDGMENT

We would like to thank A. Maas and J.B. Daalhuisen (Department of Experimental Internal Medicine) for expert biotechnical assistance, M.C. Platvoet-ter Schiphorst for her analytical assistance, and the technicians of the Laboratory of Endocrinology for the measurement of thyroid hormones.

REFERENCES

1. Martino E, Safran M, Aghini-Lombardi F, et al: Environmental iodine intake and thyroid dysfunction during chronic amiodarone therapy. *Ann Intern Med* 101:28-34, 1984
2. de Jong M, Docter R, van der Hoek J: Different effects of amiodarone on transport of T_4 and T_3 into perfused rat liver. *Am J Physiol* 266:E44-E49, 1994
3. Wiersinga WM, Trip MD, van BM, et al: An increase in plasma cholesterol independent of thyroid function during long-term amiodarone therapy. A dose-dependent relationship. *Ann Intern Med* 114:128-132, 1991
4. Kannan R, Pollak A, Singh BN: Elevation of serum lipids after chronic administration of amiodarone in rabbits. *Atherosclerosis* 44:19-26, 1982
5. Wiersinga WM, Broenink M: Amiodarone induces a dose-dependent increase of plasma cholesterol in the rat. *Horm Metab Res* 23:94-95, 1991
6. Hudig F, Bakker O, Wiersinga WM: Amiodarone-induced hypercholesterolemia is associated with a decrease in liver LDL receptor mRNA. *FEBS Lett* 341:86-90, 1994
7. Yamamoto T, Davis CG, Brown MS, et al: The human LDL receptor: A cysteine-rich protein with multiple Alu sequences in its mRNA. *Cell* 39:27-38, 1984
8. Chomczynski P, Sacchi N: Single step method of RNA isolation by acid guanidinium thiocyanate phenol chloroform extraction. *Anal Biochem* 162:156-159, 1987
9. Kingston RE, Chomczynski P, Sacchi N: Guanidine methods for total RNA preparations, in Ausubel FM, Brent R, Kingston RE, et al (eds): *Current Protocols in Molecular Biology*. New York, NY, Wiley, 1995, pp 4.2.4-4.2.8
10. Greenberg ME, Bender TP: Identification of newly transcribed RNA, in Ausubel FM, Brent R, Kingston RE, et al (eds): *Current*

Protocols in Molecular Biology. New York, NY, Wiley, 1995, pp 4.10.1-4.10.10

11. Merscher S, Hanselmann R, Welter C, et al: Nuclear runoff transcription analysis using chemiluminescent detection. *Biotechniques* 16:1024-1026, 1994

12. Segelen PO: Preparation of isolated rat liver cells. *Methods Cell Biol* 13:29-83, 1976

13. Redgrave TG, Roberts DC, West CE: Separation of plasma lipoproteins by density gradient centrifugation. *Anal Biochem* 65:42-49, 1975

14. McFarlane AS: Efficient trace-labeling of proteins with iodine. *Nature* 182:53-54, 1958

15. van Berkel TJ, Kruijt JK, van Gent T, et al: Saturable high affinity binding, uptake and degradation of rat plasma lipoproteins by isolated parenchymal and nonparenchymal cells from rat liver. *Biochim Biophys Acta* 665:22-33, 1980

16. Wiersinga WM, Chopra IJ: Radioimmunoassays of thyroxine (T₄), 3,5,3'-triiodothyronine (T₃) and 3,3'-diodothyronine (T₂). *Meth Enzymol* 84:272-303, 1982

17. Hollemans HJ, Bertina RM: Scatchard Plot and heterogeneity in binding affinity of labeled and unlabeled ligand. *Clin Chem* 21:1769-1773, 1975

18. Hollemans HJ, Toubert JL: Plot for exact calculation of specific activity in saturation analysis. *Clin Chim Acta* 56:305-306, 1974

19. Kasim SE, Bagchi N, Brown TR, et al: Effect of amiodarone on serum lipids, lipoprotein lipase and hepatic triglyceride lipase. *Endocrinology* 120:1991-1995, 1987

20. Hartong R, Wiersinga WM, Lamers WH, et al: Dose-dependent effect of amiodarone on thyroid hormone-responsive gene expression in rat liver, in Hartong R (ed): *Nuclear Thyroid Hormone Receptors and Gene Expression*. Thesis. Amsterdam, The Netherlands, Rodopi, 1989, pp 78-92

21. Koo P, Nagai MK, Farber E: Multiple sites of control of glutathione S-transferase P1-1 in rat liver. *J Biol Chem* 269:14601-14606, 1994

22. Wilson GM, Roberts EA, Deeley RG: Modulation of LDL receptor mRNA stability by phorbol esters in human liver cell culture models. *J Lipid Res* 38:437-466, 1997

23. Huang Y, Ghosh MJ, Lopes-Virella MF: Transcriptional and post-transcriptional regulation of LDL receptor gene expression in PMA-treated THP-1 cells by LDL-containing immune complexes. *J Lipid Res* 38:110-200, 1997

24. Stopeck AT, Nicholson AC, Mancini FP, et al: Cytokine regulation of low density lipoprotein receptor gene transcription in HepG2 cells. *J Biol Chem* 268:17489-17944, 1993

25. Ness GC, Zhao Z: Thyroid hormone rapidly induces hepatic LDL receptor mRNA levels in hypophysectomized rats. *Arch Biochem Biophys* 199-202, 1994

26. Bhattacharya S, Balasubramaniam S, Simons LA: Regulation of low-density-lipoprotein metabolism in the rat. *Biochem J* 234:493-496, 1986

27. Goldstein JL, Basu SK, Brown MS: Receptor-mediated endocytosis of low-density-lipoprotein in cultured cells. *Meth Enzymol* 98:241-260, 1983

28. Kovanen PT, Brown MS, Goldstein JL: Increased binding of low density lipoprotein to liver membranes from rats treated with 17 α -ethinyl estradiol. *J Biol Chem* 254:11367-11373, 1979

29. Windler EE, Kovanen PT, Chao YS, et al: The estradiol-stimulated lipoprotein receptor of rat liver. A binding site that membrane mediates the uptake of rat lipoproteins containing apoproteins B and E. *J Biol Chem* 255:10464-10471, 1980

30. Hoogerbrugge N, Jansen H, Staels B, et al: Growth hormone normalizes low-density lipoprotein receptor gene expression in hypothyroid rats. *Metabolism* 45:680-685, 1996

31. Rudling M, Angelin B: Loss of resistance of dietary cholesterol in the rat after hypophysectomy: Importance of the presence of growth hormone for hepatic low density lipoprotein-receptor expression. *Proc Natl Acad Sci USA* 90:8851-8855, 1993

32. Yin Y, Nicolas P, Perret GY, et al: Antagonism between T₃ and amiodarone on the contractility and density of β -adrenoceptors of chicken cardiac myocytes. *Eur J Pharmacol* 261:97-104, 1994

33. Salter AM, Hayashi R, al-Seeni M, et al: Effects of hypothyroidism and high-fat feeding on mRNA concentrations for the low-density-lipoprotein receptor and on acyl-CoA:cholesterol acyltransferase activities in rat liver. *Biochem J* 276:825-832, 1991

34. Ness GC, Pendleton LC, Li YC, et al: Effect of thyroid hormone on hepatic cholesterol 7 α hydroxylase, LDL receptor, HMG-CoA reductase, farnesyl pyrophosphate synthetase and apolipoprotein A-I mRNA levels in hypophysectomized rats. *Biochem Biophys Res Com* 172:1150-1156, 1990

35. Schroder-van der Elst JP, van der Heide D: T₄, T₃ and rT₃ concentrations in several tissues of rat: Effect of amiodarone and desethylamiodarone on thyroid hormone metabolism. *Endocrinology* 127:1656-1664, 1990

36. Bakker O, van Beeren HC, Wiersinga WM: Desethylamiodarone is a noncompetitive inhibitor of the binding of thyroid hormone to the thyroid hormone beta1-receptor protein. *Endocrinology* 134:1665-1670, 1994

37. van Beeren HC, Bakker O, Wiersinga WM: Desethylamiodarone is a competitive inhibitor of the binding of thyroid hormone to the thyroid hormone α 1-receptor protein. *Mol Cel Endocrinol* 112:15-19, 1995

38. Hudig F, Bakker O, Wiersinga WM: Tri-iodothyronine prevents the amiodarone-induced decrease in the expression of the liver low-density lipoprotein receptor gene. *J Endocrinol* 152:413-421, 1997

39. Chatelain P, Ferreira J, Laruel R, et al: Amiodarone induced modifications of phospholipid physical state. A fluorescence polarization study. *Biochem Pharmacol* 35:3007-3013, 1986

40. Limberd MJ, Lefkowitz RJ: Adenylate cyclase coupled beta-adrenergic receptors effect of membrane lipid-perturbing agents on receptor binding and enzyme stimulation by catecholamines. *Mol Pharmacol* 12:559-567, 1976