



UvA-DARE (Digital Academic Repository)

(Sub)-millimetre Observations of MAXI J1820+070 (ASASSN-18ey) Suggest Jet Quenching on July 6

Tetarenko, A.J.; Petitpas, G.; Sivakoff, G.R.; Miller-Jones, J.C.A.; Russell, T.D.; Schieven, G.; JACPO T XRB collaboration

Publication date

2018

Document Version

Final published version

Published in

The astronomer's telegram

License

Unspecified

[Link to publication](#)

Citation for published version (APA):

Tetarenko, A. J., Petitpas, G., Sivakoff, G. R., Miller-Jones, J. C. A., Russell, T. D., Schieven, G., & JACPO T XRB collaboration (2018). (Sub)-millimetre Observations of MAXI J1820+070 (ASASSN-18ey) Suggest Jet Quenching on July 6. *The astronomer's telegram*, 11831. <http://www.astronomerstelegam.org/?read=11831>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

The Astronomer's Telegram

[Post](#) | [Search](#) | [Policies](#)
[Credential](#) | [Feeds](#) | [Email](#)

8 Mar 2019; 08:07 UT

This space for free for your conference.



Outside
 GCN
 IAUCs

Other
 ATel on [Twitter](#) and [Facebook](#)
[ATELstream](#)
[ATel Community Site](#)

[[Previous](#) | [Next](#) | [ADS](#)]

(Sub)-millimetre Observations of MAXI J1820+070 (ASASSN-18ey) Suggest Jet Quenching on July 6

ATel #11831; *A. J. Tetarenko (Alberta), G. Petitpas (Cfa), G. R. Sivakoff (Alberta), J. C. A. Miller-Jones (Curtin), T. D. Russell (UvA), G. Schieven (NRC), and the JACPOT XRB Collaboration*

on 9 Jul 2018; 15:18 UT

Credential Certification: *Alexandra Tetarenko (tetarenk@ualberta.ca)*

Subjects: Radio, Millimeter, Sub-Millimeter, Black Hole, Transient

Referred to by ATel #: [11887](#), [11899](#), [11936](#), [12157](#)

Following the report of a hard to soft accretion state transition (ATel #[11820](#)) in the currently outbursting candidate black hole X-ray binary, MAXI J1820+070 (ATel #[11399](#), #[11400](#), #[11404](#), #[11406](#), #[11418](#), #[11420](#), #[11440](#)), we performed target-of-opportunity observations with the Atacama Large Millimeter/Submillimeter Array (ALMA) and the Submillimeter Array (SMA). We last observed MAXI J1820+070 with both ALMA and the SMA on April 12.

ALMA observations were taken on 2018 April 12 between 08:12-10:36 UTC and on 2018 July 6 between 01:31-06:18 UTC (MJD~58305.063-58305.262), in the 335.5-339.4 GHz and 347.5-351.5 GHz frequency bands. On April 12, the ALMA flux density at 343 GHz was very bright, ~120 mJy. Highly preliminary quality assurance checks on the July 6 observations detected MAXI J1820+070 with peak flux densities of 7.2 mJy (near 03 UT) and 4.9 (near 05 UT) mJy at a representative frequency of 343.5 GHz. We estimate that these preliminary measurements have absolute fluxes only good to ~25%, but that the drop in the flux density over two hours is likely real.

SMA observations occurred on 2018 April 12 between 11:00-20:00 UT and on 2018 July 6 between 09:05 - 11:40 UTC (MJD=58305.378-58305.486). During our observations, the two SMA receivers were tuned to local oscillator frequencies of 223/231 GHz (April 12) and 223/275 GHz (July 6). On April 12, the SMA flux density was ~100 mJy at 227 GHz (where the flux was highly variable during the observation). Stacking the two frequency bands together in the image plane on July 6, we detect a 3 sigma source coincident with the position measured from our previous mm/sub-mm observations (ATel #[11440](#)). Fitting a point source in the image plane we measure a preliminary flux density of 2.6 +/-0.8 mJy at 249 GHz.

We see clear evidence for reduced jet flux density values at mm/sub-mm frequencies between April 12 and July 6 observations. Over this same time, Swift/BAT only measured a moderate drop in hard X-rays (15-50 keV; 0.66+/-0.03 counts/cm²/s on April 12 dropping to 0.21 +/-0.02 counts/cm²/s on July 5/6 near 23:40 - 07:45 UTC). All measurements of the jets flux density on July 6 are significantly fainter than that predicted by the standard radio/X-ray correlation ($f_r \propto$

Related

- [12534](#) MAXI J1820+070 is close to quiescence
- [12157](#) MAXI 1820+070 has completed the decline from the recent optical re-brightening following the soft to hard transition
- [12128](#) Optical brightening of MAXI J1820+070 over the soft to hard transition observed with LCO and the AI Sadeem Observatory
- [12068](#) MAXI J1820+070 continuing its rapid evolution toward the hard state
- [12064](#) Swift observes MAXI J1820+070 in transition from the soft to the hard-intermediate state
- [12061](#) AMI radio detection of the black hole candidate MAXI J1820+070 during the soft to hard transition.
- [12057](#) MAXI/GSC detection of a rapid increase in the hard X-ray flux of MAXI J1820+070
- [11960](#) The 55 Hz signal we detected in MAXI J 1820+070 is not a QPO
- [11951](#) Detection of a 55 Hz high-frequency QPO in MAXI J1820+070 with NICER
- [11936](#) Optical timing observations of MAXI J1820+070 with IFI+IQUEYE and AQUEYE+ soon after state transition
- [11899](#) Short-lived episodes of emission line splitting in the candidate black hole X-ray binary MAXI 1820+070
- [11887](#) LOFAR observations of MAXI J1820+070 (ASASSN-18ey) during its recent state transition
- [11855](#) Polarimetric monitoring of the MAXI J1820+070 in optical and near-infrared wavelengths
- [11833](#) Declining near-infrared flux from the black-hole candidate MAXI J1820+070 (ASASSN-18ey) in transition
- [11831](#) (Sub)-millimetre Observations of MAXI J1820+070 (ASASSN-18ey) Suggest Jet Quenching on July 6

$f_x^{0.7}$) based on their April 12 values. This strongly suggests that the radio jet was quenching on July 6 as the hard intermediate - soft intermediate state transition occurred (ATel #11823).

Radio through mm/sub-mm detections of black hole X-ray binaries typically arise from a compact synchrotron-emitting jet, characterized by a flat to slightly inverted, optically thick spectrum, that can break to a steep, optically thin spectrum at higher frequencies. The spectral break of the jet has been seen to move to lower frequencies as the compact jet is quenched during transitions to softer accretion states (e.g., Russell et al. 2014, MNRAS, 439, 1390). The ALMA and SMA flux densities on July 6 are increasingly more discrepant from those expected from the hard X-ray flux. Moreover, the measurement of the early ALMA flux density is nearly simultaneous with the recently reported ~ 6 mJy AMI detection towards the end of its monitoring on July 5/6 (ATel #11827). Our best explanations for these data requires that both the jet-power drop (relative to that expected from the accretion rate) and that either: (a) the radio--sub-mm spectrum has transformed from being inverted on April 12 to nearly flat on July 6 or (b) the spectral break moved from above 350 GHz on April 12 to below 350 GHz (possibly to frequencies below the SMA band) on July 6.

We thank the SMA and ALMA staff for rapidly scheduling our observations.

- 11827 AMI-LA 15.5 GHz observations of radio flaring from the black hole candidate MAXI J1820+070 in transition
- 11824 Other low-frequency optical QPO-like features in MAXI J1820+070 detected with IFI+IQUEYE@Galileo
- 11823 Continuing NICER observations of the state transition in ASASSN-18ey/MAXI J1820+070
- 11820 A rapid state transition in MAXI J1820+070
- 11723 Low-frequency optical QPO in MAXI J1820+070 detected with IFI+IQUEYE@Galileo
- 11661 KEGS Discovery of 9 Supernova Candidates in the K2 Campaign 17 field with Pan-STARRS PS1
- 11609 Simultaneous LOFAR and AMI-LA observations of MAXI J1820+070
- 11596 MAXI J1820+070 has optical period of 3.4 hours
- 11591 Further detection of the optical low frequency QPO in the black hole transient MAXI J1820+070
- 11578 Exponential increase in X-ray QPO frequency with time in MAXI J1820+070
- 11576 NICER observations of MAXI J1820+070: Continuing evolution of X-ray variability properties
- 11574 Optical/X-ray Flux Decoupling in MAXI J1820+070
- 11540 VLITE meter-wavelength detection of MAXI J1820+070 at 339 MHz
- 11539 The 30-day monitoring of MAXI J1820+070 at 4.7 GHz
- 11533 A bright mid-infrared excess in MAXI J1820+070
- 11510 Detection of optical and X-ray QPOs at similar frequencies in MAXI J1820+070
- 11490 INTEGRAL observations of MAXI J1820+070: public data products
- 11488 Low-frequency QPOs in MAXI J1820+070 as seen by INTEGRAL/SPI
- 11482 Palomar 60-inch SEDM classification of optical transients
- 11481 MAXI J1820+070: VLT and GTC spectroscopic follow-up shows a significant spectral evolution from the early stages of the outburst
- 11480 ePESSTO spectroscopic classification of optical transients
- 11478 INTEGRAL observations of MAXI J1820+070
- 11462 Near Infrared JHKs observations of the transient MAXI J1820+070 / ASASSN-18ey: Erratum on 2MASS counterpart designation
- 11458 Near Infrared JHKs observations of the transient MAXI J1820+070 / ASASSN-18ey
- 11451 Fast infrared photometry of the black-hole candidate MAXI J1820+070

11445	First measurements of linear polarization of MAXI J1820+070
11440	NOEMA Sub-millimetre Detection of MAXI J1820+070
11439	A flat radio spectrum of MAXI J1820+070
11437	Red sub-second optical flaring in MAXI J1820+070 observed by ULTRACAM/NTT
11432	Correlated Optical/X-ray Timing Variations in MAXI J1820+070 found by Swift UVOT and XRT
11427	The hard X-ray spectrum of MAXI J1820+070 observed by Swift/BAT
11426	Detection of 10-msec scale optical flares in the black-hole binary candidate MAXI J1820+070 (ASASSN-18ey)
11425	Optical Spectra of MAXI J1820+070 with Keck
11424	SOAR/Goodman optical spectroscopy of MAXI J1820+070
11423	NICER observations of MAXI J1820+070 suggest a rapidly-brightening black hole X-ray binary in the hard state
11421	Fast optical flaring in the suspected black-hole binary MAXI J1820+070 (ASASSN-18ey)
11420	AMI radio observations of the black hole candidate MAXI J1820+070
11418	Optical observations of MAXI J1820+070 suggest it is a black hole X-ray binary
11406	MAXI J1820+070: Errata and updated XRT Position
11404	MAXI J1820+070: Swift/UVOT counterpart correction
11403	Swift detection of MAXI J1820+070
11400	Optical follow-up of MAXI J1820+070 and possible identity with ASASSN-18ey
11399	MAXI/GSC detection of a probable new X-ray transient MAXI J1820+070
10459	Ongoing radio monitoring of Cyg X-1 with the RATAN-600 radio telescope

[**Telegram Index**]

R. E. Rutledge, Editor-in-Chief
Derek Fox, Editor
Mansi M. Kasliwal, Co-Editor

rrutledge@astronomerstelegam.org
dfox@astronomerstelegam.org
mansii@astronomerstelegam.org