



UvA-DARE (Digital Academic Repository)

Re-brightening of the black hole candidate MAXI J1535-571 as it transitioned back to the soft state

Parikh, A.S.; Russell, T.D.; Wijnands, R.; Bahramain, A.; Miller-Jones, J.C.A.; Tetarenko, A.J.; Sivakoff, G.R.

Publication date

2018

Document Version

Final published version

Published in

The astronomer's telegram

License

Unspecified

[Link to publication](#)

Citation for published version (APA):

Parikh, A. S., Russell, T. D., Wijnands, R., Bahramain, A., Miller-Jones, J. C. A., Tetarenko, A. J., & Sivakoff, G. R. (2018). Re-brightening of the black hole candidate MAXI J1535-571 as it transitioned back to the soft state. *The astronomer's telegram*, 11652.
<http://www.astronomerstelegam.org/?read=11652>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Outside

GCN
IAUCs

Other

ATel on [Twitter](#) and [Facebook](#)
ATELstream
ATel Community Site[[Previous](#) | [Next](#) | [ADS](#)]

Re-brightening of the black hole candidate MAXI J1535-571 as it transitioned back to the soft state

ATel #11652; *A. S. Parikh, T. D. Russell, R. Wijnands (UvA), A. Bahramain, J. C. A. Miller-Jones (ICRAR-Curtin), A. J. Tetarenko, G. R. Sivakoff (U. Alberta)*

on 18 May 2018; 18:19 UT

Credential Certification: *Aastha Parikh (A.S.Parikh@uva.nl)*

Subjects: Radio, Millimeter, Sub-Millimeter, X-ray, Black Hole, Transient

Referred to by ATel #: [11682](#)

MAXI J1535-571 is a black-hole candidate in a low-mass X-ray binary that was detected independently by Swift and MAXI in outburst in early 2017 September (GCN #[21792](#), ATel #[10699](#)). The source transitioned to a soft state in late 2017 November (ATel #[11020](#)) and was reported to transition back to its hard state around 2018 April 30 (ATel #[11611](#)).

More recent Swift/X-ray Telescope (XRT) observations of the source in the hard state indicated that the source was decaying. On 2018 May 11 the source exhibited a minimum observed flux of $\sim 1.2E-12$ erg/cm²/s (0.5-10 keV; unabsorbed). However, the following XRT observation, carried out on 2018 May 14 (intermittently between 08:42 and 13:37 UT), showed that the flux had increased by a factor of ~ 20 to $\sim 2.4E-11$ erg/cm²/s (0.5-10 keV; unabsorbed). We fitted the spectrum from this observation with an absorbed power-law model. We fixed the equivalent hydrogen column density N_H to $4.4E22$ cm⁻² (obtained by fitting data with higher signal-to-noise ratio; see ATel #[11611](#)). The spectrum corresponded to a photon index of $\Gamma = 1.7 \pm 0.3$, indicating that the source was still in the hard state.

Around the same time as this initial X-ray re-brightening, we observed MAXI J1535-571 with the Australia Telescope Compact Array (ATCA) on 2018 May 14 (between 16:11 and 21:48 UT). These observations were taken at 5.5 and 9 GHz (simultaneously) with a bandwidth of 2 GHz at each frequency. We used PKS 1934-638 for primary calibration and 1520-58 for phase calibration. Following standard procedure in CASA (v4.7.2; McMullin et al. 2007, ASPC, 376, 127) the data were reduced then imaged (using natural weighting). We significantly detected the source at both frequencies. Fitting a point source in the image plane we measure flux densities of 0.89 ± 0.02 mJy at 5.5 GHz and 1.04 ± 0.02 mJy at 9 GHz. The radio spectral index of $\alpha = 0.3 \pm 0.1$, where $S_{\nu} \propto \nu^{\alpha}$, is consistent with emission from a compact jet during a hard X-ray state.

The most recent XRT observation, observed on 2018 May 18 (intermittently between 02:28 and 12:04 UT), indicated that the source had brightened further, by a factor of ~ 400 corresponding to

Related

- [11884](#) INTEGRAL detects re-brightening of the black hole candidate MAXI J1535-571
- [11682](#) MAXI/GSC detection of an undergoing soft-to-hard state transition of MAXI J1535-571
- [11652](#) Re-brightening of the black hole candidate MAXI J1535-571 as it transitioned back to the soft state
- [11611](#) X-ray spectral hardening and radio non-detection of MAXI J1535-571
- [11568](#) Rapidly fading the black hole X-ray nova MAXI J1535-571 from MAXI/GSC observations
- [11020](#) MAXI J1535-571 currently shows softest spectrum
- [10899](#) Radio re-brightening of MAXI J1535-571 as it transitions back towards the hard state
- [10816](#) Brackett-Gamma emission in MAXI J1535-571
- [10768](#) Initial NICER observations of a broadened iron line and QPOs in MAXI J1535-571
- [10761](#) MAXI/GSC observations indicate further softening of MAXI J1535-571 toward the high/soft state
- [10745](#) Extremely Bright Radio and (Sub-)Millimetre Detections of MAXI J1535-571
- [10734](#) Detection of the low-frequency QPOs in MAXI J1535-571
- [10731](#) Swift/XRT sees MAXI J1535-571 brightening and softening
- [10729](#) MAXI/GSC detects an X-ray spectral softening of the X-ray transient MAXI J1535-571
- [10716](#) The near-infrared counterpart of X-ray transient MAXI J1535-571
- [10714](#) MAXI J1535-571 (Candidate Black Hole X-ray Binary): Coordination
- [10711](#) ATCA radio detection of MAXI J1535-571 indicates it is a strong black hole X-ray binary candidate
- [10708](#) Further brightening of the X-ray transient MAXI J1535-571, suggesting the

a flux of $\sim 9.8 \times 10^{-9}$ erg/cm²/s (0.5-10 keV; unabsorbed). The spectrum was fit by a photon index of $\Gamma = 4.7 \pm 0.2$ (N_{H} fixed to 4.4×10^{22} cm⁻²), indicating that the source may have transitioned to a soft state. Close in time to this second XRT observation, we observed MAXI J1535-571 with ATCA and the Atacama Large Millimeter/submillimeter Array (ALMA). The ATCA observations were taken on 2018 May 17 (between 06:54 and 11:00 UT) and followed the same setup and procedures described above. With these radio observations, we do not detect the source at either 5.5 or 9 GHz, with 3-sigma upper-limits of 0.06 mJy/beam at 5.5 GHz and 0.1 mJy/beam at 9 GHz. The source was also not detected in our quasi-simultaneous ALMA observations, which were taken at 97, 140 and 230 GHz, with 8 GHz of bandwidth at each frequency. All three bands were observed on May 17, where the 97 GHz observations were taken between 01:42 and 02:02 UT, the 140 GHz observations between 08:03 and 08:30 UT, and the 230 GHz observations between 01:09 and 01:41 UT. All three ALMA observations provide preliminary 3-sigma upper limits of 0.075 mJy/beam in all three bands. These non-detections imply the compact jet was quenched, as expected during the soft X-ray spectral state.

Our current XRT campaign will continue to observe MAXI J1535-571 until 2018 May 30. We will request further XRT observations if the source remains X-ray active. We will also continue to monitor this source at radio frequencies.

We thank Swift, ATCA, and ALMA for scheduling these observations.

	presence of a black hole
10704	Erratum to ATel #10702: MAXI J1535-571 optical counterpart coordinates
10702	Discovery of the optical counterpart of MAXI J1535-571
10700	MAXI J1535-571: Swift detection and localization
10699	MAXI/GSC discovery of a new hard X-ray transient MAXI J1535-571

[**Telegram Index**]

R. E. Rutledge, Editor-in-Chief

Derek Fox, Editor

Mansi M. Kasliwal, Co-Editor

rrutledge@astronomerstelegam.org

dfox@astronomerstelegam.org

mansi@astronomerstelegam.org