



UvA-DARE (Digital Academic Repository)

Corrigendum to: Abscisic acid influences tillering by modulation of strigolactones in barley

Wang, H.; Chen, W.; Eggert, K.; Charnikhova, T.; Bouwmeester, H.; Schweizer, P.; Hajirezaei, M.R.; Seiler, C.; Sreenivasulu, N.; von Wirén, N.; Kuhlmann, M.

DOI

[10.1093/jxb/ery306](https://doi.org/10.1093/jxb/ery306)

Publication date

2018

Document Version

Final published version

Published in

Journal of Experimental Botany

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Wang, H., Chen, W., Eggert, K., Charnikhova, T., Bouwmeester, H., Schweizer, P., Hajirezaei, M. R., Seiler, C., Sreenivasulu, N., von Wirén, N., & Kuhlmann, M. (2018). Corrigendum to: Abscisic acid influences tillering by modulation of strigolactones in barley. *Journal of Experimental Botany*, 69(21), 5307. <https://doi.org/10.1093/jxb/ery306>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)



CORRIGENDUM

Corrigendum to: Abscisic acid influences tillering by modulation of strigolactones in barley

Hongwen Wang¹, Wanxin Chen², Kai Eggert¹, Tatsiana Charnikhova³, Harro Bouwmeester⁴,
Patrick Schweizer², Mohammad R. Hajirezaei¹, Christiane Seiler⁵, Nese Sreenivasulu⁶,
Nicolaus von Wirén^{1,*} and Markus Kuhlmann^{5,*}

¹ Department of Physiology and Cell Biology, Leibniz Institute of Plant Genetics and Crop Plant Research (IPK) Gatersleben, Corrensstrasse 3, 06466 Stadt Seeland, Germany

² Department of Breeding Research, Leibniz Institute of Plant Genetics and Crop Plant Research (IPK) Gatersleben, Corrensstrasse 3, 06466 Stadt Seeland, Germany

³ Laboratory of Plant Physiology, Wageningen University, Droevendaalsesteeg 1, 6708 PB Wageningen, The Netherlands

⁴ Plant Hormone Biology Group, Swammerdam Institute for Life Sciences, University of Amsterdam, 1098 XH Amsterdam, The Netherlands

⁵ Department of Molecular Genetics, Leibniz Institute of Plant Genetics and Crop Plant Research (IPK) Gatersleben, Corrensstrasse 3, 06466 Stadt Seeland, Germany

⁶ International Rice Research Institute (IRRI), Grain Quality and Nutrition Center, Metro Manila, Philippines

* Correspondence: vonwiren@ipk-gatersleben.de or kuhlmann@ipk-gatersleben.de

Journal of Experimental Botany, Vol. 69, No. 16 pp. 3883–3898, doi:10.1093/jxb/ery200

The authors of this article would like to give due credit to a colleague, Dr. Götz Hensel, whose assistance was of great value to this study. The online version of this article has now been updated to reflect that contribution.

The authors would like to apologise for the original omission.