



UvA-DARE (Digital Academic Repository)

PSR J1740+1000: A Young Pulsar Well Out of the Galactic Plane

McLaughlin, M.A.; Arzoumanian, Z.; Cordes, J.M.; Backer, D.C.; Lommen, A.; Lorimer, D.R.; Zepka, A.F.

Published in:
Astrophysical Journal

DOI:
[10.1086/324151](https://doi.org/10.1086/324151)

[Link to publication](#)

Citation for published version (APA):

McLaughlin, M. A., Arzoumanian, Z., Cordes, J. M., Backer, D. C., Lommen, A., Lorimer, D. R., & Zepka, A. F. (2002). PSR J1740+1000: A Young Pulsar Well Out of the Galactic Plane. *Astrophysical Journal*, 564(1), 333-342. DOI: 10.1086/324151

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

PSR J1740+1000: A YOUNG PULSAR WELL OUT OF THE GALACTIC PLANE

M. A. McLAUGHLIN,¹ Z. ARZOUMANIAN,² J. M. CORDES,¹ D. C. BACKER,³ A. N. LOMMEN,³ D. R. LORIMER,⁴
AND A. F. ZEPKA⁵

Received 2001 June 20; accepted 2001 August 30

ABSTRACT

We discuss PSR J1740+1000, one of five pulsars recently discovered in a search of 470 deg² at 430 MHz during the upgrade of the 305 m Arecibo Telescope. The period $P = 154$ ms and period derivative $\dot{P} = 2.1 \times 10^{-14}$ s s⁻¹ imply a spin-down age $\tau_s = P/2\dot{P} = 114$ kyr that is smaller than 95% of all known pulsars. The youth and proximity of this pulsar make it a good candidate for detection at X-ray and gamma-ray energies. Its high Galactic latitude ($b = 20^\circ 4'$) suggests a very high velocity if the pulsar was born in the midplane of the Galaxy and if its kinematic age equals its spin-down age. Interstellar scintillations, however, suggest a much lower velocity. We discuss possible explanations for this discrepancy, taking into account (1) possible birth sites away from the midplane; (2) contributions from the unmeasured radial velocity; (3) a kinematic age different from the spin-down age; and (4) biasing of the scintillation velocity by enhanced scattering from the North Polar Spur.

Subject headings: ISM: individual (North Polar Spur) — pulsars: general — pulsars: individual (J1740+1000)

1. INTRODUCTION

PSR J1740+1000, discovered in an Arecibo survey at 430 MHz (McLaughlin et al. 2000), is a young pulsar at high Galactic latitude. It may be representative of a class of high-velocity pulsars that have thus far eluded detection. Determining the velocity and origin of this object is important as pulsar velocities represent fossil information about evolution of close binary systems and core collapse processes in supernovae. The largest velocities, in excess of 1000 km s⁻¹, provide the greatest constraints on such processes. Measuring pulsar velocities is also crucial for pulsar population statistics, determining the birthrate and birth places of pulsars, and for planning future pulsar searches. Unfortunately, pulsar surveys have been strongly biased against the detection of high-velocity pulsars. While the average pulsar speed is about 500 km s⁻¹, the tail of the velocity distribution extends to at least 1600 km s⁻¹, with 10%–20% of all pulsars having speeds greater than 1000 km s⁻¹ (Lyne & Lorimer 1994; Cordes & Chernoff 1998; Arzoumanian, Chernoff, & Cordes 2001). At these speeds, within a typical 10 Myr pulsar lifetime many high-velocity pulsars will have moved beyond the volume of detectability of previous surveys at low Galactic latitudes (Lorimer, Bailes, & Harrison 1997).

In § 2 of this paper we describe the search and follow-up observations. In § 3 we present a timing model and discuss the pulsar's single pulse and interstellar scintillation (ISS) properties. In § 4 we explore ways to reconcile the kinematic and ISS velocities of the pulsar. In §§ 5 and 6 we describe prospects for astrometry and high-energy detection, respectively. In § 7 we present our main conclusions and plans for future study.

2. OBSERVATIONS

PSR J1740+1000 was discovered during the upgrade of the Arecibo Observatory, when several collaborations jointly surveyed the Arecibo sky (covering declinations roughly between -1° and 39°). As the telescope was not able to track sources, data were taken as the sky drifted through the beam at the sidereal rate. A point source drifts through the Arecibo 430 MHz 10' beam in approximately 40 s. The feed was moved once a day to allow successive declination strips to be surveyed. The nominal sensitivity of this search, which employed 32 channels covering a bandwidth of 8 MHz, was approximately 0.5 mJy. We estimate that the techniques used for radio frequency interference excision and candidate evaluation raised our sensitivity to roughly 1 mJy. With a sampling interval of 250 μ s, we were sensitive to pulsars with periods ~ 1 ms up to dispersion measures (DMs) of ~ 20 pc cm⁻³ and to pulsars with periods ~ 30 ms up to DMs of ~ 600 pc cm⁻³.

The Berkeley/Cornell collaboration covered 470 deg² between 1994 October and 1995 February. We detected five new pulsars in the search. Three (J0030+0451, J0711+0931, and J1313+0931) were reported in Lommen et al. (2000), and two (J1740+1000 and J1849+2423) were reported in McLaughlin et al. (2000). With a search-derived period of 154 ms, a dispersion measure (DM) of ~ 20 pc cm⁻³, and a signal-to-noise ratio of ~ 20 , J1740+1000 was one of several pulsar candidates that were reobserved after the upgrade was completed. Confirming observations were done in 1998 May with the Arecibo Observatory Fourier Transform Machine (AOFTM), a Fourier transform-based spectrometer with 1024 10 kHz frequency channels and 102.4 μ s sampling.⁶ The pulsar was timed from 1999 May onward with the Penn State Pulsar Machine (PSPM), a filter bank with 128 60 kHz frequency channels and 80 μ s sampling. We have also observed this pulsar with the Wideband Arecibo Pulsar Processor (WAPP), a fast-dump digital correlator covering up to a 100 MHz bandwidth

¹ Astronomy Department, Cornell University, Ithaca, NY 14853.

² Laboratory for High-Energy Astrophysics, NASA Goddard Space Flight Center, Code 662, Greenbelt, MD 20771. NAS/NRC Research Associate.

³ Astronomy Department, University of California, Berkeley, CA 94720.

⁴ Arecibo Observatory, HC3 Box 53995, PR 00612.

⁵ Currently at Numerical Technologies, 70 West Plumeria Drive, San Jose, CA 95134-2134.

⁶ <http://www.naic.edu/~aofm>.

with a programmable number of lags and time resolution (Dowd, Sisk, & Hagen 2000).⁷

We mapped the field of this pulsar with the Very Large Array (VLA) in A array at 1.4 GHz using a 50 MHz bandwidth and 12 minute integration. At the time of these observations in 1999 July, the pulsar's position was known to approximately 3' (i.e., one 1.4 GHz beam at Arecibo). We detected a continuum source in this observation with a flux density of 2.26 ± 0.17 mJy at a position (J2000.0 = MJD 51385) of R.A. = $17^{\text{h}}40^{\text{m}}25^{\text{s}}.958 \pm 0.003$, decl. = $+10^{\circ}00'05''.91 \pm 0.05$ (derived using the AIPS task *jmfitt*). Subsequent timing confirmed that this source was indeed the pulsar. The quoted 0.17 mJy error on the flux density does not account for fluctuations owing to interstellar scintillation, which are significant. No other point sources were within 3' of the position of PSR J1740+1000, but two sources, both listed in the NRAO VLA Sky Survey catalog (Condon et al. 1998), were within 5'. No filamentary or bow-shock structures were visible in the radio maps.

3. ANALYSIS AND RESULTS

3.1. Pulsar Properties

We began regular timing observations of PSR J1740+1000 in 1999 May at an observing frequency of 430 MHz. After we established an initial timing model and a more accurate position, timing observations were carried out predominantly at 1.4 GHz, where higher signal-to-noise profiles could be acquired. Pulse times of arrival (TOAs) were calculated by cross-correlating dedispersed profiles with templates, using separate templates made from the addition of many individual pulses at both 430 MHz and 1.4 GHz. These TOAs were then analyzed using the TEMPO software package⁸ (Taylor & Weisberg 1989). A timing solution for PSR J1740+1000, derived from 684 TOAs spanning almost 2 yr, is presented in Table 1. With a spin-down age τ_s of 114 kyr, this pulsar is younger than

⁷ <http://www.naic.edu/~wapp>.

⁸ <http://pulsar.princeton.edu/tempo>.

TABLE 1

OBSERVED AND DERIVED PARAMETERS FOR PSR J1740+1000

Parameter	Value
Right ascension (J2000)	17 40 25.950(5)
Declination (J2000)	+10 00 06.3(2)
Galactic longitude (deg)	34.01067(7)
Galactic latitude (deg)	+20.26828(5)
Barycentric period (s)	0.154087174313(2)
Period derivative ($10^{-15} \text{ s s}^{-1}$)	21.4651(2)
Epoch (MJD)	51662
Dispersion measure (pc cm^{-3})	23.85(5)
Rotation measure (rad m^{-2})	23.8(2.8)
Timing data span (MJD)	51310–52014
Number of TOAs	684
Postfit rms timing residual (μs)	952
Flux density at 0.4 GHz (mJy)	3.1(2)
Flux density at 1.4 GHz (mJy)	9.2(4)
Mean spectral index	0.9(1)
Distance ^a (kpc)	1.4
Dipole magnetic field strength ^b B (10^{12} G)	1.8
Characteristic age ^b τ_s (kyr)	114

NOTE.—Units of right ascension are hours, minutes, and seconds, and units of declination are degrees, arcminutes, and arcseconds. Figures in parentheses represent 1σ uncertainties in least-significant digits quoted.

^a Calculated using the TC93 model.

^b Calculated using the standard magnetic dipole formulae namely, $B = 3.2 \times 10^{19} \sqrt{\dot{P}}$ Gauss; $\tau = P/2\dot{P}$ (Manchester & Taylor 1977).

95% of all pulsars listed in the Princeton Pulsar Catalog (Taylor, Manchester, & Lyne 1993). Given the 23.85 pc cm^{-3} DM of this pulsar, we use the Taylor & Cordes (1993, hereafter TC93) model for Galactic electron density to derive a distance of 1.4 ± 0.4 kpc.

Repeated observations have shown that this pulsar's average flux density appears to be *higher* at 1.4 GHz than at 430 MHz, with a spectral index α (where $S_\nu \propto \nu^\alpha$ for an observing frequency ν) of 0.9 ± 0.1 . We note that the spectral index estimate is highly contaminated by interstellar scintillation, especially refractive interstellar scintillation (RISS) that is uncorrelated between the two frequencies. We




FIG. 1.—Folded pulse profiles for PSR J1740+1000 at 430 MHz (*left*) and 1.4 GHz (*right*). These profiles were formed by co-adding 17,000 and 11,000 s of PSPM data at 430 MHz and 1.4 GHz, respectively. For all observations, data were taken across an 8 MHz bandwidth and dedispersed off-line using a DM of 23.85 pc cm^{-3} .

note in passing that flat spectral indices may be characteristic of young pulsars (Lorimer et al. 1995), though the Crab is a counterexample to this.

As shown in Figure 1, the 1.4 GHz profile is composed predominantly of two conal pulse components, while at 430 MHz a core component is apparent. In Figure 2 we present a polarization profile based on WAPP observations carried out at 1475 MHz. The degree of linear polarization is 96%. Other pulsars with degrees of linear polarization greater than 80% include J0134–2937, B0136+57, J0631+1036, B0656+14, J0742–2822, B0740–28, B0833–45, B0906–49, B1259–63, J1359–6038, J1643–1224, B1737–30, B1913+167, and B1929+10 (McCulloch et al. 1978; Wu et al. 1993; Manchester & Johnston 1995; Xilouris & Kramer 1996; Zepka et al. 1996; Gould & Lyne 1998; Manchester, Han, & Qiao 1998; Weisberg et al. 1999). With some exceptions (most notably B1643–12 and B1913+167), these pulsars have similar ages and/or periods to PSR J1740+1000, supporting a possible correlation between the degree of linear polarization and period and/or age, as has been noted before (Manchester 1971; McCulloch et al. 1978; von Hoensbroech, Kijak, & Krawczyk 1998). From data taken over a 100 MHz bandwidth at 1.475 GHz, we derive a rotation measure of $23.8 \pm 2.8 \text{ rad m}^{-2}$. For a DM of 23.85 pc cm^{-3} , this translates to an average magnetic field along the line of sight of $1.2 \pm 0.1 \mu\text{G}$.

Because the pulsar’s young age, estimated distance, and high Galactic latitude suggest a high space velocity, a proper motion measurement is very desirable. However, the pulsar’s large timing residuals (see Fig. 3) currently prohibit this. The scatter in these residuals is likely dominated by the




FIG. 2.—Polarization profile of PSR J1740+1000 at 1475 MHz. The upper panel shows the total intensity I (solid line) and linearly polarized intensity L (dashed line). These lines are nearly indistinguishable owing to the high degree of linear polarization. Because this profile was formed from only 4 minutes of data, the pulse shape has not yet converged to that shown in Fig. 1. The inset plot shows L vs. I across the pulse. The lower panel shows the polarization position angle. We do not show the circularly polarized power because it is influenced by the unknown cross-coupling of the feed system.




FIG. 3.—Timing residuals for 684 TOAs spanning MJDs 51310 to 52014 for the timing model in Table 1, which includes a fit for P , \dot{P} , DM, R.A., and decl. Fitting for proper motion or a frequency second derivative improves the fit somewhat, but since we are unsure of the origin of the long-term trend, we do not include it here. Open and closed circles denote measurements at 430 MHz and 1.4 GHz, respectively. TOA uncertainties range from 0.5 to 5 ms at 430 MHz and from 0.3 to 3 ms at 1.4 GHz, varying by an order of magnitude or more because of scintillation and profile shape variations. Note that, despite the variation in TOA uncertainty, all TOAs were given equal weight in the least-squares fit to minimize biasing from profile shape variations between individual scans.

considerable pulse-to-pulse jitter exhibited by this pulsar, as shown in Figure 4. At 1.4 GHz, both single-peaked and multiple-peaked pulse shapes are recognizable, with the single-peaked modes generally being stronger. Sometimes a precursor, arriving ~ 10 ms before the main pulse, is present. More infrequently, a stronger postcursor, arriving ~ 10 ms after the main pulse, is apparent. An autocorrelation analysis shows that pulses are intrinsically modulated on timescales ~ 1 s (i.e., 6–7 pulses), as shown in Figure 5. This timescale is similar at both 1.4 GHz and 430 MHz. The modulation index m , corrected for additive off-pulse noise,

$$m = \frac{(\sigma_{\text{on}}^2 - \sigma_{\text{off}}^2)^{1/2}}{\langle I_{\text{on}} \rangle - \langle I_{\text{off}} \rangle}, \quad (1)$$

where σ_{on} and σ_{off} are the on- and off-pulse rms and $\langle I_{\text{on}} \rangle$ and $\langle I_{\text{off}} \rangle$ are the average on- and off-pulse intensities, is calculated across the pulse and shown in Figure 6. The shape is similar to that calculated for other strongly modulated pulsars, with greater modulation on the outer edges of the pulse (Backer 1973; Bartel, Sieber, & Wolszczan 1980).

3.2. Velocities

We define the z -velocity of an object of “kinematic” age t born at height z_0 above the Galactic plane as

$$V_z = \frac{z - z_0}{t}, \quad (2)$$

for a present-day height of $z = D \sin b$. For PSR J1740+1000, assuming $z_0 = 0$, $D = 1.4$ kpc and $t = \tau_s$, we find $V_z = 4160 \text{ km s}^{-1}$. Even for a z_0 of 0.3 kpc, over twice the best-fit scale height of the pulsar progenitor population (Cordes & Chernoff 1998; Arzoumanian et al. 2001),




FIG. 4.—Sequence of 100 single pulses (i.e., ~ 15 s of data) from PSR J1740+1000. These data were taken using the WAPP at 1.475 GHz across a 100 MHz bandwidth. The time resolution is 0.8 ms.

a velocity of 1500 km s^{-1} is required for PSR J1740+1000 to have reached its present-day z of 0.48 kpc in its 114 kyr lifetime. Note that, even for velocities as large as 4000 km s^{-1} , contamination of τ_s due to the Shklovskii effect (Shklovskii 1970) is insignificant, producing an apparent \dot{P} of roughly 1% of the true value. We later discuss the possibility that the kinematic age is much larger than the spin-down age. In Figure 7 we show V_z (for z_0 of zero) for all pulsars in the Princeton Pulsar Catalog (Taylor et al. 1993) with latitude greater than 10° . We chose this latitude cutoff to exclude pulsars within the progenitor scale height and pulsars seen through much of the Galactic disk, where uncertainties in TC93 are greater.




FIG. 5.—Autocorrelation function (ACF) for a dedispersed time series of period-averaged intensities for PSR J1740+1000. These data were taken with the AOFM at 1.4 GHz across a 10 MHz bandwidth. The lower plot shows only the first 16 s. The variation with timescale ~ 200 s is caused by diffractive interstellar scintillation (DISS), while the ~ 1 s variation is due to intrinsic pulse-to-pulse fluctuations.

To relate an object's z -velocity to its transverse velocity along of the line of sight, we adopt an (x, y, z) coordinate system such that x, y define the Galactic plane, with x toward longitude $l = 90^\circ$ and y toward $l = 180^\circ$, and z is toward the North Galactic pole. We then let \hat{n} , \hat{l} , and \hat{m} be unit vectors, with \hat{n} along the line of sight to the pulsar, \hat{l} and \hat{m} transverse to the line of sight, and $\hat{l} \cdot \hat{z} = 0$, as in equation (28) of Cordes & Rickett (1998, hereafter CR98).




FIG. 6.—Inner 20% of the 1.475 GHz pulse profile, along with the corresponding modulation index, for the full 1200 s observation from which the sequence in Fig. 4 was taken.




FIG. 7.— V_z for all pulsars in the Princeton Pulsar Catalog (Taylor et al. 1993) with Galactic latitudes $|b| > 10^\circ$. Note the logarithmic scaling of the y-axis. The value for PSR J1740+1000 is anomalously high. Pulsar distances were calculated using TC93, except in cases of a more accurate parallax measurement. We note that V_z can be quite different from measured transverse velocities. For instance, PSR B0919+06 has a V_z of 1400 km s^{-1} but an astrometrically determined transverse velocity of $505 \pm 80 \text{ km s}^{-1}$ (Chatterjee et al. 2001). We also note that imposing a latitude cutoff omits pulsars such as PSR B0531+21 and PSR B1509–58, which have V_z s of 1.6×10^5 and $5.6 \times 10^4 \text{ km s}^{-1}$, respectively, but, with known supernova remnant associations, were certainly born at $|z_0| > 0$.

The pulsar’s true space velocity in the z direction is composed of radial and perpendicular components such that

$$V_z = V_{r,z} + V_{\perp,z} = V_r \sin b + V_{\perp,m} \cos b, \quad (3)$$

where $V_{r,z}$ and $V_{\perp,z}$ are the z components of the pulsar’s radial and transverse velocity, respectively, and $V_{\perp,m}$ is the transverse velocity in the \hat{m} direction. Assuming no radial velocity and $z_0 = 0$, we find $V_{\perp,m} = 4430 \text{ km s}^{-1}$ for PSR J1740+1000. This is much higher than the transverse velocity of 1700 km s^{-1} estimated for the fastest known pulsar, PSR B2224+65 (Cordes, Romani, & Lundgren 1993). However, it is entirely possible that the space velocity is much less than 4430 km s^{-1} , as we discuss below.

3.3. Interstellar Scintillation

One way to get an independent constraint on a pulsar’s velocity is through measurement of its interstellar scintillations (ISS). Because scattering by density inhomogeneities in the ionized interstellar medium causes multiple rays to interfere, it produces frequency structure in pulsar spectra (Scheuer 1968; Rickett 1990). These spectra change significantly on timescales of minutes to hours owing to the high transverse velocities of most pulsars. When the spectrum is monitored over time, the resulting two-dimensional array or “dynamic spectrum” is often dominated by a random pattern of scintillation maxima with characteristic bandwidth and timescale (Rickett 1990). Intensity variations in time and frequency with scales of a few minutes and a few MHz are due to diffractive ISS (DISS), caused by small-scale irregularities in the interstellar plasma. Longer term




FIG. 8.—Two-dimensional ACF for a dynamic spectrum for PSR J1740+1000 taken at a center frequency of 1410 MHz in 1999 May. We also plot one-dimensional slices along the two axes, used to calculate Δv_d (as the half-width at half-max of the cut at zero time lag) and Δt_d (as the half width at e^{-1} of the cut at zero frequency lag). For this epoch, we calculate $\Delta v_d = 85 \text{ s}$ and $\Delta t_d = 1.8 \text{ MHz}$.

intensity variations are caused by RISS, owing to variations in electron density on much larger scales than those responsible for DISS.

A two-dimensional correlation analysis can be used to calculate the scintillation bandwidth Δv_d and the scintillation timescale Δt_d , which quantify the interstellar scintillation properties of a pulsar. In Figure 8 we show an ACF of a dynamic spectrum for PSR J1740+1000. In Table 2 we list diffractive bandwidths and timescales measured from ACFs at various epochs. The average values of these parameters are $\Delta t = 271 \text{ s}$ and $\Delta v = 8.0 \text{ MHz}$, with standard deviations of $\sigma_{\Delta t} = 146 \text{ s}$ and $\sigma_{\Delta v} = 6.6 \text{ MHz}$. The large variance in the scintillation parameters is likely due to a combination of measurement uncertainties (from small numbers of scintillation maxima in a given measurement) and modulation from RISS, which may be enhanced by refraction from the North Polar Spur (see below).

We may use these measurements to estimate the speed of the ISS diffraction pattern with respect to the observer. For

TABLE 2
DISS PARAMETERS MEASURED AT 1.4 GHz
FOR PSR J1740+1000

MJD	Δv_d (MHz)	Δt_d (s)	$V_{\text{ISS},5/3,u}$ (km s^{-1})
51314.....	1.8	85	329
51674.....	2.4	226	147
51722.....	2.0	193	157
51851.....	16.0	372	226
51968.....	5.4	211	231
51986.....	16.4	537	157
52022.....	11.7	273	262

a statistically uniform, Kolmogorov scattering medium, this speed can be calculated as (CR98, eq. [13])

$$V_{\text{ISS},5/3,u} = A_{\text{ISS},5/3,u} \frac{\sqrt{D\Delta v_d}}{v\Delta t_d}, \quad (4)$$

where $A_{\text{ISS},5/3,u} = 2.53 \times 10^4 \text{ km s}^{-1}$ and D , Δv_d , v , and Δt_d have units of kpc, MHz, GHz, and s, respectively. We list $V_{\text{ISS},5/3,u}$ for all epochs in Table 2 and, for a distance of 1.4 kpc, find an average value of 216 km s^{-1} , with a standard deviation of 67 km s^{-1} .

For a uniform, Kolmogorov medium, $V_{\text{ISS},5/3,u}$ will be equal to the pulsar's transverse speed. For other cases, making the reasonable assumption that the speeds of the observer and the medium are small compared to the pulsar's, we may express the transverse speed of the pulsar as (CR98, eq. [23])

$$V_{\perp} = W_C W_{D,\text{PM}} V_{\text{ISS},5/3,u}, \quad (5)$$

where the weighting factors W_C and $W_{D,\text{PM}}$ relate the measured scintillation speed to the pulsar's actual transverse speed. The contribution of W_C , which depends on both the wavenumber spectrum and the distribution of the medium (CR98, eq. [18]), is small (maximum range of 0.9–1.3). We therefore assume it is unity throughout the remaining analysis. The other factor, $W_{D,\text{PM}}$, is determined by the distribution of the scattering material along the line of sight to the pulsar. This is proportional to C_n^2 , the coefficient of the electron-density wavenumber spectrum. An exact expression for $W_{D,\text{PM}}$ for a square law phase structure function is given in CR98 (eq. [25]) as

$$\begin{aligned} W_{D,\text{PM}}(D) &= \left[\frac{2 \int_0^D ds (s/D)(1-s/D) C_n^2(s)}{\int_0^D ds (1-s/D)^2 C_n^2(s)} \right]^{1/2} \\ &= \frac{\text{SM}_{\tau}^{1/2}}{[3\text{SM} - (\text{SM}_{\tau} + \text{SM}_{\theta})]^{1/2}}, \end{aligned} \quad (6)$$

where s , the distance measured from the pulsar to the observer, varies from 0 to the pulsar distance D and

$$\text{SM} = \int_0^D ds C_n^2(s), \quad (7)$$

$$\text{SM}_{\tau} = 6 \int_0^D ds (s/D)(1-s/D) C_n^2(s), \quad (8)$$

$$\text{SM}_{\theta} = 3 \int_0^D ds (s/D)^2 C_n^2(s). \quad (9)$$

define the scattering measure, the scattering measure weighted for pulse broadening, and the scattering measure weighted for angular broadening, respectively.

If we use TC93 to calculate SM , SM_{τ} , and SM_{θ} , and thus the weight factor $W_{D,\text{PM}}$, we find $W_{D,\text{PM}} = 0.85$ and $V_{\perp} = 184 \text{ km s}^{-1}$, far smaller than the kinematic z -velocity calculated using $z_0 = 0$ and $t = \tau_s$. In the following, we consider several ways to reconcile the various constraints on the pulsar's velocity. Some reduce the value of the kinematic z -velocity, while others increase the scintillation estimate of the transverse velocity. We note that, if the actual distance to the pulsar is much less than the distance estimated by TC93, the difference between the velocity estimated from ISS (proportional to \sqrt{D}) and the kinematic velocity (proportional to D) becomes smaller.

4. INTERPRETATION OF J1740+1000'S KINEMATICS

4.1. Large Radial Velocity

An unrealistically large radial velocity is required if we assume that it accounts solely for the difference between the V_z calculated assuming $V_r = 0$ and the ISS speed, which reflects only the transverse components. Letting z be the pulsar's height above the Galactic plane, z_{TC} the height above the plane given the TC93 model distance, and ignoring any contribution to V_z from V_{\perp} , we have

$$\begin{aligned} V_r &\approx \left(\frac{D_{\text{TC}}}{\tau_s} \right) \left(\frac{\tau_s}{t} \right) \left(\frac{z - z_0}{z_{\text{TC}}} \right) \\ &\approx 1.2 \times 10^4 \text{ km s}^{-1} \left(\frac{\tau_s}{t} \right) \left(\frac{z - z_0}{z_{\text{TC}}} \right). \end{aligned} \quad (10)$$

Assuming that a value $V_r \approx 10^3 \text{ km s}^{-1}$ is “reasonable” in comparison with the highest known velocities, it appears that either the pulsar is kinematically much older than the spin-down age such that $t \gg \tau_s$, or the pulsar was born at a relatively high z_0 such that $|z - z_0| \ll z_{\text{TC}}$. However, we later discuss other interpretations where the transverse velocity makes a sizeable contribution to V_z and V_r need not be so large.

4.2. Large Kinematic Age

Depending on the spin-down law for the pulsar and on the evolutionary history of the neutron star's progenitor, the spin-down age τ_s may or may not be a good estimate for the kinematic age t . First, the spin-down age may only approximate the true age of the pulsar. Young pulsars show estimated braking indices $n \equiv \Omega\dot{\Omega}/\Omega^2 < 3$, where $\Omega = 2\pi/P$, implying ages greater than the spin-down age τ_s , which is calculated for $n = 3$. However, a pulsar born with spin period not significantly smaller than its present-day period could be much younger than τ_s . In fact, a statistical comparison of kinematic z -velocities and proper motion derived velocities suggests that, on average, pulsars are about 50% younger than their spin-down ages (Cordes & Chernoff 1998).

A pulsar could be older than τ_s if the magnetic field was not constant, but has been growing over the lifetime of the pulsar, as suggested in Blandford, Applegate, & Hernquist (1983). A measurement of PSR J1740+1000's braking index could provide support for this hypothesis. Recently, Gaensler & Frail (2000) argued that the pulsar PSR B1757–24 was significantly older than τ_s by comparing the pulsar's location to its birthplace, estimated through identification of the geometric center of the associated supernova remnant. The upper bound on the proper motion of the synchrotron nebula around the pulsar suggests an older age. However, we believe that the errors associated with estimating ages based on remnant morphology are too large to allow any general statement about pulsar ages.

Second, even if the pulsar age is well approximated by τ_s , the kinematic age could be significantly larger. For example, the progenitor of J1740+1000 may have been a member of a binary system whose primary exploded long before the observed pulsar was produced, imparting significant space velocities to its remnant star and to the progenitor of J1740+1000, either as an intact binary or as individual objects. A scenario of this type was proposed by Gott, Gunn, & Ostriker (1970) to account for the current

locations of the Crab pulsar (B0531+21) and the nearby, older, and longer period pulsar, B0525+21, with the binary producing both pulsars originating in the I Gem OB association.

4.3. Halo Progenitor

It is conceivable that PSR J1740+1000 may have been born well out of the Galactic plane from a halo-star progenitor, perhaps through a rare accretion-induced collapse (AIC) or merger event (Bailyn & Grindlay 1990; Rasio & Shapiro 1995). If the neutron star formed through accretion induced collapse (AIC) of a halo white dwarf, the expected transverse speed would be $\sim 250 \text{ km s}^{-1}$ (i.e., the characteristic speed of a halo star), in the absence of a supernova-induced kick. This is consistent with the scintillation estimate. Given one object in 114 kyr, a naive estimate of the birthrate of such objects would be $\sim 10^{-5} \text{ yr}^{-1}$. This rate is about 10^3 times smaller than that of disk-born pulsars and about 3–10 times larger than the estimated birthrate of millisecond pulsars (Lorimer et al. 1995; Cordes & Chernoff 1997). Therefore, if AIC produces only large magnetic field objects, the birthrate implied by PSR J1740+1000 would be compatible with birthrates of other neutron star subpopulations. Because we have not accounted for any selection effects in detecting objects produced through AIC, this is of course a very rough estimate.

4.4. Interstellar Medium Structure and DISS

The local ISM has been sculpted by supernovae and stellar winds into shells and loops that could perturb both the pulsar's distance estimate, which is based on the dispersion measure, and the scintillation parameters. If only the distance is in error, reconciliation of the DISS and kinematic velocities implies $D \approx 4 \text{ pc}$ (for $z_0 = 0$), which requires an unrealistically large H II column density within a few parsec of the Sun. For $z_0 \neq 0$ the distance could be larger and a smaller fraction of DM would be needed from an H II region.

As shown in Figure 9, the line of sight to PSR J1740+1000 passes through the North Polar Spur (NPS) and the Gould Belt, an expanding disk of gas and young stars (Stothers & Frogel 1974). The North Polar Spur is the brightest part of the larger radio feature called Loop I, a 116° diameter circular feature on the sky centered at longitude 330° and latitude $+18^\circ$. The NPS rises from the Galactic plane at a longitude of 30° and covers latitudes of 15° – 30° , appearing as a bright, narrow ridge in radio and X-ray maps. Sofue (1977) postulates that the NPS is part of a shock front induced by an explosive event at the Galactic center. However, a local, supernova remnant origin for the NPS is usually assumed (Heiles et al. 1980; Salter 1983; Egger & Aschenbach 1995; Heiles 1998). Distance estimates given this assumption range from 50 to 200 pc (Berkhuijsen




FIG. 9.— $105^\circ \times 105^\circ$ image of the Galaxy at 408 MHz, showing the NPS rising from the Galactic plane (Haslam et al. 1982). The position of PSR J1740+1000 is denoted with a cross. The positions of the three extreme scattering event sources associated with the NPS are denoted with asterisks.

1973; Bingham 1967), while the thickness of the NPS is estimated to be ~ 20 pc (Berkhuijsen 1973). The distance of the Gould Belt in the direction of PSR J1740+1000 is 100–300 pc, also much less than the TC93 distance for PSR J1740+1000.

The discovery of PSR J1740+1000 may have implications for the long-standing question of how the NPS was formed. While it is generally accepted that such shells are produced through the release of energy into the interstellar medium, the age of the NPS is uncertain. Its slow rate of expansion indicates that it is fairly old, but the production of X-rays in the interior poses an apparent contradiction. If these X-rays are due to reheating processes, the NPS could be roughly 2×10^6 yr old and produced by one normal supernova explosion. However, if the X-ray emission is due to just a single event, the age of the NPS could be only 2×10^5 yr (Heiles et al. 1980), roughly the spin-down age of PSR J1740+1000.

Structure in the local ISM like that seen in the NPS and the Gould Belt perturbs both DM and SM for the line of sight to the pulsar. Strong evidence for the diffracting and refracting power of the NPS comes from examining the distribution of extreme scattering event (ESE) sources, extragalactic sources that show extreme variations in their radio light curves (Fiedler et al. 1987). ESEs are generally believed to be caused by strong diffraction and refraction due to enhanced electron density turbulence in the ISM. As shown in Figure 9, three of the 10 known ESE sources (1749+096, 1821+107, and 1741–038) have lines of sight that pass near or through the North Polar Spur (Fey, Clegg, & Fiedler 1996).

To proceed, we follow Chatterjee et al. (2001) by superposing a thin screen at distance D_s from the pulsar on a smoothly varying medium such as is modeled by TC93. This yields a DISS velocity correction factor

$$W_{D,PM}(D) = \frac{[6(D_s/D)(1 - D_s/D)\Delta SM + SM_\tau]^{1/2}}{[3(1 - D_s/D)^2\Delta SM + 3SM - (SM_\tau + SM_\theta)]^{1/2}}, \quad (11)$$

where ΔSM is the additional scattering measure contributed by the screen. Given the standard definitions of SM and DM (i.e., eq. [9] and $DM = \int_0^D n_e(s)ds$, where n_e is the electron density in the medium), we may relate ΔSM and ΔDM , the differential DM contributed by the screen. For a screen of width Δs ,

$$\Delta SM = (1/3)(2\pi)^{-1/3} K_u F_s \frac{(\Delta DM)^2}{\Delta s}, \quad (12)$$

where $K_u = 10.2 \times 10^{-3} \text{ m}^{-20/3} \text{ cm}^6 \text{ kpc pc}^{-1}$ is a unit conversion factor, and the units of ΔSM , ΔDM , and Δs are $\text{kpc m}^{-20/3}$, pc cm^{-3} , and pc, respectively. As defined in TC93, F_s is a dimensionless “fluctuation parameter” [i.e., $C_n^2(s) \propto F_s n_e^2(s)$], which is found to range from 0.1 to 100 for the various Galactic components of the TC93 model. We note that while DM and SM are correlated, the excess electron density from a screen could cause a large fractional change in scattering measure but a much smaller fractional change in dispersion measure.

Given the SM, SM_τ , and SM_θ estimated by TC93 along the line of sight to the pulsar, $W_{D,PM}$ will increase with increasing ΔSM , asymptoting to a value of $[2(D_s/D)/(1 - D_s/D)]^{1/2}$. For the measured ISS velocity to be made

consistent with $z_0 = 0$ (assuming $t = \tau_s$ and ignoring the radial velocity contribution to V_z), a $W_{D,PM}$ of 20.5 is needed. This requires an unreasonable $D_s/D = 0.995$, or, for a pulsar distance of 1.4 kpc, a screen that is only 7 pc away from the Earth. To calculate the minimum z_0 for a range of realistic model parameters, we constrain ΔSM such that the resultant ΔDM is less than half of the pulsar’s DM. We note that $\Delta DM = 11.9 \text{ pc cm}^{-3}$ corresponds to a screen electron density of 0.6 cm^{-3} for a screen thickness of 20 pc, roughly consistent with the constraint $n_e \leq 0.4 \text{ cm}^{-3}$ derived by Heiles et al. (1980) from NPS rotation and emission measures. Assuming Δs , D_s , D , and F of 20 pc, 130 pc, 1.4 kpc, and 10, we calculate a minimum z_0 of 0.34 kpc. Allowing screen distances as small as 50 pc, a pulsar distance as low as 0.6 kpc, a screen thickness as small as 10 pc, and a fluctuation parameter F_s as large as 1000, the lowest height at which the pulsar could have been born is 0.10 kpc, consistent with estimated distances to the NPS. Note that these minimum values of z_0 depend on the assumption that there is no radial contribution to the transverse velocity and that the kinematic age is equal to the spin-down age.

4.5. Refractive Modulation of DISS

Another effect of a strong phase screen is that it can refract the radiation into a cone of received directions that is not centered on the direct ray path between pulsar and observer. The result is to decrease the DISS bandwidth while leaving the DISS timescale unchanged (Cordes et al. 1986). It is therefore possible that strong RISS is contaminating our measured Δv_d , causing an underestimate of $V_{ISS,5/3,u}$. For the $V_{ISS,5/3,u}$ estimate to agree with the V_\perp estimated assuming $V_r = 0$, $z_0 = 0$ and $t = \tau_s$, the actual bandwidth, unperturbed by RISS, would have to be an impossibly high 3360 MHz. However, while RISS may not be solely responsible for the discrepancy between the pulsar’s age and height above the plane and the measured scintillation parameters, it could be biasing our results. The distribution of ESE sources provides strong evidence for the refracting properties of the NPS.

5. PROSPECTS FOR ASTROMETRY

The most robust way to determine the velocity and possible origin of PSR J1740+1000 is through measurement of a proper motion and parallax. An estimated distance of 1.4 kpc corresponds to a parallax of 0.71 mas. A transverse speed of 10^3 km s^{-1} for a pulsar at that distance corresponds to a proper motion $\sim 50 \text{ mas yr}^{-1}$.

Because the timing residuals of the pulsar are large, the prospects for measuring the proper motion through VLBI are much better than through timing. Given the current rms timing residuals, we estimate that we could expect to measure a 150 mas yr^{-1} proper motion to about 10σ with 2 more yr of timing data. However, if there is significant timing noise, as is expected for a young pulsar, a proper motion may be unmeasurable. A parallax measurement through timing is not feasible.

The current Very Large Baseline Array (VLBA) has sufficient angular resolution to easily measure the anticipated proper motion. However, the $\sim 2 \text{ mJy}$ 1.4 GHz average flux density of this pulsar necessitates more collecting area than is available with the VLBA alone. With the addition of Arecibo and the Green Bank Telescope to the array, a proper motion should be measurable within several months. Moreover, we have seen episodes of scintillation

enhancement of the pulsar's flux density for sustained times (~ 1 hr) by a factor of 10, which will boost the astrometric sensitivity of VLBI measurements. As in the case of PSR B0919+06 (Chatterjee et al. 2001), the situation may also be aided by the presence of two nearby calibrators with separations and flux densities of 4' and 5' and 2.7 and 2.6 mJy, respectively (Condon et al. 1998). The in-beam calibrators may also allow measurement of the pulsar's parallax, especially if the pulsar is associated with the nearby NPS or Gould Belt.

6. HIGH-ENERGY EMISSION

Because of its youth and proximity, PSR J1740+1000 is an excellent candidate for detection at high energies. Assuming that X-ray flux is proportional to \dot{E}/D^2 (Becker & Trümper 1997), where \dot{E} is the spin-down energy loss rate, PSR J1740+1000 has the 10th highest predicted X-ray flux out of 547 pulsars in the Princeton Pulsar Catalog (Taylor et al. 1993). Upcoming observations with *Chandra* may reveal the existence of thermal or magnetospheric radiation from the star. Nebular emission from a compact synchrotron nebula or a pulsar wind nebula bow shock, especially likely if the velocity of the pulsar is high, should be detectable.

This pulsar should also be a strong gamma-ray source. Given the model of McLaughlin & Cordes (2000), PSR J1740+1000 ranks 20th out of 547 pulsars for OSSE ($50 \leq E \leq 200$ keV) flux and 10th out of 547 pulsars for EGRET ($E > 100$ MeV) flux. Because this pulsar was not detected as an EGRET point source (Hartman et al. 1999), we can place an upper limit on its gamma-ray flux of 8.7×10^{32} ergs s^{-1} kpc^{-2} , just above the flux of 8.1×10^{32} ergs s^{-1} kpc^{-2} predicted by the McLaughlin & Cordes model. Unfortunately, even in the closest EGRET viewing period, PSR J1740+1000 is 20° off-axis. Still, we did search this and other viewing periods for periodicities over a range of periods and period derivatives close to that predicted by our current timing model. To do this, we first created a barycentered time series of photon arrival times given the known position of the pulsar. Photons are then individually weighted by the EGRET point-spread function and folded for each trial ephemeris. The point-spread function weighting procedure produces a dramatic improvement in signal-to-noise ratio. For each profile, we use the Bayesian procedure presented in McLaughlin et al. (1999) to calculate the probability of a pulse of unknown width and phase. Using this method, we detect the known EGRET pulsars with high signal-to-noise. While, not surprisingly, we find

no convincing pulsed signatures for PSR J1740+1000, this pulsar should be easily detectable, modulo any beaming effects, with the *Gamma-Ray Large Area Space Telescope* (*GLAST*), with a projected sensitivity of 3×10^{31} ergs s^{-1} kpc^{-2} . Gamma-ray observations of this pulsar will be aided greatly by its high latitude, where the Galactic gamma-ray background is much smaller than in the Galactic plane.

7. CONCLUSIONS

PSR J1740+1000 is a young pulsar whose study will yield important information about pulsar velocities and the local interstellar medium. As a young object at a high Galactic latitude, its existence may signify a complex progenitor history, either from a Galactic disk population or from a halo population. Alternatively, it may simply have an especially high velocity.

Using simple models, it is difficult to reconcile the measured scintillation properties of this pulsar with its age and height above the Galactic plane. While it is of course possible that this pulsar was indeed born out of the Galactic plane, there are several ways in which the measured ISS parameters could be reconciled with a birth in the midplane of the Galaxy. Clearly the best way to resolve this issue is through measurement of the proper motion and parallax. With the upcoming addition of Arecibo and the Greenbank Telescope to the VLBA, a proper motion will be measurable within a few months. Measurement of the parallax will rely heavily on the quality of nearby calibrator sources and exploitation of strong scintillation maxima that boost the apparent flux density of the pulsar by a factor of 10 or more. We are also planning high-energy observations with *Chandra* and *GLAST*, as, because of its youth and proximity, the pulsar should be a strong high-energy source.

We thank Shami Chatterjee for useful discussions and the anonymous referee for an insightful comment. This work was partially supported by NSF grant AST 98-19931. The work was also supported by the National Astronomy and Ionosphere Center, which is operated by Cornell University under cooperative agreement with the National Science Foundation (NSF). Part of this work was performed while Z. A. held a National Research Council Research Associateship Award at NASA/GSFC. We acknowledge the use of NASA's SkyView facility⁹ located at NASA Goddard Space Flight Center.

⁹ <http://skyview.gsfc.nasa.gov>.

REFERENCES

- Arzoumanian, Z., Chernoff, D. F., & Cordes, J. M. 2001, ApJ, submitted (astro-ph/0106159)
 Backer, D. C. 1973, ApJ, 182, 245
 Bailyn, C. D., & Grindlay, J. E. 1990, ApJ, 353, 159
 Bartel, N., Sieber, W., & Wolszczan, A. 1980, A&A, 90, 58
 Becker, W., & Trümper, J. 1997, A&A, 326, 682
 Berkhuysen, E. M. 1973, A&A, 24, 143
 Bingham, R. G. 1967, MNRAS, 137, 157
 Blandford, R. D., Applegate, J. H., & Hernquist, L. 1983, MNRAS, 204, 1025
 Chatterjee, S., Cordes, J. M., Lazio, T. J. W., Goss, W. M., Fomalont, E. B., & Benson, J. M. 2001, ApJ, 550, 287
 Condon, J. J., Cotton, W. D., Greisen, E. W., Yin, Q. F., Perley, R. A., Taylor, G. B., & Broderick, J. J. 1998, AJ, 115, 1693
 Cordes, J. M., & Chernoff, D. F. 1997, ApJ, 482, 971
 Cordes, J. M., & Chernoff, D. F. 1998, ApJ, 505, 315
 Cordes, J. M., Pidwerbetsky, A., & Lovelace, R. V. E. L. 1986, ApJ, 310, 737
 Cordes, J. M., & Rickett, B. J. 1998, ApJ, 507, 846
 Cordes, J. M., Romani, R. W., & Lundgren, S. C. 1993, Nature, 362, 133
 Dowd, A., Sisk, W., & Hagen, J. 2000, in IAU Colloq. 177, Pulsar Astronomy—2000 and Beyond, ed. M. Kramer, N. Wex, and R. Wielebinski (ASP Conf. Ser. 202; San Francisco: ASP), 275
 Egger, R. J., & Aschenbach, B. 1995, A&A, 294, L25
 Fey, A. L., Clegg, A. W., & Fiedler, R. L. 1996, ApJ, 468, 543
 Fiedler, R. L., Dennison, B., Johnston, K. J., & Hewish, A. 1987, Nature, 326, 675
 Gaensler, B. M., & Frail, D. A. 2000, Nature, 406, 158
 Gott, J. R. I., Gunn, J. E., & Ostriker, J. P. 1970, ApJ, 160, L91

- Gould, D. M., & Lyne, A. G. 1998, *MNRAS*, 301, 235
Hartman, R. C., et al. 1999, *ApJS*, 123, 79
Haslam, C. G. T., Stoffel, H., Salter, C. J., & Wilson, W. E. 1982, *A&AS*, 47, 1
Heiles, C. 1998, in *IAU Colloq. 506, The Local Bubble and Beyond*, ed. D. Breitschwendt, M. J. Freyberg, & J. Trümper (Berlin: Springer), 229
Heiles, C., Chu, Y.-H., Troland, T. H., Reynolds, R. J., & Yegingil, I. 1980, *ApJ*, 242, 533
Lommen, A. N., Zepka, A., Backer, D. C., McLaughlin, M., Cordes, J. M., Arzoumanian, Z., & Xilouris, K. 2000, *ApJ*, 545, 1007
Lorimer, D. R., Bailes, M., & Harrison, P. A. 1997, *MNRAS*, 289, 592
Lorimer, D. R., Yates, J. A., Lyne, A. G., & Gould, D. M. 1995, *MNRAS*, 273, 411
Lyne, A. G., & Lorimer, D. R. 1994, *Nature*, 369, 127
Manchester, R. N. 1971, *ApJS*, 23, 283
Manchester, R. N., Han, J. L., & Qiao, G. J. 1998, *MNRAS*, 295, 280
Manchester, R. N., & Johnston, S. 1995, *ApJ*, 441, L65
McCulloch, P. M., Hamilton, P. A., Manchester, R. N., & Ables, J. G. 1978, *MNRAS*, 183, 645
McLaughlin, M. A., Arzoumanian, Z., & Cordes, J. M. 2000, in *ASP Conf. Ser. 202, Pulsar Astronomy—2000 and Beyond*, ed. M. Kramer, N. Wex, & R. Weilebinksi (San Francisco: ASP), 41
McLaughlin, M. A., & Cordes, J. M. 2000, *ApJ*, 538, 818
McLaughlin, M. A., Cordes, J. M., Hankins, T. H., & Moffett, D. A. 1999, *ApJ*, 512, 929
Rasio, F. A., & Shapiro, S. L. 1995, *ApJ*, 438, 887
Rickett, B. J. 1990, *ARA&A*, 28, 561
Salter, C. J. 1983, *Bull. Astron. Soc. India*, 11, 1
Scheuer, P. A. G. 1968, *Nature*, 218, 920
Shklovskii, I. S. 1970, *Soviet Astron.*, 13, 562
Sofue, Y. 1977, *A&A*, 60, 327
Stothers, R., & Frogel, J. A. 1974, *AJ*, 79, 456
Taylor, J. H., & Cordes, J. M. 1993, *ApJ*, 411, 674
Taylor, J. H., Manchester, R. N., & Lyne, A. G. 1993, *ApJS*, 88, 529
Taylor, J. H., & Weisberg, J. M. 1989, *ApJ*, 345, 132
von Hoensbroech, A., Kijak, J., & Krawczyk, A. 1998, *A&A*, 334, 571
Weisberg, J. M., et al. 1999, *ApJS*, 121, 171
Wu, X., Manchester, R. N., Lyne, A. G., & Qiao, G. 1993, *MNRAS*, 261, 630
Xilouris, K. M., & Kramer, M. 1996, in *IAU Colloq. 160, Pulsars: Problems and Progress*, ed. S. Johnson, M. A. Walker, & M. Bailes, (ASP Conf. Ser. 105; San Francisco: ASP), 245
Zepka, A., Cordes, J. M., Wasserman, I., & Lundgren, S. C. 1996, *ApJ*, 456, 305