



UvA-DARE (Digital Academic Repository)

MAXI J1807+132 is fading in both X-rays and optical wavelengths

Armas Padilla, M.; Wijnands, R.; Degenaar, N.; Munoz-Darias, T.; Jimenez-Ibarra, F.; Mata Sanchez, D.; Casares, J.; Charles, P.A.

Publication date

2017

Document Version

Final published version

Published in

The astronomer's telegram

License

Unspecified

[Link to publication](#)

Citation for published version (APA):

Armas Padilla, M., Wijnands, R., Degenaar, N., Munoz-Darias, T., Jimenez-Ibarra, F., Mata Sanchez, D., Casares, J., & Charles, P. A. (2017). MAXI J1807+132 is fading in both X-rays and optical wavelengths. *The astronomer's telegram*, 10224.
<http://www.astronomerstelegam.org/?read=10224>

General rights


It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Outside
 GCN
 IAUCs

Other
 ATel on [Twitter](#) and [Facebook](#)
[ATELstream](#)
[ATel Community Site](#)



[[Previous](#) | [Next](#) | [ADS](#)]

MAXI J1807+132 is fading in both X-rays and optical wavelengths

ATel #10224; *M. Armas Padilla (IAC-Tenerife), R. Wijnands, N. Degenaar (U. of Amsterdam), T. Munoz-Darias, F. Jimenez-Ibarra, D. Mata Sanchez, J. Casares (IAC-Tenerife), P. A. Charles (Southampton)*
 on 29 Mar 2017; 13:46 UT
 Credential Certification: [Rudy Wijnands \(radwijnands@gmail.com\)](mailto:radwijnands@gmail.com)

Subjects: Optical, Ultra-Violet, X-ray, Request for Observations, Black Hole, Neutron Star, Transient

Referred to by ATel #: [10245](#)

We have analyzed a new Swift observation (performed on 06:44UT on March 29, 2017) of the recently discovered X-ray transient MAXI J1807+132 [ATEL #[10208](#), #[10215](#), #[10216](#), #[10217](#), #[10221](#), #[10222](#), #[10223](#)].

The average spectrum obtained during this observation has been fitted with a power-law model affected by absorption. The hydrogen column has been fixed to the value of $N_H \sim 2.3 \times 10^{21} \text{ cm}^{-2}$ obtained during the analysis of the previous Swift-XRT observation [ATel #[10216](#)]. The photon index is 2.5 ± 0.2 and the resulting absorbed (unabsorbed) 0.3-10 keV flux is $(9.2 \pm 0.1) \times 10^{-12}$ ($(1.9 \pm 0.1) \times 10^{-11}$) $\text{ergs cm}^{-2} \text{ s}^{-1}$.

This represent a $\sim 75\%$ drop in X-ray flux (0.3 - 10 keV) with respect to the previous Swift-XRT observation (2 days ago; ATel #[10216](#)). Similarly, the UVOT data show also a flux drop (the below magnitudes are given in the AB system):

March 27	March 29
$v > 17.56$	$v > 18.02$
$b = 18.1 \pm 0.3$	$b = 18.6 \pm 0.2$
$u = 18.43 \pm 0.19$	$u = 18.76 \pm 0.16$
$uvw1 = 18.56 \pm 0.13$	$uvw1 = 19.10 \pm 0.13$
$uvm2 = 18.72 \pm 0.13$	$uvm2 = 19.61 \pm 0.14$
$uvw2 = 18.75 \pm 0.09$	$uvw2 = 19.59 \pm 0.11$

The above results suggest that the source may be decaying to quiescence after a very short outburst. If confirmed this behavior is reminiscent of the X-ray transient MAXI J1957+032 (Mata

- Related**
- [10245](#) MAXI J1807+132: Lulin observations find substantial optical variability during the X-ray fading phase
 - [10227](#) ASAS-SN Optical Observations of MAXI J1807+132
 - [10224](#) MAXI J1807+132 is fading in both X-rays and optical wavelengths
 - [10223](#) MAXI J1807+132: MITSuME Akeno Optical Observation
 - [10222](#) Swift/XRT spectral analysis of MAXI J1807+132: possibility of a faint low mass X-ray binary
 - [10221](#) MAXI J1807+132: GTC spectroscopy reveals a LMXB-like optical spectrum.
 - [10217](#) MAXI J1807+132: archival optical variability of Swift counterpart
 - [10216](#) MAXI J1807+132: UV/Optical counterpart, further XRT observations
 - [10215](#) MAXI J1807+132: Swift/XRT localization
 - [10208](#) MAXI/GSC detection of a long-term soft X-ray flaring event MAXI J1807+132

Sanchez et al. 2017, arXiv:1701.03381) and that of the accreting millisecond X-ray pulsar NGC 6440 X-2 (Altamirano et al. 2010, ApJ, 712, L58; Heinke et al. 2010, ApJ, 714, 894). Both those sources have shown recurrent, very short (~week long) outbursts. However, we urge more X-ray observations of MAXI J1807+132 to investigate the further evolution of the outburst of this source.

[**Telegram Index**]

R. E. Rutledge, Editor-in-Chief

`rrutledge@astronomerstelegam.org`

Derek Fox, Editor

`dfox@astronomerstelegam.org`

Mansi M. Kasliwal, Co-Editor

`mansi@astronomerstelegam.org`