



UvA-DARE (Digital Academic Repository)

The neutron-star low-mass X-ray binary H 1658-298 back in quiescence

Parikh, A.; Wijnands, R.; Bahramian, A.; Degenaar, N.; Heinke, C.

Publication date

2017

Document Version

Final published version

Published in

The astronomer's telegram

License

Unspecified

[Link to publication](#)

Citation for published version (APA):

Parikh, A., Wijnands, R., Bahramian, A., Degenaar, N., & Heinke, C. (2017). The neutron-star low-mass X-ray binary H 1658-298 back in quiescence. *The astronomer's telegram*, 10169. <http://www.astronomerstelegam.org/?read=10169>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).


Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Outside
 GCN
 IAUCs

Other
 ATel on [Twitter](#) and [Facebook](#)
[ATELstream](#)
[ATel Community Site](#)

This space for free for your conference.



[[Previous](#) | [Next](#) | [ADS](#)]

The neutron-star low-mass X-ray binary H 1658-298 back in quiescence

ATel #10169; *Aastha Parikh, Rudy Wijnands, Arash Bahramian, Nathalie Degenaar, Craig Heinke*

on 14 Mar 2017; 13:06 UT

Credential Certification: Rudy Wijnands (rudy@space.mit.edu)

Subjects: X-ray, Binary, Neutron Star, Transient

The transient and eclipsing neutron-star low-mass X-ray binary H 1658-298 began its most recent outburst in August 2015 as determined using MAXI (ATel #7943) and we continued to monitor the outburst using Swift/XRT (e.g., ATel #7957, #8046). On 2017 February 15, the source was clearly detected at a luminosity of $L_X \sim 1e36$ erg/s (0.5 - 10 keV; assuming a distance of 10 kpc) but during the subsequent pointing on March 7 the source was not detected in a ~ 1.7 ksec observation. It indicated a count rate upper limit of $2.1e-3$ c/s (0.5 - 10 keV; the exposure time was corrected by 900 seconds to account for the expected eclipse during the observation; determined using the linear ephemeris of Oosterbroek et al. 2001). This strong drop in flux suggested that the outburst had possibly ceased after ~ 1.5 yr.

To confirm that the source truly transitioned into quiescence, we requested 6 ksec of Swift/XRT data, with a 2 ksec observation every day between March 8 and March 10. During each observation the source was not or only marginally detected. We combined the 4 obtained observations (obs ID 00034002072 - 00034002075, March 7 - 10) and the stacked image shows that H 1658-298 was detected, showing the presence of 15 photons within a circular region having a radius 20 arcsec centred on the source. This shows a count rate of $\sim 2.2e-3$ c/s (0.5 - 10 keV) for the source (having corrected the exposure time for the several eclipses expected to have occurred during our observations). To understand what this implies for the temperature of the accretion-heated neutron star, we extracted a spectrum from the stacked data and fitted it with a neutron star atmosphere model (nstamos; Heinke et al. 2006), with a fixed value of $N_h = 2.2e21$ cm⁻² implemented using tbabs. The obtained effective temperature as seen by an observer at infinity is 96 +/- 10 eV and we detected an unabsorbed 0.5 - 10 keV flux of $\sim 1.1e-13$ erg/cm²/s resulting in an associated luminosity (for 10 kpc) of $\sim 1.3e33$ erg/s. The last outburst detection of the source was on February 15 and therefore our temperature estimate was obtained at a maximum of ~ 20 d after the end of the outburst. However, it is likely that this quiescent temperature was obtained closer to the end of the outburst since the source had to decay from $\sim 1e36$ erg/s to $\sim 1e33$ erg/s during that time span. This effective temperature is consistent with the initial temperature measurement of H 1658-298 ~ 40 d after the end of the 1999 - 2001 outburst (Wijnands et al. 2003).

We have triggered our XMM-Newton observation and will request additional Swift and Chandra observations to study the cooling of the accretion-heated neutron star in quiescence, similar to

Related	
10169	The neutron-star low-mass X-ray binary H 1658-298 back in quiescence
8699	Continuing outburst from H 1658-298
8046	Swift follow-up observations of outburst from H 1658-298
7957	Swift/XRT confirmation of activity from H 1658-298, no detection of MAXI J1327-627
7947	Swift/XRT follow up on X-ray burst from SAX J1324.5-63.13
7946	INTEGRAL confirms the detection of renewed activity from the NS transient H 1658-298
7943	MAXI/GSC detection of renewed X-ray activities of SAX J1324.5-6313/MAXI J1327-627 and H 1658-298/MAXI J1702-301

what has been done after its previous outburst (Wijnands et al. 2003, Cackett et al. 2013). Currently the first XMM-Newton observation is scheduled on March 23. More multi-wavelength coverage of the source, especially in optical would aid this study to determine if accretion has indeed ceased and settled into a cold, quiescent disk.

Bahramian, A, et al. 2015, ATel #[7957](#)
Bahramian, A, et al. 2015, ATel #[8046](#)
Cackett, E, et al. 2013, ApJ 774, 131
Heinke, C, et al. 2006, ApJ 644, 1090
Negoro, H, et al. 2015, ATel #[7943](#)
Oosterbroek, T, et al. 2001, A&A, 376, 532
Wijnands, R, et al. 2003, ApJ, 594, 952

[[Telegram Index](#)]

R. E. Rutledge, Editor-in-Chief

rrutledge@astronomerstelegam.org

Derek Fox, Editor

dfox@astronomerstelegam.org

Mansi M. Kasliwal, Co-Editor

mansi@astronomerstelegam.org