



**UvA-DARE (Digital Academic Repository)**

**Chemokine receptor signals: role in migration, invasion and cancer metastasis**

Zeelenberg, I.S.

[Link to publication](#)

*Citation for published version (APA):*

Zeelenberg, I. S. (2004). Chemokine receptor signals: role in migration, invasion and cancer metastasis

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

## Contents

<b>Chapter 1</b>	Introduction	<b>9</b>
<b>Chapter 2</b>	Retention of CXCR4 in the endoplasmic reticulum blocks dissemination of a T cell hybridoma. <i>J. Clin. Invest.</i> 108:269-277 (2001)	<b>49</b>
<b>Chapter 3</b>	Stromal cell-derived factor-1-induced LFA-1 activation during in vivo migration of T cell hybridoma cells requires Gq/11, RhoA, and myosin, as well as Gi and Cdc42. <i>J. Immunol.</i> 166(7):4293-4301 (2001)	<b>61</b>
<b>Chapter 4</b>	Blocking G-protein $\beta\gamma$ dimers reduces desensitization of CXCR4, causing enhanced and persistent migration, independent of PI3-kinase. <i>Submitted for publication</i>	<b>73</b>
<b>Chapter 5</b>	Fusion of docked vesicles, driven by synaptotagmin, is required for chemokine-induced T-cell migration. <i>To be submitted</i>	<b>85</b>
<b>Chapter 6</b>	The chemokine receptor CXCR4 is required for outgrowth of colon carcinoma micrometastases. <i>Cancer Res.</i> 63:3833-3839 (2003)	<b>97</b>
<b>Chapter 7</b>	A putative role for the chemokine receptor CXCR5 in tumor growth of colon carcinoma cells. <i>Preliminary report</i>	<b>107</b>
<b>Summary</b>		<b>117</b>
<b>Samenvatting</b>		<b>121</b>
<b>Dankwoord</b>		<b>127</b>
<b>Curriculum vitae</b>		<b>133</b>
<b>List of publications</b>		<b>135</b>

