



UvA-DARE (Digital Academic Repository)

Chemokine receptor signals: role in migration, invasion and cancer metastasis

Zeelenberg, I.S.

[Link to publication](#)

Citation for published version (APA):

Zeelenberg, I. S. (2004). Chemokine receptor signals: role in migration, invasion and cancer metastasis

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

List of publications

- Van Vugt M.J., Reefman E., **Zeelenberg I.**, Boonen G., Leusen J.H., van de Winkel J.G. The alternatively spliced CD64 transcript FcγRIIb2 does not specify a surface-expressed isoform. *Eur. J. Immunol.* 29(1):143-149 (1999)
- Van Vugt M.J., Kleijmeer M.J., Keler T., **Zeelenberg I.**, van Dijk M.A., Leusen J.H., Geuze H.J., van de Winkel J.G. The FcγRIa (CD64) ligand binding chain triggers major histocompatibility complex class II antigen presentation independently of its associated FcR gamma-chain. *Blood* 94(2):808-817 (1999).
- Johnsson A., **Zeelenberg I.**, Min Y., Hilinski J., Berry C., Howell S.B., Los G. Identification of genes differentially expressed in association with acquired cisplatin resistance. *Br. J. Cancer* 83(8):1047-1054 (2000).
- Soede R.D., **Zeelenberg I.S.**, Wijnands Y.M., Kamp M., Roos E. Stromal cell-derived factor-1-induced LFA-1 activation during in vivo migration of T cell hybridoma cells requires Gq/11, RhoA, and myosin, as well as Gi and Cdc42. *J. Immunol.* 166(7):4293-4301 (2001).
- Zeelenberg I.S.**, Ruuls-Van Stalle L., Roos E. Retention of CXCR4 in the endoplasmic reticulum blocks dissemination of a T cell hybridoma. *J. Clin. Invest.* 108(2):269-277 (2001).
- Zeelenberg I.S.**, Ruuls-Van Stalle L., Roos E. The chemokine receptor CXCR4 is required for outgrowth of colon carcinoma micrometastases. *Cancer Res.* 63(13):3833-3839 (2003).
- Zeelenberg, I.S.**, Roos, E. The role of chemokine receptors, in particular CXCR4, in lymphoma and carcinoma metastasis. In: **CANCER GROWTH AND PROGRESSION** (Dr. H. Kaiser, ed.). Vol. 15, *Integration/Interaction of Oncologic Growth*. (Dr. G.G. Meadows, ed.), Kluwer Acad. Publ. (in press)

