UvA-DARE (Digital Academic Repository)

Effects of gemfibrozil or simvastatin on apolipoprotein-B-containing lipoproteins, apolipoprotein-CIII and lipoprotein(a) in familial combined hyperlipidaemia

Bredie, S.J.H.; Westerveld, H.T.; Knipscheer, H.C.; Bruin, T.; Kastelein, J.J.P.; Stalenhoef, A.F.H.

DOI

[10.1016/0300-2977\(96\)00015-0](https://doi.org/10.1016/0300-2977(96)00015-0)

Publication date

1996

Published in

The Netherlands Journal of Medicine

[Link to publication](#)

Citation for published version (APA):

Bredie, S. J. H., Westerveld, H. T., Knipscheer, H. C., Bruin, T., Kastelein, J. J. P., & Stalenhoef, A. F. H. (1996). Effects of gemfibrozil or simvastatin on apolipoprotein-B-containing lipoproteins, apolipoprotein-CIII and lipoprotein(a) in familial combined hyperlipidaemia. *The Netherlands Journal of Medicine*, 49, 59-67.
[https://doi.org/10.1016/0300-2977\(96\)00015-0](https://doi.org/10.1016/0300-2977(96)00015-0)

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Download date: 26 Apr 2023

ELSEVIER

Netherlands Journal of Medicine 49 (1996) 59–67

The Netherlands
JOURNAL OF
MEDICINE

Original article

Effects of gemfibrozil or simvastatin on apolipoprotein-B-containing lipoproteins, apolipoprotein-CIII and lipoprotein(a) in familial combined hyperlipidaemia¹

S.J.H. Bredie^a, H.T. Westerveld^b, H.C. Knipscheer^c, T.W.A. de Bruin^b,
J.J.P. Kastelein^c, A.F.H. Stalenhoef^{a,*}

^a Department of Medicine, Division of General Internal Medicine, University Hospital Nijmegen, PO Box 9101, 6500 HB Nijmegen, Netherlands

^b Departments of Internal Medicine and Endocrinology, University Hospital Utrecht, Utrecht, Netherlands

^c Centre for Vascular Medicine, Academic Medical Centre, University of Amsterdam, Amsterdam, Netherlands

Received 9 November 1995; revised 19 February 1996; accepted 22 February 1996

Abstract

Background: Familial combined hyperlipidaemia (FCH), characterized by elevated very-low-density lipoprotein (VLDL) and/or low-density lipoprotein (LDL), is associated with an increased prevalence of premature cardiovascular disease. Therefore, lipid-lowering is frequently indicated. **Methods:** We evaluated in a parallel, double-blind randomized fashion the effect of gemfibrozil (1200 mg/day) ($n = 40$) or simvastatin (20 mg/day) ($n = 41$) on lipids, apolipoprotein-B (apo-B)-containing lipoproteins, apo-CIII and lipoprotein(a) [Lp(a)], in 81 well-defined FCH patients. **Results:** While both drugs lowered plasma cholesterol and triglyceride levels, gemfibrozil lowered plasma triglycerides more effectively by reduction of triglycerides in VLDL and LDL, whereas simvastatin was more effective in its reduction of total plasma cholesterol by exclusively decreasing LDL cholesterol. High-density lipoprotein (HDL) increased to an equal extent on both therapies. Total serum apo-B levels were reduced with both drugs; however, gemfibrozil decreased apo-B only in VLDL + IDL, whereas simvastatin decreased apo-B in both VLDL + IDL and LDL. In keeping with a more effective reduction of VLDL particles, a more pronounced reduction of apo-CIII also was observed after gemfibrozil, which correlated with the reduction in plasma triglycerides. Baseline concentrations of Lp(a) showed a wide range in both treatment groups. Median Lp(a) levels increased after simvastatin, but were not affected by gemfibrozil. **Conclusion:** Both therapies exhibited their specific effects, although none of the drugs alone completely normalized the lipid profiles of these patients with FCH. Therefore, the choice of treatment should be based on the most elevated lipoprotein fraction, and in some cases a combination of the two drugs may be indicated.

Keywords: Lipids; Therapeutics; Clinical trial

1. Introduction

Familial combined hyperlipidaemia (FCH) is the most common of the hereditary lipid disorders, with

* Corresponding author. Tel. (+31-24) 3614782; fax: (+31-24) 3541734.

¹ Part of this work was presented at the "Internistendagen" Veldhoven, The Netherlands, 1995 (Neth J Med 1995;46:A68).

an estimated prevalence of 0.5–1.0% in the general population and characterized by a strong predisposition to premature cardiovascular diseases in patients and first-degree relatives [1]. Affected individuals have elevated very-low-density lipoprotein (VLDL) concentrations, low-density lipoprotein (LDL) concentrations, or both. Furthermore, a low high-density lipoprotein (HDL) cholesterol concentration, an elevated apolipoprotein-B (apo-B) concentration [2], and an increased prevalence of atherogenic small dense LDL subfractions are observed [3–5]. Because of the absence of a specific clinical or metabolic marker for the disorder, and because of characteristic variability in the presenting lipid phenotype, family investigation is pivotal to establish the diagnosis FCH in a single patient [2]. FCH seems to be a metabolically and genetically heterogeneous lipid disorder [6] which, in general, is explained by an overproduction of VLDL particles [7,8]. Furthermore, an impaired hydrolytic capacity of lipoprotein lipase (LpL) itself [9], or elevated plasma levels of a LpL inhibitor apo-CIII, possibly linked to genetic polymorphisms in the AI-CIII-AIV gene cluster on chromosome 11 [10,11], may be related to the FCH phenotypes.

In general, elevated concentrations of lipoprotein(a) [Lp(a)] are related to an increased risk for cardiovascular disease, which is independent of the cardiovascular risk associated with elevated lipid levels [12]. Although FCH is not associated with elevated Lp(a) levels [13], Lp(a) concentrations correlate positively with LpL-mediated metabolism of triglyceride-rich lipoproteins [14]. Reduction of the elevated pool of triglyceride-rich lipoproteins, as seen in FCH, may influence plasma Lp(a) levels [15].

Because of the high incidence of cardiovascular diseases in these patients, lipid-lowering treatment should be initiated. If diet and lifestyle changes have insufficient effect, drug therapy may be indicated. For this purpose, the HMG-CoA reductase inhibitor, simvastatin, is highly effective in patients with primary hypercholesterolaemia [16], and fibrates like gemfibrozil with a primary triglyceride-lowering effect [17] are useful.

To date, few comparative data are available on the different effects of these agents in FCH. In the present study we describe the baseline levels of lipids, apo-B-containing lipoproteins, apo-CIII and Lp(a) in well-defined patients with FCH, and com-

pare the results of treatment with either gemfibrozil or simvastatin on these parameters.

2. Materials and methods

2.1. Subjects

In total, 81 FCH probands from 3 centres participated in this study. A subgroup of this population has been studied regarding LDL subfraction profiles and LDL oxidizability [5]. In the present study the effects of treatment with either gemfibrozil or simvastatin on serum lipids, lipoproteins, apo-CIII and Lp(a) in all 81 FCH probands are presented.

At the end of the screening period, all FCH patients met the following criteria: (1) a total serum cholesterol > 6.5 mmol/l and triglyceride level between 2.3 and 5.6 mmol/l, without hypolipidaemic drugs and on a standard lipid-lowering diet (30 energy % fat, P/S ratio of 2:1 and a cholesterol intake < 300 mg/day); (2) at least 1 first-degree relative with significant hypercholesterolaemia, hypertriglyceridaemia or both; (3) a positive family history of premature coronary heart disease; (4) total apo-B levels above 1200 mg/l. Furthermore, all patients were > 30 years old and patients with secondary causes for dyslipidaemia or with apolipoprotein phenotype E 2/2 were excluded. The study protocol was approved by the local medical ethical committees, and all patients participated after informed consent was obtained.

2.2. Experimental design

This study was a double-blind trial with a double dummy design, divided into a screening period (weeks –8 to –5), a baseline placebo period (weeks –4 to day 0) and an active treatment period (day 0 to week 12). In the screening period, patients were taken off all hypolipidaemic drugs, receiving a standard lipid-lowering diet. When total plasma cholesterol concentration exceeded 6.5 mmol/l, and plasma triglyceride concentration remained between 2.3 and 5.6 mmol/l, the patients entered the baseline placebo period of the study. During this period, each patient received 2 bottles, one containing placebo matching gemfibrozil and one containing placebo matching

simvastatin. During the active treatment period, patients were randomly assigned to receive either simvastatin 20 mg/day together with a placebo matching gemfibrozil ($n = 41$) or gemfibrozil 1200 mg/day together with a placebo matching simvastatin ($n = 40$). In the present study we compared concentrations of total plasma cholesterol and plasma triglycerides obtained at the end of the entry period, at the end of the placebo period and at two time points (6 and 12 weeks) during the period of active treatment. Furthermore, analyses of lipoproteins, apoproteins and Lp(a) at the end of the placebo-controlled period were compared with the same data at the end of the active treatment period.

2.3. Plasma

Blood samples were obtained after an overnight fast and collected in ethylenediaminetetraacetic acid (EDTA) (1 mg/ml)-containing vacutainers. Plasma was isolated immediately and a saccharose solution (600 mg/ml H₂O) was added to prevent denaturation of VLDL and LDL during freezing; samples were stored at -80°C for 4 to 15 weeks. The analysis of apo-CIII and Lp(a) was performed in the Lipid Research Laboratory, University Hospital of Utrecht; apo-B was determined in the Lipid Research Laboratory, University Hospital of Amsterdam; all other lipid determinations were performed in the Lipid Research Laboratory, University Hospital of Nijmegen.

2.4. Analytical methods

Very-low-density lipoproteins (VLDL) and intermediate-density lipoproteins (IDL) were isolated together by ultracentrifugation at density ≤ 1.019 g/ml for 16 h at 40 000 rpm in a fixed-angle rotor (TFT 45.6 rotor, Kontron, Zürich), in a Beckman L7-55 ultracentrifuge (Beckman, Palo Alto, USA) [18]. After removal of VLDL + IDL, we measured cholesterol and triglycerides in the remaining plasma and in total plasma. HDL was determined from whole plasma by the polyethylene glycol method [19]. All cholesterol and triglyceride measurements were determined by enzymatic methods (Boehringer Mannheim, Mannheim, Germany, cat. no. 237574 and Sera Pak, Tournai, Belgium, cat. no. 6669,

respectively) with a centrifugal analyzer (Multistat III; Instrumentation Laboratory, Lexington, MA). VLDL + IDL cholesterol and triglyceride were calculated by subtraction. Apo-B and apo-AI concentrations in total plasma and apo-B concentrations in the fraction that remained after VLDL + IDL removal were determined by nephelometry [20]. VLDL + IDL-apo-B was calculated by subtraction. Apo-CIII was determined quantitatively by radial immunodiffusion using plates and apo-CIII standards, according to the manufacturer's instructions (Daiichi Pure Chemicals, Ltd, Japan). The diameter of the precipitation ring was measured by an investigator unaware of the specimen's identity. Serial samples from each subject were measured in the same assay. Lp(a) concentrations were determined by measuring the apoprotein(a) moiety in a commercially available solid-phase two-site immunoradiometric assay (IRMA) using 2 different specific anti-apoprotein(a) monoclonal antibodies (Pharmacia, Uppsala, Sweden).

2.5. Statistics

Results are expressed as mean $\pm$ SD and as median for Lp(a). Statistical analysis of alterations within one group of treatment was performed with Wilcoxon's Signed Rank test. Differences in terms of percentage between the two groups of treatment were analyzed with the Mann-Whitney test. A two-tailed probability value of less than 0.05 was considered to be significant. The statistical analysis were performed with procedures available in the SPSS PC + (Statistical Package for the Social Sciences) software package Version 4.0 (SPSS Inc. Chicago, IL, USA)

3. Results

3.1. Participants

After the screening period (4 weeks), participants who exhibited a IIb phenotype (total plasma cholesterol concentration ≥ 6.5 mmol/l and plasma triglyceride concentration ≥ 2.3 mmol/l) were allowed to enter the placebo period. Due to variability in presenting lipid phenotype during the placebo

Fig. 1. Change in plasma concentrations (mmol/l) of total cholesterol (TC) and triglycerides (TG) during the placebo period (week -4 to day 0), and at two time-points (week 6 and week 12) during the period of active treatment with either gemfibrozil (□) ($n = 40$) or simvastatin (■) ($n = 41$) of patients with familial combined hyperlipidaemia. * $P < 0.001$, ** $P < 0.0001$, compared with the baseline level (week 0).

period (4 weeks), 55 patients exhibited a phenotype IIb, 12 patients exhibited a phenotype IIa (total plasma cholesterol concentration ≥ 6.5 and plasma triglyceride concentrations < 2.3 mmol/l), and 14 patients exhibited a phenotype IV (total plasma cholesterol concentration < 6.5 mmol/l and plasma triglyceride concentration ≥ 2.3 mmol/l, respectively) at the end of the baseline placebo period. Description of the two treatment groups, the body

mass index (BMI), and the blood pressure are presented in Table 1.

3.2. Effect of treatment on lipid and lipoprotein concentrations

In Fig. 1 the changes in the mean total plasma cholesterol and plasma triglyceride concentrations during the placebo period and the period of treatment

Table 2

Lipid and lipoprotein concentrations at the end of the placebo period (= before) and after treatment (= after) with either gemfibrozil ($n = 40$) or simvastatin ($n = 41$) of patients with familial combined hyperlipidaemia

	Drug	Before	After	Delta (%)	P
TC	G	7.50 $\pm$ 0.98	6.73 $\pm$ 0.91	-9.5 $\pm$ 13.2	0.0001
	S	7.30 $\pm$ 0.96	5.67 $\pm$ 1.00	-22.4 $\pm$ 9.2 ^{aa}	< 0.0001
TG	G	3.22 $\pm$ 1.22	1.64 $\pm$ 0.59	-45.5 $\pm$ 19.7	< 0.0001
	S	3.20 $\pm$ 1.20	2.63 $\pm$ 0.88	-14.6 $\pm$ 27.8 ^{aa}	0.0003
HDL-C	G	0.91 $\pm$ 0.20	1.07 $\pm$ 0.26	17.4 $\pm$ 17.6	< 0.0001
	S	0.84 $\pm$ 0.19	0.95 $\pm$ 0.21	13.9 $\pm$ 13.1	< 0.0001
VLDL + IDL-C	G	2.43 $\pm$ 0.84	1.15 $\pm$ 0.84	-50.6 $\pm$ 20.6	< 0.0001
	S	2.60 $\pm$ 1.02	1.56 $\pm$ 0.62	-37.6 $\pm$ 20.1 ^{aa}	< 0.0001
VLDL + IDL-TG	G	2.61 $\pm$ 1.18	1.43 $\pm$ 1.15	-44.7 $\pm$ 28.0	< 0.0001
	S	2.59 $\pm$ 1.04	2.08 $\pm$ 0.79	-14.1 $\pm$ 32.6 ^{aa}	0.0008
LDL-C	G	4.08 $\pm$ 0.99	4.29 $\pm$ 1.26	8.3 $\pm$ 41.9	n.s.
	S	3.87 $\pm$ 0.98	3.16 $\pm$ 0.87	-16.3 $\pm$ 21.0 ^{aa}	< 0.0001
LDL-TG	G	0.67 $\pm$ 0.20	0.49 $\pm$ 0.13	-22.9 $\pm$ 19.9	< 0.0001
	S	0.60 $\pm$ 0.12	0.53 $\pm$ 0.12	-10.9 $\pm$ 16.8 ^a	0.0003

Values are presented in mmol/l as mean $\pm$ SD. TC = total plasma cholesterol; TG = plasma triglycerides; VLDL = very-low-density lipoproteins; IDL = intermediate-density lipoproteins; HDL = high-density lipoproteins; LDL = low-density lipoproteins; G = gemfibrozil, S = simvastatin; n.s., not significant. ^a $P < 0.001$, ^{aa} $P < 0.0001$, S versus G.

Table 1

Age, gender, blood pressure and body mass index of 81 patients with familial combined hyperlipidaemia

	Gemfibrozil	Simvastatin
<i>n</i>	40	41
Age (years)	53.4 ± 9.4	50.4 ± 10.8
Gender M/F	26/14	32/9
Systolic blood pressure (mmHg)	131 ± 18	127 ± 13
Diastolic blood pressure (mmHg)	84 ± 9	81 ± 8
Body mass index (kg/m ²)	27.2 ± 3.0	26.6 ± 2.7

Values are presented as mean ± SD.

are presented. At entry and at the end of the placebo period there were no significant differences in mean lipid and lipoprotein concentrations between the two groups, and as shown in Fig. 1, mean lipid levels were unaffected during the placebo period. Lipoprotein levels at the end of the baseline period and after therapy are summarized in Table 2. During the treatment period, the BMI appeared to be constant in both groups. Gemfibrozil affected total triglyceride levels in plasma as well as in the VLDL + IDL and LDL fraction significantly more than simvastatin, whereas simvastatin induced the largest reduction in total plasma cholesterol with an exclusive reduction of LDL-cholesterol. Gemfibrozil tended to increase LDL-cholesterol (+8.3%, not significant). In both groups, the reduction of cholesterol and triglycerides in the VLDL + IDL fraction was responsible for the major decrease on total plasma cholesterol and triglycerides. With both drugs, HDL-cholesterol concentrations increased significantly to an equal extent.

3.3. Effect of treatment on apo-B-containing lipoproteins

Alterations in the apo-B concentrations in total plasma, the VLDL + IDL fraction and the LDL fraction are presented in Fig. 2. Simvastatin and gemfibrozil reduced total plasma apo-B (−25.6 and −13.7%, respectively, but simvastatin was more effective if measured in percentage of change, $P = 0.02$). Simvastatin reduced apo-B in both the VLDL + IDL and the LDL fraction (−35.4 and 16.7%, respectively). Gemfibrozil, on the contrary, reduced apo-B only in the VLDL + IDL fraction (−42.4%; difference simvastatin vs. gemfibrozil in terms of percentage change, $P < 0.0001$), without affecting the LDL fraction (+5.5%, not significant; difference simvastatin vs. gemfibrozil in terms of percentage $P < 0.0001$).

3.4. Effect of treatment on apo-AI and apo-CIII

Changes in apo-AI and apo-CIII are summarized in Table 3. The increase in HDL-cholesterol on both treatments was not accompanied by an increase of apo-AI in both groups. Apo-CIII decreased significantly with both therapies, but more markedly after gemfibrozil. In general, the absolute decrease in Apo-CIII concentrations coincided with the decrease in VLDL + IDL particles, as measured by apo-B concentrations in this fraction. There was a significant correlation between the changes in apo-CIII and the change in plasma triglycerides after gemfibrozil

Fig. 2. Effect of treatment with either gemfibrozil ($n = 40$, left) or simvastatin ($n = 41$, right) on apolipoprotein-B-containing lipoproteins in total plasma, the LDL fraction and the VLDL + IDL fraction of patients with familial combined hyperlipidaemia, before (solid columns) and after (hatched columns) treatment. * $P = 0.001$, ** $P = 0.0001$.

Table 3

Lipoprotein (a) [Lp(a)], apolipoprotein-AI (apo-AI) and apolipoprotein-CIII (apo-CIII) at the end of the placebo period (= before) and after treatment (= after) with either gemfibrozil ($n=40$) or simvastatin ($n=41$) of patients with familial combined hyperlipidaemia

		Drug Before	After	Delta (%)	P
Lp(a)	G	18.0	20.7	10.2 ± 38.9	n.s.
	S	19.6	26.1	14.9 ± 31.1	0.015
Apo-A I	G	131.5 ± 24.7	134.4 ± 24.1	2.9 ± 12.2	NS
	S	136.7 ± 25.4	133.7 ± 22.2	-0.1 ± 18.1	NS
Apo-CIII	G	14.8 ± 3.4	10.6 ± 2.3	-26.0 ± 17.2	< 0.0001
	S	14.7 ± 4.7	13.6 ± 3.6	-6.4 ± 20.3 ^a	0.005

Values are presented in mg/dl as median for Lp(a) and as mean ± SD for apo-AI and apo-CIII. S = simvastatin; G = gemfibrozil; n.s. = not significant. ^a $P < 0.0001$, S versus G.

and simvastatin ($r = 0.69$ and $r = 0.67$; $P = 0.001$, respectively).

3.5. Effect of treatment on Lp(a)

Before and after therapy Lp(a) values showed a wide distribution. Effects on median levels are summarized in Table 3. The effect of gemfibrozil was not significant, 18 out of 40 patients showed a decrease, but conversely, 27 out of 41 patients showed an increase in Lp(a) levels on simvastatin.

4. Discussion

In spite of often mildly elevated lipid levels compared to other lipid disorders, there is a high prevalence of cardiovascular diseases in FCH families. Consequently, family screening and pharmacological intervention in affected individuals may be indicated [21]. The increased tendency to premature cardiovascular disease in these patients must be related to the increased pool of triglyceride-rich apo-B-containing lipoproteins in conjunction with elevated LDL-cholesterol levels [22,23]. In addition, hypertriglyceridaemia, as found in FCH with type IIb and IV phenotypes, is associated with augmented exchange of lipids between VLDL, HDL and LDL, mediated by cholesteryl transfer protein (CETP), which may result in more small dense atherogenic LDL particles in total LDL [24,25]. Elevated apo-CIII concentra-

tions may be associated with hypertriglyceridaemia, because of a possible inhibition of LpL activity by apo-CIII [26]. Furthermore, decreased HDL-cholesterol, apo-AI concentrations and elevated concentrations of Lp(a) may also be involved in the atherogenesis in FCH [13,27].

Recently, large prevention trials and studies with solid cardiovascular end-points have now established that cholesterol-lowering results in a decrease in mortality from cardiovascular disease, and regression of atherosclerotic lesions [28–31]. Although the study populations did not meet the FCH diagnostic criteria, it can be deduced from these results that lipid-lowering may affect mortality and morbidity from cardiovascular events in FCH patients as well, because lipid parameters associated with an increase in cardiovascular risk were also influenced in the present study.

The observed reduction of total plasma cholesterol and plasma triglyceride concentrations with simvastatin (–22.4 and –9.5%, respectively) and gemfibrozil (–14.6 and –45.5%, respectively) after 12 weeks of treatment in our study are in accordance with previous reports [32,33]. To investigate the effect of both therapies on triglyceride-rich apo-B-containing lipoproteins, VLDL and IDL particles were isolated together. This explains the relatively large contribution of VLDL + IDL-cholesterol and the relatively small contribution of LDL-cholesterol to total plasma cholesterol, both before and after treatment. The increase in HDL-cholesterol with gemfibrozil therapy agrees with previous studies as well [32,34], but the increase in HDL on simvastatin, despite a less pronounced reduction of triglycerides, is larger than previously reported [32]. The mean apo-AI concentration was unaffected, despite an increase in HDL-cholesterol and may reflect HDL-cholesterol particles containing less protein [35]. The effects on LDL-cholesterol are comparable with other reports [32,33]. The depletion of triglycerides in the VLDL + IDL fraction, leading to more small dense particles, which are more likely to be converted into LDL particles, is suggested to be a cause of the observed increase in LDL-cholesterol after gemfibrozil [36]. On the other hand, a triglyceride reduction in the LDL particle, more pronounced after gemfibrozil, yields more buoyant LDL subfractions, as we have demonstrated before [5,37]. These buoy-

ant LDL subfractions may be less atherogenic because an increased resistance against oxidative modification has been described [18]. We observed that simvastatin most effectively reduced the total LDL-cholesterol concentration, but did not affect the LDL subfraction profile of FCH patients [5], which was recently demonstrated for pravastatin as well [38].

All patients had moderate to severe elevations of apo-B levels. Both therapies reduced total apo-B, but in accordance with the reduction of total apo-B-containing particles, simvastatin reduced apo-B in both the VLDL + IDL and LDL fraction more effectively than gemfibrozil, which only reduced VLDL + IDL-apo-B. Since VLDL + IDL and LDL particles contain only one molecule of apo-B per particle, a reduction of apo-B observed in these fractions reflects a reduction of the number of potential atherogenic particles per isolated fraction.

It has been suggested that an elevated apo-CIII concentration may be a metabolic marker for FCH. The decrease in the apo-CIII concentrations in this study, however, significantly correlated with the decrease in triglyceride-rich lipoproteins and paralleled the reduction of the particle numbers in the VLDL + IDL fraction. Recently, Patsch et al. found a lower apo-CII/apo-CIII ratio, due to an elevated apo-CIII concentration, in VLDL particles of patients with type IIb hyperlipidaemia, which was related to sequence variations in the minor allele of the AI/CIII/AIV gene cluster on chromosome 11 [39]. From our results, it remains to be seen whether increased apo-CIII concentrations have some role in the pathophysiology of FCH and can serve as a metabolic marker.

So far, the effects of hypolipidaemic drugs on Lp(a) levels have not been very successful. Studies with simvastatin demonstrated either no effect [40] or an increase in Lp(a) concentrations [15]. Although it has been shown that Lp(a) may be bound to triglyceride-rich lipoproteins [41] and that LpL-mediated clearance of triglyceride-rich lipoproteins influences Lp(a) concentrations [14], the mechanism behind this association remains unclear. Synthesis of Lp(a) in the liver may share some characteristics with VLDL synthesis, which is dependent on the amount of free fatty acids to the liver cell [42]. It is still unclear which causes underlie this possible increased supply of free fatty acids to the liver cell in

FCH. Gemfibrozil, however, which may have some effect on the release of fatty acids from adipose tissue, had no significant effect on Lp(a) concentrations.

In conclusion, we directly compared the effects of two drugs with different working mechanisms in a well-defined large group of FCH patients in order to support the choice of treatment for this heterogeneous lipid disorder. None of these agents completely normalized the lipid and lipoprotein profiles. However, each agent has its specific effect in the treatment of subjects with both elevated total plasma cholesterol and triglyceride concentrations. The use of drugs should be based on which lipoprotein fraction is elevated the most during several visits to the out-patient clinic, and probably a combination of gemfibrozil and simvastatin may be the therapy of choice in selected FCH patients with a high risk of cardiovascular disease.

Acknowledgements

We gratefully acknowledge the expert technical assistance of Mrs. Janine Vogelaar, Heidi Hak Lemmers, Anneke Hijmans and Margreet van Linde-Sibenius-Trip.

Drs T.W.A. de Bruin and J.J.P. Kastelein are senior clinical investigators of the Dutch Heart Foundation.

This work was supported by research grant No. 92.056 from the Dutch Heart Foundation and by a grant from Merck Sharp and Dohme, Haarlem, The Netherlands.

References

- [1] Goldstein JL, Schrott HG, Hazzard WR, Bierman EL, Motulsky AG. Hyperlipidemia in coronary heart disease. II. Genetic analysis of lipid levels in 176 families and delineation of a new inherited disorder, combined hyperlipidemia. *J Clin Invest* 1973;52:1544–1568.
- [2] Grundy SM, Chait A, Brunzell JD. Familial combined hyperlipidemia workshop. *Arteriosclerosis* 1987;7(2):203–207.
- [3] Hokanson JE, Austin MA, Zambon A, Brunzell JD. Plasma triglyceride and LDL heterogeneity in familial combined hyperlipidemia. *Arterioscler Thromb* 1993;13:427–434.

- [4] Bredie SJH, Kiemeny LA, De Haan AFJ, Demacker PMN, Stalenhoef AFH. Inherited susceptibility determines the distribution of dense low density lipoprotein subfraction profiles in familial combined hyperlipidemia. *Am J Hum Genet* 1996;58:812–822.
- [5] Bredie SJH, de Bruin TWA, Demacker PNM, Kastelein JJP, Stalenhoef AFH. Comparison of gemfibrozil versus simvastatin in familial combined hyperlipidemia and effects on apolipoprotein-B-containing lipoproteins, low-density lipoprotein subfraction profile, and low-density lipoprotein oxidizability. *Am J Cardiol* 1995;75:348–353.
- [6] Kwiterovich POJ. Genetics and molecular biology of familial combined hyperlipidemia. *Curr Opin Lipidol* 1993;4:133–143.
- [7] Kissebah AH, Alfarsi S, Adams PW. Integrated regulation of very low density lipoprotein triglyceride and apolipoprotein-B kinetics in man: normolipemic subjects, familial hypertriglyceridemia and familial combined hyperlipidemia. *Metabolism* 1981;30(9):856–868.
- [8] Stalenhoef AFH, Demacker PNM, Lutterman JA, van't Laar A. Plasma lipoproteins, apolipoproteins, and triglyceride metabolism in familial hypertriglyceridemia. *Arteriosclerosis* 1986;6:387–394.
- [9] Babirak SP, Brown BG, Brunzell JD. Familial combined hyperlipidemia and abnormal lipoprotein lipase. *Arterioscler Thromb* 1992;12:1176–1183.
- [10] Wojciechowski AP, Farrall M, Cullen P, et al. Familial combined hyperlipidaemia linked to the apolipoprotein AI-CIII-AIV gene cluster on chromosome 11q23-q24. *Nature* 1991;349:161–164.
- [11] Ginsberg HN, Le NA, Goldberg IJ, et al. Apolipoprotein B metabolism in subjects with deficiency of apolipoproteins CIII and AI. Evidence that apolipoprotein CIII inhibits catabolism of triglyceride-rich lipoproteins by lipoprotein lipase in vivo. *J Clin Invest* 1986;78:1287–1295.
- [12] Loscalzo J. Lipoprotein(a). A unique risk factor for atherothrombotic disease. *Arteriosclerosis* 1990;10:672–679.
- [13] Fonda M, Da Col PG, La Verde R, et al. Lipoprotein(a) serum concentration in familial combined hyperlipidemia. *Clin Chim Acta* 1993;223:121–127.
- [14] Williams KJ, Flees GM, Petrie KA, Snyder ML, Brocia RW, Swenson TL. Mechanisms by which lipoprotein lipase alters cellular metabolism of lipoprotein(a), low density lipoprotein, and nascent lipoproteins. Roles for low density lipoprotein receptors and heparan sulfate proteoglycans. *J Biol Chem* 1992;267:13284–13292.
- [15] Castro Cabezas M, de Bruin TW, van Linde-Sibenius-Trip M, Kock LA, Jansen H, Erkelens DW. Lipoprotein(a) plasma concentrations associated with lipolytic activities in eight kindreds with familial combined hyperlipidemia and normolipidemic subjects. *Metabolism* 1993;42:756–761.
- [16] Grundy SM. HMG-CoA reductase inhibitors for treatment of hypercholesterolemia. *N Engl J Med* 1988;319:24–33.
- [17] Grundy SM, Vega GL. Fibric acids: effects on lipids and lipoprotein metabolism. *Am J Med* 1987;83:9–20.
- [18] de Graaf J, Hak-Lemmers HLM, Hectors MPC, Demacker PNM, Hendriks JCM, Stalenhoef AFH. Enhanced susceptibility to in vitro oxidation of the dense low density lipoprotein subfraction in healthy subjects. *Arterioscler Thromb* 1991;11:298–306.
- [19] Demacker PNM, Hijmans AG, Vos-Janssen HE, van't Laar A, Jansen AP. A study of the use of polyethylene glycol in estimating cholesterol in high-density lipoprotein. *Clin Chem* 1980;26:1775–1779.
- [20] Lopes-Virella MF, Virella G, Malenkos SB, Colwell JA. Immunonephelometry assay of human apolipoprotein AI. *Clin Chem* 1980;26:1205–1208.
- [21] The recognition and management of hyperlipidaemia in adults: a policy statement of the European Atherosclerosis Society. *Eur Heart J* 1988;9:571–600.
- [22] Assmann G, Schulte H. Relation of high-density lipoprotein cholesterol and triglycerides to incidence of atherosclerotic coronary artery disease (the PROCAM experience). Prospective Cardiovascular Munster study. *Am J Cardiol* 1992;70:733–737.
- [23] Manninen V, Huttunen JK, Heinonen OP, Tenkanen L, Frick MH. Relation between baseline lipid and lipoprotein values and the incidence of coronary heart disease in the Helsinki Heart Study. *Am J Cardiol* 1989;63:42H–47H.
- [24] Karpe F, Tornvall P, Olivecrona T, Steiner G, Carlson LA, Hamsten A. Composition of human low density lipoprotein: effects of postprandial triglyceride-rich lipoproteins, lipoprotein lipase, hepatic lipase and cholesteryl ester transfer protein. *Atherosclerosis* 1993;98:33–49.
- [25] Austin MA, Breslow JL, Hennekens CH, Buring JE, Willett WC, Krauss RM. Low-density lipoprotein subclass patterns and risk of myocardial infarction. *J Am Med Assoc* 1988;260:1917–1921.
- [26] Ito Y, Azrolan N, O'Connell A, Walsh A, Breslow JL. Hypertriglyceridemia as a result of human apo CIII gene expression in transgenic mice. *Science* 1990;249:790–793.
- [27] Tall AR. Plasma high density lipoproteins; metabolism and relationship to atherosclerosis. *J Clin Invest* 1990;86:379–384.
- [28] Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet* 1994;344:1383–1389.
- [29] Shepherd J, Cobbe SM, Ford I, et al. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. West of Scotland Coronary Prevention Study Group. *N Engl J Med* 1995;333:1301–1307.
- [30] Effect of simvastatin on coronary atheroma: the Multicentre Anti-Atheroma Study (MAAS). *Lancet* 1994;344:633–638.
- [31] Jukema JW, Bruschke AV, van-Boven AJ, et al. Effects of lipid lowering by pravastatin on progression and regression of coronary artery disease in symptomatic men with normal to moderately elevated serum cholesterol levels. The Regression Growth Evaluation Statin Study (REGRESS). *Circulation* 1995;91:2528–2540.
- [32] Ojala JP, Helve E, Tikkanen MJ. Treatment of combined hyperlipidemia with lovastatin versus gemfibrozil: a comparison study. *Cardiology* 1990;77(suppl 4):39–49.

- [33] East C, Bilheimer DW, Grundy SM. Combination drug therapy for familial combined hyperlipidemia. *Ann Intern Med* 1988;109:25–32.
- [34] Odman B, Ericsson S, Lindmark M, Berglund L, Angelin B. Gemfibrozil in familial combined hyperlipidaemia: effect of added low-dose cholestyramine on plasma and biliary lipids. *Eur J Clin Invest* 1991;21:344–349.
- [35] Vega GL, Grundy SM. Management of primary mixed hyperlipidemia with lovastatin. *Arch Intern Med* 1990;150:1313–1319.
- [36] Griffin BA, Packard CJ. Metabolism of VLDL and LDL subclasses. *Curr Opin Lipidol* 1994;5:200–206.
- [37] de Graaf J, Hendriks JCM, Demacker PNM, Stalenhoef AFH. Identification of multiple dense LDL subfractions with enhanced susceptibility to in vitro oxidation among hypertriglyceridemic subjects. Normalization after clofibrate treatment. *Arterioscler Thromb* 1993;13:712–719.
- [38] Franceschini G, Cassinotti M, Vecchio G, et al. Pravastatin effectively lowers LDL cholesterol in familial combined hyperlipidemia without changing LDL subclass pattern. *Arterioscler Thromb* 1994;14:1569–1575.
- [39] Patsch W, Sharrett AR, Chen IY, et al. Associations of allelic differences at the A-I/C-III/A-IV gene cluster with carotid artery intima-media thickness and plasma lipid transport in hypercholesterolemic-hypertriglyceridemic humans. *Arterioscler Thromb* 1994;14:874–883.
- [40] Fieseler HG, Armstrong VW, Wieland E, et al. Serum Lp(a) concentrations are unaffected by treatment with the HMG-CoA reductase inhibitor pravastatin: results of a 2-year investigation. *Clin Chim Acta* 1991;204:291–300.
- [41] Cohn JS, Lam CW, Sullivan DR, Hensley WJ. Plasma lipoprotein distribution of apolipoprotein(a) in the fed and fasted states. *Atherosclerosis* 1991;90:59–66.
- [42] Sniderman AD, Cianflone KM. Substrate delivery as a determinant of hepatic apoB secretion. *Arterioscler Thromb* 1993;13:629–636.