



UvA-DARE (Digital Academic Repository)

Study of the Circinus X-1 Broadband Spectrum at Orbital Phases Close to the Apoastron

Iaria, R.; di Salvo, T.; Robba, N.R.; Burderi, L.

Published in:
Astrophysical Journal

DOI:
[10.1086/338507](https://doi.org/10.1086/338507)

[Link to publication](#)

Citation for published version (APA):

Iaria, R., di Salvo, T., Robba, N. R., & Burderi, L. (2002). Study of the Circinus X-1 Broadband Spectrum at Orbital Phases Close to the Apoastron. *Astrophysical Journal*, 567(1), 503-509. DOI: 10.1086/338507

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

STUDY OF THE CIRCINUS X-1 BROADBAND SPECTRUM AT ORBITAL PHASES CLOSE TO THE APOASTRON

R. IARIA,¹ T. DI SALVO,² N. R. ROBBA,¹ AND L. BURDERI³

Received 2001 October 9; accepted 2001 November 6

ABSTRACT

We report on the results of a *BeppoSAX* (1.8–200 keV) observation of the peculiar X-ray binary source Circinus X-1 (Cir X-1) at the orbital phases between 0.61 and 0.63. We find that three components are needed to fit the broadband spectrum: a blackbody component at a temperature of ~ 0.6 keV; a Comptonized component with a seed-photon temperature of ~ 1.2 keV, electron temperature of ~ 6 keV, and optical depth of ~ 1.7 ; and a power-law component dominating the spectrum at energies higher than 20 keV. We interpret the blackbody as the emission from the accretion disk, while the Comptonized component probably comes from a corona surrounding the inner part of the system. This spectrum is different from that observed at the periastron (in our previous work) because of the presence of the blackbody component. We discuss the implications of this difference and the presence of the power-law component.

Subject headings: accretion, accretion disks — stars: individual (Circinus X-1) — stars: neutron — X-rays: general — X-rays: stars

1. INTRODUCTION

Circinus X-1 (Cir X-1) is a peculiar low-mass X-ray binary (LMXB). Generally in LMXBs the companion star has a mass $\lesssim 1 M_{\odot}$ and the orbit is almost circular; contrarily, Cir X-1 has a companion star of $\sim 3\text{--}5 M_{\odot}$ (probably a subgiant; see Johnston et al. 1999), an orbital period of ~ 16.6 days, and an eccentric orbit with $e \sim 0.7\text{--}0.9$ (Murdin et al. 1980; Tauris et al. 1999). It is a widely accepted idea that Cir X-1 is a runaway binary because of its association with the supernova remnant G321.9–0.3 (Clark et al. 1975). The most recent estimation of the distance to the supernova remnant G321.9–0.3, and hence to Cir X-1, is 5.5 kpc (Case & Bhattacharya 1998). Therefore we adopt 5.5 kpc as the distance to the source.

The spectral and fast-timing properties of accreting low-magnetic field neutron stars permit distinguishing them into two classes, Z and atoll sources, using the pattern each source describes in an X-ray color-color diagram (Hasinger & van der Klis 1989). The Z sources are very bright and could have a moderate magnetic field, while the atoll sources have a lower luminosity and probably a weaker magnetic field. The timing analysis of *Rossi X-Ray Timing Explorer* data (Shirey et al. 1996) evidenced that Cir X-1 is more similar to LMXBs of the Z class. However, no quasi-periodic oscillations at kilohertz frequencies were observed in Cir X-1 (Shirey 1996), contrary to all the other Z sources.

Brandt et al. (1996), using *ASCA*/GIS data taken on 1994 August 4–5, found a sudden variation (with a timescale of ~ 20 minutes) of the flux at the periastron phase, where the count rate increased from ~ 30 to ~ 300 counts s^{-1} . In the low-count rate state they obtained a good fit to the 0.6–10 keV spectrum using a partial-covering component, in addition to a two-blackbody model, with a corresponding

hydrogen column of $\sim 10^{24}$ cm^{-2} ; in the high-count rate state the partial covering of the X-ray spectrum was not needed. Because of the similarity of the spectra of Seyfert 2 galaxies to Compton-thin tori, Brandt et al. (1996) suggested that matter at the outer edge of the accretion disk, together with an edge-on disk orientation, could explain the partial covering of the spectrum in Cir X-1.

Iaria et al. (2001a), using *BeppoSAX* data, analyzed the 0.1–100 keV spectrum of Cir X-1 at orbital phases near the periastron. They found that the continuum could be well described by either a two-blackbody model or a Comptonization model. In the first case, the “Eastern model” proposed by Mitsuda et al. (1984), the soft blackbody emission is interpreted as the emission from the inner region of the accretion disk while the hard blackbody comes from the neutron star surface. In the case of Cir X-1, the radius of the soft blackbody was ~ 140 km, which should correspond to the inner radius of the accretion disk, while the radius of the harder blackbody component was ~ 30 km—too large to be interpreted as the neutron star radius. Therefore the Comptonization model was preferred. A hard power-law tail, dominating the spectrum at energies higher than 15 keV, was also observed. A similar feature was observed in other Z-source LMXBs (GX 17+2, Di Salvo et al. 2000; GX 349+2, Di Salvo et al. 2001; GX 5–1, Asai et al. 1994; Sco X-1, D’Amico et al. 2000), suggesting that Cir X-1 shows a behavior similar to that of other Z sources. Finally, a strong absorption edge at $\sim 8.4\text{--}8.7$ keV, produced by highly ionized iron, was needed to fit the spectrum of Cir X-1.

Brandt & Schulz (2000), using *Chandra* data, observed Cir X-1 near the periastron. They found, for the first time in the X-ray band, P Cygni profiles of lines emitted by highly ionized matter. The outflow velocity necessary to explain the P Cygni profiles was ~ 2000 $km s^{-1}$. Under the hypothesis that the source is seen nearly edge-on, they supposed that the model of a thermally driven wind (Begelman et al. 1983) could explain this outflow of ionized matter from the system. They also suggested that the absorption edge observed by Iaria et al. (2001a) could be connected to the outflow.

¹ Dipartimento di Scienze Fisiche ed Astronomiche, Università di Palermo, Via Archirafi 36, 90123 Palermo, Italy; iaria@gifco.fisica.unipa.it.

² Astronomical Institute “Anton Pannekoek,” University of Amsterdam and Center for High-Energy Astrophysics, Kruislaan 403, NL 1098 SJ Amsterdam, the Netherlands.

³ Osservatorio Astronomico di Roma, Via Frascati 33, 00040 Monteporzio Catone, Rome, Italy.

Iaria et al. (2001b), using *ASCA* data, studied the energy spectrum of Cir X-1 along its orbit. They distinguished three different X-ray states of the source as a function of its orbital phase. At first, when the source is near the periastron (according to the ephemeris of Stewart et al. 1991), the luminosity is super-Eddington and a large X-ray flaring activity is present. In the second state, corresponding to the orbital phase interval 0.2–0.6, the luminosity is sub-Eddington and the light curve does not show flaring activity. In the third state, in the orbital phase interval 0.7–0.9, an excess of neutral matter (probably the atmosphere of the companion star and/or the accreting matter) partially covers the emitting region. In all these states, the continuum was well fitted using a Comptonization model. Unfortunately, because of the relatively narrow band of *ASCA* (0.6–10 keV), the spectral evolution of the hard tail along the orbit could not be studied. On the other hand, it was possible to study the evolution of the absorption edge along the orbit of Cir X-1. At the phase of the flaring activity, the hydrogen column derived from the edge was $\sim 10^{24} \text{ cm}^{-2}$. When the source moved away from the periastron, the hydrogen column decreased; the absorption edge (from ionized iron) was no longer detected at orbital phases from 0.78 to the periastron. It is possible that the outflowing wind is connected to the huge flaring activity at the periastron.

In this work we present the broadband (1.8–100 keV) spectrum of Cir X-1 from a *BeppoSAX* observation taken at orbital phases 0.61–0.63. The broad band allows us to constrain the continuum components of the spectrum and to detect the presence of a hard tail at energies higher than 20 keV.

2. SPECTRAL ANALYSIS OF CIR X-1 AT THE APOASTRON

A pointed observation of Cir X-1 was carried out between 1999 February 7 03:15:19 UT and 1999 February 8 19:25:33 UT with the narrow field instruments on board *BeppoSAX* (Boella et al. 1997). These consist of four co-aligned instruments covering the 0.1–200 keV energy range: a Low Energy Concentration Spectrometer (LECS; operating in the range 0.1–10 keV), two Medium Energy Concentration Spectrometers (MECS; 1.3–10 keV), a High-Pressure Gas Scintillation Proportional Counter (HPGSPC; 7–60 keV), and a Phoswich Detector System (PDS; 13–200 keV). The LECS and the HPGSPC were not active during this observation. The effective exposure time was 25 ks in the MECS and 11 ks in the PDS. In the MECS image, data were extracted from a circular region of 4' radius centered on the source centroid. Data extracted from the same detector region during blank field observations were used for background subtraction. Background subtraction for the PDS data was obtained from data accumulated during off-source intervals. We rebinned the energy spectra in order to have at least 25 counts channel^{-1} and to oversample the instrument resolution with the same number of channels at all energies⁴.

Using the orbital parameters reported by Stewart et al. (1991), we have calculated that this observation corresponds to the orbital phase interval 0.61–0.63, just after the apoastron. In Figure 1 we plotted the hardness-intensity diagram, where the hardness is the ratio of the count rate in




FIG. 1.—Hardness-intensity diagram of Cir X-1. We show in the same box the data at the periastron binary phase (Iaria et al. 2001a) and the new data taken near the apoastron phase, as indicated in the figure. The hardness is defined by the ratio $[3-7 \text{ keV}]/[1-3 \text{ keV}]$ and the intensity of the source is calculated in the energy band $[1-7 \text{ keV}]$. The bin time is 137 s. The data at the apoastron correspond to lower intensity and higher hardness (top left in the diagram).

the 3–7 keV energy band to the count rate in the 1–3 keV energy band, and the intensity is the count rate in the 1–7 keV energy band. In this figure we show both the data of this observation and those of a previous *BeppoSAX* observation at the periastron (see Iaria et al. 2001a). The data at the apoastron show larger hardness and lower intensity with respect to the data at the periastron. In Figure 2 we plotted the color-color diagram of Cir X-1 data, where the hard color is the ratio of the count rate in the energy band 7–10 keV to that in the energy band 3–7 keV and the soft color is the hardness defined above. The data at the apoastron are at the top right in this figure while the data obtained at the periastron are at the bottom left, implying that the spectrum at the apoastron is harder in both the soft and hard color. In Figure 3 we plotted the light curve of the data at the apoastron in the energy band 1.8–10 keV (upper panel) together with the soft color as a function of time (lower panel). The count rate smoothly decreases from $\sim 300 \text{ counts s}^{-1}$ at the beginning of the observation to $\sim 280 \text{ counts s}^{-1}$ at the end of the observation. No flaring activity is present.

Because no significant spectral variation is observed in the hardness ratios described above, we produced a unique spectrum in the band 1.8–200 keV from this observation. A systematic error of 1% was added to each spectrum. As customary, in the spectral-fitting procedure we allowed for

⁴ See the *BeppoSAX* cookbook at <http://www.sdc.asi.it/software/index.html>.




FIG. 2.—Color-color diagram of Cir X-1. We report here the same data shown in Fig. 1. The hard color is the hardness ratio $[7-10 \text{ keV}]/[3-7 \text{ keV}]$ and the soft color is the hardness ratio $[3-7 \text{ keV}]/[1-3 \text{ keV}]$. The bin time is 137 s. The data at the apoastron are at the top right of the box showing that the spectrum is harder in both the soft and the hard color.




FIG. 3.—Cir X-1 light curve in the energy band 1.8–10 keV (MECS data, upper panel) and ratio of the count rate in the energy band 3–7 keV to that in 1.8–3 keV (lower panel). The bin time is 137 s.




FIG. 4.—Residuals in units of σ with respect to the Comptonization model (upper panel), to model 1 (middle panel), to model 2 (lower panel). The models are described in the text and in Table 1.

a different normalization of the PDS spectrum relative to the MECS spectrum, always checking that derived values were in the standard range for that instrument.⁵ The energy ranges used for the spectral analysis are 1.8–10 keV for the MECS and 15–200 keV for the PDS. Because of the lack of low-energy data, we fixed the value of the average photoelectric absorption by cold matter at $1.7 \times 10^{22} \text{ cm}^{-2}$ (see, e.g., Iaria et al. 2001a, 2001b). We started fitting the data to the Comptonization model Comptt (Titarchuk 1994). This model gave an unacceptable fit with $\chi^2/\text{dof} = 1161/100$; the residuals with respect to this model are shown in Figure 4 (upper panel). From the residuals, it is evident that there is a feature between 8–10 keV and an excess at energies higher than 15 keV. Note that because this simple model does not fit the high-energy part of the spectrum, we had to fix the PDS normalization factor to 0.85 (its typical value) to avoid unreasonably high values of this parameter.

Then, in agreement with the model proposed by Iaria et al. (2001a) for the spectrum of Cir X-1 at the periastron, we added a power law to fit the high energy excess, an absorption edge at ~ 8.4 keV, and a Gaussian emission line at ~ 6.7 keV (with the Gaussian σ fixed at 0.25 keV), keeping fixed the PDS normalization factor. The addition of each of these components is statistically significant and in this way the fit improves significantly, giving $\chi^2/\text{dof} = 117/94$. In Table 1 (model 1) we present the best-fit parameters obtained for this model. Looking at the residuals with

⁵ See the *BeppoSAX* cookbook at <http://www.sdc.asi.it/software/index.html>.

TABLE 1
RESULTS OF THE FIT OF CIR X-1 1.8–200 keV SPECTRUM AT THE BINARY PHASE 0.61–0.63

Parameter	Model 1	Model 2	Model 3
	Comptt + Power Law + Line + Edge	Comptt + Modified Power Law + Line + Edge	BB + Comptt + Power Law + Line + Edge
$N_{\text{H}} (\times 10^{22} \text{ cm}^{-2})$	1.7 (fixed)	1.7 (fixed)	1.7 (fixed)
$E_{\text{edge}} (\text{keV})$	$8.40_{-0.11}^{+0.13}$	$8.41_{-0.15}^{+0.21}$	$8.47_{-0.24}^{+0.29}$
τ_{edge}	0.175 ± 0.025	$0.125_{-0.047}^{+0.023}$	$0.071_{-0.035}^{+0.016}$
$kT_{\text{bb}} (\text{keV})$	$0.589_{-0.021}^{+0.036}$
N_{bb}	$0.1710_{-0.0098}^{+0.0067}$
$kT_0 (\text{keV})$	$0.637_{-0.023}^{+0.027}$	$0.569_{-0.047}^{+0.037}$	$1.182_{-0.030}^{+0.042}$
$kT_e (\text{keV})$	1.641 ± 0.014	$1.510_{-0.046}^{+0.027}$	$6.0_{-2.6}^{+11.1}$
τ	$16.38_{-0.53}^{+0.31}$	$19.14_{-0.69}^{+0.86}$	$1.72_{-0.33}^{+1.96}$
N_{comp}	$5.71_{-0.15}^{+0.18}$	$6.37_{-0.28}^{+0.64}$	$0.75_{-0.23}^{+1.59}$
$f_{\text{bol}} (\times 10^{-8} \text{ ergs cm}^{-2} \text{ s}^{-1})$	3.2	3.2	2.3
$E_{\text{Fe}} (\text{keV})$	$6.702_{-0.100}^{+0.084}$	$6.757_{-0.097}^{+0.087}$	$6.800_{-0.100}^{+0.094}$
$\sigma_{\text{Fe}} (\text{keV})$	0.25 (fixed)	< 0.16	< 0.15
$I_{\text{Fe}} (\times 10^{-2} \text{ photons cm}^{-2} \text{ s}^{-1})$	$0.92_{-0.23}^{+0.20}$	$0.73_{-0.22}^{+0.36}$	$0.69_{-0.24}^{+0.29}$
$\text{EQW}_{\text{Fe}} (\text{eV})$	$47.7_{-12.1}^{+10.1}$	$39.6_{-11.7}^{+19.9}$	$42.2_{-13.3}^{+17.9}$
Photon index	$3.318_{-0.066}^{+0.020}$	$3.593_{-0.146}^{+0.065}$	$2.96_{-0.53}^{+0.30}$
N_{PL}	$6.15_{-0.93}^{+0.57}$	$14.1_{-3.9}^{+3.9}$	$0.89_{-0.77}^{+1.92}$
Low-energy cutoff (keV)	$1.510_{-0.046}^{+0.027}$...
PDS normalization	0.85 (fixed)	$1.049_{-0.033}^{+0.156}$	$0.98_{-0.10}^{+0.15}$
χ^2/dof	117/94	96/92	71/92

NOTE.—The best fit (model 3) is given by a blackbody, a Comptonized spectrum modeled by Comptt, a power law, an absorption edge, and a Gaussian emission line. Uncertainties are at 90% confidence level for a single parameter. kT_{bb} and N_{bb} are, respectively, the blackbody temperature and normalization in units of L_{39}/D_{10}^2 , where L_{39} is the luminosity in units of 10^{39} ergs s^{-1} and D_{10} is the distance in units of 10 kpc. kT_0 is the temperature of the seed photons for the Comptonization, kT_e is the electron temperature, τ is the optical depth of the scattering cloud using a spherical geometry, N_{comp} is the normalization of the Comptt model in XSPEC v.11 units, and f_{bol} is the bolometric (unabsorbed) flux of the Comptonization component. EQW_{Fe} indicates the equivalent width of the Gaussian emission line E_{Fe} , its centroid energy and I_{Fe} its intensity. The power-law normalization, N_{PL} , is in units of photons $\text{keV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}$ at 1 keV.




FIG. 5.—Energy spectrum (1.8–200 keV) of Cir X-1 at the orbital phase 0.61–0.63. Data and the corresponding best-fit model (see Table 1, model 3) are shown in the upper panel; residuals in units of σ with respect to the best-fit model are shown in the lower panel.

respect to model 1 (Fig. 4, *middle panel*), we still see modulations between 1.8 and 5 keV and between 20 and 40 keV, indicating that the continuum emission could be more complex.

The modulation in the residuals could be due to the large contribution at low energies of the power law, which has a large normalization and a photon index of ~ 3.3 . We modified the power law, multiplying it for an exponential cutoff at low energy (*Expfac* in XSPEC), fixing the low-energy cutoff at the electron temperature of the Comptonized component. In this way we found a stable (and reasonable) fit, with the PDS normalization factor and the width of the Gaussian emission line as free parameters (see Table 1, model 2). The residuals (see Fig. 4, *lower panel*) show that the modulation between 1.8 and 5 keV is now smaller. We obtain $\chi^2/\text{dof} = 96/92$ and an *F*-test with respect to the previous model gives a probability of chance improvement of the fit of $\sim 1.1 \times 10^{-4}$. However, the modulation between 20 and 40 keV is still visible.

We therefore tried an alternative model, consisting of model 1, to which we added a blackbody component at low energies. The addition of the blackbody component improved the fit significantly, giving $\chi^2/\text{dof} = 71/91$ with a probability of chance improvement with respect to model 1 of $\sim 6.5 \times 10^{-10}$. Hence we consider this the best-fit model. The *F*-test for the addition of the power-law component in this case gives a probability of chance improvement of $\sim 9.8 \times 10^{-10}$. In Table 1 (model 3) we report the parameters of the best-fit model. The blackbody temperature is $kT_{\text{bb}} \sim 0.59$ keV. The parameters of the Comptonized component are as follows: the seed-photon temperature is $kT_0 \sim 1.2$ keV, the electron temperature is $kT_e \sim 6$ keV, and the optical depth of the Comptonizing cloud is $\tau \sim 1.7$. The




FIG. 6.—Unfolded spectrum of Cir X-1 and the best-fit model, shown in this figure as a solid line. The single components of the model are also shown, namely the blackbody (*dashed line*), Comptt (*dotted line*), power-law (*double-dot-dashed line*), and iron emission line (*dot-dashed line*). The absorption edge at 8.45 keV is visible.

centroid of the Gaussian emission line is at ~ 6.8 keV, with FWHM ~ 0.39 keV, and the absorption edge is at ~ 8.47 keV, with an optical depth $\tau_{\text{edge}} \sim 0.07$. The PDS normalization factor is ~ 1 .

In Figure 5 we show the data and the residuals with respect to model 3. Note that in this case the modulations between 1.8 and 5 keV and between 20 and 40 keV disappear, confirming that model 3 gives the best fit to these data. In Figure 6 we show the corresponding unfolded spectrum. The extrapolated unabsorbed flux in the energy band 0.1–200 keV is $\sim 2.3 \times 10^{-8}$ ergs $\text{s}^{-1} \text{cm}^{-2}$ for the Comptonized component, 1.4×10^{-8} ergs $\text{s}^{-1} \text{cm}^{-2}$ for the blackbody component, and 1.35×10^{-8} ergs $\text{s}^{-1} \text{cm}^{-2}$ for the power-law component.

3. DISCUSSION

We analyzed data of Cir X-1 from a *BeppoSAX* observation (in the orbital phase range 0.61–0.63) in the energy range 1.8–200 keV. The light curve and the hardness ratio do not show large variations during the observation. From Figures 1 and 2 it is clear that the spectrum, in the range 1–10 keV, is different from that observed at the periastron (Iaria et al. 2001a); in fact, the spectrum at the apoastron is harder in both the soft and hard colors and the intensity is lower than that at the periastron. The spectrum of Cir X-1 during this observation is very similar to that of other Z sources (see Di Salvo et al. 2000; Di Salvo et al. 2001). The soft X-ray continuum in the energy range 1.8–10 keV is well fitted by a blackbody and a Comptonized component, while

a power-law component, with photon index ~ 3 , is present at energies higher than 20 keV.

The blackbody temperature is ~ 0.6 keV and the corresponding radius of the emitting region is ~ 58 km. This radius is too large to be associated with the emission from the neutron star surface; therefore, the blackbody component is probably emitted by the inner part of the accretion disk. The seed-photon and electron temperature of the Comptonized component are ~ 1.2 keV and ~ 6 keV, respectively, and the optical depth is ~ 2 . We calculated the radius, R_w , of the seed-photon emitting region using the parameters reported in Table 1 (model 3) following in 't Zand et al. (1999); we obtain $R_w \sim 17$ km. This component could be produced by a corona surrounding the neutron star and occulting the inner region of the accretion disk.

We found an absorption edge at ~ 8.5 keV with $\tau_{\text{edge}} \sim 0.07$. The energy of the absorption edge is compatible with iron ionization levels of Fe xxiii–xxiv (see Turner et al. 1992 for a correspondence between iron edge energy and ionization level). The best-fit value for the optical depth τ_{edge} , considering the photoionization cross section for the K-shell of Fe xxiii (Krolik & Kallman 1987), corresponds to a hydrogen column density of $\sim 1.2 \times 10^{23} \text{cm}^{-2}$. A Gaussian emission line is also detected at ~ 6.8 keV. The energy of the emission line corresponds to iron ionization levels Fe xxiv–xxv, compatible with that of the absorption edge (see Turner et al. 1992). This suggests that the iron emission line could be produced in the same region where the absorption edge is produced. Note that the *ASCA* spectrum at orbital phase 0.54–0.56, the nearest to the orbital phase of the *BeppoSAX* observation analyzed here, shows both the iron emission line and the absorption edge, and their parameters are in agreement with the values we found from this *BeppoSAX* observation (compare with Iaria et al. 2001b).

The best-fit model obtained here is different from that used by Iaria et al. (2001a) for the *BeppoSAX* spectrum of Cir X-1, taken at the periastron when a flaring activity was present. In that case only the Comptonized component was needed to fit the continuum from 0.12 to 10 keV and the blackbody component was not needed. The seed-photon and electron temperatures at the periastron were ~ 0.4 and ~ 1 keV, respectively, with a radius of the seed-photon emitting region of ~ 140 km.

One possibility to explain the large seed-photon radius observed at the periastron, in the hypothesis that it is the inner radius of the accretion disk, might be the presence of a large neutron star magnetic field with a strength of $\sim 4 \times 10^{10}$ G, more than one order of magnitude higher than the typical value estimated for the Z sources. In light of the new results on Cir X-1 presented in this work, we can exclude this possibility. In fact, the flux from the source was larger than 5.4×10^{-8} ergs $\text{cm}^{-2} \text{s}^{-1}$ during the observation at the periastron and, for a distance to the source of 5.5 kpc (Case & Bhattacharya 1998), the corresponding luminosity is super-Eddington. During the observation analyzed here, where Cir X-1 is near the apoastron, the estimated flux is 3.8×10^{-8} ergs $\text{cm}^{-2} \text{s}^{-1}$ and the luminosity is sub-Eddington. Because the X-ray luminosity is connected to the accretion rate, the ram pressure of the disk should be lower by a factor of ~ 1.4 at the apoastron than at the periastron. Consequently the magnetospheric radius (see, e.g., Burderi et al. 1998) should be ~ 154 km at the apoastron, assuming the same strength of the magnetic field

of $\sim 4 \times 10^{10}$ G. However we find that the inner radius of the accretion disk is ~ 58 km.

Another possibility to explain the large inner radius of the accretion disk observed at the periastron can be the occultation of the inner region by a corona surrounding the neutron star. This corona should therefore be more extended at the periastron and cover a larger part of the accretion disk. This hypothesis is also supported by the differences between the spectrum at the periastron and at the apoastron. The seed-photon temperature is much lower at the periastron (~ 0.4 keV), with a comparatively higher accretion rate, than at the apoastron (~ 1.2 keV), implying that at the periastron the seed photons for the Comptonization come from a further region in the accretion disk. In particular at the periastron the seed photons probably come from a region which extends up to a radius of ~ 140 km in the accretion disk (see Iaria et al. 2001a). On the other hand, at the apoastron, the seed photons probably come from a region which extends up to ~ 20 km in the accretion disk. This also explains why we observe a blackbody component at the apoastron, which we interpret as the emission of the part of the accretion disk not covered by the corona, i.e., from radii larger than ~ 60 km, while this component is not observed at the periastron. In fact, the disk emission from radii larger than 150 km would probably be too soft to be significantly detected by the *BeppoSAX*/LECS.

A power-law component, dominating the spectrum of Cir X-1 at energies higher than 20 keV, is also detected with high statistical significance. This component was observed two times in Cir X-1: at the periastron (Iaria et al. 2001a) and near the apoastron (this paper). At the periastron of Cir X-1, the photon index was ~ 3.3 with a flux in the band 10–200 keV of $\sim 1.4 \times 10^{-10}$ ergs s $^{-1}$ cm $^{-2}$. At the apoastron the photon-index is ~ 2.9 with a flux in the band 10–200 keV of $\sim 1.1 \times 10^{-10}$ ergs s $^{-1}$ cm $^{-2}$. Note that the radio emission in Cir X-1 is maximum near the periastron (Stewart et al. 1993; Fender 1997). Also, the observation of Cir X-1 presented in this work is at the orbital phase between 0.61–0.63; at the same orbital phases Fender (1997) detected a secondary radio-flaring activity with a maximum intensity of ~ 25 mJy, lower than at the periastron (where the maximum intensity is ~ 40 mJy). Although a systematic study of the presence of the hard power-law component along the entire orbit is needed to draw a conclusion, the coincidence between the presence of this component and periods of radio flares suggests a connection between these two phenomena. In particular the power-law component might have a nonthermal origin and might be related to the presence of the high-velocity electrons responsible for the radio emission.

As mentioned above, there is evidence that a jet is present at the periastron (Stewart et al. 1993; Fender 1997). Also, the prominent absorption edge from highly ionized matter observed at the periastron of Cir X-1 is probably connected with the outflowing wind observed by *Chandra* (Brandt & Schulz 2000; Iaria et al. 2001b). These observations could be explained using a model similar to that proposed for young stellar objects (YSOs). The source HH 30 presents both a jet and a wind from the disk. Goodson et al. (1999) explain the presence of these two phenomena by postulating that the interaction between the stellar magnetosphere and the surrounding accretion disk drives a two-component outflow, the jet and the disk wind. The magnetic loops connecting the accretion disk to the star expand because of the

differential rotation between the star and the disk (Hayashi et al. 1996; Lovelace et al. 1995). The plasma trapped in the loops is driven toward the rotation axis, producing a collimated jet and an outflowing wind outward along the disk surface. Magnetic reconnections can produce large X-ray flaring activity and allow the process to repeat.

In this interpretation, the P Cygni profiles and the absorption edge observed near the periastron could be explained by the disk wind expected in this model, the radio emission by the collimated jet, and the flaring activity at the periastron by the magnetic reconnections. The Comptonized emission observed at the periastron could also come from the outflowing wind. The outflow and then the Comptonizing region, because of the large magnetic activity, probably extends far from the neutron star above the disk, and, as mentioned above, this might explain why we do not observe the blackbody emission from the accretion disk at the periastron. However, note that the work of Goodson et al. (1999) is based on YSOs and possibly the results might not be applied in a straightforward way to the case of Cir X-1 because of the different strength of the neutron star magnetic field. A study of the consequences of this model in the more extreme case of magnetized neutron stars would be interesting in this context. Following this scenario, we can suppose that at the apoastron there are no magnetic reconnections and consequently the source is not highly variable. However, the reason for this difference between the periastron and the apoastron is not clear, although we can suppose it can be related to the difference in the accretion rate and/or tidal interactions, and should be investigated with a proper model.

4. CONCLUSIONS

In this work we presented the analysis of the broadband (1.8–200 keV) spectrum of Cir X-1 at the orbital phase 0.61–0.63 using a *BeppoSAX* observation. The luminosity of the source is sub-Eddington and the spectrum is harder than that at the periastron (Iaria et al. 2001a). The model used to fit this spectrum is the same that has been used for other Z sources (see Di Salvo et al. 2000; Di Salvo et al. 2001); we find that the continuum is well fitted using three components. The first is a blackbody with a temperature of ~ 0.6 keV, probably produced by the inner region of the accretion disk. The corresponding blackbody radius is ~ 58 km. The second is a Comptonized component with a seed-photon temperature of ~ 1.2 keV, an electron temperature of ~ 6 keV, and an optical depth of ~ 1.7 . We interpret this component as the emission from a corona (or an outflowing wind) around the neutron star. In this case there is a weaker outflow of matter, suggested by the relatively low optical depth of the absorption edge, which is now $\tau_{\text{edge}} \sim 0.07$, one order of magnitude lower than at the periastron. The third component is a power-law that could be produced by Comptonization due to nonthermal electrons or by synchrotron emission in a jet. An emission line is also present at ~ 6.8 keV, with FWHM ~ 0.39 keV. The ionization level inferred from the line is Fe XXIV–XXV. The last observed feature is an absorption edge from highly ionized iron (Fe XXIII–XXIV), probably produced by an outflowing wind along the disk surface.

This work was partially supported by the Italian Space Agency and the Ministero della Ricerca Scientifica e Tecnologica.

REFERENCES

- Asai, K., et al. 1994, PASJ, 46, 479
Begelman, M. C., McKee, C. F., & Shields, G. A. 1983, ApJ, 271, 70
Boella, G., Butler, R. C., Perola, G. C., Piro, L., Scarsi, L., & Blecker, J. 1997, A&AS, 122, 299
Brandt, W. N., Fabian, A. C., Dotani, T., Nagase, F., Inoue, H., Kotani, T., & Segawa, Y. 1996, MNRAS, 283, 1071
Brandt, W. N., & Schulz, N. S. 2000, ApJ, 544, L123
Burderi, L., Di Salvo, T., Robba, N. R., del Sordo, S., Santangelo, A., & Segreto, A. 1998, ApJ, 498, 831
Case, G. L., & Bhattacharya, D. 1998, ApJ, 504, 761
Clark, D. H., Parkinson, J. H., & Caswell, J. L. 1975, Nature, 254, 674
D'Amico, F., Heindl, W. A., Rothschild, R. E., & Gruber, D. E. 2001, ApJ, 547, L147
Di Salvo, T., et al. 2000, ApJ, 544, L119
———. 2001, ApJ, 554, 49
Fender, R. P. 1997, in AIP Conf. Proc. 410, Proc. 4th Compton Symposium, ed. C. D. Dermer, M. S. Strickman, & J. D. Kurfess (Melville: AIP), 798
Goodson, A. P., Bohm, K. H., & Winglee, R. M. 1999, ApJ, 524, 142
Hasinger, G., & van der Klis, M. 1989, A&A, 225, 79
Hayashi, M. R., Shibata, K., & Matsumoto, R. 1996, ApJ, 468, L37
Iaria, R., Burderi, L., Di Salvo, T., La Barbera, A., & Robba, N. R. 2001a, ApJ, 547, 412
Iaria, R., Di Salvo, T., Burderi, L., & Robba, N. R. 2001b, ApJ, 561, 321
in 't Zand, J. J. M., et al. 1999, A&A, 345, 100
Johnston, H. M., Fender, R., & Wu, K. 1999, MNRAS, 308, 415
Krolik, J. H., & Kallman, T. R. 1987, ApJ, 320, L5
Lovelace, R. V. E., Romanova, M. M., & Bisnovatyi-Kogan, G. S. 1995, MNRAS, 275, 244
Mitsuda, K., et al. 1984, PASJ, 36, 741
Murdin, P., Jauncey, D. L., Haynes, R. F., Lerche, I., Nicolson, G. D., Holt, S. S., & Kaluziński, L. J. 1980, A&A, 87, 292
Shirey, R. E., Bradt, H. V., Levine, A. M., & Morgan, E. H. 1996, ApJ, 469, L21
Stewart, R. T., Caswell, J. L., Haynes, R. F., & Nelson, G. J. 1993, MNRAS, 261, 593
Stewart, R. T., Nelson, G. J., Penninx, W., Kitamoto, S., Miyamoto, S., & Nicolson, G. D. 1991, MNRAS, 253, 212
Tauris, T. M., Fender, R. P., van den Heuvel, E. P. J., Johnston, H. M., & Wu, K. 1999, MNRAS, 310, 1165
Titarchuk, L. 1994, ApJ, 434, 570
Turner, T. J., Done, C., Mushotzky, R., & Madejski, G. 1992, ApJ, 391, 102