



**UvA-DARE (Digital Academic Repository)**

**The lives and deaths of star clusters near the Galactic Center?**

McMillan, S.L.W.; Portegies Zwart, S.F.

*Published in:*  
Astrophysical Journal

*DOI:*  
[10.1086/324600](https://doi.org/10.1086/324600)

[Link to publication](#)

*Citation for published version (APA):*

McMillan, S. L. W., & Portegies Zwart, S. F. (2001). The lives and deaths of star clusters near the Galactic Center? *Astrophysical Journal*, 565, 256. DOI: 10.1086/324600

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

## THE LIVES AND DEATHS OF STAR CLUSTERS NEAR THE GALACTIC CENTER

SIMON F. PORTEGIES ZWART,<sup>1</sup> JUNICHIRO MAKINO,<sup>2</sup> STEPHEN L. W. MCMILLAN,<sup>3</sup> AND PIET HUT<sup>4</sup>

Received 2001 February 14; accepted 2001 August 29

### ABSTRACT

We study the evolution and observability of young, compact star clusters near the Galactic center, such as the Arches and Quintuplet systems. The clusters are modeled by integrating the equations of motion of all stars while accounting for the internal evolution of stars and binaries, as well as the effect of the Galactic tidal field. We find that clusters within 150 pc of the Galactic center dissolve within  $\sim 55$  Myr, but their projected densities drop below the background density in the direction of the Galactic center within only a few megayears, effectively making these clusters undetectable after that time. Detailed observations of the Arches cluster, taken at face value, suggest that its mass function is unusually flat and that the cluster contains an overabundance of stars more massive than  $20 M_{\odot}$ . Our dynamical analysis, however, shows that the observed characteristics of the Arches cluster are consistent with a perfectly normal initial mass function. The observed anomalies are then caused by a combination of observational selection effects and the dynamical evolution of the cluster. We calibrate the current parameters of the Arches cluster using a normal initial mass function and conclude that the cluster is more massive than  $40,000 M_{\odot}$ , has a half-mass radius of about 0.35 pc, and is located between 50 and 90 pc from the Galactic center.

*Subject headings:* binaries: close — blue stragglers — globular clusters: general — open clusters and associations: individual (Arches, Quintuplet cluster) — stars: evolution

### 1. INTRODUCTION

A number of young, dense star clusters have been observed within the inner few hundred parsecs of the Galactic center. Best known are the Arches cluster (Object 17) (Nagata et al. 1995) and the Quintuplet cluster (AFGL 2004) (Nagata et al. 1990; Okuda et al. 1990). However, it is likely that others exist, as these clusters lie behind thick layers of obscuring material (Portegies Zwart et al. 2001a). Most are expected to be invisible at optical wavelengths, but should be readily detectable in the infrared (Vrba et al. 2000) or in the 2MASS survey (Dutra & Bica 2000).

The Arches and the Quintuplet clusters are the Galactic counterparts of R136, the central star cluster in NGC 2070: the 30 Doradus region in the Large Magellanic Cloud (Massey & Hunter 1998). The structural parameters of these clusters—masses, radii, and density profiles—are quite similar, as are their ages. R136, however, is located far from the perturbing influence of the Galaxy, and the tidal effect of the LMC is negligible (Portegies Zwart et al. 1999). The Arches and the Quintuplet clusters, on the other hand, lie at projected distances of  $\lesssim 40$  pc of the Galactic center, and their evolution is strongly affected by the Galactic tidal field.

We study the evolution of clusters like Arches and Quintuplet using a fully self-consistent star-cluster model in which the dynamics of stars are followed by direct  $N$ -body integration and the evolution of individual stars and binaries are followed using a stellar and binary evolution program. The importance of the tidal field of the Galaxy is

studied by repeating each model calculations at several distances from the Galactic center. We compare the results of our calculations with detailed observations of the Arches cluster and with model calculations performed by others.

We find that, while all our models start with a mass function comparable to that in the solar neighborhood, to an observer the current mass function may well appear to be much flatter. The unusually flat mass functions observed in both the Arches and the Quintuplet clusters may therefore in part be attributed to a combination of observational selection effect, the age of the cluster, and the location at which the measurements were taken.

The star clusters studied in this paper experience core collapse within a few million years. During this phase and at later time many stars experience collisions with other stars. The clusters finally dissolve in the tidal field of the Galaxy. The disruption of the cluster is driven mostly by tidal stripping and two-body relaxation. Stellar evolution plays only a minor role in the disruption of the clusters.

The clusters expand as they age, causing their surface densities to decline. Clusters older than about 5 Myr often have surface densities below that of their surroundings, making these clusters virtually undetectable. Since the modeled clusters disrupt at ages greater than 5 Myr, we estimate that there may be many clusters like the Arches and Quintuplet, but only the youngest are observable. This suggests that the formation of Arches- and Quintuplet-like clusters may be a continuous process. We cannot exclude, however, that these clusters were formed in a recent burst of star formation and that older counterparts do not exist.

The numerical methods and selection of the initial conditions are discussed in § 2. The results are presented in § 3 and discussed in § 4. We summarize and conclude in § 5.

### 2. THE MODEL

The  $N$ -body portion of the simulations is carried out using the *kira* integrator operating within the STARLAB

<sup>1</sup> Massachusetts Institute of Technology, Cambridge, MA 02139. S. P. Z. is a Hubble Fellow.

<sup>2</sup> Department of Astronomy, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan.

<sup>3</sup> Department of Physics, Drexel University, Philadelphia, PA 19104.

<sup>4</sup> Institute for Advanced Study, Princeton, NJ 08540.

software environment. Time integration of stellar orbits is accomplished using a fourth-order Hermite scheme (Makino & Aarseth 1992). *Kira* also incorporates block time steps (McMillan 1986a, 1996b; Makino 1991), special treatment of close two-body and multiple encounters of arbitrary complexity, and a robust treatment of stellar and binary evolution and stellar collisions. The special-purpose GRAPE-4 (Makino et al. 1997) system is used to accelerate the computation of gravitational forces between stars. The treatment of stellar mass loss is described by Portegies Zwart et al. (1998). A concise description of the STARLAB environment is given by Portegies Zwart et al. (2001b, hereafter Paper IV).<sup>5</sup>

Evolution of stars and binaries is handled using the prescription given by Portegies Zwart & Verbunt (1996, § 2.1). However, some changes are made to the treatment of main-sequence mass loss in massive stars (see Portegies Zwart et al. 1999, hereafter Paper III), and we incorporate a more extended prescription for stellar collisions, as described by Portegies Zwart, Hut & Verbunt (1997).

The system of units used internally in the  $N$ -body models is defined by  $M = G = -4E = 1$ , where  $E$  is the initial internal energy of the stellar system,  $M$  is the total mass in stars, and  $G$  is the gravitational constant (Heggie & Mathieu 1986).

The time required for a star to cross the clusters' virial radius  $r_{\text{vir}}$  is

$$t_{\text{vir}} = \left( \frac{GM}{r_{\text{vir}}^3} \right)^{-1/2}. \quad (1)$$

For the runs considered here, the  $N$ -body and physical length scales are connected by the requirement that initially the star cluster is in virial equilibrium and exactly fills its zero-velocity surface in the Galactic tidal field (see § 2.2 below). Once the cluster mass and distance to the Galactic center are known, the clusters' size scale is determined. The half-mass crossing time ( $t_{\text{hm}} = 2\sqrt{2}t_{\text{vir}}$ ) can then be

expressed in convenient physical units as

$$t_{\text{hm}} \simeq 42.2 \left( \frac{M_{\odot}}{M} \right)^{1/2} \left( \frac{r_{\text{hm}}}{\text{pc}} \right)^{3/2} \text{ (Myr)}. \quad (2)$$

For most density profiles the half-mass radius  $r_{\text{hm}}$  is slightly ( $\lesssim 25\%$ ) smaller than  $r_{\text{vir}}$ .

The half-mass relaxation time is calculated as (Spitzer 1987)

$$t_{\text{hrx}} = \left( \frac{r_{\text{hm}}^3}{GM} \right)^{1/2} \frac{N}{8 \log \Lambda}. \quad (3)$$

Here  $\Lambda \equiv \gamma N$  is the coulomb logarithm;  $\gamma \sim 0.4$  is a scaling factor introduced to model the effects of the cutoff in the long range Coulomb logarithm (see Giersz & Heggie 1994, 1996). We will also use the relaxation time at the tidal radius  $t_{\text{trx}}$ , for which we use equation (3), but with the tidal radius  $r_{\text{tide}}$  substituted for  $r_{\text{hm}}$ .

### 2.1. Selection of Initial Conditions

The two clusters on which we concentrate here, the Arches and the Quintuplet, lie in projection within  $\sim 40$  pc of the Galactic center. Table 1 lists some observed parameters of these clusters, along with some other clusters (and cluster candidates) having comparable characteristics. All have masses of  $\sim 10^4 M_{\odot}$ , are very compact,  $r_{\text{hm}} \lesssim 1$  pc, and are only a few million years old. However, of the systems listed, only the Arches and Quintuplet are significantly perturbed by external tidal fields.

### 2.2. The Tidal Field Near the Galactic Center

A star cluster embedded in the Galactic tidal field is not spherically symmetric. Rather, it is flattened and the stellar velocity distribution is anisotropic, particularly in the outer regions. The initial models that best describe such a cluster are the anisotropic density profiles described by Heggie & Ramamani (1995). As in the usual spherically symmetric King (1966) models, the density profile is described by the dimensionless parameter  $W_0$ . Higher values of  $W_0$  indicate a more centrally concentrated cluster (see the illustration in Fig. 1).

TABLE 1  
OBSERVED PARAMETERS FOR SOME YOUNG, DENSE CLUSTERS

Name	Reference	Age (Myr)	$M$ ( $10^3 M_{\odot}$ )	$r_{\text{GC}}$ (pc)	$r_{\text{tide}}$ (pc)	$r_{\text{hm}}$ (pc)	$\rho_{\text{hm}}$ ( $10^5 M_{\odot} \text{pc}^{-3}$ )	$t_{\text{hrx}}$ (Myr)
R136 .....	1	2–4	21–79	LMC	$\gtrsim 20$	$\sim 0.5$	0.4–1.5	70
Arches .....	2	1–2	12–50	30	1	0.2	3.6–15	12
Quintuplet .....	3	3–5	10–16	35	1	$\sim 0.5$	0.14–0.31	12
NGC 3603 .....	4	2–3	3–13	few k	4–5	0.23	0.59–2.6	44
W43 (= G30.8-0.2).....	5	$\lesssim 10$	? <sup>a</sup>					
Norma .....	6		? <sup>b</sup>	few k				
Nameless .....	7	$\lesssim 10$	? <sup>c</sup>	12k–15k	$> 1$	0.2–0.4	0.45–4.5	

NOTE.—Columns give the cluster name, reference, age, mass, distance to the Galactic center (the cluster R136 is located in the Large Magellanic Cloud), the tidal radius  $r_{\text{tide}}$ , and the half-mass radius  $r_{\text{hm}}$ . The last two columns give the density within the half-mass radius and the half-mass relaxation time.

<sup>a</sup> Contains approximately 100 spectral type O and WN stars, but further information is not available.

<sup>b</sup> Contains a dozen embedded early-type stars surrounding the M3 Ia star HD 143183.

<sup>c</sup> Contains at least 13 early type O stars. The cluster does not have a name but is located near the soft gamma-ray repeater SGR 1900+14 (also SGR 1806-20 may be associated with a young and compact star cluster [van Kerkwijk et al. 1995; Kulkarni et al. 1995]).

REFERENCES—(1) Brandl et al. 1996; Campbell et al. 1992; Massey & Hunter 1998. (2) Figer et al. 1999a; (3) Glass, Catchpole & Whitelock 1987; Nagata et al. 1990; Figer et al. 1999b. (4) Brandl 1999; (5) Blum, Daminelli & Conti 1999; (6) Moffat 1976; (7) Vrba et al. 2000.

<sup>5</sup> Additional information is available at <http://manybody.org>.




FIG. 1.—Illustration of the structure of star clusters with  $W_0 = 1$  (left),  $W_0 = 4$  (middle) and  $W_0 = 7$  (right). The outer ellipse represents the zero velocity surface of the cluster in the Galactic field. The two inner circles represent the half-mass and core radii. The dashed circle shows the “tidal radius” that would be obtained if the initial model were described by King’s spherically symmetric density profile. The Galactic center is to the top of the figure, at a distance of 150 pc. The length scale is indicated by the horizontal bar.

Following Heggie & Ramamani, we model the tidal potential  $\phi_T$  by the quadrupole formula

$$\phi_T(x, y, z) = -\frac{1}{2} (\alpha_1 x^2 + \alpha_3 z^2). \quad (4)$$

The cluster is taken to move on a circular orbit around the Galactic center. The  $x$ -axis in the rotating frame of reference centered on the cluster always points toward the Galactic center; the  $z$ -axis is perpendicular to the orbital plane. The quantities  $\alpha_1 (> 0)$  and  $\alpha_3 (< 0)$  are conveniently expressed in terms of the kinematic Oort constants  $A$  and  $B$  (Oort 1927), and the local Galactic density  $\rho_G$  is expressed as

$$\begin{aligned} \alpha_1 &= -4A(A - B), \\ \alpha_3 &= 4\pi G\rho_G + 2(A^2 - B^2). \end{aligned} \quad (5)$$

Loosely speaking, we can think of  $\alpha_1$  as determining the overall strength of the tidal field, while the ratio  $\alpha_1/\alpha_3$  determines its geometry.

The Oort constants  $A$  and  $B$  are defined as

$$\begin{aligned} A &= \frac{1}{2} \left( \frac{v_c}{r} - \frac{dv_c}{dr} \right), \\ B &= -\frac{1}{2} \left( \frac{v_c}{r} + \frac{dv_c}{dr} \right), \end{aligned} \quad (6)$$

where  $v_c$  is the circular rotational velocity:

$$v_c = \sqrt{\frac{GM_{\text{Gal}}(r)}{r}}. \quad (7)$$

Taking the mass of the Galaxy within the cluster’s orbit at distance  $r_{\text{GC}}$  ( $\lesssim 100$  pc) from the Galactic center to be (Mezger et al. 1999)

$$M_{\text{Gal}}(r_{\text{GC}}) = 4.25 \times 10^6 \left( \frac{r_{\text{GC}}}{\text{pc}} \right)^{1/2} M_{\odot}, \quad (8)$$

we find

$$\begin{aligned} v_c &= 136.8 \left( \frac{r}{\text{pc}} \right)^{0.1} \text{ km s}^{-1} \\ \frac{dv_c}{dr} &= 13.7 \left( \frac{r}{\text{pc}} \right)^{-0.9} \text{ km s}^{-1} \text{ pc}^{-1}, \end{aligned} \quad (9)$$

and hence

$$\begin{aligned} A &\simeq 61.5 r_{\text{GC}}^{-0.9} \text{ km s}^{-1} \text{ pc}^{-1} \\ B &\simeq -75.3 r_{\text{GC}}^{-0.9} \text{ km s}^{-1} \text{ pc}^{-1} \\ \rho_G &\simeq 4.06 \times 10^5 r_{\text{GC}}^{-1.8} M_{\odot} \text{ pc}^{-3}. \end{aligned} \quad (10)$$

Table 2 lists these parameters at selected Galactocentric distances.

The distance from the center of the star cluster (of mass  $M$ ) to the first Lagrange point (the Jacobi radius) is

$$r_{\text{L1}} \equiv \left( \frac{-M}{\alpha_1} \right)^{1/3}. \quad (11)$$

Substitution of equation (8) into equation (11) yields

$$r_{\text{L1}} \simeq 4.90 \times 10^{-3} \left( \frac{M}{M_{\odot}} \right)^{1/3} \left( \frac{r_{\text{GC}}}{\text{pc}} \right)^{0.6} \text{ pc}. \quad (12)$$

### 2.3. Initial Cluster Structure

We adopt initial models comparable to those observed for the Arches and the Quintuplet clusters. By varying the density profile and the distance to the Galactic center we study how the cluster evolution depends on the initial conditions.

Our calculations start with 12k (12,288) stars at zero age. We assign stellar masses  $m$  in the range  $0.1 M_{\odot} < m < 100 M_{\odot}$  from the mass function suggested for the solar neighborhood by Scalo (1986). The median mass of this mass function is about  $0.3 M_{\odot}$ ; the mean mass is  $\langle m \rangle \simeq 0.6 M_{\odot}$ . For models with 12k stars this results in a total cluster mass of  $\sim 7500 M_{\odot}$ . We adopt a mass function that is applicable to the solar neighborhood rather than the flatter mass spectrum suggested by observations (Figer et al. 1999a) in order to determine whether the flat mass spectrum can be attributed to cluster dynamical evolution. Initially all stars are single, although binaries do form via three-body encounters

TABLE 2  
PARAMETERS FOR THE GALACTIC TIDAL FIELD OF THE CALCULATED STAR CLUSTERS AT SELECTED DISTANCES FROM THE GALACTIC CENTER

$r_{\text{GC}}$ (pc)	$\rho_G(r)$ ( $M_{\odot} \text{ pc}^{-3}$ )	$v_c$ ( $\text{km s}^{-1}$ )	$A$ ( $\text{km s}^{-1} \text{ pc}^{-1}$ )	$B$ ( $\text{km s}^{-1} \text{ pc}^{-1}$ )
Sun.....	0.11	$184 \pm 8$	$11.3 \pm 1.1 \times 10^{-3}$	$-13.9 \pm 0.9 \times 10^{-3}$
34 .....	711	193	2.55	-3.12
90 .....	123	212	1.06	-1.30
150 .....	49	224	0.67	-0.82

NOTE.— Each row gives the distance to the Galactic center, the local stellar density, and the Oort  $A$  and  $B$  constants. The first row gives that value for the local stellar density from Hill, Hiditch, & Barnes 1985 and the Oort constants in the solar vicinity from Olling & Merrifield 1998.

in which one star carries away the excess energy and angular momentum necessary for two other stars to become bound.

We adopt three standard distances from the Galactic center: 34, 90, and 150 pc. The shape of the zero-velocity surface in the tidal field of the Galaxy only depends on  $\alpha_3/\alpha_1$ , which is independent of  $r_{\text{GC}}$ . However, this does not mean that the models can be scaled with respect to distance from the Galactic center using the relaxation time alone. The timescale on which the stars in the cluster evolve depends on the size of the cluster via equation (2); when the cluster is located further from the Galactic center the crossing time increases and the stars evolve relatively more quickly compared to the dynamical evolution of the cluster. This results in a more active mass loss by stellar evolution for clusters which are farther from the Galactic center.

Initial density profiles and velocity dispersions are taken from Heggie & Ramamani (1995) with  $W_0 = 1$ ,  $W_0 = 4$ , and  $W_0 = 7$ , for a total of nine models. At birth the clusters are in virial equilibrium and exactly fill their critical zero-velocity surfaces (Roche lobes) in the Galactic tidal field. Table 3 presents a summary of the adopted initial models. Figure 1 indicates the shape and structure of models with  $W_0 = 1$ , 4, and 7 at  $r_{\text{GC}} = 150$  pc; models closer to the Galactic center are identical in shape, but the scale is different. The zero-velocity surfaces of the various models are represented as ellipses.

We test the reproducibility of our calculations by performing several calculations per set of selected initial conditions. Each calculation was performed twice with a different random seed. In addition, we performed a set of calculations without stellar evolution for models with  $W_0 = 1$ , 4, and 7. These same initial realizations were rerun with stellar evolution switched on, for  $r_{\text{GC}} = 34, 90$ , and 150 pc. In total, 30 simulations were performed.

Stars are removed from the  $N$ -body system when their distance from the center of the cluster exceeds  $3r_{\text{L1}}$ . In  $N$ -body calculations it is not always trivial to determine the moment the cluster dissolves, as a few stars may remain bound for an extended period of time (Portegies Zwart et al 1998). In our models we identify the cluster's disruption as the moment when no stars lie within the zero velocity surface. Typically, a few hundred stars remain in the  $N$ -body system (within  $3r_{\text{L1}}$ ) at this time.

### 3. RESULTS

We first discuss the global parameters of all models. Later (in §§ 3.5 and 3.6) we will concentrate on a few representative models and compare these with the observed clusters.

#### 3.1. The Evolution of the Cluster

Figure 2 (left) shows the mass evolution of the models listed in Table 3. Not surprisingly, clusters located at larger distances from the Galactic center tend to live longer than those closer to the Galactic center.

Figure 2b gives the evolution of the number of stars in several of our models with  $W_0 = 4$  at various distances from the Galactic center and compares the results with a model in which stellar evolution was not taken into account and in which stars were not allowed to collide. All the calculations presented in this figure were started with identical initial conditions. The only difference between the various runs is the distance from the Galactic center (varying from 34 to 150 pc) and whether stellar evolution is included in the calculation. In this figure, time is scaled with respect to each model's initial relaxation time, i.e.,

$$t_{\text{rlx}} \propto N/8 \log(0.4N). \quad (13)$$

The various lines in Figure 2 do not overlap perfectly, mainly owing to stochastic differences in the moments when binaries formed and interacted. The small deviations from the model in which stellar evolution was not taken into account indicate that stellar evolution in these models is unimportant. If stellar evolution were important, clusters at greater distances from the Galactic center should dissolve more rapidly. However, we see the opposite trend: clusters farther from the Galactic center tend to dissolve slightly more slowly; the model without stellar evolution dissolves fastest. We have no ready explanation for this trend. Note that for the models with stellar evolution this trend may be attributed in part to collisions, as they tend to heat the cluster; so clusters with higher collision rates (closer to the Galactic center) would dissolve more quickly in terms of their initial relaxation time. Collisional heating also causes the cluster to become less compact, again reducing the cluster lifetime.

Alternatively, the trend of longer lifetime at greater Galactocentric distance may also be explainable in terms of

TABLE 3  
OVERVIEW OF INITIAL CONDITIONS FOR THE SIMULATIONS PERFORMED

Model	$r_{\text{GC}}$ (pc)	$W_0$	$t_{\text{trx}}$ (Myr)	$t_{\text{hrx}}$ (Myr)	$t_{\text{hm}}$ (Kyr)	$r_{\text{core}}$ (pc)	$r_{\text{vir}}$ (pc)	$r_{\text{tide}}$ (pc)	$n_{\text{coll}}$ (Myr)	$t_{\text{cc}}$ (Myr)	$t_{\text{end}}$ (Myr)	$n_{\text{coll}}/t_{\text{end}}$ (Myr $^{-1}$ )
R34W1 .....	34	1	53	5.5	43	0.12	0.20	0.76	5	1.9	7.4	0.68
R34W4 .....	34	4	53	3.2	27	0.05	0.14	0.76	8	1.2	12.7	0.63
R34W7 .....	34	7	53	0.36	9	0.018	0.077	0.80	24	0.4	12.0	2.00
R90W1 .....	90	1	134	14.6	105	0.19	0.36	1.4	7	1.9	25.6	0.27
R90W4 .....	90	4	134	8.1	68	0.09	0.26	1.4	15	1.2	32.6	4.02
R90W7 .....	90	7	134	1.0	25	0.032	0.14	1.4	14	1.4	32.4	32.4
R150W1 .....	150	1	218	23.6	169	0.30	0.50	1.9	5	2.0	45.8	1.94
R150W4 .....	150	4	218	13	110	0.14	0.36	1.9	8	2.0	53.4	4.08
R150W7 .....	150	7	218	4.5	40	0.044	0.19	2.0	14	0.4	55.0	12.2

NOTE.—Each row lists the model name, the distance to the Galactic center, the initial King parameter  $W_0$ , the initial tidal relaxation time, the initial half-mass relaxation time (see eq. [3]), and see Paper IV for the version in more usual astronomical units), the half-mass crossing time (see eq. [2]), and the initial core radius, virial radius, and the tidal radius. The final four columns give the number of stellar collisions observed in each calculation, the time of core collapse, the time at which the cluster mass dropped below 5% of the initial mass (about  $375 M_{\odot}$ ), and the collision rate in number of collisions per million years.




FIG. 2.—(left) Evolution of total cluster mass for models with  $W_0 = 1$  (solid lines),  $W_0 = 4$  (dashed lines), and  $W_0 = 7$  (dotted line) at distances of 34 pc (left lines) and 150 pc (right lines) from the Galactic center. (right) The evolution of the number of stars (renormalized to the initial total of 12k) for the models with  $W_0 = 4$  as a function of time in units of the initial half-mass relaxation time (see eq. [13]) for  $r_{GC} = 34$  (solid line),  $r_{GC} = 90$  (dashed line) and  $r_{GC} = 150$  (dotted line). The dash-dotted line gives the evolution of the number of stars for the model without stellar evolution. These models were all calculated using the same realization of the initial conditions.

the interplay among stellar evolution, binary formation, and stellar collisions. These possibilities require more detailed study, which is beyond the scope of this paper; however, work is in progress to address this point. We emphasize that, although the trend looks like a subtlety, it may play an important role in the dynamical evolution of compact star clusters.

Figure 3 shows the time variation of the relaxation time for the models at 34 and 90 pc from the Galactic center. Note that the relaxation time measured some time after zero age bears little information about the clusters' initial conditions. The behavior of the half-mass relaxation time is qualitatively similar to that found in less compact clusters at greater distances (6–12 kpc) from the Galactic center (Portegies Zwart et al. 2001b)—it first rises by a factor of a few as the cluster expands and then slowly decreases as the cluster loses mass.

Figure 4 (left) shows the evolution of the mean densities within the half-mass radius in models R150W1, R90W4, and R34W7. The initial half-mass densities for these models

were very different, ranging about 3 orders of magnitude. At later times the differences in the half-mass densities become much smaller; after five million years the difference has decreased by about a factor of 5.

As in Figure 2, 4 (right) also shows the densities of the models as functions of time scaled to the initial relaxation time (see Portegies Zwart et al 2001a and Baumgardt 2001). The trend visible in Figure 2 is somewhat obscured by random fluctuations in the density.

Figure 5 shows the evolution of the core radius, selected Lagrangian radii, and the Jacobi radius  $r_{L1}$ , for model R90W4. Core collapse occurs within  $\sim 1.2$  Myr followed by a gradual overall expansion of the cluster. Similar behavior is evident in the isolated clusters considered by Portegies Zwart et al. (1999). The postcollapse expansion stops after a few million years, by which time the relaxation time has reached its maximum value (see Fig. 3). Subsequently, the Lagrangian radii decrease as the Jacobi radius shrinks and the cluster dissolves.

### 3.2. Evolution of the Binding Energy

Figure 6 shows the evolution of the binding energy ( $-E_{tot}$ , in scaled  $N$ -body units) for models R90W4 and R90W7, which are selected for their similar lifetimes. (Some deviations between the solid and the dotted lines are a result of our definition of a binary. Here we consider a binary to be a pair of bound stars for which the internal forces are a hundred times higher than those due to the nearest neighbor.) Initially, each cluster has  $E_{tot} = -0.25$  (see § 2), but because of escapers, binary activity, stellar mass loss, mergers, supernovae, etc., the total energy changes. A cluster ceases to exist when its binding energy becomes positive. The two clusters exhibit quite similar global evolution. Model R90W4 shows an initial “plateau” of a few megayears during which mass segregation occurs, driving the cluster into core collapse (see Fig. 5). For model R90W7 this phase is hardly noticeable as the cluster experiences several strong encounters and a few collisions early in its evolution.

During the first deep core collapse the binding energy fluctuates rapidly because binaries frequently harden and single stars are ejected following three-body encounters.




FIG. 3.—Evolution of the half-mass relaxation time for clusters lying 34 and 150 pc from the Galactic center, and with  $W_0 = 1$  (solid lines) and  $W_0 = 7$  (dotted lines; the  $W_0 = 7$  models live longer than the  $W_0 = 1$  models). The models with  $W_0 = 4$  and those at  $R_{GC} = 90$  pc are not given.




FIG. 4.—Evolution of the half-mass density for clusters. (left) Panel in astronomical units ( $M_{\odot} \text{pc}^{-3}$  for the models R150W1 (solid line), R90W4 (dashed line), and R34W7 (dotted line). Error bars give the values for age and core density listed in Table 1. (right) The density evolution of the models with  $W_0 = 4$ , but now time is in units of the half-mass relaxation time and density in units of star per half-mass radius cubed (see also Fig. 2b).

The first collisions happen during this phase (Fig. 6, *filled circles*) and in both models mass transfer occurs in a dynamically formed binary, giving rise to an excursion in the binding energy (Fig. 6, *dotted line*). Core collapse is followed by the ejection of a hard binary (Fig. 6, *arrows*).

After the first collapse of the core the binding energies of both clusters decrease steadily with some strong fluctuations caused by close two- and three-body encounters (Fig. 6, *triangles*), high-velocity escapers (*arrows*), stellar collisions (*filled circles*), mass transfer in close binaries, and supernovae (*stars*).

Model R90W7 experiences much more binary activity than model R90W4. This can be seen from the many departures of the dotted line in Figure 6 from the solid curve. Such excursions are the result of binary activity, such as mass transfer. A stable phase of mass transfer generally results in an increase in the binding energy of the binary. Such binaries first become very hard as the orbital period decreases, then soften when the mass of the donor drops below that of the accreting star. One example of this is clearly visible in Figure 6 (*right*), around  $t = 11$  Myr. The

phase of mass transfer stops after about 3 Myr, indicating that it is a case A (Kippenhahn & Weigert 1967), i.e. stable on a nuclear timescale (see also Portegies Zwart et al. 2001b). This episode stops when the donor leaves the main sequence, becomes an envelope-depleted helium-burning star and finally explodes in a supernova (*filled star*). The mass loss in the supernova, forming a black hole, causes the binary to be ejected from the cluster (*up-pointed arrow*).

The spikes to lower binding energy in both models R90W4 and R90W7 are the result of stars that receive a high velocity following a strong encounter. Such a high-energy encounter may decrease the binding energy of the entire cluster for as long as the star is considered a cluster member (within  $3r_{L1}$  from the cluster center). Model R90W7 produces many more high-velocity escapers than model R90W4, consistent with the greater dynamical activity within the more compact cluster.

### 3.3. Evolution of the Total Luminosity

The total luminosities of the model clusters decrease with time as the clusters dissolve in the Galactic tidal field and the most massive stars evolve off the main sequence and become dark remnants.

Figure 7 presents the integrated visual luminosity for the models R90W4 (see also Fig. 6 (*left*)) and R150W4 (both models had identical initial stellar masses [in  $M_{\odot}$ ], positions, and velocities [in scaled  $N$ -body units]). Initially the luminosity greatly exceeds  $10^6 L_{\odot}$ , then it drops steadily by more than 2 orders of magnitude as the cluster ages. After about 4 Myr the luminosity suddenly drops because of to the explosion of the most massive stars (for model R150W4) or because a few massive stars escape from the cluster as the binary in which they reside is ejected (for model R90W4, see also Fig. 6).

Occasionally the luminosity again exceeds  $10^6 M_{\odot}$  as a runaway collision product leaves the main sequence and becomes a luminous blue variable or a Wolf-Rayet star. (Paper III discusses the collision rate in models with comparable initial conditions. A summary is given in § 4.1.) In model R90W4 (Fig. 7 (*left*)) this happens around 9.5 Myr and 23–25 Myr, and for model R150W4 (Fig. 7 (*right*)) near  $\sim 6$  Myr and 16–22 Myr. In these episodes the total luminosity of the entire star cluster is dominated by a single




FIG. 5.—Evolution of the core radius (lower solid line) and the 25%, 50%, and 75% Lagrangian radii (dotted, dashed, and upper dotted lines, respectively) for model R90W4. The top line shows the instantaneous Jacobi radius of the cluster.




FIG. 6.—Evolution of the binding energy for models (*left*) R90W4 and (*right*) R90W7. The solid lines include the binding energies of all stars (only the center of mass energy of binaries and higher order systems are included), and the dotted lines include the binding energies of the binaries. Initially  $E_{\text{tot}} = -0.25$  (see § 2). The two lines with symbols in the horizontal bar above the figure indicate important happenings in the cluster lifetime. Triangles indicate the moment at which a binary or triple is formed by a three- (or more) body interaction, the filled and open stars indicate the formation of a black hole and neutron star, respectively. The filled circles indicate the moment a collision occurs and arrows pointing upward indicate the moment then a hard binary is ejected from the cluster. All these events are reflected in the evolution of the binding energy.

star. This episode suddenly stops when the star explodes or escapes from the cluster. The regular rises and sudden drops in luminosity (for example near  $t = 33$  Myr and  $t = 47$  Myr in Fig. 7 (*right*)) are caused by single stars that ascend the giant branch and then explode in supernovae (*open and filled stars*) or form a white dwarf (*circles*). The essential point, however, is that the cluster is brightest at early times. Later increases in brightness may be caused by the rejuvenation of stars in collisions, but in general the total luminosity decreases steadily until the cluster dissolves completely.

### 3.4. Fraction of Massive Stars in the Cluster Center

Figure 8 shows the evolution of the fraction of stars having masses greater than  $1 M_{\odot}$ , within the 5% Lagrangian

radius and near the 50% and 75% Lagrangian radii of model R90W4. Mass segregation causes the central mass function to flatten rapidly, but it takes considerably longer before the outer region of the cluster is significantly affected. After  $\sim 4$  million years, the mass function within the inner 5% Lagrangian radius contains about 4 times as many stars with masses exceeding  $1 M_{\odot}$  (relative to the local number density) than does the rest of the cluster. The mass function near the half-mass radius remains comparable to that of the cluster as a whole (see also Vesperini & Heggie 1997). The outer 75% of the cluster becomes slightly depleted of high-mass stars. Within one initial half-mass relaxation time, the fraction of stars having  $m > 1 M_{\odot}$  in this region falls by about a factor of 3. At later times, the fraction of high-mass stars throughout the cluster increases as a result of the pref-




FIG. 7.—Evolution of the visual luminosity of all stars within  $3r_{L1}$  for the model (*left*) R90W4 and model (*right*) R150W4. Both models were started with the same random seed and therefore have identical stellar masses, positions and velocities (in scaled  $N$ -body units). The solid lines include the luminosity of all stars. The two lines with symbols in the horizontal bar above the figure indicate important happenings in the cluster lifetime. Fig. 6 gives the explanation of the symbols (we omitted the information about close encounters since they bear little information, and in [*right*] we also omitted the escapers). The open circles in the upper bar of (*right*) indicate the moment when a super giant forms a white dwarf.





FIG. 8.—Fraction of stars having masses greater than  $M_{\odot}$  within the 0%–5% (upper line), 10%–50% (middle line), and 50%–90% (lower line) Lagrangian radii for model R90W4.

erential escape of low-mass stars; at disruption, the cluster is rich in high-mass stars, while low-mass stars are depleted. This is in agreement with the findings of Takahashi & Portegies Zwart (2000), who concluded that the observed flat mass function in the globular cluster NGC 6712 indicates that this cluster is close to dissolution.

### 3.5. Has the Arches Cluster an Unusually Flat Mass Function?

Figer, McLean, & Morris (1999b) studied the mass function of the Arches cluster in two projected annuli, the inner one spanning  $2''.5$  to  $4''.5$  from the cluster center ( $0.098$ – $0.18$  pc, assuming a distance of 8 kpc) and the outer one extending from  $4''.5$  to  $7''.5$  ( $0.18$ – $0.29$  pc). They found that the cluster mass function is much flatter than the Salpeter distribution (a power law with exponent  $x = -2.35$ ) and that the mass function in the inner annulus is even flatter ( $x = +1$  to  $-1.5$ ) than that in the outer parts ( $x = -1.5$  to  $-2.0$ ). They estimate completeness down to  $20 M_{\odot}$  in the inner annulus and down to  $10 M_{\odot}$  in the outer one. The total number of stars in the inner annulus is 50 down to  $20 M_{\odot}$ , for a total mass of  $2173 M_{\odot}$ . The total number of stars



in the outer annulus is 122 down to  $10 M_{\odot}$ , resulting in a total mass of  $3164 M_{\odot}$ .

Figure 9 shows the observed mass function in the inner and outer annuli and compares them with the mass functions of our models with  $W_0 = 4$  at 34 pc (dotted line), 90 pc (dashed line), and 150 pc (solid line) from the Galactic center. The projected distance was kept constant at 34 pc. The comparison in Figure 9 is performed at an age of 3 Myr. Each calculation was repeated three times to improve statistics. We further compensate for the uncertainty in cluster ages and distances from the Galactic center by using annuli that are somewhat narrower (from  $0.098$  to  $0.139$  pc and from  $0.18$  to  $0.235$  pc for the inner and outer annuli, respectively) than the observed annuli (from  $0.098$  to  $0.18$  pc for the inner annulus and from  $0.18$  to  $0.29$  pc for the outer annulus). We normalize to match a star cluster with  $10^5$  stars.<sup>6</sup>

The mass function for the cluster farther from the Galactic center is in better agreement with the observations; the models at  $r_{GC} = 34$  pc do not reproduce the observations well. In addition, the inner annulus is much better represented by both sets of models than is the outer annulus, probably because the latter is much more sensitive to the dynamical evolution of the cluster and the adopted distance from the Galactic center. The top end of the mass function ( $\geq 50 M_{\odot}$ ) is underrepresented in both annuli. This discrepancy is caused in part by the steep slope at the upper end of our selected initial mass function (which has an exponent  $-2.8$ , where Salpeter is  $-2.35$ ). We simply have not enough massive stars in our simulations. For stars more massive than  $\geq 50 M_{\odot}$ , the initial mass function in young star clusters may well be flatter than adopted by Scalo (1986).

This method of comparing mass functions is very sensitive to the total mass of the cluster, its age, and the actual distance between the cluster and the Galactic center. These three parameters are coupled via the relaxation time, and

<sup>6</sup> When comparing two stellar systems with a different number of stars a choice must be made about the time at which the comparison is carried out. The time can be expressed in units of the initial crossing time (or in million years, as in Fig. 9), or in terms of the initial relaxation time of the cluster. For a purely dynamical comparison the initial relaxation time may be more natural (for a discussion see Portegies Zwart et al. 1998).




FIG. 9.—Mass function for models R150W4 (solid line), R90W4 (dashed line), and R34W4 (dotted line) at an age of 3 Myr. The initial mass function of the whole cluster at birth is given by the dotted line (Scalo 1986). (left) Mass functions at the projected inner annulus ( $2''.5 < r < 4''.5$ ). (right) Mass function in the outer annulus ( $4''.5 < r < 7''.5$ ) from the cluster center. These annuli correspond to those in which Figer et al. (1999a) observed the mass function for this cluster (bullets). Arrows indicate lower limits due to incompleteness.

small changes in any of them may have a profound effect on the derived mass function for the real cluster.

In order to compare our model calculations more directly with the observations we introduce the ratio of the mean mass in the outer annulus to that in the inner annulus, which for the observations is  $f_{o/i} \equiv \langle m \rangle_{\text{out}} / \langle m \rangle_{\text{in}} \simeq 25.9 M_{\odot} / 43.5 M_{\odot} = 0.597$ . By presenting this value as a function of time (measured in units of the initial relaxation time of the cluster), we remove much of the observational bias.

This mean mass ratio  $f_{o/i}$  is a characteristic of the cluster. For a Scalo (1986) mass function at zero age and in a system with a homogeneous density the value of this fraction is  $f_{o/i} = 0.501$ .<sup>7</sup> As more massive stars sink to the cluster center owing to mass segregation the value of  $f_{o/i}$  changes. The value of  $f_{o/i}$  also changes as a result of the variation of the density profile in the cluster, and therefore is also a function of the distances from the cluster center at which  $f_{o/i}$  is measured. The shapes of these curves, however, are rather insensitive to the initial density profile because the evolution of the cluster mass (Fig. 2 (right)) and density (Fig. 4 (right)) do not depend sensitively on the adopted density profile. Because of the uncertainties in the masses and hence the length scales of our models, we compute  $f_{o/i}$  at various distances from the centers of our model ( $W_0 = 4$ ) clusters as functions of time.

We selected a total of six sets of inner and outer annuli, starting close to the cluster center and moving progressively outward. For the innermost annulus we selected  $r_{\text{in}} = 0.003125$ ,  $r_{\text{mid}} = 0.05625$ , and  $r_{\text{out}} = 0.09375$  (in units of the initial virial radius), and we increased the radii by factors of 2 for subsequent annuli out to  $r_{\text{in}} = 1.0$ ,  $r_{\text{mid}} = 1.8$  and  $r_{\text{out}} = 3.0$ . For each of our  $N$ -body calculations we computed  $f_{o/i}$  for all combinations of six inner and outer annuli. To improve statistics we then combined the values for the inner two (Fig. 10, *solid line*), the middle two (*dashed line*), and the outer two (*dotted line*) annuli.

Figure 10 plots these values of  $f_{o/i}$  as a functions of time for our model calculations with  $W_0 = 4$  and compares them with the observed mass function. We combined all models with  $W_0 = 4$  to improve the statistics at the top end of the mass function. We computed a total of 14 models with these parameters, of which eight were run for only 0.91 initial relaxation times and six continued until they dissolved (more than  $3 t_{\text{hrx}}$ ). The various models were scaled to the initial relaxation time before being superimposed, as explained in § 3. At later times ( $t \gtrsim t_{\text{hrx}}$ ) some effect from stellar evolution may be influencing this scaling, but since we are mostly concerned here with earlier times this does not affect our conclusions. The vertical bar representing the observations may be shifted along the dotted line. The best fit for all annuli is obtained in the interval  $0.05 \lesssim t_{\text{hrx}} \lesssim 0.2$ .

The value of  $f_{o/i}$  for the inner annulus in Figure 10 is much lower than for the outer annuli, but  $f_{o/i}$  steadily increases to match the outer annuli near  $t \simeq 0.05 t_{\text{hrx}}$ . In the same time interval  $f_{o/i}$  for the outer annuli decreases somewhat. This effect is caused by mass segregation, as the most massive stars sink to the cluster center more rapidly than somewhat less massive stars. On their way to the cluster center the most massive stars pass through the various annuli. The




FIG. 10.—Values for  $f_{o/i}$  (the mass ratio of the mean mass in the outer annulus divided by the mean mass in the inner annulus) as a function of the time in units of the initial half-mass relaxation time. Each of the various lines combines the results for two of the measured values for  $f_{o/i}$ . The solid line gives the sum of the inner two values (with  $r_{\text{mid}} = 0.06$  and  $0.11$ ), the dashed line gives the sum for the intermediate range ( $r_{\text{mid}} = 0.2$  and  $0.5$ ), and the dotted line gives the value of  $f_{o/i}$  for the outer annuli ( $r_{\text{mid}} = 0.9$  and  $1.8$ ). The horizontal dotted line gives the measured value of  $f_{o/i}$  as measured by Figer et al. (1999a) with the Poissonian error bar (right).

loss of a few massive stars has little effect on the value of  $f_{o/i}$  in the outer part of the cluster as these annuli contains many more stars than the inner annuli. The arrival of a few massive stars in the cluster center, however, carries a relatively larger contribution to the total mass in high-mass stars to the inner annulus. The inner  $f_{o/i}$  rises therefore more quickly than the value in the outer annuli decreases.

At later time  $t \gtrsim 0.2 t_{\text{hrx}}$  the value of  $f_{o/i}$  rises sharply for the inner annulus, while for the outer annuli  $f_{o/i}$  continues to decrease. The reason for this is the continuing effect of mass segregation; the more massive stars pile up in the cluster center, while the clusters' outer parts become depleted of massive stars. After  $t \simeq 0.4 t_{\text{hrx}}$ ,  $f_{o/i}$  for the inner annuli (Fig. 10, *solid line*) starts to decrease rapidly. The sharp peak in this curve is the result of the two competing effects: mass segregation causing  $f_{o/i}$  to increase and stellar evolution resulting in its decline. This strong decline in  $f_{o/i}$  is caused by the onset of stellar evolution as the model with the longest relaxation time ( $t_{\text{hrx}} = 13$  Myr at  $r_{\text{GC}} = 150$  pc; see Table 3) starts to run out of massive stars. The first decline starts at  $t \simeq 0.4 t_{\text{hrx}}$ , corresponding to an age of 5.2 Myr for the  $t_{\text{hrx}} = 13$  Myr model. This time corresponds to the nuclear lifetime of a  $34 M_{\odot}$  star. The lower mass limit used for the outer annuli is  $20 M_{\odot}$ . Such a star has a nuclear burning lifetime of about 9.1 Myr i.e., by the time  $t \gtrsim 0.70 t_{\text{hrx}}$  all stars more massive than  $20 M_{\odot}$  have experienced a supernova and  $f_{o/i}$  has consequently diminished. (The slightly later time at which the outer annuli run out of stars is caused by the inhomogeneous mixture of models used to make Fig. 10.)

The observed value of  $f_{o/i}$  (vertical bar) may be moved along the horizontal dotted line until it matches any of the curves representing the model calculations. A matching result indicates that the observed mass function at these annuli agrees with a cluster at the appropriate age in units of the initial relaxation time. We illustrate this with a few examples.

At later times ( $t \gtrsim 0.1 t_{\text{hrx}}$ ) the observations fit quite well with the solid line in Figure 10, representing the innermost

<sup>7</sup> The initial value of  $f_{o/i} < 1$  because of the different lower cutoffs in the nominator and the denominator. The selected cutoffs are identical to those in the observations, which are more affected by crowding in the inner annulus.

annuli. Initially, high-mass stars are undersampled in the inner part of the cluster, and only at later times does mass segregation cause them to sink to center of the potential well of the system, explaining the initial rise in  $f_{oi}$  for the innermost annuli. If indeed the observed value of  $f_{oi}$  coincides with the model calculations at the innermost part of the cluster, the observed cluster must be older than about  $t \gtrsim 0.1t_{\text{hrx}}$ . At an observed age of the Arches cluster of 1–2 Myr the initial relaxation time must then have been  $t_{\text{hrx}} \lesssim 20$  Myr. Using the intermediate annulus of  $r_{\text{mid}} = 0.18$  pc to set the size scale of the models, the half-mass radius of the Arches cluster is then  $r_{\text{hm}} \gtrsim 1.6$ –3.0 pc. A cluster with a half-mass radius of  $r_{\text{hm}} \gtrsim 1.6$  pc and a relaxation time  $t_{\text{hrx}} \lesssim 20$  Myr contains fewer than a hundred stars. Since the number of stars observed easily exceeds this number we can firmly reject this solution: the observed radius of  $r_{\text{mid}} = 0.18$  pc then corresponds to  $\lesssim 2r_{\text{vir}}$ .

Alternatively, we may observe the cluster when it is younger in terms of its relaxation time. In the very early evolution of the cluster  $t \lesssim 0.05t_{\text{hrx}}$  the observations are quite consistent with  $0.2 \lesssim r_{\text{mid}} \lesssim 1.8$  (Fig. 10, *dashed and dotted lines*), but is inconsistent with  $r_{\text{mid}} \lesssim 0.11$ . The intermediate radius in the observations corresponds to  $r = 4''.5$  (or about 0.18 pc at a distance of 8 kpc). With these values the cluster would be described accurately by  $r_{\text{vir}} = 0.1$ –0.9 pc at an age  $\lesssim 0.05t_{\text{hrx}}$ . With an age of the Arches cluster of 1–2 Myr (see Table 1) the initial relaxation time of the cluster must be larger than 20–40 Myr. With a half-mass radius  $r_{\text{vir}} \lesssim 0.9$  (corresponding to  $r_{\text{hm}} \simeq 0.75$  pc), the cluster then has a mass of about  $4000 M_{\odot}$  in order to produce a half-mass relaxation time of  $t_{\text{hrx}} \lesssim 40$  Myr. If  $r_{\text{vir}} = 0.1$  pc, the total mass is  $2.4 \times 10^6 M_{\odot}$  and the relaxation time is 20 Myr.

Taking the observed half-mass radius of  $r_{\text{hm}} \sim 0.2$  pc and correcting it for the effects of mass segregation and expansion during its early dynamical evolution to  $r_{\text{hm}} \sim 0.35$  pc, a relaxation time of 30 Myr implies a cluster of about 64k stars and a total mass of about  $40,000 M_{\odot}$ . These numbers would also agree with the observed number of massive stars (50 stars with  $m > 10 M_{\odot}$  in the inner annulus) if a normal Scalo (1986) mass function were adopted. We thus see no reason to invoke a flatter than usual mass function to explain the observations. This is consistent with the suggestion of Serabyn, Shupe, & Figer (1998) that the Arches contains  $\sim 10^5$  stars, far more than the 12k adopted in our models or as suggested by Figer et al. (1999a), as we discuss in § 4.

The limits on the initial parameters for the Arches cluster are summarized in Figure 11. The shaded ellipse shows the likely initial parameters for this star clusters.

Assuming that the density profile of the cluster is consistent with a King model with  $W_0 = 4$ –6 and adopting  $r_{\text{hm}} = 0.35$ , we derive a Jacobi radius of 1.6 pc–2.5 pc. Consequently the distance from the Arches cluster to the Galactic center is between 43 pc and 91 pc, for models with  $W_0 = 4$  and  $W_0 = 6$ , respectively. We conclude that the Arches cluster is likely to be at a distance from the Galactic center somewhat, but not much, greater than its projected distance. Table 4 reviews our conclusions about the Arches system.

### 3.6. The Density Profile

Figure 12 shows the projected density profile of our model clusters with  $W_0 = 4$  for stars having masses  $m \gtrsim 20$




FIG. 11.—Graphical representation of the constraints for our model of the Arches cluster. The horizontal and vertical axis give the half-mass radius and initial cluster mass, respectively. The diagonal lines are of constant relaxation time at the half-mass radius (eq. [3]). The two error crosses locate the observed parameters for the Arches and Quintuplet star clusters. The ellipse identifies the parameter space in which our model calculations are consistent with the observed mass function, number of high-mass stars and density structure of the observations for the Arches cluster (see discussion in § 3.5).

$M_{\odot}$  at various instants. The data are presented in virial units (horizontal axis in units of  $r_{\text{vir}}$ , and the vertical axis in units of stars per  $r_{\text{vir}}^2$ ). The various curves (and symbols) represent the density distribution of the model clusters at zero age (*solid line*), at  $t = 0.05$  initial relaxation times (*dashed line*), and at  $t = 0.5t_{\text{hrx}}$  (*dotted line*). Mass segregation causes the density profile of the older clusters to steepen with time, and tidal stripping causes the global density to decrease.

The open squares in Figure 12 present the observed density distribution for the Arches cluster (Figer et al. 1999a). These data are given in arcseconds (or parsec when scaled to the appropriate distance). We plot them on the same figure assuming that the observed median radius of 0.18 pc ( $4''.5$ ) corresponds to  $0.18r_{\text{vir}}$ , i.e., that the virial radius of the Arches cluster is 1 pc. This may be an overestimate, as discussed in the previous section. There is some

TABLE 4

DERIVED PARAMETERS FOR THE ARCHES CLUSTER RESULTING FROM OUR BEST VALUES FROM OUR MODEL CALCULATIONS ASSUMING A NORMAL MASS FUNCTION COMPARABLE TO THAT OF THE SOLAR NEIGHBORHOOD<sup>a</sup>

Parameter	Value
$M (M_{\odot})$ .....	$\sim 40,000$
$N$ .....	$\sim 65,000$ stars
$r_{\text{GC}}$ (pc) .....	43–91
$t_{\text{hrx}}$ (Myr) .....	20–40
Age ( $t_{\text{hrx}}$ ) .....	$0.05 \pm 0.02$
$r_{\text{hm}}$ (pc) .....	$0.35 \pm 0.05$
$r_{\text{vir}}$ (pc) .....	$0.42 \pm 0.05$
$W_0$ .....	4–6
$r_{\text{tide}}$ (pc) .....	1.6–2.5

<sup>a</sup> These numbers are derived in § 3.5 and discussed in the §§ 3.6 and 3.7.




FIG. 12.—Projected density profile of the Arches cluster for stars with  $m > 20 M_{\odot}$  from the observations of Figer et al. (1999a; data from Kim et al. 2000 [open squares]). These data were scaled to a virial radius of 1 pc. The arrow starting from the upper left most square indicates the direction in which to shift these symbols when the size scaling (in arcseconds) is increased by  $1 \sigma$ . They adopted the inner annulus between  $2.5 < r < 4.5$  ( $0.098 < r < 0.18$  pc) and an outer annulus between  $2.5 < r < 4.5$  arcseconds ( $0.18 < r < 0.29$  pc). The various other lines and symbols give the results of our  $N$ -body calculations with  $W_0 = 4$ . The various lines represent the density profile at the moment indicated (top right-hand corner). These moments are presented in units of the initial relaxation time of our models, ranging from zero age (solid line) to  $0.5 t_{\text{hrx}}$ . The densities for our model calculations were increased with a factor 64/12 to account for the larger number of stars expected in these clusters. In the lower left-hand corner is a  $1 \sigma$  Poissonian error bar for reference.

freedom in shifting the observed points along the arrow, which then changes the scaling for the cluster.

The observed points fit best with the dashed line and are not consistent with the solid (zero age) or the dotted ( $t = 0.5 t_{\text{hrx}}$ ) lines. The projected density profile of the observed Arches cluster thus suggests an age of about  $t \simeq 0.05 t_{\text{hrx}}$ , in agreement with our findings in § 3.5.

### 3.7. Distances to the Galactic Center

The Arches cluster is located at a projected distance of  $\sim 30$  pc from the Galactic center, the Quintuplet at  $\sim 35$  pc. These measurements provide lower limits to the true distances of these clusters from the center.

The mean density within the Jacobi radius of a tidally limited cluster is proportional to the local stellar density. (This is not a matter of definition—it happens to be true for point-mass fields if one averages the point mass over  $r_{\text{GC}}$ , and for the power-law density profiles we consider here, since the tidal field  $\sim M/R^3 \sim M_G/r_{\text{GC}}^3 \sim r_{\text{GC}}^{-2} dM_G/dr_{\text{GC}} \simeq \rho_g$ , but it is not in general the case.) Thus, a cluster closer to the Galactic center is more compact, has a shorter relaxation time, and therefore evolves more rapidly than a similar cluster at a greater distance.

Most of our 34 pc models have densities higher than any of the observed clusters listed in Table 4 (see Fig. 4). Only the model with an extremely shallow initial density profile (R34W1) has density comparable to the observed systems. Clusters at greater distances from the Galactic center (but with the same projected distance) have lower densities. Since the densities of the 34 pc models are too high by at least a factor of 2, this suggests that the real clusters are somewhat farther out, at  $r_{\text{GC}} \gtrsim 50$  pc; a factor of 2 in density corresponds to a factor of  $2^{1/3}$  in radius, so the true Galactocentric distance should exceed 43 pc.

Figure 13 plots the evolution of the mean surface density within the projected half-mass radius for models R34W7 (dotted line), R90W4 (dashed line), and R150W1 (solid line), which brackets the central densities of the models. We view the clusters along the  $x$ -axis, so we look through the second and first Lagrangian points toward the Galactic center, giving the highest possible cluster density and hence an overestimate of the true density contrast (see Fig. 1). The two error bars give the projected half-mass densities for the Arches (left) and the Quintuplet (right) clusters. The horizontal dotted lines give the background surface densities a projected distances of 34, 90, and 150 pc from the Galactic center.

The surface density at projected distance  $d$  from the Galactic center can be calculated by integrating the local stellar density  $\rho_G$  (eq. [10]) along the line of sight to the cluster:

$$\Sigma(d) = 4.06 \times 10^5 \int (d^2 + z^2)^{-0.9} dz M_{\odot} \text{ pc}^{-2}. \quad (14)$$

Integrating equation (14) from  $-100$  to  $100$  pc (the range of validity of the Mezsger's equation) gives the projected surface densities toward the Arches (at  $d = 30$  pc) and Quintuplet cluster (at  $d = 35$  pc). We integrate equation (14) numerically. Figures 14 and 15 illustrate this. An extra correction for stars between 100 pc from the Galactic center and Earth adds little ( $\lesssim 10\%$ ) to the total (Bahcall & Soneira 1980). The projected densities of the observed clusters are comparable to the background. Clusters with lower background densities may well remain unnoticed among the background stars. Note, however, that the background stellar population is probably older than the star clusters studied here, and may therefore have a smaller mass-to-light ratio, making the clusters stand out somewhat better. This may be the reason that the Quintuplet was found even though its projected density is smaller than that of the background.

Figure 14 shows equations (14) and (10) both normalized to their value at  $r_{\text{GC}} = 5$  pc, which are  $\Sigma(r_{\text{GC}} = 5 \text{ pc}) = 1.6 \times 10^5 M_{\odot} \text{ pc}^{-2}$  and  $\rho_G(r_{\text{GC}} = 5 \text{ pc}) = 1.6 \times 10^4 M_{\odot} \text{ pc}^{-3}$ .




FIG. 13.—Evolution of the projected density within the half-mass radius for models R34W7 (dotted line), R90W4 (dashed line), and R150W1 (solid line). The dotted horizontal lines give the surface density at a projected distance of 34, 90, and 150 pc (as indicated) from the Galactic center. The selected cases are quite extreme; R34W7 being the most concentrated model and R150W1 the least concentrated. The two error bars (to the left) indicate the location of the Arches (left) and the Quintuplet (right) clusters.




FIG. 14.—Projected (solid line; eq. [14]) and three dimensional (dashed line; eq. [10]) density as a function of the distance to the Galactic center. Both densities are normalized to a distance of  $r_{GC} = 5$  pc,

The projected density of model R150W1 remains well below the background, and such a cluster could easily remain unseen throughout its entire lifetime. The two initially more concentrated models, R90W4 and R34W7, have projected densities above the background, at least initially. The cluster farther from the Galactic center has a lower initial density because it is more extended; after the first few million years, it may become hard to see. The observation of the Quintuplet provides an upper limit on the critical contrast below which the cluster cannot be detected. We arbi-




FIG. 15.—Cartoon of the location of the Arches (left) and the Quintuplet (far left) clusters with respect to the Galactic center. Earth is below. The large partial circle has a radius of 100 pc. The ellipses representing the clusters are exaggerated by a factor of 10.

trarily adopt a minimum contrast equals to the surface density of the Quintuplet cluster (i.e.,  $\sim 10^4 M_{\odot} \text{pc}^{-2}$ ) as a threshold for distinguishing a star cluster from the background. In that case, the 150 pc cluster would be invisible for its entire lifetime, while the 34 and 90 pc clusters remain visible for about  $\sim 8$  million years ( $\sim 65\%$  and  $\lesssim 25\%$  of their respective lifetimes). Thus, although the clusters farther from the Galactic center live much longer, their visible lifetime is actually less than that of a cluster at smaller Galactocentric radius.

Although not shown in Figure 13, the surface density evolution of model R34W1 is also consistent with the densities of the observed clusters. On these grounds we therefore cannot exclude the possibility that the Arches and Quintuplet systems lie at Galactocentric distances of 34 pc, but in that case they must have been born with very shallow density profiles (which, however, would contradict our earlier analysis from § 3.5; see Table 4).

The surface mass density as presented in Figure 13 may not be the ideal way to express the visibility of these clusters; a luminosity density contrasted with the background would be more appropriate. This, however, requires a detailed study of the observational selection effects, which goes beyond the scope of this paper. Another way to illustrate the visibility of clusters like Arches is by looking at the integrated visual luminosity of these clusters (see Fig. 7).

#### 4. DISCUSSION

The Arches and the Quintuplet are among the youngest star clusters known in the Galaxy. They are not as rich as small globular clusters, but are considerably more compact. Their central densities are comparable to those of the densest—postcollapse—globulars, but their life expectancies are only a few tens of millions of years. These clusters are destroyed by tidal forces, accelerated by impulsive mass loss from supernovae (after  $\sim 3$  Myr) and strong binary activity. In the following we first focus on comparison with other calculations; we then discuss the consequences of our results for the inferred distances of the Arches and Quintuplet clusters from the Galactic center and for the number of such clusters that may be hidden near the Galactic center.

##### 4.1. Comparison with Other Work

Portegies Zwart et al. (Paper III) studied the evolution of R136 in the LMC using  $N$ -body simulations. Their initial conditions were comparable to those adopted here, although small differences exist. They adopted King models with  $W_0 = 6$ , whereas we selected  $W_0 = 4$  and  $W_0 = 7$ .

The calculations in Paper III included the effects of stellar evolution and physical collisions between stars, but neglected the effect of the Galactic tidal field and the possible presence of primordial binaries. Both may have a profound effect on a cluster's dynamical evolution. For R136 the neglect of a tidal field may be appropriate, but for the Arches and Quintuplet clusters the tidal field is crucial. Our clusters therefore dissolve much more rapidly than the calculations in Paper III. The latter models lost only a few percent of their mass within the first relaxation time, where our models lose up to  $\sim 50\%$  on this timescale, confirming the strong influence of the Galactic tidal field.

Portegies Zwart et al. (1999) found that physical collisions between stars in their models were frequent and that the evolution of the most massive stars and the dynamical

evolution of the cluster were closely coupled. In all cases, a single star grew steadily in mass through mergers with other stars, forming a very massive ( $\gtrsim 100 M_{\odot}$ ) star in less than 3–4 Myr. The growth rate of this runaway merger was much larger than estimates based on simple cross section arguments, mainly because the star was typically found in the core and tended to form binaries with other massive stars there. We observe the same general behavior in our calculations, in the sense that collisions tend to occur between high-mass ( $m \gtrsim 10 M_{\odot}$ ) stars and that most collisions occur repeatedly with the same star. Although not discussed in detail in this paper, we find the same trend in our calculations and confirm the findings of Portegies Zwart et al. (1999). Kim, Morris, & Lee (1999, hereafter KML99) performed two-dimensional Fokker-Planck calculations of dense star clusters near the Galactic center and studied the lifetimes of the Arches and the Quintuplet clusters. Their calculations are somewhat different than those presented in this paper in the sense that they only solve the two-dimensional Fokker-Planck equation instead of the equation of motions for all stars in the stellar system. KML99 include standard binary heating (Lee, Fahlman, & Richer 1991) in their calculations and neglect stellar collisions.

We selected the most representative cases of KML99 and recalculated these with our model. The Fokker Planck models dissolve much more rapidly (often more than a factor 2 in lifetime) than the  $N$ -body models (see Fig. 16). Similar discrepancies are observed in the comparison between the Fokker-Planck calculations of Chernoff & Weinberg (1990) and the  $N$ -body calculations of Fukushima & Heggie (1995) and Portegies Zwart et al. (1998). The discrepancies are the result of the rather simple treatment of escapers in the Fokker-Planck models, which can drive the evaporation of the cluster on a shorter timescale than when a self-consistent tidal field is used (see Takahashi & Portegies Zwart 1998 for details). An extra complication is introduced in the calculations of KML99 by selecting a maximum mass of  $150 M_{\odot}$  in the initial mass function. Such

stars dominate the early evolution of the star cluster, while their behavior cannot accurately be represented with a statistical average as is done in a Fokker-Planck solver. Part of the spread in the Fokker-Planck results in Figure 16 may be attributed to these effects.

Kim et al. (2000) also perform  $N$ -body calculations of several of the models explored by KML99 using Aarseth's (1999) NBODY6. They also find that the  $N$ -body models live longer than the Fokker-Planck models, but arrive at the same conclusion as KML99: the Arches and Quintuplet star clusters must have unusually flat initial mass function and dissolve within about 10 Myr. The results of KML99 are completely understandable from their choice of initial conditions. Many of their calculations start with a power-law initial mass function with an exponent as flat as  $-1.5$  (Salpeter =  $-2.35$ ) and omitting all stars with a mass smaller than  $1 M_{\odot}$ . Together with a small number of stars of only a few thousand, their models have an initial relaxation time at the tidal radius of only  $t_{\text{trx}} = 5$  to 60 Myr. The lifetime of these clusters is therefore expected to be only a few million years.

Figure 16 compares the results of our calculations (*filled circles and triangles*) with the Fokker-Planck calculations of KML99 (*crosses*) and the  $N$ -body calculations of Kim et al. (2000) (*squares*). For comparison we added the results of the analytic model presented by Portegies Zwart et al. (2001a) as a solid line. The lowest values of our models (*lower filled circles for each  $t_{\text{trx}}$* ) are computed with  $W_0 = 1$ , which dissolve somewhat quicker than the more concentrated models. The lifetime of the  $N$ -body models start to depend on  $N$  via the Coulomb logarithm (see eq. [3]) when the number of stars in the models drops below a few thousand. In this small  $N$  limit the scaling proposed by Portegies Zwart et al. (2001a) probably breaks down. This effect may cause our models with the shortest initial relaxation times to live longer than expected (leftmost triangles in Fig. 16). A similar effect may cause the models of Kim et al. (2000; *open squares*) to have somewhat shorter lifetimes than expected, as most of their models contain less than 3000 stars. Part of the discrepancy at small  $N$  and therefore at small relaxation times may also be caused by statistical fluctuations, which become most noticeable at small  $N$ .

#### 4.2. How Many Clusters are Still Hidden Near the Galactic Center

Although the Arches and Quintuplet clusters are very dense, it may still be difficult to distinguish them near the Galactic center. The three-dimensional density contrast is several orders of magnitude, as can be seen in Figure 4. In projection on the sky, however, the density contrast is reduced, mainly because of the accumulation of stars along the line of sight (see Fig. 13). The cluster density within the projected half-mass radius also decreases with time, so these clusters are more easily seen at early ages than at later times. As outlined in § 3.7, to estimate the time over which our model clusters would be observable, we simply compare the integrated stellar density  $\Sigma(d)$  along the line of sight to the cluster (eq. [14]) with the projected stellar densities of the models.

In § 3.7 we introduced a limiting density contrast above which a cluster can be discriminated from the background stars and used this to estimate the time during which our models would be detectable. At a Galactocentric distance of 34 pc, our most compact models (R34W7) remain visible for




FIG. 16.—Cluster lifetime as a function of the initial relaxation time at the tidal radius for the model calculations presented in Table 3 (nine models; *filled circles*), in Kim, Morris, & Lee (*crosses*), and in Kim et al. (2000; *open squares*). The solid line ( $t_{\text{end}} = 0.25t_{\text{trx}}$ ) presents the scaling of the models as proposed by Portegies Zwart et al. (2001a). The triangles are calculated as an extra check on the behavior of the models when  $N \lesssim 3000$ , which represents the parameter space covered by Kim et al. (2000; their models 101, 111, 112, and 141).

$\sim 9$  Myr; at 90 pc they become invisible after about 7 Myr, and at a distance of 150 pc from the Galactic center they remain invisible for their entire lifetime. Models at greater distances from the Galactic center, or those born with shallower density profiles, remain undetectable for their entire life times. It is therefore not surprising that the clusters we observe near the Galactic center are extremely compact and very young. Less compact or older clusters are unobservable owing to their low surface density contrasts.

Portegies Zwart et al. (2001a) studied the timescale over which clusters such as Arches and the Quintuplet systems remain visible. The results of their detailed  $N$ -body calculations are used to calibrate a simple analytical model that is applicable over a wider range of cluster initial conditions. They conclude that clusters within 200 pc of the Galactic center dissolve within  $\sim 70$  Myr. However, their projected densities drop below the background density in the direction of the Galactic center within  $\sim 20$  Myr, effectively making these clusters undetectable after that time. Clusters farther from the Galactic center but at the same projected distance are more strongly affected by this selection effect, and may go undetected for their entire lifetimes. Based on these findings, they conclude that the region within 200 pc of the Galactic center could easily harbor some 50 clusters with properties similar to those of the Arches or the Quintuplet systems. The results of our more extended parameter study is consistent with their findings. Given the higher mass we derive for the Arches cluster (see Table 4) compared to what was adopted by Portegies Zwart et al. (2001a), we argue that their results may even be somewhat conservative.

Another effect which may contribute to the difficulty in detecting clusters like the Arches is the enormous range in luminosity of the brightest stars. CCD cameras have a dynamic range of less than  $2^{16} = 64k$ , causing the brightest stars to saturate the detector and preventing faint stars from being detected. However, these brightest stars are also the least common; only with very deep exposures is the rest of the cluster revealed. Thus, wherever two or more bright blue stars are seen together, there may be an entire star cluster lurking in the background. An example of such a cluster may be the small group of stars identified as R140.<sup>8</sup>

Figure 7 presented the integrated luminosity for two representative models (R90W4 and R150W4). In these models the total luminosity dropped by more than 2 orders of magnitude over the life span of the cluster. The individual high peaks are the results of single stars that become extremely bright as a result of their rejuvenation in several collisions. Such stars can temporarily outshine the rest of the star cluster. Of course, the luminosity and the lifetime of such a multiple collision product is quite sensitive to details in modeling the collision process.

## 5. CONCLUSION

We have studied the evolution of young, dense star clusters near the Galactic center, taking the Arches and Quintuplet clusters as specific examples. These clusters are generally referred to as “young,” because they are only a few million years old, and indeed, even in a dynamical sense

both clusters are quite young, having lifetimes considerably smaller than their initial relaxation times. However, by observing only the most massive stars in these clusters the dynamical image sketched by the observers is biased toward greater age, as the massive stars evolve dynamically on shorter timescales than average cluster members. By observing only stars with  $m \gtrsim 10 M_{\odot}$ , one selects a part of the cluster that is dynamically rather mature, being comparable in age to the local initial relaxation time. This makes the Arches cluster (and probably also the Quintuplet) appear dynamically older than it really is. We cannot test this hypothesis for the Quintuplet system because the available data are of lower quality than for the Arches.

The modeled clusters lose mass at a more-or-less constant rate, inversely proportional to the initial relaxation time, i.e.  $\propto r_{\text{hm}}^{-3/2} \propto r_{\text{GC}}^{-0.9}$  (see also Portegies Zwart et al. 2001a). Star clusters which are born farther from the Galactic center live therefore considerably longer than those closer in. The relaxation time derived from observation, however, contains little information about the initial relaxation time.

Mass segregation in our model clusters quickly causes the most massive stars to sink to the cluster center. As a result core collapse occurs within about two million years, even for models as shallow as  $W_0 = 1$ . During core collapse close binaries are formed and collisions between stars are frequent. The collision rate is much higher than would be expected from simple cross section arguments, owing to mass segregation and binary formation. We confirm the finding of Portegies Zwart et al. (1999) that the most massive stars are generally involved in a collision runaway, in which few low-mass stars participate. This process continues until the runaway product is ejected by a supernova or a strong encounter with a binary or the cluster dissolves in the tidal field of the Galaxy.

By comparing mass functions from our models at various distances from the cluster center with observations of the Arches mass function, we conclude that the age of the Arches cluster must be about  $0.05 t_{\text{hrx}}$ . With an age of about 1.5 Myr the initial relaxation time of the cluster is then about 30 Myr. To reconcile the observed mass function with the observed density distribution of the Arches, we conclude that the cluster must be about  $4 \times 10^4 M_{\odot}$  and lie some 50–90 pc from the Galactic center. With these parameters we can reproduce the observed density distribution and the observed mass function for stars between 10 and  $50 M_{\odot}$  without requiring an unusually flat initial mass function. The top end of the mass function contains more stars than our adopted mass function and this may require a flatter  $\lesssim -2.8$  (Salpeter =  $-2.35$ ) initial mass function for stars  $\gtrsim 50 M_{\odot}$ . However, we still conclude that the Arches cluster can be explained with a “normal” mass function. Most of the observed flattening of the mass function can be attributed to selection effects caused by the limited range in distance from the cluster center in which the observations were made.

The young clusters we discussed expand as they become older, causing their surface densities and total luminosity to drop. Clusters which are older than about 5 Myr often have surface densities below that of their surrounding, making these clusters virtually undetectable. By comparing the projected density of the model clusters with that of the field stars in the direction of the Galactic center we conclude that the region within 200 pc of the Galactic center could easily

<sup>8</sup> R140 contains at least two WN stars and one WC star (Moffat et al. 1987) located at a projected distance of 11.5 pc from R136. One of the WN stars (R140a) is a bright  $\log L_{0.2-3.5 \text{ keV}} = 10^{35.2}$  ergs  $\text{s}^{-1}$  X-ray source, possibly a colliding wind binary (Portegies Zwart, Pooley, & Lewin. 2001c).

harbor some 50 clusters with properties similar to those of the Arches or the Quintuplet systems. To find more clusters like Arches and Quintuplet systems, we urge observers to look for single extremely bright stars. Closer study of the region near such a star may well reveal the underlying star cluster.

Finally, with our estimate of about 50 clusters with a mass of  $\sim 40,000 M_{\odot}$  each and dissolving within 100 Myr, we derive a star formation rate near the Galactic center of  $\sim 0.02 M_{\odot} \text{ yr}^{-1}$ . Clusters like the Arches and Quintuplet may therefore contribute significantly to the total star formation rate in the Galaxy.

We thank Rob Olling and Mark Morris for discussions, and the excellent comments of the anonymous referee. This work was supported by NASA through Hubble Fellowship

grant HF-01112.01-98A awarded by the Space Telescope Science Institute, by ATP grants NAG 5-6964 and NAG 5-9264, and by the Research for the Future Program of Japan Society for the Promotion of Science (JSPS-RFTP97P01102). S. P. Z. is grateful to Drexel University, Tokyo University, and to the Institute for Advanced Study for their hospitality and the use of their fabulous hardware. Part of this letter was written while we were visiting the American Museum of Natural History. We acknowledge the hospitality of their astrophysics department and visualization group. We also acknowledge the expert visualization help offered to us by Stuart Levy, from the Virtual Director Group, National Center for Supercomputing Applications, University of Illinois at Urbana-Champaign. We thank the Alfred P. Sloan Foundation for a grant to Hut for observing astrophysical computer simulations in the Hayden Planetarium at the Museum.

## REFERENCES

- Aarseth, S. J. 1999, *PASP*, 111, 1333  
 Bahcall, J. N., & Soneira, R. M. 1980, *ApJS*, 44, 73  
 Baumgardt, H. 2000, in *MNRAS*, 325, 1323  
 Blum, R. D., Damiani, A., & Conti, P. S. 1999, *AJ*, 117, 1392  
 Brandl, B., Brandner, W., Eisenhauer, F., Moffat, A. F. J., Palla, F., & Zinnecker, H. 1999, *A&A*, 352, L69  
 Brandl, B., et al. 1996, *ApJ*, 466, 254  
 Campbell, B., et al. 1992, *AJ*, 104, 1721  
 Chernoff, D. F., & Weinberg, M. D. 1990, *ApJ*, 351, 121  
 Dutra, C. M., & Bica, E. 2000, *A&A*, 359, L9  
 Figer, D. F., Kim, S. S., Morris, M., Serabyn, E., Rich, R. M., & McLean, I. S. 1999a, *ApJ*, 525, 750  
 Figer, D. F., McLean, I. S., & Morris, M. 1999b, *ApJ*, 514, 202  
 Fukushige, T., & Heggge, D. C. 1995, *MNRAS*, 276, 206  
 Giersz, M., & Heggge, D. C. 1994, *MNRAS*, 268, 257  
 ———. 1996, *MNRAS*, 279, 1037  
 Glass, I. S., Catchpole, R. M., & Whitelock, P. A. 1987, *MNRAS*, 227, 373  
 Heggge, D. C., & Mathieu, R. 1986, in *The Use of Supercomputers in Stellar Dynamics*, ed. S. McMillan (Berlin: Springer), 233  
 Heggge, D. C., & Ramamani, N. 1995, *MNRAS*, 272, 317  
 Hill, G., Hilditch, R. W., & Barnes, J. V. 1979, *MNRAS*, 186, 813  
 Kim, S. S., Figer, D. F., Lee, H. M., & Morris, M. 2000, *ApJ*, 545, 301  
 Kim, S. S., Morris, M., & Lee, H. M. 1999, *ApJ*, 525, 228  
 King, I. R. 1966, *AJ*, 71, 64  
 Kippenhahn, R., & Weigert, A. 1967, *Z. Astrophys.*, 65, 251  
 Kulkarni, S. R., Matthews, K., Neugebauer, G., Reid, I. N., van Kerkwijk, M. H., & Vasisht, G. 1995, *ApJ*, 440, L61  
 Lee, H. M., Fahlman, G. G., & Richer, H. B. 1991, *ApJ*, 366, 455  
 Makino, J. 1991, *ApJ*, 369, 200  
 Makino, J., & Aarseth, S. J. 1992, *PASJ*, 44, 141  
 Makino, J., Taiji, M., Ebisuzaki, T., & Sugimoto, D. 1997, *ApJ*, 480, 432  
 Massey, P., & Hunter, D. A. 1998, *ApJ*, 493, 180  
 McMillan, S. L. W. 1986a, *ApJ*, 306, 552  
 ———. 1986b, *ApJ*, 307, 126  
 Mezger, P. G., Zylka, R., Philipp, S., & Launhardt, R. 1999, *A&A*, 348, 457  
 Moffat, A. F. J. 1976, *A&A*, 50, 429  
 Moffat, A. F. J., Niemela, V. S., Phillips, M. M., Chu, Y., & Seggewiss, W. 1987, *ApJ*, 312, 612  
 Nagata, T., Woodward, C. E., Shure, M., & Kobayashi, N. 1995, *AJ*, 109, 1676  
 Nagata, T., Woodward, C. E., Shure, M., Pipher, J. L., & Okuda, H. 1990, *ApJ*, 351, 83  
 Okuda, H., et al. 1990, *ApJ*, 351, 89  
 Olling, R. P., & Merrifield, M. R. 1998, *MNRAS*, 297, 943  
 Oort, J. 1927, *Bull. Astron. Inst. Netherlands*, 3, 275  
 Portegies Zwart, S. F., Hut, P., Makino, J., & McMillan, S. L. W. 1998, *A&A*, 337, 363  
 Portegies Zwart, S. F., Hut, P., & Verbunt, F. 1997, *A&A*, 328, 130  
 Portegies Zwart, S. F., Makino, J., McMillan, S. L. W., & Hut, P. 1999, *A&A*, 348, 117  
 ———. 2001a, *ApJ*, 546, L101  
 Portegies Zwart, S. F., McMillan, S. L. W., Hut, P., & Makino, J. 2001b, *MNRAS*, 321, 199  
 Portegies Zwart, S. F., Pooley, D., & Lewin, W. 2001c, *ApJ*, in press (astro-ph/0106109)  
 Portegies Zwart, S. F., & Verbunt, F. 1996, *A&A*, 309, 179  
 Scalo, J. M. 1986, *Fundam. Cosmic Phys.*, 11, 1  
 Serabyn, E., Shupe, D., & Figer, D. F. 1998, *Nature*, 394, 448  
 Spitzer, L. 1987, *Dynamical Evolution of Globular Clusters* (Princeton: Princeton Univ. Press), 35  
 Takahashi, K., & Portegies Zwart, S. F. 1998, *ApJ*, 503, L49  
 ———. 2000, *ApJ*, 535, 759  
 van Kerkwijk, M. H., Kulkarni, S. R., Matthews, K., & Neugebauer, G. 1995, *ApJ*, 444, L33  
 Vesperini, E., & Heggge, D. C. 1997, *MNRAS*, 289, 898  
 Vrba, F. J., Henden, A. A., Luginbuhl, C. B., Guetter, H. H., Hartmann, D. H., & Klose, S. 2000, *ApJ*, 533, L17