



## UvA-DARE (Digital Academic Repository)

### Temporal aggregation in a periodically integrated autoregressive process

Franses, P.H.; Boswijk, H.P.

**DOI**

[10.1016/0167-7152\(95\)00225-1](https://doi.org/10.1016/0167-7152(95)00225-1)

**Publication date**

1996

**Published in**

Statistics & Probability Letters

[Link to publication](#)

**Citation for published version (APA):**

Franses, P. H., & Boswijk, H. P. (1996). Temporal aggregation in a periodically integrated autoregressive process. *Statistics & Probability Letters*, 30(3), 235-240.  
[https://doi.org/10.1016/0167-7152\(95\)00225-1](https://doi.org/10.1016/0167-7152(95)00225-1)

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.



A078352398  
NCC/IBL AANVRAAGBON

KOPIE PERIODIEK EGB

(7)  
18-04-2005

Datum indienen : 15-04-2005 18:36                    5493-1    Clearing House  
Datum plaatsen : 15-04-2005 18:36  
Aanvrager : 0004/9998  
Aanvraagident :  
Aanvragerident : 0004/9999  
Eindgebruiker : 041631433

Telefoonnummer : 050-3635057  
Cooperatiecode : R

Leverwijze : Elektronisch  
Fax :  
Ftp :  
E-Mail : m.s.van.delden@rug.nl  
Ariel :

Plaatscode : 559301545 ; WK 3296 ; ; 1985 V3 - 2000 V50

- |                            |                                |
|----------------------------|--------------------------------|
| (1) [ ] Origineel gestuurd | (6) [ ] Niet beschikbaar       |
| (2) [ ] Copie gestuurd     | (7) [ ] Uitgeleend             |
| (3) [ ] Overige            | (8) [ ] Wordt niet uitgeleend  |
| (4) [ ] Nog niet aanwezig  | (9) [ ] Bibliografisch onjuist |
| (5) [ ] Niet aanwezig      | (0) [ ] Bij de binder          |




A078352398  
NCC/IBL AANVRAAGBON

Verzamelnota volgt.  
KOPIE PERIODIEK EGB

18-04-2005

Datum indienen : 15-04-2005 18:36                    9560-1    UvA Keur  
Datum plaatsen : 15-04-2005 18:36                    UB Groningen  
Aanvrager : 0004/9998                                    Broerstraat 4  
Aanvraagident :  
Aanvragerident : 0004/9999                                9700 AN Groningen  
Eindgebruiker : 041631433                                tav

Aantal



PPN Titel : 559301545  
Titel : Statistics & probability letters  
Auteur :  
Deel/Supplem. :  
Corporatie :    Externe database:  
Jaar/Editie : 1982                                        Extern nummer :  
Uitgave : Amsterdam                                    North-Holland Publishing Company  
Serie/Sectie :  
Pag-ISSN/ISBN :    0167-7152

Plaatscode : 559301545 ; WK 3296 ; ; 1985 V3 - 2000 V50

Jaar : 1996-00-00  
Volume : 30  
Aflevering : 3  
Eindgebruiker : 041631433                    Aanvraagident. :  
Auteur : Franses, P.H. (ed.)nvragerident. : UVA KEUR (UB GRONINGEN)  
Artikel : Temporal aggregation in a periodically integrated autoregressiv  
Bladzijden : 235-240  
Bron :  
Opmerking : ARNO:19934



ELSEVIER

Statistics & Probability Letters 30 (1996) 235–240

**STATISTICS &  
PROBABILITY  
LETTERS**

## Temporal aggregation in a periodically integrated autoregressive process

Philip Hans Franses<sup>a,\*</sup>, H. Peter Boswijk<sup>b</sup>

<sup>a</sup>*Econometric Institute, Erasmus University Rotterdam, P.O. Box 1738, NL-3000 DR Rotterdam, Netherlands*

<sup>b</sup>*Department of Econometrics, University of Amsterdam, Netherlands*

Received July 1995; revised October 1995

---

### Abstract

A periodically integrated autoregressive process for a time series which is observed  $S$  times per year assumes the presence of  $S - 1$  cointegration relations between the annual series containing the seasonal observations, with the additional feature that these relations are different across the seasons. This means that there is a single unit root in the vector autoregression for these annual series. In this paper it is shown that temporally aggregating such a process does not affect the presence of this unit root, i.e. the aggregated series is also periodically integrated.

---

### 1. Introduction

Temporal aggregation in flow-type univariate time series processes is studied, e.g. in Amemiya and Wu (1972) and Weiss (1986) with respect to the order of the ARMA process for the aggregated time series. Recently, the focus is also on the effects of temporal aggregation on unit roots. For example, Hylleberg et al. (1993) investigate to what extent seasonal unit roots in monthly series are preserved in quarterly time series. In this paper we consider the effects of temporally aggregating a so-called periodically integrated process.

Section 2 of the present paper is dedicated to a brief review of several concepts for periodically integrated autoregressions. Section 3 presents the main results.

### 2. Periodic integration

Consider a time series  $y_t$  which is observed  $S$  times a year during  $N$  years. The index  $t$  runs from 1 to  $n = SN$ . Assume that  $y_t$  can be described by a periodic autoregressive process of order  $p$  [PAR( $p$ )],

$$y_t = \mu_s + \phi_{1s}y_{t-1} + \dots + \phi_{ps}y_{t-p} + \varepsilon_t \quad (s = 1, 2, \dots, S), \quad (1)$$

---

\* Corresponding author.



were  $\varepsilon_t$  denotes a standard white noise process with variance  $\sigma^2$ . The index  $s$  indicates that the parameters are assumed to be time-varying, cf. Jones and Brelsford (1967) and Troutman (1979). Note that the mean of  $y_t$  may be  $\delta$ , i.e. nonperiodic, since (1) can originate from a PAR( $p$ ) process for  $y_t - \delta$ . The AR order  $p$  corresponds to the maximum order, i.e.  $p = \max(p_s)$ , where  $p_s$  is the AR order per season.

The process in (1) can be rewritten in vector notation by stacking the observations  $y_t$  in the  $(S \times 1)$  vector of annual observations per season, to be denoted as  $Y_T$  with  $T = 1, \dots, N$ , and representing (1) by

$$A_0 \begin{bmatrix} Y_{1,T} \\ Y_{2,T} \\ Y_{3,T} \\ \vdots \\ Y_{S,T} \end{bmatrix} = \mu + A_1 \begin{bmatrix} Y_{1,T-1} \\ Y_{2,T-1} \\ Y_{3,T-1} \\ \vdots \\ Y_{S,T-1} \end{bmatrix} + A_2 \begin{bmatrix} Y_{1,T-2} \\ Y_{2,T-2} \\ Y_{3,T-2} \\ \vdots \\ Y_{S,T-2} \end{bmatrix} + \dots + A_p \begin{bmatrix} Y_{1,T-p} \\ Y_{2,T-p} \\ Y_{3,T-p} \\ \vdots \\ Y_{S,T-p} \end{bmatrix} + \omega_T, \quad (2)$$

see e.g. Osborn (1991). The  $\mu$  and  $\omega_T$  are partitioned conformily with  $Y_T = (Y_{1,T}, Y_{2,T}, \dots, Y_{S,T})'$ . Model (2) corresponds with a VAR( $P$ ) process for  $Y_T$ , with  $P = 1 + [(p-1)/4]$ , where  $[\cdot]$  denoting the entire-function.  $\omega_T$  is a vector white noise process with covariance matrix  $\sigma^2 I_S$ . The matrices  $A_0, \dots, A_p$  are  $(S \times S)$  parameter matrices.

The vector process for  $Y_T$ , which can be described by (2), is stationary if the roots of the characteristic equation

$$|A_0 - A_1 z - \dots - A_p z^p| = 0 \quad (3)$$

are outside the unit circle. In that case we say that the univariate series  $y_t$  is periodically stationary. If a single solution to (3) equals unity, while the other solutions are outside the unit circle, there are  $S-1$  so-called cointegrating relationships between the  $S$  elements of  $Y_T$ , see Engle and Granger (1987) and Johansen (1991). Cointegration relations correspond to stationary linear combinations of the  $Y_{s,T}$  ( $s = 1, 2, \dots, S$ ) variables. A common representation of a cointegration model for (2) is given by rewriting it as

$$\Delta Y_T = \mu^* + \Gamma_1 \Delta Y_{T-1} + \dots + \Gamma_{p-1} \Delta Y_{T-(p-1)} + \Pi Y_{T-p} + \omega_T^*, \quad (4)$$

where  $\Delta$  is the first differencing filter for annual time series, where  $\mu^*$ ,  $\Gamma_1$  to  $\Gamma_{p-1}$  and  $\omega_T^*$  are functions of  $\mu$ ,  $A_0^{-1}$  and the  $A_1$  to  $A_{p-1}$  matrices, and where

$$\Pi = A_0^{-1} \left( \sum_{i=1}^p A_i - I_S \right). \quad (5)$$

When  $\Pi$  has reduced rank  $S-1$ , it can be decomposed as  $\alpha\beta'$ , where  $\alpha$  and  $\beta$  are  $S \times (S-1)$  matrices. The matrix  $\beta$  contains the cointegration parameters. It can be shown that in case of this reduced rank  $S-1$ , the  $S-1$  variables  $\beta' Y_T$  are stationary, i.e. do not contain a unit root. Hence, model (4) contains time series variables which are all stationary. The  $\alpha$  parameters can be interpreted as dynamic adjustment parameters. There are several methods to test for cointegration between the  $Y_{s,T}$  variables ( $s = 1, 2, \dots, S$ ). An elegant method is developed in Johansen (1991), which involves formal tests of the rank of  $\Pi$ . In Franses (1994) this method is applied to the case of periodic time series as (1).

When there are  $S-1$  cointegrating relations between the  $S$  variables, these relations are given by  $Y_{j,T} - \delta_j Y_{j-1,T}$  for  $j = 2, \dots, S$ . In case the  $\delta_j$  values are not equal across the seasons, the  $y_t$  process is called periodically integrated. Boswijk and Franses (1995) propose tests for periodic integration.

An illustrative example of the expressions in (1)–(5) is given by the periodic first-order autoregressive model for a quarterly observed series  $y_t$  with zero mean, or

$$y_t = \phi_s y_{t-1} + \varepsilon_t, \quad s = 1, 2, 3, 4, \quad (6)$$

where  $\phi_s$  is unequal to zero, with  $\phi_s \neq \phi$  for all  $s = 1, 2, 3, 4$ , and where some of the  $\phi_s$  can be larger than 1. Stacking  $y_t$  into  $Y_T = (Y_{1,T}, Y_{2,T}, Y_{3,T}, Y_{4,T})'$  ensures that (6) can be written as

$$A_0 Y_T = A_1 Y_{T-1} + \varepsilon_T, \quad (7)$$

with

$$A_0 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -\phi_2 & 1 & 0 & 0 \\ 0 & -\phi_3 & 1 & 0 \\ 0 & 0 & -\phi_4 & 1 \end{bmatrix}, \quad A_1 = \begin{bmatrix} 0 & 0 & 0 & \phi_1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

The characteristic equation for (7) reads

$$|A_0 - A_1 z| = (1 - (\phi_1 \phi_2 \phi_3 \phi_4) z) = 0. \quad (8)$$

The process  $Y_T$  is periodically stationary if  $|\phi_1 \phi_2 \phi_3 \phi_4| < 1$ , and periodically integrated if  $\phi_1 \phi_2 \phi_3 \phi_4 = 1$ . To investigate the presence of cointegration relations between the  $Y_{s,T}$  in (7), it is convenient to rewrite (7) as

$$\Delta_1 Y_T = \Pi Y_{T-1} + \omega_T, \quad (9)$$

where  $\omega_T = A_0^{-1} \varepsilon_T$ , with

$$\Pi = \begin{bmatrix} -1 & 0 & 0 & \phi_1 \\ 0 & -1 & 0 & \phi_1 \phi_2 \\ 0 & 0 & -1 & \phi_1 \phi_2 \phi_3 \\ 0 & 0 & 0 & \phi_1 \phi_2 \phi_3 \phi_4 - 1 \end{bmatrix}. \quad (10)$$

Under the restriction  $\phi_1 \phi_2 \phi_3 \phi_4 = 1$ , this matrix has rank 3. Hence, for this quarterly time series, there are three cointegrating relations between the elements of  $Y_T$ , and these are  $Y_{2,T} - \phi_2 Y_{1,T}$ ,  $Y_{3,T} - \phi_3 Y_{2,T}$  and  $Y_{4,T} - \phi_4 Y_{3,T}$ .

### 3. Temporal aggregation

Consider the temporal aggregation of a  $y_t$  series, which is observed  $S$  times per year and which can be described by a PAR process as in (1), to a series  $ya_t$ , that is observed  $S^*$  times, with  $kS^* = S$  and  $k$  usually is 6, 4, 3 or 2. In general, the  $ya_t$  series can be described by a periodic ARMA process [PARMA], see also Weiss (1984) for nonperiodic processes.

For example, consider the aggregation of the quarterly series  $y_t$ , which is generated from (6) into a bi-annual series  $z_t$ , where  $\tau$  runs from 1 through  $n/2$ . The observations  $z_t$  are generated as  $z_1 = y_1 + y_2$ ,  $z_2 = y_3 + y_4$ , and so on. Substituting the appropriate expressions for  $y_t$ ,  $i = 1, \dots, n$ , one easily obtains that



$z_t$  follows a periodic ARMA process of order (1, 1),

$$z_t = \alpha_s z_{t-1} + v_t + \beta_s v_{t-1} \quad (11)$$

with

$$\alpha_1 = ((\phi_2 + 1)\phi_1\phi_4)/(\phi_4 + 1), \quad (12)$$

$$\alpha_2 = ((\phi_4 + 1)\phi_2\phi_3)/(\phi_2 + 1), \quad (13)$$

where the  $v_t + \beta_s v_{t-1}$  is a moving average process of order 1 with parameter  $\beta_s$ , which also varies with the season. Note that this assumes that  $\phi_2$  and  $\phi_4$  are not equal to  $-1$ . If this is the case,  $z_t$  becomes a white noise process. From (12) and (13) it is clear that  $\alpha_1 = \alpha_2$  only when a specific nonlinear parameter restriction holds. This suggests that temporal aggregation of a PAR generically yields a PARMA process.

Aggregating the  $y_t$  series in (6) to an annually observed time series  $x_T$  yields

$$x_T = \phi_1\phi_2\phi_3\phi_4 x_{T-1} + v_T - \theta v_{T-1}, \quad (14)$$

which is a nonperiodic ARMA(1, 1) process.

**Proposition.** Let  $\{Y_T\}$  be an  $(S \times 1)$  cointegrated process with cointegrating rank  $S - 1$  and  $S \times (S - 1)$  matrix of cointegrating vectors  $\beta$ . Define  $\beta_\perp \neq 0$  to be an  $S$ -vector that spans the null space of  $\beta$ , i.e.  $\beta' \beta_\perp = 0$ . Let  $A$  be an arbitrary  $(S \times S^*)$  matrix,  $S^* < S$ , and let  $\{X_T = A' Y_T\}$  be an  $S^*$ -vector process. Then  $X_T$  is a stationary process iff  $M(A) \subset M(B)$ , where  $M(\cdot)$  denotes the column space, or equivalently iff  $A' \beta_\perp = 0$ ; otherwise,  $\{X_T\}$  is a cointegrated process with cointegrating rank  $S^* - 1$ .

**Proof.** Johansen's version of the Granger representation theorem (Theorem 4.1) states that

$$Y_T = \beta_\perp (\alpha'_\perp \Psi \beta_\perp)^{-1} \alpha'_\perp S_T + U_T, \quad (15)$$

where  $\alpha_\perp$  and  $\Psi$  are as defined in Johansen (1991), where  $S_T$  is an  $S$ -vector random walk, and where  $U_T$  is an  $(S \times 1)$  stationary process. Because  $\beta_\perp$  and  $\alpha_\perp$  are  $(S \times 1)$  vectors, there is one common trend in the system:  $\alpha'_\perp S_T$  is a scalar random walk. The same single common trend appears in the system

$$X_T = A' \beta_\perp (\alpha'_\perp \Psi \beta_\perp)^{-1} \alpha'_\perp S_T + A' U_T, \quad (16)$$

unless  $A' \beta_\perp = 0$ , so that the linear combinations  $A' Y_T$  annihilates the common trend and thus is stationary. Finally, Johansen's Theorem 4.1 implies that  $S^*$ -vector process with a single common trend is cointegrated with cointegrating rank  $S^* - 1$ .  $\square$

For the periodic AR process  $y_t$  in (1) with a single unit root in the multivariate system (2), it applies that  $\beta'$  is the  $(S - 1) \times S$  matrix

$$\beta' = \begin{bmatrix} -\delta_2 & 1 & 0 & \cdot & \cdot & \cdot & 0 \\ 0 & -\delta_2 & 1 & & & & 0 \\ 0 & 0 & -\delta_3 & 1 & \cdot & \cdot & 0 \\ \vdots & \vdots & 0 & & \ddots & & \vdots \\ \vdots & \vdots & \vdots & & & & \vdots \\ 0 & 0 & 0 & & & -\delta_S & 1 \end{bmatrix}, \quad (17)$$

and that  $\beta_\perp$  is given by

$$\beta'_\perp = [1, \delta_2, \delta_2 \delta_3, \dots, \delta_2 \delta_3 \dots \delta_{S-1}, \delta_2 \delta_3 \dots \delta_S]. \quad (18)$$

For temporal aggregation of the process  $Y_T$  in (2), the matrix  $A'$  in the proposition is given by the  $(S^* \times S)$  matrix

$$A' = \begin{bmatrix} 1 \dots 1 & 0 & 0 & \dots & 0 \\ 0 & 0 & 1 \dots 1 & 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \dots & 1 \dots 1 & \dots & \dots \end{bmatrix} \tag{19}$$

where the  $(1 \dots 1)$  parts are  $(1 \times k)$  vectors. The event that all  $S^*$  elements of  $A'\beta_{\perp}$  are equal to zero, where  $A$  and  $\beta_{\perp}$  are given by (18) and (19), has measure zero. Hence, if a periodically integrated process for an  $S$  times observed  $y_t$  series, i.e. a process with  $S - 1$  cointegration relations, is aggregated to an  $S^*$  times observed series  $ya_t$ , this  $ya_t$  series generically has  $S^* - 1$  cointegrating relations between its annually observed components.

An additional intuitive argument is that the only way to get rid of the unit root  $z = 1$  is to use some differencing filter. Temporal aggregation of  $y_t$  does not involve such a filter, and hence the unit root is not removed.

For the periodically integrated AR(1) process in (6) the cointegration vectors are  $(-\phi_2, 1, 0, 0)'$ ,  $(0, -\phi_3, 1, 0)'$  and  $(0, 0, -\phi_4, 1)'$ , and the  $(4 \times 1)$  vector  $\beta_{\perp}$  is  $(\phi_1, \phi_1\phi_2, \phi_1\phi_2\phi_3, 1)$ . Temporal aggregation to the  $z_{\tau}$  in (11) implies that

$$A' = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix} \tag{20}$$

and hence  $z_{\tau}$  is periodically integrated unless  $A'\beta_{\perp} = 0$ , or  $\phi_2 = \phi_4 = -1$ , see below (12) and (13). The process in (11) can be represented in vector notation as  $B_0 Z_T = B_1 Z_{T-1} + \psi_T$ , where  $\psi_T$  is a moving average type error process. The characteristic equation is  $|B_0 - B_1 z| = (1 - (\phi_1\phi_2\phi_3\phi_4)z) = 0$ , which is the same as that in (8). An expression for the cointegration vector can be found from

$$B_0^{-1} B_1 - I = \begin{bmatrix} -1 & ((\phi_2 + 1)\phi_1\phi_4)/(\phi_4 + 1) \\ 0 & \phi_1\phi_2\phi_3\phi_4 - 1 \end{bmatrix}. \tag{21}$$

The rank of the matrix in (21) is at least one, and the cointegration relation between  $Z_{1,T}$  and  $Z_{2,T}$  is easily found.

To obtain the annual series  $X_T$  in (14), one uses  $A' = [1, 1, 1, 1]$ . Hence, the  $X_T$  is an integrated series provided that  $\phi_1 + \phi_1\phi_2 + \phi_1\phi_2\phi_3 \neq -1$ .

In summary, in this paper it is shown that temporally aggregating a univariate periodically integrated autoregressive time series [PIAR], i.e. a process of the multivariate representation contains a single unit root, yields a PIAR again.

**Acknowledgements**

Both the authors thank the Royal Netherlands Academy of Arts and Sciences for its financial support and a referee for useful comments.

**References**

Amemiya, T. and R.Y. Wu (1972), The effect of aggregation on prediction in the autoregressive model, *J. Amer. Statist. Assoc.* **67**, 628-632.

- Boswijk, H.P. and P.H. Franses (1995), Unit roots in periodic autoregressions, *J. Time Series Anal.*, to appear.
- Engle, R.F. and C.W.J. Granger (1987), Cointegration and error correction: representation, estimation, and testing, *Econometrica* **55**, 251–276.
- Franses, P.H. (1994), A multivariate approach to modeling univariate seasonal time series, *J. Econometrics* **63** 133–151.
- Hylleberg, S., C. Jørgensen and N.K. Sørensen (1993), Aggregation and seasonal unit roots, unpublished manuscript.
- Johansen, S. (1991), Estimation and hypothesis testing of cointegration vectors in Gaussian vector autoregressive models, *Econometrica* **59**, 1551–1580.
- Jones, R.H. and W.M. Brelsford (1967), Time series with periodic structure, *Biometrika* **54**, 403–408.
- Osborn, D.R. (1991), The implications of periodically varying coefficients for seasonal time-series processes, *J. Econometrics* **28**, 373–384.
- Troutman, B.M. (1979), Some results in periodic autoregression, *Biometrika* **66**, 365–373.
- Weiss, A.A. (1986), Systematic sampling and temporal aggregation in time series models, *J. Econometrics* **26**, 271–281.