



UvA-DARE (Digital Academic Repository)

A minor-monotone graph parameter based on oriented matroids

Edmonds, J.; Laurent, M.; Schrijver, A.

Published in:
Discrete mathematics

DOI:
[10.1016/S0012-365X\(96\)00172-0](https://doi.org/10.1016/S0012-365X(96)00172-0)

[Link to publication](#)

Citation for published version (APA):
Edmonds, J., Laurent, M., & Schrijver, A. (1997). A minor-monotone graph parameter based on oriented matroids. *Discrete mathematics*, 165-166, 219-226. [https://doi.org/10.1016/S0012-365X\(96\)00172-0](https://doi.org/10.1016/S0012-365X(96)00172-0)

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.



A minor-monotone graph parameter based on oriented matroids

Jack Edmonds^a, Monique Laurent^{b,*}, Alexander Schrijver^{c,d}

^a *Department of Combinatorics and Optimization, University of Waterloo, Waterloo, Ont., Canada N2L 3G1*

^b *LIENS, Ecole Normale Supérieure, 45 rue d'Ulm, 75230 Paris Cedex 05, France*

^c *CWI, Kruislaan 413, 1098 SJ Amsterdam, The Netherlands*

^d *Department of Mathematics, University of Amsterdam, Plantage Muidergracht 24, 1018 TV Amsterdam, The Netherlands*

Abstract

For an undirected graph $G = (V, E)$ let $\lambda'(G)$ be the largest d for which there exists an oriented matroid M on V of corank d such that for each nonzero vector (x^+, x^-) of M , x^+ is nonempty and induces a connected subgraph of G .

We show that $\lambda'(G)$ is monotone under taking minors and clique sums. Moreover, we show that $\lambda'(G) \leq 3$ if and only if G has no K_5 - or V_8 -minor; that is, if and only if G arises from planar graphs by taking clique sums and subgraphs.

1. Introduction

In [5] the following invariant $\lambda(G)$ for a graph $G = (V, E)$ was introduced: $\lambda(G)$ is equal to the largest dimension of any linear subspace X of \mathbb{R}^V with the property that for any nonzero $x \in X$ the graph $\langle \text{supp}_+(x) \rangle$ induced by $\text{supp}_+(x)$ is nonempty and connected. (Here $\text{supp}_+(x)$ denotes the *positive support* of x ; that is, the set $\{v \in V \mid x(v) > 0\}$. Similarly, $\text{supp}_-(x)$ denotes the *negative support* of x ; that is, the set $\{v \in V \mid x(v) < 0\}$. Moreover, for any $U \subseteq V$, $\langle U \rangle$ denotes the subgraph of G induced by U ; that is, the subgraph with vertex set U and edges all edges of G contained in U . In this paper, all graphs are assumed to be simple.)

This graph parameter can be easily seen to be monotone under taking minors. That is, if G is a minor of H , then $\lambda(G) \leq \lambda(H)$. So for each natural number d the class of graphs G with $\lambda(G) \leq d$ is closed under taking minors.

*Corresponding author. CWI, P.O. Box 94079, 1090 GB Amsterdam, Netherlands. E-mail: monique@cwi.nl.

In [5] it is also shown that $\lambda(G) = \max\{\lambda(G_1), \lambda(G_2)\}$ if G is a *clique sum* of G_1 and G_2 (that is, arises by identifying two cliques of equal size in G_1 and G_2). It was shown that

- (i) $\lambda(G) \leq 1$ if and only if G is a forest;
- (ii) $\lambda(G) \leq 2$ if and only if G is series-parallel;
- (iii) $\lambda(G) \leq 3$ if and only if G arises by taking subgraphs and clique sums from planar graphs. (1)

The function $\lambda(G)$ was motivated by the graph invariant $\mu(G)$ introduced by Colin de Verdière [2] (cf. [3]), although we do not know a relation between the two numbers. (It might be that $\lambda(G) \leq \mu(G)$ holds for each graph G .)

In the discussion after presenting the results above at the *5ème Colloque International Graphes et Combinatoire* in Marseille Luminy (September 1995), the first author of the present paper raised the question of extending these results to oriented matroids. The present paper shows that indeed most results of [5] are maintained under such an extension.

We first give the definition of oriented matroid (see [1] for background). To this end it is convenient to introduce, for any ordered pair $x = (a, b)$, the notation $x^+ := a$ and $x^- := b$.

Let $M = (V, X)$ be an oriented matroid, where X is the set of ‘vectors’ of M . That is, X is a collection of ordered pairs $x = (x^+, x^-)$ of subsets of V satisfying

- (i) for each $x \in X$, $x^+ \cap x^- = \emptyset$;
- (ii) $\mathbf{0} := (\emptyset, \emptyset) \in X$;
- (iii) if $x \in X$ then $-x := (x^-, x^+) \in X$;
- (iv) if $x, y \in X$, then $x \cdot y := (x^+ \cup (y^+ \setminus x^+), x^- \cup (y^- \setminus x^-)) \in X$;
- (v) if $x, y \in X$ and $u \in x^+ \cap y^-$, then there exists a $z \in X$ such that $u \notin z^+ \cup z^-$, $(x^+ \setminus y^+) \cup (y^+ \setminus x^+) \subseteq z^+ \subseteq x^+ \cup y^+$, and $(x^- \setminus y^-) \cup (y^- \setminus x^-) \subseteq z^- \subseteq x^- \cup y^-$. (2)

The elements of X are called the *vectors* of the oriented matroid ($\mathbf{0}$ is the *zero*). Any linear subspace Y of \mathbb{R}^V gives an oriented matroid (V, X) , by taking $X := \{(\text{supp}_+(x), \text{supp}_-(x)) \mid x \in Y\}$.

For any oriented matroid $M = (V, X)$, the minimal nonempty subsets of $\{x^+ \cup x^- \mid x \in X\}$ form the circuit collection of a matroid, again denoted by M . Thus matroid terminology applies to oriented matroids. We give the concepts we need below, expressed in terms of the circuits of M .

The *rank* of a subset U of V is the size of a largest subset U' of U not containing a circuit of M . The *rank* $\text{rank}(M)$ of M is the rank of V .

A *cobase* is a base of the dual matroid M^* ; that is, it is an inclusionwise minimal subset intersecting each circuit of M . The *cospan* $\text{cospan}(U)$ of a subset U of V is the set of elements $v \in V$ such that there is no circuit containing v and not intersecting U (so $U \subseteq \text{cospan}(U)$). The *corank* $\text{corank}(U)$ of a subset U of V is the size of a minimal subset U' of U such that $U \subseteq \text{cospan}(U')$. A basic matroid theory

formula is

$$\text{corank}(U) = |U| + \text{rank}(V \setminus U) - \text{rank}(V). \tag{3}$$

The *corank* $\text{corank}(M)$ of M is equal to $\text{corank}(V)$, which is equal to $|V| - \text{rank}(V)$.

Finally, we denote the deletion and contraction of U by $M \setminus U$ and M/U , respectively. In terms of oriented matroids, if $M = (V, X)$ is an oriented matroid and $U \subseteq V$, then $M \setminus U$ is the oriented matroid $(V \setminus U, X')$ with $X' := \{x \in X \mid (x^+ \cup x^-) \cap U = \emptyset\}$, and M/U is the oriented matroid $(V \setminus U, X'')$ with $X'' := \{(x^+ \setminus U, x^- \setminus U) \mid x \in X\}$.

We next describe our graph parameter based on oriented matroids. Let $G = (V, E)$ be an undirected graph. A *valid representation* for G is any oriented matroid $M = (V, X)$ with the property that for each nonzero $x \in X$, the subgraph $\langle x^+ \rangle$ of G induced by x^+ is nonempty and connected. Let $\lambda'(G)$ be the largest corank of any valid representation for G .

As each subspace of \mathbb{R}^V gives an oriented matroid, with corank equal to the dimension of the subspace, we have for each graph G

$$\lambda(G) \leq \lambda'(G). \tag{4}$$

One of the consequences of this paper is that there are no graphs with $\lambda(G) \leq 3$ and $\lambda(G) < \lambda'(G)$. In fact, we do not know any graph G with strict inequality in (4).

2. λ' is minor-monotone

We now first show:

Theorem 1. *If G is a minor of H then $\lambda'(G) \leq \lambda'(H)$.*

Proof. Let $M = (V, X)$ be a valid representation of $G = (V, E)$ with $\text{corank}(M) = \lambda'(G)$. If G arises from H by deleting an edge of G , then M is also a valid representation for H . So $\lambda'(H) \geq \text{corank}(M) = \lambda'(G)$.

If G arises from H by contracting an edge $e = uv$ of H to vertex w of G , then replacing in any $x \in X$, x^+ by $(x^+ \setminus \{w\}) \cup \{u, v\}$ if $w \in x^+$, and similarly, x^- by $(x^- \setminus \{w\}) \cup \{u, v\}$ if $w \in x^-$, gives a valid representation M' for H , with $\text{corank}(M') = \text{corank}(M) = \lambda'(G)$. \square

This theorem implies, by Robertson and Seymour’s theorem [4], that for each fixed n there is a finite class \mathcal{F}_n of graphs with the property that for any graph G : $\lambda'(G) \geq n$ if and only if G has no minor in \mathcal{F}_n .

We note that for the complete graph K_n one has

Theorem 2. $\lambda'(K_n) = n - 1$.

Proof. Let $M = (V, X)$ be a valid representation for K_n . If $\text{corank}(M) = n$, then $\text{rank}(M) = 0$, and therefore $\{v\}$ is a circuit for each $v \in V$. So $\{v\}$ contains $x^+ \cup x^-$

for some nonzero $x \in X$. This contradicts the fact that both x^+ and x^- are non-empty.

On the other hand, the set X of all pairs (U, W) with $U = \emptyset = W$ or $U \neq \emptyset \neq W$ and $U \cap W = \emptyset$, gives a valid representation for K_n of corank $n - 1$. \square

So Hadwiger’s conjecture implies the conjecture that $\gamma(G) \leq \lambda'(G) + 1$ for each graph G , where $\gamma(G)$ is the chromatic number of G . (Hadwiger’s conjecture states that $\gamma(G) \leq n$ if G does not have any K_{n+1} -minor.)

It is useful to note:

Theorem 3. *If graph G' arises from graph G by deleting one vertex, then $\lambda'(G) \leq \lambda'(G') + 1$.*

Proof. Let $M = (V, X)$ be a valid representation for $G = (V, E)$, of corank $\lambda'(G)$. Let G' arise from G by deleting vertex v . Then the matroid $M' := M \setminus \{v\}$ obtained from M by deleting v is a valid representation for G' . Moreover $\text{rank}(M') \leq \text{rank}(M)$, and hence $\text{corank}(M') = |V| - 1 - \text{rank}(M') \geq |V| - 1 - \text{rank}(M) = \lambda'(G) - 1$. \square

3. Clique sums

In this section we show that the function $\lambda'(G)$ does not increase by taking clique sums, and from this we directly derive characterizations of the classes of graphs G satisfying $\lambda'(G) \leq 1$ and $\lambda'(G) \leq 2$. We first prove a lemma on oriented matroids.

Lemma 1. *Let $M = (V, X)$ be an oriented matroid and let $x, y \in X$ with $\emptyset \neq x^+ \subseteq y^-$ and $x \neq -y$. Then there is a nonzero $z \in X$ such that $z^+ \subseteq y^+$ and $x^+ \not\subseteq z^-$.*

Proof. Choose a nonzero $z \in X$ such that (i) $x^+ \not\subseteq z^-$, (ii) $z^+ \subseteq x^+ \cup y^+$, (iii) $x^- \setminus y^+ \subseteq z^- \subseteq x^- \cup y^-$, and (iv) $|y^+ \cup z^+|$ as small as possible. Such a z exists, since $z = x$ satisfies (i)–(iii).

Assume $z^+ \not\subseteq y^+$, and choose $u \in z^+ \setminus y^+$. So $u \in x^+$, and hence $u \in y^-$. Therefore, applying (2)(v) to y, z , there is a $z' \in X$ such that $u \notin z'^+ \cup z'^-$, $z'^+ \subseteq y^+ \cup z^+$, $z'^- \subseteq y^- \cup z^-$, $(y^+ \setminus z^-) \cup (z^+ \setminus y^-) \subseteq z'^+$, and $(z^- \setminus y^+) \cup (y^- \setminus z^+) \subseteq z'^-$. Then $x^+ \not\subseteq z'^-$ (as $u \notin z'^-$), $z'^+ \subseteq y^+ \cup z^+ \subseteq x^+ \cup y^+$, $x^- \setminus y^+ \subseteq z^- \setminus y^+ \subseteq z'^- \subseteq y^- \cup z^- \subseteq x^- \cup y^-$, and $y^+ \cup z'^+ \subset y^+ \cup z^+$ (as $u \notin y^+ \cup z'^+$). Since $|y^+ \cup z^+|$ is minimal it follows that $z' = \mathbf{0}$. Hence $y^+ \subseteq z^-$, $z^+ \subseteq y^-$, $z^- \subseteq y^+$, and $y^- \subseteq z^+$. So $z = -y$, and therefore $y^- \subseteq x^+$ and $y^+ \subseteq x^-$. Moreover, $x^- \setminus y^+ \subseteq y^+$, and hence $x^- \subseteq y^+$. So $x = -y$, contradicting our assumption. \square

The lemma is used to prove

Theorem 4. *Let $M = (V, X)$ be a valid representation for $G = (V, E)$ and let $y, z \in X$. If $y \neq -z$ then $\langle y^+ \cup z^+ \rangle$ is connected.*

Proof. Suppose $y \neq -z$ and $\langle y^+ \cup z^+ \rangle$ is disconnected. So y and z are nonzero and $y^+ \cap z^+ = \emptyset$. Consider $z \cdot y = (z^+ \cup (y^+ \setminus z^-), z^- \cup (y^- \setminus z^+))$. Since $\langle z^+ \cup (y^+ \setminus z^-) \rangle$ is connected, $y^+ \setminus z^- = \emptyset$, that is, $y^+ \subseteq z^-$. This implies by Lemma 1 that there is a nonzero $w \in X$ such that $w^+ \subseteq z^+$ and $y^+ \not\subseteq w^-$. Consider $w \cdot y = (w^+ \cup (y^+ \setminus w^-), w^- \cup (y^- \setminus w^+))$. Then w^+ is a nonempty subset of z^+ and $y^+ \setminus w^-$ is a nonempty subset of y^- , contradicting the fact that $\langle w^+ \cup (y^+ \setminus w^-) \rangle$ is connected. \square

This theorem does not apply if $y = -z$. This case can be described as follows.

Theorem 5. *Let $M = (V, X)$ be a valid representation for $G = (V, E)$. Then for all $y \in X$ with $\langle y^+ \cup y^- \rangle$ not connected, there exist corank(M) pairwise openly vertex-disjoint paths connecting y^+ and y^- , except if corank(M) = 1 and y^+ and y^- are contained in different components of G .*

Proof. Suppose not. Then by Menger’s theorem there exists a subset U of V such that y^+ and y^- are contained in different components of $G - U$ and such that $|U| < \text{corank}(M)$. By Theorem 4, $y^+ \cup y^-$ is the unique circuit of M contained in $V \setminus U$. (Indeed, by Theorem 4, for any nonzero $x \in X$ and $x^+ \cup x^- \subseteq V \setminus U$ one has $x \in \{y, -y\}$.) Therefore $\text{rank}(V \setminus U) = |V \setminus U| - 1$.

If $U = \emptyset$ then $\text{rank}(M) = |V| - 1$, and hence $\text{corank}(M) = 1$. If $U \neq \emptyset$, we can choose some $u \in U$. Let $x \in X$ be such that $x^+ \cup x^- \subseteq (V \setminus U) \cup \{u\}$. If $u \notin x^+$ then $x \in \{y, -y\}$ (by Theorem 4). Similarly, if $u \notin x^-$ then again $x \in \{y, -y\}$. Concluding, $y^+ \cup y^-$ is the unique circuit contained in $(V \setminus U) \cup \{u\}$ and hence $\text{rank}(V \setminus U) \cup \{u\} = |V \setminus U|$. Hence $\text{rank}(M) \geq \text{rank}((V \setminus U) \cup \{u\}) = |V \setminus U|$. This contradicts the fact that $\text{rank}(M) = |V| - \text{corank}(M) = |V| - d < |V \setminus U|$. \square

We use Theorems 4 and 5 to investigate the behaviour of $\lambda'(G)$ upon taking a ‘clique sum’, which is defined as follows. Let $G = (V, E)$ be a graph and let V_1 and V_2 be subsets of V such that $V = V_1 \cup V_2, K := V_1 \cap V_2$ is a clique in G and such that there is no edge connecting $V_1 \setminus K$ and $V_2 \setminus K$. Then G is called a *clique sum* of $G_1 := \langle V_1 \rangle$ and $G_2 := \langle V_2 \rangle$.

Theorem 6. *If G is a clique sum of G_1 and G_2 then $\lambda'(G) = \max\{\lambda'(G_1), \lambda'(G_2)\}$, except if $G = \bar{K}_2$.*

Proof. Since G_1 and G_2 are subgraphs of G , we have $\lambda'(G) \geq \max\{\lambda'(G_1), \lambda'(G_2)\}$. So it suffices to show that $\lambda'(G) = \lambda'(G_i)$ for $i = 1$ or 2 . Assume that $\lambda'(G) > \max\{\lambda'(G_1), \lambda'(G_2)\}$. Let $d := \lambda'(G), G = (V, E), G_1 = (V_1, E_1), G_2 = (V_2, E_2), K := V_1 \cap V_2$, and $t := |K|$. We may assume that we have chosen this counterexample so that t is as small as possible.

Then $\langle V_1 \setminus K \rangle$ has a component L such that each vertex in K is adjacent to at least one vertex in L . Otherwise G would be a repeated clique sum of subgraphs of G_1 and G_2 with common clique sum smaller than t . In that case $\lambda'(G) = \max\{\lambda'(G_1), \lambda'(G_2)\}$

would follow by the minimality of t . Concluding, G_1 has a K_{t+1} -minor, and therefore $\lambda'(G_1) \geq t$. Hence $\lambda'(G) > t$.

Let $M = (V, X)$ be a valid representation for G with $\text{corank}(M) = d$. There exists a nonzero $y \in X$ such that $y^+ \cup y^- \subseteq V \setminus K$ (otherwise $\text{rank}(M) \geq |V \setminus K|$, contradicting the fact that $d > t$).

By Theorem 5 both y^+ and y^- are contained in the same component of $G - K$. Hence we may assume that $y^+ \cup y^- \subseteq V_1 \setminus K$. Hence by Theorem 4 we have that there is no nonzero $x \in X$ with $x^+ \subseteq V_1 \setminus K$. So $M/(V_1 \setminus K)$ has corank equal to $\text{corank}(M)$. Moreover, for each nonzero $x \in X$, $x^+ \cap V_2$ induces a nonempty connected subgraph of G_2 . Hence $\lambda'(G_2) \geq \text{corank}(M) = \lambda'(G)$, contradicting our assumption that $\lambda'(G_2) < d$. \square

This theorem directly implies characterizations of those graphs G satisfying $\lambda'(G) \leq 1$ and $\lambda'(G) \leq 2$.

Corollary 6a. *For any graph G , $\lambda'(G) \leq 1$ if and only if G does not have a K_3 -minor; that is, if and only if G is a forest.*

Proof. If $\lambda'(G) \leq 1$ then G has no K_3 -minor, as $\lambda'(K_3) = 2$. Conversely, if G is a forest, then G arises by taking clique sums and subgraphs from the graph K_2 . As $\lambda'(K_2) = 1$, Theorem 6 gives the corollary. \square

Corollary 6b. *For any graph G , $\lambda'(G) \leq 2$ if and only if G does not have a K_4 -minor; that is, if and only if G is a series-parallel graph.*

Proof. If $\lambda'(G) \leq 2$ then G has no K_4 -minor, as $\lambda'(K_4) = 3$. Conversely, if G is a series-parallel graph, then G arises by taking clique sums and subgraphs from the graph K_3 . As $\lambda'(K_3) = 2$, Theorem 6 gives the corollary. \square

4. Graphs satisfying $\lambda'(G) \leq 3$

We characterize in this section the graphs G satisfying $\lambda'(G) \leq 3$. The main step consists in proving that $\lambda'(G) \leq 3$ if G is planar.

Theorem 7. *If G is planar then $\lambda'(G) \leq 3$.*

Proof. Suppose $G = (V, E)$ is a planar graph with $\lambda'(G) \geq 4$ and $|V|$ minimal. We assume that we have an embedding of G in the sphere. For each face f of G let V_f be the set of vertices incident with f . Note that G is 4-connected, since otherwise it would be a subgraph of clique sums of smaller planar graphs, and hence we would have $\lambda'(G) \leq 3$ by Theorem 6.

Let $M = (V, X)$ be a valid representation for G with $\text{corank}(M) \geq 4$. Then $\text{corank}(\{u\}) = 1$ for each $u \in V$; that is, u is contained in at least one circuit of M . Otherwise, we can delete u from G and M .

We may assume that, for each edge uv , $\text{corank}(\{u, v\}) = 2$; that is, there is a circuit containing u but not v . Otherwise, either for each $x \in X$ one has $u \in x^+ \Leftrightarrow v \in x^-$, in which case we can delete the edge $\{u, v\}$ from G , or for each $x \in X$ one has $u \in x^+ \Leftrightarrow v \in x^+$, in which case we can contract the edge $\{u, v\}$ in G and identify elements u and v in M .

Note that this implies that if f and f' are adjacent faces (that is, have an edge in common) and $\text{corank}(V_f) = 2 = \text{corank}(V_{f'})$, then $\text{cospan}(V_f) = \text{cospan}(V_{f'})$.

Fixing V we choose E maximal under the condition that $\text{corank}(\{u, v\}) = 2$ for each edge $\{u, v\}$. Then $\text{corank}(V_f) \in \{2, 3\}$ for each face f . Indeed, $\text{corank}(V_f) \geq 2$, as each edge e has $\text{corank}(e) \geq 2$. Moreover, if $\text{corank}(V_f) \geq 4$, V_f contains at least two nonadjacent vertices u, v with $\text{corank}(\{u, v\}) = 2$. This contradicts the maximality of E .

For $x \in X$ let \mathcal{F}_x be the set of faces f for which $V_f \cap x^+ \neq \emptyset$ and $V_f \cap x^- \neq \emptyset$. Then:

$$\begin{aligned} &\text{Let } f \text{ and } f' \text{ be two faces with } \text{corank}(V_f \cup V_{f'}) \geq 4. \\ &\text{Then there is an } x \in X \text{ with } f, f' \in \mathcal{F}_x. \end{aligned} \tag{5}$$

As $\text{corank}(V_f) \geq 2$, $\text{corank}(V_{f'}) \geq 2$, and $\text{corank}(V_f \cup V_{f'}) \geq 4$, there exist $u, v \in V_f$, $u', v' \in V_{f'}$ with $\text{corank}(\{u, v, u', v'\}) = 4$. Therefore, we can find $x \in X$ such that $u, u' \in x^+$ and $v, v' \in x^-$. So $f, f' \in \mathcal{F}_x$, proving (5).

For $x \in X$ let $W_x := \bigcup \{V_f \mid f \in \mathcal{F}_x\}$. We show:

$$\text{corank}(W_x) \leq 3 \text{ for all } x \in X. \tag{6}$$

Note that (6) implies an immediate contradiction with (5), as $\text{corank}(V) \geq 4$.

We show that (6) holds. It suffices to show the result for $x \in X$ such that $x^+ \cup x^- = V$. (Indeed, if there exists $u \notin x^+ \cup x^-$, let $y \in X$ with $u \in y^+$ and set $z := x \cdot y$. Then, $z^+ \supseteq x^+ \cup \{u\}$, $z^- \supseteq x^-$ and $W_z \supseteq W_x$. Hence validity of the result for z will imply validity for x .)

Let $x \in X$ with $x^+ \cup x^- = V$ be given. Observe that if f and f' are faces with $\text{corank}(V_f) = \text{corank}(V_{f'}) = 2$ and having a common edge, e say, then $\text{cospan}(V_f) = \text{cospan}(V_{f'})$, as it is equal to $\text{cospan}(e)$. Similarly, $\text{cospan}(V_f) \subseteq \text{cospan}(V_{f'})$ if $\text{corank}(V_f) = 2$, $\text{corank}(V_{f'}) = 3$ and f, f' share a common edge.

As both $\langle x^+ \rangle$ and $\langle x^- \rangle$ are connected, the cut $\delta(x^+)$ corresponds in the dual graph of G to a circuit C which traverses exactly two edges in each face $f \in \mathcal{F}_x$.

Suppose, to obtain a contradiction, that $\text{corank}(W_x) \geq 4$. Then there exist faces $f, f' \in \mathcal{F}_x$ with $\text{corank}(V_f) = \text{corank}(V_{f'}) = 3$ and such that $\text{cospan}(V_f) \neq \text{cospan}(V_{f'})$. They correspond to two nodes on C . Denote by f_1, \dots, f_t the faces between f and f' when traveling from f to f' along C (in a given direction). Then we may assume that $\text{corank}(V_{f_i}) = 2$ for all $i = 1, \dots, t$. For $i = 0, 1, \dots, t$, let $u_i v_i$ be the edge common to the faces f_i and f_{i+1} , setting $f_0 := f$ and $f_{t+1} := f'$. So each $u_i v_i$ belongs to $\delta(x^+)$ (as G is 4-connected). We may assume that $u_i \in x^+$ and $v_i \in x^-$ for each i .

Now choose $w \in V_f \setminus \text{cospan}(V_{f'})$ and $w' \in V_{f'} \setminus \text{cospan}(V_f)$. Then the set $\{u_0, v_0, w, w'\}$ has corank 4. Hence, there exists $y \in X$ such that $w, w' \in y^+$ and $u, v \notin y^+ \cup y^-$. Hence, the set $y^+ \cup y^-$ contains none of the vertices on the faces f_1, \dots, f_t (since $V_{f_i} \subseteq \text{cospan}(\{u_0, v_0\})$ for all $i = 1, \dots, t$). In particular, $u_i, v_i \notin y^+ \cup y^-$ for $i = 1, \dots, t$. By connectivity of $\langle y^+ \rangle$ there exists a path P from w to w' which is entirely contained in y^+ .

Consider the region $R := \bigcup_{i=0}^{t+1} f_i$ (where faces are assumed to be topologically closed). As P joins two nodes on the boundary of R , $R \cup P$ partitions the rest of the sphere into two regions R_1 and R_2 . We choose indices such that R_1 has the vertices u_0, \dots, u_t on its boundary, while R_2 has the vertices v_0, \dots, v_t on its boundary.

By the connectivity of $\langle y^- \rangle$, y^- is contained either in \bar{R}_1 or in \bar{R}_2 . Suppose first that y^- is contained in \bar{R}_1 . Consider the element $z := y \cdot x$ of X . Then, $z^- \supseteq \{v_0, \dots, v_t\} \cup y^-$, while $u_0, \dots, u_t \in z^+$. Then there is no path joining v_0 and y^- which is entirely contained in z^- , contradicting the connectivity of $\langle z^- \rangle$.

Suppose next that y^- is contained in \bar{R}_2 . Set $z := y \cdot (-x)$. Then we arrive similarly at a contradiction. \square

We can now characterize the graphs G satisfying $\lambda'(G) \leq 3$. It follows from Theorems 6 and 7 that $\lambda'(G) \leq 3$ if G can be obtained from planar graphs by taking clique sums and subgraphs. On the other hand, it follows from a result by Wagner [6] that the graphs that can be obtained from planar graphs by taking clique sums and subgraphs are precisely the graphs with no K_5 - or V_8 -minor. (V_8 is the graph with vertices v_1, \dots, v_8 , where v_i and v_j are adjacent if and only if $|i - j| \in \{1, 4, 7\}$.) It is shown in [5] that $\lambda(V_8) = 4$. Hence $\lambda'(V_8) \geq 4$. As deleting any vertex of V_8 gives a planar graph, Theorem 3 implies that $\lambda'(V_8) = 4$. Moreover, by Theorem 2 $\lambda'(K_5) = 4$. Therefore,

Theorem 8. *A graph G satisfies $\lambda'(G) \leq 3$ if and only if G has no K_5 - or V_8 -minor; that is, if and only if G can be obtained from planar graphs by taking clique sums and subgraphs.*

References

- [1] A. Björner, M. Las Vergnas, B. Sturmfels, N. White and G. Ziegler, *Oriented Matroids* (Cambridge University Press, Cambridge, 1993).
- [2] Y. Colin de Verdière, Sur un nouvel invariant des graphes et un critère de planarité, *J. Combin. Theory Ser. B* 50 (1990) 11–21.
- [3] Y. Colin de Verdière, On a new graph invariant and a criterion for planarity, in: N. Robertson and P. Seymour, eds., *Graph Structure Theory, Contemporary Mathematics* (American Mathematical Society, Providence, RI, 1993) 137–147.
- [4] N. Robertson and P.D. Seymour, Graph minors. XX. Wagner's conjecture, preprint, 1988.
- [5] H. van der Holst, M. Laurent and A. Schrijver, On a minor-monotone graph invariant, *J. Combin. Theory Ser. B* 65 (1995) 291–304.
- [6] K. Wagner, Über eine Eigenschaft der ebene Komplexe, *Mathematische Annalen* 114 (1937) 570–590.