UvA-DARE (Digital Academic Repository)

Superfluid Fermi gas in a 1D optical lattice.

Orso, G.; Shlyapnikov, G.V.

Publication date
2005

Published in
Physical Review Letters

[Link to publication](#)

Citation for published version (APA):

Orso, G., & Shlyapnikov, G. V. (2005). Superfluid Fermi gas in a 1D optical lattice. *Physical Review Letters*, 95, 260402.

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Superfluid Fermi Gas in a 1D Optical Lattice

G. Orso¹ and G. V. Shlyapnikov^{2,3}

¹*BEC-INFM and Dipartimento di Fisica, Università di Trento, I-38050 Povo, Italy*

²*Laboratoire Physique Theorique et Modeles Statistiques, Université Paris Sud, Bâtiment 100, 91405 Orsay Cedex, France*

³*Van der Waals-Zeeman Institute, University of Amsterdam, Valckenierstraat 65/67, 1018 XE Amsterdam, The Netherlands*

(Received 6 August 2005; published 19 December 2005)

We calculate the superfluid transition temperature for a two-component 3D Fermi gas in a 1D tight optical lattice and discuss a dimensional crossover from the 3D to quasi-2D regime. For the geometry of finite size discs in the 1D lattice, we find that even for a large number of atoms per disc the critical effective tunneling rate for a quantum transition to the Mott insulator state can be large compared to the loss rate caused by three-body recombination. This allows the observation of the Mott transition, in contrast to the case of Bose-condensed gases in the same geometry.

DOI: [10.1103/PhysRevLett.95.260402](https://doi.org/10.1103/PhysRevLett.95.260402)

PACS numbers: 03.75.Ss, 03.75.Kk, 03.75.Lm

The observation of a BCS superfluid transition remains a challenging goal in the studies of ultracold Fermi gases. It was recently suggested that gases confined to low dimensions are promising candidates for achieving superfluidity as the confinement enhances interaction effects [1]. Adding a tunable periodic potential allows one to combine the benefit of the reduced dimensionality with the advantage of working with large yet coherent samples. Importantly, in the presence of a 1D periodic potential plus a superimposed weak harmonic trapping, the superfluid transition is signaled by a marked change in the behavior of the center of mass oscillations of the atomic cloud *along* the lattice direction [2].

In this Letter, we obtain the BCS transition temperature T_c for a two-component 3D Fermi gas in a 1D optical lattice. We reveal how the presence of the lattice renormalizes the effective coupling constant, leading to a density-dependent coupling and introducing a crossover to the quasi-2D regime. These findings form a basis for studying superfluid transport for atomic fermions in 1D lattices. For the geometry of finite size discs in the 1D lattice, we also discuss the possibility of achieving the superfluid-Mott insulator quantum transition by tuning the lattice depth above a critical value [3]. In this peculiar phase, the gas is superfluid in each separate disc, but the coherence along the lattice direction is completely lost. We show that for Fermi superfluids the critical effective tunneling rate can be large compared to the loss rate of all inelastic processes and therefore the Mott transition can be achieved. This result is a direct consequence of the Fermi statistics and is in marked contrast with the case of Bose-Einstein condensates in the same geometry, where the Mott transition can be hardly observed [4] unless the number of atoms per disc is very small.

We consider a two-component atomic Fermi gas in the presence of a 1D optical potential [5]:

$$V_{\text{opt}} = sE_R \sin^2 q_B z, \quad (1)$$

where s is a dimensionless factor fixed by the intensity of the laser beam and $E_R = \hbar^2 q_B^2 / 2m$ is the recoil energy, with $\hbar q_B$ being the Bragg momentum and m the atom mass. The potential (1) has periodicity $d = \pi / q_B$ along the z axis. The weak attraction between atoms in different internal states is modeled by a s -wave pseudo-potential $U(\mathbf{r}) = g\delta(\mathbf{r})\partial_r(r\cdot)$ with coupling constant $g = 4\pi\hbar^2 a / m$, where $a < 0$ is the 3D scattering length.

We will discuss the situation where the laser intensity is sufficiently large ($s \gtrsim 5$) and the Fermi energy ϵ_F is small compared to the interband gap ϵ_g . We thus confine ourselves to the lowest Bloch band where the physics is governed by the ratio of the Fermi energy to the bandwidth $4t$, where t is the hopping rate between neighboring wells. For $\epsilon_F < 4t$ the Fermi surface is closed and the system retains a 3D behavior, whereas in the case of $\epsilon_F > 4t$ the Fermi surface is open and the system undergoes a dimensional crossover. Hence, one has two distinct regimes: an anisotropic 3D regime ($\epsilon_F \ll t$) and a quasi-2D regime ($\epsilon_g \gg \epsilon_F \gg t$). This is clearly different from the case of a 3D lattice [6] where the Fermi energy scales with the bandwidth and can therefore be much smaller than the corresponding value in free space for a given atom density.

The mean field transition temperature T_c^0 is the highest temperature at which the Gorkov equation for the gap parameter has a nontrivial solution [7]. This gives

$$\frac{1}{g_{\text{eff}}} = \text{P} \int \frac{d^3 q}{(2\pi)^3} \frac{1}{\xi_{\mathbf{q}}} \frac{1}{\exp(\xi_{\mathbf{q}} / T_c^0) + 1}, \quad (2)$$

where g_{eff} is an effective coupling constant. The symbol P stands for the principal value and $\xi_{\mathbf{q}} = \hbar^2 \mathbf{q}_{\perp}^2 / 2m + \epsilon_1(q_z) - \mu$, where $\mathbf{q}_{\perp}$ is the momentum in the direction perpendicular to the lattice, $\epsilon(q_z)$ is the band dispersion, and $\mu \simeq \epsilon_F$ is the chemical potential. A straightforward integration of Eq. (2) yields

$$T_c^0 = \frac{2\gamma}{\pi} \mu \exp\left(\frac{1}{g_{\text{eff}}} \frac{1}{\nu(\mu)} - F(\mu)\right), \quad (3)$$

with $\gamma = 1.781$ and $\nu(\mu) = \int \delta(\xi_{\mathbf{q}}) d\mathbf{q} / (2\pi)^3$ being the density of states per internal state at the Fermi level. The function F is defined as

$$F = - \frac{\int_{-q_B}^{q_B} dq_z \ln(1 - \epsilon(q_z)/\mu) \Theta(\mu - \epsilon(q_z))}{\int_{-q_B}^{q_B} dq_z \Theta(\mu - \epsilon(q_z))}, \quad (4)$$

where $\Theta(x)$ is the unit-step function.

The effective coupling constant is related to the scattering amplitude $f(E)$ for Cooper pairs by $g_{\text{eff}}^{-1} = (m/4\pi\hbar^2) \text{Re}[1/f(E = 2\mu)]$ [8]. This requires us to solve the two-body problem for finding the scattering amplitude in the presence of the 1D lattice. In this case the expression for $f(E)$ is given by

$$f(E) = a \int dZ \phi_E^*(Z, 0) \partial_r (r \Psi(Z, \mathbf{r}))_{r=0}, \quad (5)$$

where $\phi_E(Z, \mathbf{r}) = \phi_{1q_z}(z_1) \phi_{1-q_z}(z_2) e^{i\mathbf{q}_1 \cdot \mathbf{r}_\perp}$ is the incoming wave function for two atoms undergoing Cooper pairing. The center of mass and relative coordinates are $Z = (z_1 + z_2)/2$ and $\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2$, and $E = \hbar^2 \mathbf{q}_\perp^2 / m + 2\epsilon_1(q_z)$ is the total energy. The two-particle wave function $\Psi(Z, \mathbf{r})$ obeys the Schrödinger equation

$$\left(-\frac{\hbar^2}{m} \Delta - \frac{\hbar^2}{4m} \frac{\partial^2}{\partial Z^2} + V(Z, z) + g \delta(\mathbf{r}) \frac{\partial}{\partial r} r - E \right) \Psi = 0, \quad (6)$$

where $V(Z, z) = V_{\text{opt}}(z_1) + V_{\text{opt}}(z_2)$. Equation (6) can be written in integral form as

$$\Psi(Z, \mathbf{r}) = \phi_E(Z, \mathbf{r}) + g \int dZ' G_E \partial_{r'} (r' \Psi(Z', \mathbf{r}'))_{r'=0}, \quad (7)$$

where $G_E(\mathbf{r}, Z; \mathbf{0}, Z')$ is the Green function of Eq. (6) with $g = 0$. The behavior of the Green function at short distances r is governed by the Laplacian term in Eq. (6) yielding $G_E(\mathbf{r}, Z; \mathbf{0}, Z') = -\delta(Z - Z') m / 4\pi\hbar^2 r + K_E(Z, Z')$, where $K_E(Z, Z')$ is a regular function. Then, from Eq. (7) we immediately obtain an equation for the function $Y(Z) = \partial_r (r \Psi(Z, \mathbf{r}))_{r=0}$ appearing in Eq. (5):

$$Y(Z) = \phi_E(Z, 0) + g \int dZ' K_E(Z, Z') Y(Z'). \quad (8)$$

Writing the kernel of the integral Eq. (8) in the form $K_E(Z, Z') = [G_E(\mathbf{r}, Z; \mathbf{0}, Z') - G_{E=0}(\mathbf{r}, Z; \mathbf{0}, Z')]_{r=0} + K_{E=0}(Z, Z')$, we expand the Green function G_E in eigenstates of noninteracting atoms,

$$G_E(\mathbf{r}, Z; \mathbf{0}, Z') = \sum_{n_1, n_2} \int \frac{d^2 \mathbf{q}_\perp}{(2\pi)^2} \int_{-q_B}^{q_B} \frac{dq_{1z}}{2\pi} \frac{dq_{2z}}{2\pi} e^{i\mathbf{q}_1 \cdot \mathbf{r}} \frac{\phi_{n_1, q_{1z}}(Z) \phi_{n_2, q_{2z}}(Z) \phi_{n_1, q_{1z}}^*(Z') \phi_{n_2, q_{2z}}^*(Z')}{E + i0 - \epsilon_{n_1}(q_{1z}) - \epsilon_{n_2}(q_{2z}) - \hbar^2 \mathbf{q}_\perp^2 / m}, \quad (9)$$

and retain only the contribution of the lowest Bloch band. In the tight binding limit, these states can be written in terms of Wannier functions as $\phi_{1q_z}(z) \sim \sum_{\ell} e^{i\ell q_z d} w(z - \ell d)$, where $w(z) = (1/\pi^{1/4} \sigma^{1/2}) \exp(-z^2/2\sigma^2)$ is a variational Gaussian ansatz. By minimizing the energy of noninteracting lattice atoms with respect to σ , one finds $d/\sigma = \pi s^{1/4} \exp(-1/4\sqrt{s})$ [9].

We now insert the ansatz $Y(Z) = A \sum_{\ell} w^2(Z - \ell d)$ into Eq. (8) and take into account that the relation $\int dZ' K_{E=0}(Z, Z') Y(Z') dZ' = Y(Z) m / 4\pi\hbar^2 a$ gives a critical value of the scattering length $a = a_{\text{cr}}$ needed to form a two-body bound state in the lattice [10]. Then, using the dispersion relation $\epsilon_1(q_z) = 2t[1 - \cos(q_z d)]$ and obtaining the kernel $K_E(Z, Z')$ on the basis of Eq. (9), we find the coefficient A in the expression for $Y(Z)$. Equation (5) then leads to the scattering amplitude

$$f(E) = \frac{aC}{1 - a/a_{\text{cr}} + (a/\sqrt{2\pi}\sigma)\alpha(E/4t)}, \quad (10)$$

where $C = d/\sqrt{2\pi}\sigma$. The function α is defined as $\alpha(x) = i \arccos(1 - x)$ for $x < 2$ and $\alpha(x) = -\ln[x(1 + \sqrt{1 - 2/x})^2/2] + i\pi$ for $x \geq 2$.

Equation (10) is one of the key results of this Letter. It shows that the scattering amplitude undergoes a dimen-

sional crossover as a function of energy. In the anisotropic 3D regime ($E \ll 8t$), we have $f = aC/(1 - a/a_{\text{cr}} + iaC\sqrt{Em^*}/\hbar)$, where m^* is the effective mass at the bottom of the band. In the quasi-2D regime ($E \gg 8t$), the tunneling between wells is irrelevant and the two atoms are in the ground state of an effective harmonic potential of frequency $\omega_0 = \hbar/m\sigma^2$. The scattering amplitude of Eq. (10) should then reduce to $f = f_{2D}d$, where $f_{2D} = (a/\sqrt{2\pi}\sigma)/[1 + (a/\sqrt{2\pi}\sigma)(\ln[\lambda\hbar\omega_0/E] + i\pi)]$ and $\lambda = 0.915/\pi$ [11]. This provides us with the asymptotic behavior of the critical value of the scattering length

$$a_{\text{cr}} = -\sqrt{2\pi}\sigma \ln^{-1}(\lambda\hbar\omega_0/2t), \quad (11)$$

which agrees with numerics of Ref. [10] already for $s \gtrsim 5$.

We see that the 1D lattice affects $f(E)$ and the coupling constant g_{eff} in a *nontrivial* way. For $\epsilon_F < 4t$, the coupling is density independent and from Eq. (3) we get

$$T_c^0 = \frac{2\gamma}{\pi} e^{-F} \epsilon_F \exp\left[-\frac{\pi}{2q_{zF}C} \left(\frac{1}{|a|} - \frac{1}{|a_{\text{cr}}|}\right)\right], \quad (12)$$

where the function F is given by Eq. (4), and $q_{zF} = \arccos(1 - \epsilon_F/2t)/d$ is the Fermi wave vector along the z axis. Equation (12) is valid provided $T_c^0 \ll \epsilon_F$, which

implies $|a| < |a_{\text{cr}}|$. In the low density limit $\epsilon_F \ll 4t$, the temperature T_c^0 reduces to the mean field transition temperature [12] for a homogeneous gas of atoms with an anisotropic quadratic dispersion and a renormalized 3D inverse scattering length $a_{\text{eff}}^{-1} = C^{-1}(|a|^{-1} - |a_{\text{cr}}|^{-1})$. The presence of the lattice causes an effective shift of the resonance from $1/a = 0$ to $1/a = 1/a_{\text{cr}} < 0$, which in turn gives rise to a sharp increase in T_c^0 at a fixed value of the 3D scattering length.

For $\epsilon_F > 4t$, the coupling constant g_{eff} becomes *density dependent*. Equation (4) yields $F(\mu) = 2 \ln[2/(1 + \sqrt{1 - 4t/\mu})]$, and from Eq. (3) we find

$$T_c^0 = \frac{\gamma}{2\pi} \left(1 + \sqrt{1 - \frac{4t}{\epsilon_F}} \right) \sqrt{\epsilon_F 4t} \exp \left[\sqrt{\frac{\pi}{2}} \left(\frac{\sigma}{a} - \frac{\sigma}{a_{\text{cr}}} \right) \right]. \quad (13)$$

Note that the exponent on the right-hand side of Eq. (13) does not depend on the Fermi energy, the density of states being constant for $\epsilon_F > 4t$.

Finally, for $\epsilon_F \gg 4t$ and values of the scattering length $|a| > |a_{\text{cr}}|$, but still $|a| \ll \sigma$, the BCS temperature reduces to the value $T_c^0 = \gamma \sqrt{2\epsilon_F E_b} / \pi$ found for isolated discs, where E_b is the energy of the two-body bound state. In this limit one has $t \leq T_c^0$ and the system behaves as a stack of quasi-2D superfluids weakly coupled by Josephson junctions. We emphasize that the low-energy *collective* excitations of the isolated disc are Bogolubov-Anderson phonons, so each pancake is superfluid *below* the Kosterlitz-Thouless temperature $T_{\text{KT}} \ll \epsilon_F$. Despite the fact that there is no real condensation of pairs in two dimensions at finite temperatures [13], BCS theory provides a rather good estimate for T_{KT} , namely, $T_c/T_{\text{KT}} - 1 \sim T_c/\epsilon_F \ll 1$ [14]. The inclusion of a small interplanar coupling changes the nature of the superfluid transition from Kosterlitz-Thouless to the continuous (second order) type, leading to a critical temperature T_c slightly larger than T_{KT} . This upward shift has been calculated for layered superconductors treating the hopping of Cooper pairs along the lattice in the mean field approach [15].

We next proceed to evaluate Gorkov's correction to the transition temperature due to the polarization of the medium [12]. Following Ref. [16], we introduce the static Lindhard function

$$L(\mathbf{p}) = \int \frac{d\mathbf{q}}{(2\pi)^3} \frac{f(\xi_{\mathbf{q}}) - f(\xi_{\mathbf{q}+\mathbf{p}})}{\xi_{\mathbf{q}+\mathbf{p}} - \xi_{\mathbf{q}}}, \quad (14)$$

where $f(x) = \Theta(-x)$ is the Fermi distribution at $T = 0$. Since $L(\mathbf{p} = 0) = \nu(\mu)$, we write $L(\mathbf{p}) = \nu(\mu)B(\mathbf{p})$, where B is a dimensionless *positive* function sensitive to the geometry of the Fermi surface. The critical temperature is then given by $T_c = T_c^0 e^{-(B)_{\text{FS}}}$, where

$$\langle B \rangle_{\text{FS}} = \frac{\int B(\mathbf{q} + \mathbf{q}') \delta(\xi_{\mathbf{q}}) \delta(\xi_{\mathbf{q}'}) d\mathbf{q} d\mathbf{q}'}{\int \delta(\xi_{\mathbf{q}}) \delta(\xi_{\mathbf{q}'}) d\mathbf{q} d\mathbf{q}'}. \quad (15)$$

The integration in (15) is done numerically and the corresponding Gorkov correction T_c/T_c^0 is shown in

FIG. 1. Gorkov's correction versus $\epsilon_F/4t$. The limiting value $T_c/T_c^0 = e^{-1}$ at $\epsilon_F/4t \gg 1$ is shown by the dotted line.

Fig. 1. For $\epsilon_F \ll 4t$, the system has an anisotropic quadratic dispersion and we recover the result for the homogeneous case $T_c = T_c^0/(4e)^{1/3} = 0.45T_c^0$. In the limit $\epsilon_F \gg 4t$, the band dispersion $\epsilon(k_z)$ can be neglected in Eq. (14) and we find $T_c/T_c^0 = e^{-1}$, in agreement with Ref. [1]. The cusp at $\epsilon_F = 4t$ is expected as this is the point of the Van Hove singularity [17] where the derivative of the density of states, $\partial\nu/\partial\epsilon$, diverges.

So far we have discussed the BCS superfluid transition in a 1D optical lattice. In the second part of the Letter we assume that the superfluid gas is at zero temperature and it is confined in the x, y directions by a trapping potential. Then, as the tunneling rate between neighboring discs is tuned below a critical value t_c , the system undergoes the superfluid-Mott insulator quantum transition. For a large number of atoms per well ($N \gg 1$), the critical hopping rate can be evaluated within the hydrodynamic approach [18]. Neglecting the coupling with radial degrees of freedom and the particle loss due to inelastic processes, the proper dynamical variables are the particle number fluctuation N'_ℓ and the phase Φ_ℓ of the order parameter in each disc. The hydrodynamic equations are equivalent to the classical equations of motion of the 1D phase Hamiltonian

$$H_P = \sum_\ell (E_c/2) N_\ell'^2 - E_J \cos(\Phi_{\ell+1} - \Phi_\ell), \quad (16)$$

where $E_c = 2\mu/N$ and $E_J = t^2 N/\mu$ are the charging and the Josephson energies, respectively, and $\hbar N'_\ell$ and Φ_ℓ are considered as conjugated variables.

Quantization of the classical Hamiltonian (16) is achieved by replacing these variables with operators $\hbar \hat{N}'_\ell$ and $\hat{\Phi}$ satisfying the commutation relation $[\hbar \hat{N}'_\ell, \hat{\Phi}] = i\hbar$. The quantized Hamiltonian is known to exhibit a phase transition at the critical value $E_c = \eta E_J$, with $\eta \approx 0.81$ [19]. The superfluid phase occurs for $E_c < \eta E_J$ and is characterized by an algebraic decay of the phase correlation function $\langle \cos(\Phi_\ell - \Phi_k) \rangle$ at large distances $|\ell - k| \gg 1$. The decay becomes exponential for $E_c > \eta E_J$ where one enters the Mott regime, characterized by large phase fluctuations which suppress interwell tunneling. The ground state is an insulator with a fixed number of atoms

per disc and a finite gap in the excitation spectrum. By comparing the values of the charging and the Josephson energies, we find that for BCS superfluids one has

$$\frac{t_c}{\mu} = \frac{1}{N} \sqrt{\frac{2}{\eta}}. \quad (17)$$

This result differs from the corresponding value for Bose condensates in the same geometry, $t_c^b/\mu^b \sim 1/N^2$, where $\mu^b = n g_{2D}$ is the chemical potential and g_{2D} is the 2D coupling constant. This is because the Josephson energy in the Hamiltonian (16) for the bosonic case is $E_J^b = tN$.

Equation (17) has been derived under the assumption that the effective tunneling rate $\nu_c = t_c^2/\hbar\mu \approx \mu/N^2$ is large compared to the loss rate $\tilde{\nu}$. The most severe losses come from three-body recombination. For an array of Bose-condensed atomic gases in the same geometry the corresponding loss rate is always large compared to the critical tunneling rate [4], unless the number of atoms per disc is very small as in the experiment of Ref. [3]. For Fermi superfluids the situation is completely different because the inelastic processes are *strongly inhibited* by quantum statistics. In the quasi-2D geometry, in analogy with the 3D case [20], for the 3-body loss rate one can write $\tilde{\nu} = -\dot{n}/n = Ln^2(k_F R_e)^2$, where L is the quasi-2D recombination coefficient and the small factor $(k_F R_e)^2 \ll 1$ comes from Pauli blocking, with R_e being a characteristic radius of the interatomic potential. For the ratio of the loss to critical tunneling rate, which should be small for consistency, we then find

$$\tilde{\nu}/\nu_c \approx (\hbar Ln^2/\mu)(k_F R_e)^2 N^2. \quad (18)$$

Note that compared to the case of bosons, where the coefficient L has the same order of magnitude but the statistics-induced inhibition of 3-body losses is absent, the ratio $\tilde{\nu}/\nu_c$ has an extra small factor $(k_F R_e)^2 \sim n R_e^2$.

For a weakly interacting gas the quantity $(\hbar Ln^2/\mu)$ is small, and already a crude dimensional estimate $(\hbar Ln^2/\mu) \sim n R_e^2$ indicates wide possibilities to have a small ratio (18) for a large number N of atoms per pancake. For example, let us consider $N = 10^3$ fermionic potassium atoms ($R_e \approx 5$ nm) in each disc, with density $n = 10^9$ cm $^{-2}$ corresponding to a chemical potential $\mu = \epsilon_F = 380$ nK k_B . From Eq. (17) we find $t_c/\hbar \sim 70$ s $^{-1}$ corresponding to $s \sim 25$ for a lattice period $d = 400$ nm, and $\sigma \approx 60$ nm. This leads to $\nu_c \sim 0.1$ s $^{-1}$. Then, taking into account that $L \sim L_{3D}/\sigma^2$ and assuming the 3D recombination $L_{3D} \sim 10^{-28}$ cm 6 /s, from Eq. (18) we obtain $\tilde{\nu}/\nu_c \sim 0.1$. Hence, owing to quantum statistics, in Fermi superfluids one can easily have $\tilde{\nu} \ll \nu_c$ and achieve the Mott insulator transition. It is important to emphasize that, in the given example, the suppression of recombination processes by a factor of $(k_F R_e)^2 \sim 10^{-3}$ originating from the Pauli principle is crucial to keep the ratio (18) small even for $N \sim 10^3$.

In conclusion, we have found the superfluid transition temperature for a two-component Fermi gas in a 1D optical

lattice and revealed that the effective coupling constant depends in a nontrivial way on both the atom density and the parameters of the optical field. For an array of finite size discs with a large number of atoms per disc, we have shown that the critical effective tunneling rate for the Mott insulator quantum transition can be larger than the rate of particle losses. Thus, the Mott phase transition in a chain of weakly coupled 2D Fermi superfluids can be observed for ultracold gases.

We acknowledge fruitful discussions with M. A. Baranov, L. P. Pitaevskii, M. Randeria, S. Stringari, and W. Zwerger. We thank F. Dalfovo for a critical reading of the manuscript. One of the authors (G. O.) wishes to thank the Van der Waals-Zeeman Institute (Amsterdam) and the LPTMS (Orsay) for their kind hospitality. This work was supported by the Ministero dell'Istruzione, dell'Università e della Ricerca (MIUR), by the Ministère de la Recherche (Grant No. ACI Nanoscience 201), and by the Nederlandse Stichting voor Fundamenteel Onderzoek der Materie (FOM).

-
- [1] D. S. Petrov, M. A. Baranov, and G. V. Shlyapnikov, Phys. Rev. A **67**, 031601(R) (2003).
 - [2] G. Orso, L. P. Pitaevskii, and S. Stringari, Phys. Rev. Lett. **93**, 020404 (2004).
 - [3] M. Greiner *et al.*, Nature (London) **415**, 39 (2002).
 - [4] Z. Hadzibabic *et al.*, Phys. Rev. Lett. **93**, 180403 (2004).
 - [5] For a review of optical lattice, see P. S. Jessen and I. H. Deutsch, Adv. At. Mol. Opt. Phys. **37**, 95 (1996); G. Grynberg and C. Robilliard, Phys. Rep. **355**, 335 (2001).
 - [6] W. Hofstetter *et al.*, Phys. Rev. Lett. **89**, 220407 (2002).
 - [7] P. G. de Gennes, *Superconductivity of Metals and Alloys* (Addison-Wesley, Reading, MA, 1994).
 - [8] A. A. Abrikosov, L. P. Gorkov, and I. E. Dzyaloshinskij, *Quantum Field Theoretical Methods in Statistical Physics* (Pergamon Press, New York, 1965).
 - [9] L. P. Pitaevskii and S. Stringari, *Bose-Einstein Condensation* (Clarendon Press, Oxford, 2003).
 - [10] G. Orso *et al.*, Phys. Rev. Lett. **95**, 060402 (2005).
 - [11] D. S. Petrov, M. Holzmann, and G. V. Shlyapnikov, Phys. Rev. Lett. **84**, 2551 (2000); D. S. Petrov and G. V. Shlyapnikov, Phys. Rev. A **64**, 012706 (2001).
 - [12] L. P. Gorkov and T. K. Melik-Barkhudarov, Sov. Phys. JETP **13**, 1018 (1961).
 - [13] N. D. Mermin and H. Wagner, Phys. Rev. Lett. **17**, 1133 (1966).
 - [14] K. Miyake, Prog. Theor. Phys. **69**, 1794 (1983).
 - [15] See C. Deutsch and S. Doniach, Phys. Rev. B **29**, 2724 (1984), and references therein.
 - [16] H. Heiselberg *et al.*, Phys. Rev. Lett. **85**, 2418 (2000).
 - [17] N. W. Ashcroft and N. D. Mermin, *Solid State Physics* (Rinehart and Winston, New York, 1976).
 - [18] M. Wouters, J. Tempere, and J. T. Devreese, Phys. Rev. A **70**, 013616 (2004); see also L. P. Pitaevskii, S. Stringari, and G. Orso, Phys. Rev. A **71**, 053602 (2005).
 - [19] R. M. Bradley and S. Doniach, Phys. Rev. B **30**, 1138 (1984).
 - [20] D. S. Petrov, Phys. Rev. A **67**, 010703(R) (2003).