UvA-DARE (Digital Academic Repository)

Virtual Compton scattering off the nucleon at low energies

Scherer, S.; Korchin, A.Y.; Koch, J.H.

DOI

[10.1103/PhysRevC.54.904](https://doi.org/10.1103/PhysRevC.54.904)

Publication date

1996

Published in

Physical Review C

[Link to publication](#)

Citation for published version (APA):

Scherer, S., Korchin, A. Y., & Koch, J. H. (1996). Virtual Compton scattering off the nucleon at low energies. *Physical Review C*, 54, 904. <https://doi.org/10.1103/PhysRevC.54.904>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Virtual Compton scattering off the nucleon at low energies

S. Scherer

Institut für Kernphysik, Johannes Gutenberg-Universität, D-55099 Mainz, Germany

A. Yu. Korchin

National Scientific Center Kharkov Institute of Physics and Technology, 310108 Kharkov, Ukraine

J. H. Koch

National Institute for Nuclear Physics and High-Energy Physics, 1009 DB Amsterdam, The Netherlands

(Received 19 March 1996)

We investigate the low-energy behavior of the four-point Green's function $\Gamma^{\mu\nu}$ describing virtual Compton scattering off the nucleon. Using Lorentz invariance, gauge invariance, and crossing symmetry, we derive the leading terms of an expansion of the operator in the four-momenta q and q' of the initial and final photon, respectively. The model-independent result is expressed in terms of the electromagnetic form factors of the free nucleon, i.e., on-shell information which one obtains from electron-nucleon scattering experiments. Model-dependent terms appear in the operator at $O(q_\alpha q'_\beta)$, whereas the orders $O(q_\alpha q_\beta)$ and $O(q'_\alpha q'_\beta)$ are contained in the low-energy theorem for $\Gamma^{\mu\nu}$, i.e., no new parameters appear. We discuss the leading terms of the matrix element and comment on the use of on-shell equivalent electromagnetic vertices in the calculation of "Born terms" for virtual Compton scattering. [S0556-2813(96)02108-5]

PACS number(s): 13.40.Gp, 13.60.Fz, 14.20.Dh

I. INTRODUCTION

Low-energy theorems (LET's) play an important role in studies of properties of particles. Based on a few general principles, they determine the leading terms of the low-energy amplitude for a given reaction in terms of global, model-independent properties of the particles. Clearly, this provides a constraint for models or theories of hadron structure: unless they violate these general principles they must reproduce the predictions of the low-energy theorem. On the other hand, the low-energy theorems also provide useful constraints for experiments. Experimental studies designed to investigate particle properties beyond the global quantities and to distinguish between different models must be carried out with sufficient accuracy at low energies to be sensitive to the higher-order terms not predicted by the theorems. Another option is, of course, to go to an energy regime where the low-energy theorems do not apply anymore and model-dependent terms in the theoretical predictions are important.

The best-known low-energy theorem for electromagnetic interactions is the theorem for "Compton scattering" (CS) of real photons off a nucleon [1–3]. Based on the requirement of gauge invariance, Lorentz invariance, and crossing symmetry, it specifies the terms in the low-energy scattering amplitude up to and including terms linear in the photon momentum. The coefficients of this expansion are expressed in terms of global properties of the nucleon: its mass, charge, and magnetic moment. In experiments, one can make the photon momentum, the kinematical variable in which one expands, small to ensure the convergence of the expansion and to allow for a direct comparison with the data. By increasing the energy of the photon one will become sensitive to terms that depend on details of the structure of the nucleon beyond the global properties. Terms of second order in the frequency, which are not determined by this theorem, can be

parametrized in terms of two new structure constants, the electric and magnetic polarizabilities of the nucleon (see, for example, Ref. [4]).

As in all studies with electromagnetic probes, the possibilities to investigate the structure of the target are much greater if virtual photons are used. A virtual photon allows one to vary the three-momentum and energy transfer to the target independently. Therefore it has recently been proposed to also use "virtual Compton scattering" (VCS) as a means to study the structure of the nucleon [5–7]. The proposed reaction is $p(e, e'p)\gamma$, i.e., in addition to the scattered electron also the recoiling proton is detected to completely determine the kinematics of the final state consisting of a real photon and a proton. It is the purpose of this work to extend the standard low-energy theorem for Compton scattering of real photons to the general case where one or both photons are virtual. The latter would be the case, e.g., in the reaction $e^- + p \rightarrow e^- + (e^- + e^+) + p$. We will refer to both possibilities as "VCS."

There are several different approaches to derive the LET for Compton scattering of real photons. One was first used by Low [1]. It made use of the fact that in terms of "unitarity diagrams" the scattering amplitude is dominated by the single-nucleon intermediate state. In such unitarity diagrams, not to be confused with Feynman diagrams, all intermediate states are on their mass shell [8]. Lorentz invariance and gauge invariance then allow a prediction for the amplitude to first order in the frequency. Another approach [2,3], first used by Gell-Mann and Goldberger [2], relies on a completely covariant description in terms of the basic building blocks of electromagnetic vertices and nucleon propagators. They split the amplitude into two classes, A and B: class A consists of one-particle-reducible contributions that can be built up from dressed photon-nucleon vertices and dressed nucleon propagators. Class B contains all one-particle-

irreducible two-photon diagrams, where the second photon couples into the dressed vertex of the first one. Application of two Ward-Takahashi identities, one relating the photon-nucleon vertex operator to the nucleon propagator, the other relating the irreducible two-photon vertex to the dressed one-photon vertex, then lead to the same result for the leading powers of the low-energy amplitude. One can use yet another technique introduced by Low [9] to describe bremsstrahlung processes. This technique relies on the observation that poles in the photon momentum can only be due to photon emission from external nucleon lines of the scattering amplitude. Low's method has, in a modified form, also been widely used in the framework of partially conserved axial-vector currents (PCAC) [10].

So far, there have been only a few investigations of the general VCS matrix element, since most calculations were restricted to the Compton scattering of real photons. In [11] electron-proton bremsstrahlung was calculated in first-order Born approximation. The photon scattering amplitude, for one photon virtual and the other one real, was analyzed in terms of 12 invariant functions of three scalar kinematical variables. In [12] it was shown that the general VCS matrix element—virtual photon to virtual photon—requires 18 invariant amplitudes depending on four scalar variables. In [13] the reaction $\gamma p \rightarrow p + e^+ e^-$ was investigated. Sizable effects on the dilepton spectrum from the timelike electromagnetic form factors of the proton were found. Very recently the low-energy behavior of the VCS matrix element was investigated [14]. Using Low's approach [9] the leading terms in the outgoing photon momentum were derived. It was shown that the virtual Compton scattering amplitude at low energies involves 10 “generalized polarizabilities” that depend on the absolute value of the three-momentum of the virtual photon. These new polarizabilities were estimated in a nonrelativistic quark model. In [15] the VCS amplitude was calculated in the framework of a phenomenological Lagrangian including baryon resonances in the s and u channels as well as π^0 and σ exchanges in the t channel. A prediction for the $|\vec{q}|$ dependence of the electromagnetic polarizabilities α and β was made.

The purpose of this work is to identify, in an analogous form to the real CS case, those terms of VCS which are determined on the basis of only gauge invariance, Lorentz invariance, crossing symmetry, and the discrete symmetries. In the following, we will refer to such terms not fixed by this LET for simplicity as “model dependent.” By introducing additional constraints, such as chiral symmetry, a statement about these terms also becomes possible. This is, however, beyond the scope of the present work. In our study of low-energy virtual Compton scattering, below the onset of pion production, we will mainly work on the operator level. This allows us to work without specifying a particular Lorentz frame or a gauge. We combine the method of Gell-Mann and Goldberger [2] with an effective Lagrangian approach. Class A is obtained in the framework of a general effective Lagrangian describing the interaction of a single nucleon with the electromagnetic field [4]. In the specific representation we choose, this turns out to be a simple covariant and gauge invariant “modified Born term” expression, involving on-shell Dirac and Pauli nucleon form factors F_1 and F_2 . The Ward-Takahashi identities then allow us to determine the

leading-order term of the unknown class B contribution in an expansion in both the initial and final photon momenta. Furthermore, with the help of crossing symmetry a definite prediction can be made concerning the order at which one expects model-dependent terms.

Our work is organized as follows. We start out in Sec. II by outlining the general structure of the VCS Green's function in the framework of Gell-Mann and Goldberger. We state the ingredients for the derivation of the LET, namely, crossing symmetry and gauge invariance. Section III derives the LET for virtual photon Compton scattering and we discuss the leading terms of the matrix element for the reaction $e^- + p \rightarrow e^- + p + \gamma$ in the center-of-mass frame. As the notion of “Born terms” is important for the LET, we comment on this aspect in Sec. IV and point out ambiguities that arise in their definition, both for real and virtual photons. Our results are summarized and put into perspective in Sec. V.

II. STRUCTURE OF THE VIRTUAL COMPTON SCATTERING TENSOR AND GAUGE INVARIANCE

In this section we will define the Green's functions and the kinematical variables relevant for the discussion of VCS off the proton. We will consider the constraints imposed by the fundamental requirements of gauge invariance, Lorentz invariance, and crossing symmetry. We do this in the framework of a manifestly covariant description, incorporating gauge invariance in its strong version, namely, in the form of the Ward-Takahashi identities [16,17]. The approach is similar to that of [3] using, however, a somewhat more modern formulation.

The electromagnetic three-point and four-point Green's functions are defined as

$$G_{\alpha\beta}^{\mu}(x, y, z) = \langle 0 | T [\Psi_{\alpha}(x) \bar{\Psi}_{\beta}(y) J^{\mu}(z)] | 0 \rangle, \quad (2.1)$$

$$G_{\alpha\beta}^{\mu\nu}(w, x, y, z) = \langle 0 | T [\Psi_{\alpha}(w) \bar{\Psi}_{\beta}(x) J^{\mu}(y) J^{\nu}(z)] | 0 \rangle, \quad (2.2)$$

where J^{μ} is the electromagnetic current operator in units of the elementary charge, $e > 0$, $e^2/4\pi = 1/137$, and where Ψ denotes a renormalized interpolating field of the proton; T denotes the covariant time-ordered product [18]. Electromagnetic current conservation, $\partial_{\mu} J^{\mu} = 0$, and the equal-time commutation relation of the charge density operator with the proton field,

$$[J^0(x), \Psi(y)] \delta(x^0 - y^0) = -\delta^4(x - y) \Psi(y), \quad (2.3)$$

are the basic ingredients for deriving Ward-Takahashi identities [16,17].

Using translation invariance, the momentum-space Green's functions corresponding to Eqs. (2.1) and (2.2) are defined through a Fourier transformation,

$$(2\pi)^4 \delta^4(p_f - p_i - q) G_{\alpha\beta}^{\mu}(p_f, p_i) = \int d^4x d^4y d^4z e^{i(p_f \cdot x - p_i \cdot y - q \cdot z)} G_{\alpha\beta}^{\mu}(x, y, z), \quad (2.4)$$

$$\begin{aligned}
(2\pi)^4 \delta^4(p_f + q' - p_i - q) G_{\alpha\beta}^{\mu\nu}(P, q, q') \\
= \int d^4w d^4x d^4y d^4z e^{i(p_f \cdot w - p_i \cdot x - q \cdot y + q' \cdot z)} G_{\alpha\beta}^{\mu\nu}(w, x, y, z),
\end{aligned} \quad (2.5)$$

where p_i and p_f refer to the four-momenta of the initial and final proton lines, respectively, $P = p_i + p_f$, and where q and $-q'$ denote the momentum transferred by the currents J^μ and J^ν , respectively. We note that $G_{\alpha\beta}^{\mu\nu}$ depends on two independent four-momenta, e.g., p_i and p_f . In particular, it is not assumed that these momenta obey the mass-shell condition $p_i^2 = p_f^2 = M^2$. Similarly, $G_{\alpha\beta}^{\mu\nu}$ depends on three four-momenta which are completely independent as long as one considers the general off-mass-shell case. This will prove to be an important ingredient below when analyzing the general structure of the VCS tensor.

Finally, the truncated three-point and four-point Green's functions relevant for our discussion of VCS are obtained by multiplying the external proton lines by the inverse of the corresponding full (renormalized) propagators,

$$\Gamma^\mu(p_f, p_i) = [iS(p_f)]^{-1} G^\mu(p_f, p_i) [iS(p_i)]^{-1}, \quad (2.6)$$

$$\Gamma^{\mu\nu}(P, q, q') = [iS(p_f)]^{-1} G^{\mu\nu}(P, q, q') [iS(p_i)]^{-1}, \quad (2.7)$$

where, for convenience, from now on we omit spinor indices. Using the definitions above, it is straightforward to obtain the Ward-Takahashi identities

$$q_\mu \Gamma^\mu(p_f, p_i) = S^{-1}(p_f) - S^{-1}(p_i), \quad (2.8)$$

$$\begin{aligned}
q_\mu \Gamma^{\mu\nu}(P, q, q') &= i(S^{-1}(p_f)S(p_f - q)\Gamma^\nu(p_f - q, p_i) \\
&\quad - \Gamma^\nu(p_f, p_i + q)S(p_i + q)S^{-1}(p_i)).
\end{aligned} \quad (2.9)$$

Following Gell-Mann and Goldberger [2], we divide the contributions to $\Gamma^{\mu\nu}$ into two classes, A and B , $\Gamma^{\mu\nu} = \Gamma_A^{\mu\nu} + \Gamma_B^{\mu\nu}$, where class A consists of the s - and u -channel pole terms; class B contains all the other contributions. We emphasize that this procedure does not restrict the generality of the approach. The separation into the two classes is such that all terms which are irregular for $q^\mu \rightarrow 0$ (or $q'^\mu \rightarrow 0$) are contained in class A , whereas class B is regular in this limit. Strictly speaking, one also assumes that there are no massless particles in the theory which could make a low-energy expansion in terms of kinematical variables impossible [1]; furthermore, the contribution due to t -channel exchanges, such as a π^0 , has not been considered.

The contribution from class A , expressed in terms of the full renormalized propagator and the irreducible electromagnetic vertices, reads

$$\begin{aligned}
\Gamma_A^{\mu\nu} &= \Gamma^\nu(p_f, p_f + q') iS(p_i + q) \Gamma^\mu(p_i + q, p_i) \\
&\quad + \Gamma^\mu(p_f, p_f - q) iS(p_i - q') \Gamma^\nu(p_i - q', p_i).
\end{aligned} \quad (2.10)$$

Note that $\Gamma_A^{\mu\nu}$ is symmetric under crossing, $q \leftrightarrow -q'$ and $\mu \leftrightarrow \nu$, i.e.,

$$\Gamma_A^{\mu\nu}(P, q, q') = \Gamma_A^{\nu\mu}(P, -q', -q). \quad (2.11)$$

Since also the total $\Gamma^{\mu\nu}$ is crossing symmetric, this must also be true for the contribution of class B separately [2]. Using the Ward-Takahashi identity, Eq. (2.8), one obtains the following constraint for class A as imposed by gauge invariance:

$$\begin{aligned}
q_\mu \Gamma_A^{\mu\nu}(P, q, q') &= i(\Gamma^\nu(p_f, p_f + q') - \Gamma^\nu(p_i - q', p_i) \\
&\quad + S^{-1}(p_f)S(p_i - q')\Gamma^\nu(p_i - q', p_i) \\
&\quad - \Gamma^\nu(p_f, p_f + q')S(p_i + q)S^{-1}(p_i)) \\
&\equiv f_A^\nu(P, q, q').
\end{aligned} \quad (2.12)$$

Similarly, contraction of $\Gamma_A^{\mu\nu}$ with q'_ν results in

$$\begin{aligned}
q'_\nu \Gamma_A^{\mu\nu}(P, q, q') &= -i(\Gamma^\mu(p_f, p_f - q) - \Gamma^\mu(p_i + q, p_i) \\
&\quad + S^{-1}(p_f)S(p_i + q)\Gamma^\mu(p_i + q, p_i) \\
&\quad - \Gamma^\mu(p_f, p_f - q)S(p_i - q')S^{-1}(p_i)) \\
&= -f_A^\mu(P, -q', -q),
\end{aligned} \quad (2.13)$$

which is, of course, the same constraint which one obtains from Eq. (2.12) using the crossing-symmetry property of $\Gamma_A^{\mu\nu}$:

$$\begin{aligned}
q'_\nu \Gamma_A^{\mu\nu}(P, q, q') &= -(-q'_\nu) \Gamma_A^{\nu\mu}(P, -q', -q) \\
&= -f_A^\mu(P, -q', -q).
\end{aligned} \quad (2.14)$$

Combining Eqs. (2.9) and (2.12) generates the following constraint for the contribution of class B :

$$\begin{aligned}
q_\mu \Gamma_B^{\mu\nu} &= q_\mu (\Gamma^{\mu\nu} - \Gamma_A^{\mu\nu}) \\
&= i[\Gamma^\nu(p_i - q', p_i) - \Gamma^\nu(p_f, p_f + q')],
\end{aligned} \quad (2.15)$$

relating it to the one-photon vertex [3]. Once again, the second gauge-invariance condition, obtained by contracting with q'_ν , is automatically satisfied due to crossing symmetry.

The 4×4 matrix $\Gamma_B^{\mu\nu}$ of class B is a function of the three independent four-momenta P , q , and q' . It is important to realize that the 12 components of these momenta are independent variables only for the complete off-shell case, i.e., if one allows for arbitrary values of p_i^2 and p_f^2 . This will be important when making use of the constraints imposed by gauge invariance. Using Lorentz invariance, gauge invariance, crossing symmetry, parity and time-reversal invariance, it was shown in [12] that the general $\Gamma^{\mu\nu}$ for VCS off a free nucleon with both photons virtual consists of 18 independent operator structures. The functions associated with each operator depend on four Lorentz scalars, e.g., $q^2, q'^2, \nu = P \cdot q = P \cdot q'$, and $t = (p_i - p_f)^2$. When allowing the external nucleon lines to be off their mass shell, one will have an even more complicated structure [19].

However, in our derivation of the low-energy behavior of the electromagnetic four-point Green's function we will not require the full structure as discussed in [12]. At low energies we expand $\Gamma_B^{\mu\nu}$ in terms of the four-momenta q^μ and q'^μ ,

$$\Gamma_B^{\mu\nu} = a^{\mu\nu}(P) + b^{\mu\nu,\rho}(P)q_\rho + c^{\mu\nu,\sigma}(P)q'_\sigma + \dots, \quad (2.16)$$

where the coefficients are 4×4 matrices and can be expressed in terms of the 16 independent Dirac matrices $1, \gamma_5, \gamma^\mu, \gamma^\mu \gamma_5, \sigma^{\mu\nu}$. An expansion of the type of Eq. (2.16) is expected to work below the lowest relevant particle-production threshold, in this case the pion-production threshold; we refer the reader to, e.g., Ref. [20], where a similar discussion for the case of pion photoproduction can be found.

So far we have considered general features of the operators entering into the description of VCS. It is clearly one of the advantages of using a covariant description of the type of Eq. (2.16) that it neither uses a particular Lorentz system nor a specific gauge. When referring to powers of q or q' , we mean the ones coming from the Dirac structures and their associated functions. This is different when one works on the level of nucleon matrix elements or the invariant amplitude, where also kinematical variables from the spinors or normalization factors enter into the power counting.

We conclude this section by noting that the above framework can easily be applied to VCS off a spin-0 particle, such as the pion. In that case $\Gamma^{\mu\nu}$, of course, has no complicated spinor structure. The building blocks for class A are simply the corresponding irreducible, renormalized electromagnetic vertex for a spinless particle and the full, renormalized propagator $\Delta(p)$ [21].

III. LOW-ENERGY BEHAVIOR OF VCS

In a two-step reaction on a single nucleon, such as $\gamma^* N \rightarrow \gamma^* N$, the intermediate nucleon lines in the s - and u -channel pole diagrams of class A are off mass shell, while the external nucleons are on shell. The early, manifestly covariant derivations of the low-energy theorem for Compton scattering [2,3] took into account that the associated half-off-shell electromagnetic vertex of the nucleon has a different and more complicated structure than the free vertex. However, it was shown that the model- and representation-dependent properties of an off-shell nucleon do not enter in the leading terms of the full Compton scattering amplitude when the irreducible two-photon amplitude of class B is included consistently. This was explicitly shown by expanding Eq. (2.10) to first order in q^μ and by constructing the leading-order term of $\Gamma_B^{\mu\nu}$ with the help of Eq. (2.15) and crossing symmetry or, equivalently, the second gauge-invariance constraint. The final result for the amplitude was given in the laboratory frame and in Coulomb gauge.

In order to obtain the LET for VCS, we will proceed on the operator level and combine the method of [2,3] with ideas of an effective Lagrangian approach to Compton scattering [4]. Let us first recall that the electromagnetic three-point and four-point Green's functions of Eqs. (2.1) and (2.2) depend on the choice of the interpolating field Ψ of the proton, i.e., are "representation dependent." Therefore, the truncated Green's functions in momentum space, Eqs. (2.6) and (2.7), are in general not directly related to observables, except for $p_i^2 = p_f^2 = M^2$. Consequently, the separation into class A and class B is necessarily representation dependent, since the total Green's function, Eq. (2.6), as well as the individual building blocks of class A are representation de-

pendent; this has of course no effect on the final on-shell result, which is representation independent. This was explicitly shown in [22] for the case of real Compton scattering off the pion. On the other hand, for any given appropriate interpolating field satisfying the equal-time commutation relation of Eq. (2.3) the Ward-Takahashi identities, Eqs. (2.8) and (2.9), hold, providing important consistency relations between the different Green's functions. In the following we will make use of these relations for arbitrary P , q , and q' , which, in particular, includes arbitrary p_i^2 and p_f^2 . Only at the end, the observable on-shell case will be considered.

A. Derivation of the low-energy theorem

We will derive this theorem by using a convenient representation, the "canonical form," of the most general and gauge invariant effective Lagrangian. In the framework of effective Lagrangians, a canonical form is defined as a representation with the minimal number of independent structures (see, e.g., [23]). For the class A terms, we need the electromagnetic vertex and the propagator of the nucleon. Below the pion-production threshold, the Lagrangian for a single proton, interacting with an electromagnetic field, can be brought into the canonical form [4]

$$\begin{aligned} \mathcal{L}_{\gamma NN} = & \bar{\Psi}(i\not{D} - M)\Psi - e \sum_{n=1}^{\infty} [(-\square)^{n-1} \partial^\nu F_{\mu\nu}] F_{1n} \bar{\Psi} \gamma^\mu \Psi \\ & - \frac{e}{4M} \left(\kappa F_{\mu\nu} + \sum_{n=1}^{\infty} [(-\square)^n F_{\mu\nu}] F_{2n} \right) \bar{\Psi} \sigma^{\mu\nu} \Psi, \end{aligned} \quad (3.1)$$

where $D_\mu \Psi = (\partial_\mu + ieA_\mu)\Psi$, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$. The electromagnetic structure of the proton is accounted for through the Dirac and Pauli form factors, F_1 and F_2 , respectively, which are expanded according to

$$\begin{aligned} F_1(q^2) &= 1 + \sum_{n=1}^{\infty} (q^2)^n F_{1n}, \\ F_2(q^2) &= \kappa + \sum_{n=1}^{\infty} (q^2)^n F_{2n}, \quad \kappa = 1.79. \end{aligned} \quad (3.2)$$

To this representation of the Lagrangian belongs, of course, a particular canonical Lagrangian of order e^2 that generates the class B terms. However, as will be seen below, we do not need to know it in detail for this derivation. Clearly, Eq. (3.1) is invariant under the gauge transformation $\Psi \rightarrow \exp[-ie\alpha(x)]\Psi$, and $A_\mu \rightarrow A_\mu + \partial_\mu \alpha$. In order to arrive at Eq. (3.1), use has implicitly been made of the method of field transformations (see, e.g., [23–26]). It should be stressed that all ingredients needed for Eq. (3.1) are on-shell quantities that can be determined model independently from electron-proton scattering. Any explicit off-shell dependence of the irreducible three-point Green's function has been transformed away and will thus not show up in the class A contribution. Such transformations, however, generate concomitant irreducible class B terms for the amplitude that must be treated consistently (see [4] for details).

Using standard Feynman rules, the irreducible vertex associated with the effective Lagrangian of Eq. (3.1) is found to be

$$\Gamma_{\text{eff}}^{\mu}(p_f, p_i) = \Gamma_{\text{eff}}^{\mu}(p_f - p_i) = \gamma^{\mu} F_1(q^2) + \frac{1 - F_1(q^2)}{q^2} q^{\mu} \not{q} + i \frac{\sigma^{\mu\nu} q_{\nu}}{2M} F_2(q^2), \quad q = p_f - p_i. \quad (3.3)$$

Since it is an important ingredient in our derivation of the LET, we emphasize that the vertex of Eq. (3.3) satisfies the Ward-Takahashi identity of Eq. (2.8); the corresponding propagator in the representation that yields Eq. (3.1) is the Feynman propagator of a point proton. As a consequence of gauge invariance, Eq. (3.1) automatically generates a term in the vertex proportional to q^{μ} .

We note that the effective Lagrangian approach provides a natural explanation for the vertex of Eq. (3.3) which has previously been used by several authors as a simple means to restore gauge invariance in the form of the Ward-Takahashi identity (see, for example, [27]). However, it is not an independent building block for any amplitude, but must be used together with the corresponding irreducible class B terms for the reaction in question.

In principle, we could now proceed to construct the most general effective Lagrangian relevant to generate the corresponding class B terms for VCS. This would allow us, together with a calculation of the pole terms involving the vertex of Eq. (3.3), to determine the model-dependent terms of $\Gamma^{\mu\nu}$ at low energies. However, at this point it is more straightforward to apply the results of the last section. We obtain for $\Gamma_{A,\text{eff}}^{\mu\nu}$ in the framework of Eqs. (3.1) and (3.3)

$$\Gamma_{A,\text{eff}}^{\mu\nu} = \Gamma_{\text{eff}}^{\nu}(-q') i S_F(p_i + q) \Gamma_{\text{eff}}^{\mu}(q) + \Gamma_{\text{eff}}^{\mu}(q) i S_F(p_i - q') \Gamma_{\text{eff}}^{\nu}(-q'). \quad (3.4)$$

Making use of Eq. (2.15) to obtain the gauge-invariance constraint for class B in our representation, we see that it has the simple form¹

$$q_{\mu} \Gamma_B^{\mu\nu} = 0. \quad (3.5)$$

This is due to the fact that the vertex of Eq. (3.3) depends on the momentum transfer, only. Note that Eq. (3.5) is still an operator equation, valid for arbitrary P , q , and q' .

Class A is defined to contain for any representation the pole terms of $\Gamma^{\mu\nu}$ which are singular in the limit $q^{\mu} \rightarrow 0$; we will come back to this point in the next section. We thus can make for class B the following ansatz for the q dependence:

$$\Gamma_B^{\mu\nu}(P, q, q') = a^{\mu,\nu} + b^{\mu\rho,\nu} q_{\rho} + c^{\mu\rho\sigma,\nu} q_{\rho} q_{\sigma} + \dots \quad (3.6)$$

The coefficients $a^{\mu,\nu}, \dots$ are 4×4 matrices which have to be constructed from the 16 Dirac matrices, the metric tensor $g^{\alpha\beta}$, the completely antisymmetric Levi-Civita pseudotensor $\epsilon^{\alpha\beta\gamma\delta}$, and the remaining independent variables P^{α} and

q'^{α} . The form of these coefficients will be constrained by Lorentz invariance, gauge invariance, crossing symmetry, and the discrete symmetries, C , P , and T . Contracting this ansatz with q_{μ} , the condition for the class B operator, Eq. (3.5), becomes

$$q_{\mu} \Gamma_B^{\mu\nu} = a^{\mu,\nu} q_{\mu} + b^{\mu\rho,\nu} q_{\rho} q_{\mu} + \dots = 0. \quad (3.7)$$

Multiple partial differentiation of Eq. (3.7) with respect to q_{α} results in the following conditions for the coefficients,

$$a^{\mu,\nu} = 0, \quad b^{\mu\rho,\nu} + b^{\rho\mu,\nu} = 0, \quad \sum_{\substack{6 \text{ perm.} \\ (\mu,\rho,\sigma)}} c^{\mu\rho\sigma,\nu} = 0, \dots \quad (3.8)$$

Since current conservation is expected to hold for arbitrary q , the technique of partial differentiation to obtain conditions for the coefficients can easily be extended to obtain constraints for higher-order coefficients. The simple constraints of Eq. (3.8) are based on the fact that P^{α} , q^{α} , and q'^{α} are independent variables; an implicit dependence would make matters more complicated.

From $a^{\mu,\nu} = 0$ we can already conclude that the operator of class B contains no contributions which involve powers of q' only, without powers of q . Taking Eq. (3.8) into account, we now expand with respect to q' ,

$$\Gamma_B^{\mu\nu}(P, q, q') = B^{\mu\rho,\nu} q_{\rho} + B^{\mu\rho,\nu\alpha} q_{\rho} q'_{\alpha} + \dots + C^{\mu\rho\sigma,\nu} q_{\rho} q_{\sigma} + C^{\mu\rho\sigma,\nu\alpha} q_{\rho} q_{\sigma} q'_{\alpha} + \dots \quad (3.9)$$

If we apply crossing symmetry to Eq. (3.9) we find, as expected, that indeed all terms vanish that involve powers of q' only. Thus we find for the leading term of the model-dependent class B ,

$$\Gamma_B^{\mu\nu}(P, q, q') = B^{\mu\rho,\nu\alpha}(P) q_{\rho} q'_{\alpha} + O(q q q', q q' q'),$$

$$B^{\mu\rho,\nu\alpha} = -B^{\rho\mu,\nu\alpha} = B^{\nu\alpha,\mu\rho}, \quad (3.10)$$

where the conditions for $B^{\mu\rho,\nu\alpha}$ result from gauge invariance and crossing symmetry, respectively. To be specific, after imposing the constraints of Eq. (3.10) and of the discrete symmetries one obtains three possible structures for $B^{\mu\rho,\nu\alpha}$ on the *operator level*:

$$\begin{aligned} B^{\mu\rho,\nu\alpha}(P) = & i(g^{\mu\nu} g^{\rho\alpha} - g^{\rho\nu} g^{\mu\alpha}) f_1(P^2) + i(g^{\mu\nu} P^{\rho} P^{\alpha} \\ & + g^{\rho\alpha} P^{\mu} P^{\nu} - g^{\rho\nu} P^{\mu} P^{\alpha} - g^{\mu\alpha} P^{\rho} P^{\nu}) f_2(P^2) \\ & + \epsilon^{\mu\rho\nu\alpha} \gamma_5 f_3(P^2), \quad \epsilon_{0123} = 1. \end{aligned} \quad (3.11)$$

In summary, we have shown that on the *operator level* the terms of order $O(q_{\alpha}^{-1})$, $O(q'^{-1}_{\alpha})$, $O(1)$, $O(q_{\alpha})$, and $O(q'_{\alpha})$ are contained in $\Gamma_{A,\text{eff}}^{\mu\nu}$, Eq. (3.4). They are therefore determined model independently through on-shell quantities; model-dependent terms first appear in the order $O(q_{\alpha} q'_{\beta})$; all operators which contain either only powers of q or only powers of q' can entirely be obtained from this $\Gamma_{A,\text{eff}}^{\mu\nu}$. In the next section we will discuss the implications of these findings for the on-shell VCS matrix element.

¹In the following we omit the subscript “eff.”

B. Application

We now want to apply the above result to the observable case where the nucleons are on mass shell, i.e., we consider the matrix element. For that purpose we first define the matrix element of $\Gamma^{\mu\nu}$ between positive-energy spinors as

$$V_{s_i s_f}^{\mu\nu}(P, q, q') = \bar{u}(p_f, s_f) \Gamma^{\mu\nu}(P, q, q') u(p_i, s_i). \quad (3.12)$$

At this point we have to keep in mind that for the variables we use, P , q , and q' , the on-shell condition $p_i^2 = p_f^2 = M^2$ is equivalent to $P \cdot (q - q') = 0$, and $P^2 + (q - q')^2 = 4M^2$. In other words, the four-momenta chosen for the description of the off-shell Green's function will no longer be independent for the on-shell invariant amplitude. In particular, the distinction between powers of q only or, respectively, of q' only in $\Gamma^{\mu\nu}$ is not valid anymore for the matrix element since $q \cdot P = q' \cdot P$.

Let us consider as an application the case where the initial photon is virtual and spacelike and the final photon is real, $\gamma^*(q, \epsilon) + p(p_i, s_i) \rightarrow \gamma(q', \epsilon') + p(p_f, s_f)$. The following discussion does not include the Bethe-Heitler terms of the physical process $p(e, e' p) \gamma$, where the real photon is radiated by the initial or final electron, since these terms are not

part of $\Gamma^{\mu\nu}$. For the final photon the Lorentz condition $q' \cdot \epsilon' = 0$ is automatically satisfied, and we are free to choose, in addition, the Coulomb gauge $\epsilon'^\mu = (0, \vec{\epsilon}')$ which implies $\vec{q}' \cdot \vec{\epsilon}' = 0$. We write the invariant amplitude in the convention of [28] as $\mathcal{M} = -ie^2 \epsilon_\mu M^\mu$, where $\epsilon_\mu = e \bar{u} \gamma_\mu u / q^2$ is the polarization vector of the virtual photon. With a suitable choice for the reference frame, $q^\mu = (q_0, 0, 0, |\vec{q}|)$, the Lorentz condition $q \cdot \epsilon = 0$ and current conservation, $q_\mu M^\mu = 0$, may be used to reexpress the invariant amplitude as [29]

$$\mathcal{M} = ie^2 \left(\vec{\epsilon}_T \cdot \vec{M}_T + \frac{q^2}{q_0} \epsilon_z M_z \right). \quad (3.13)$$

Note that as $q_0 \rightarrow 0$, both ϵ_z and M_z tend to zero such that $\mathcal{M}$ in Eq. (3.13) remains finite. Making use that we now know the general structure of the first undetermined operator, Eq. (3.11), we can now consider the matrix element up to second order in q or q' . This will enable us to explicitly show how far the amplitude is determined and where the first model-dependent terms enter. After a reduction from Dirac spinors to Pauli spinors the transverse and longitudinal parts of Eq. (3.13) may be expressed in terms of eight and four independent structures, respectively [4,14,30]:

$$\begin{aligned} \vec{\epsilon}_T \cdot \vec{M}_T = & \vec{\epsilon}'^* \cdot \vec{\epsilon}_T A_1 + i \vec{\sigma} \cdot (\vec{\epsilon}'^* \times \vec{\epsilon}_T) A_2 + (\hat{q}' \times \vec{\epsilon}'^*) \cdot (\hat{q} \times \vec{\epsilon}_T) A_3 + i \vec{\sigma} \cdot (\hat{q}' \times \vec{\epsilon}'^*) \times (\hat{q} \times \vec{\epsilon}_T) A_4 + i \hat{q} \cdot \vec{\epsilon}'^* \vec{\sigma} \cdot (\hat{q} \times \vec{\epsilon}_T) A_5 \\ & + i \hat{q}' \cdot \vec{\epsilon}_T \vec{\sigma} \cdot (\hat{q}' \times \vec{\epsilon}'^*) A_6 + i \hat{q} \cdot \vec{\epsilon}'^* \vec{\sigma} \cdot (\hat{q}' \times \vec{\epsilon}_T) A_7 + i \hat{q}' \cdot \vec{\epsilon}_T \vec{\sigma} \cdot (\hat{q} \times \vec{\epsilon}'^*) A_8, \end{aligned} \quad (3.14)$$

$$\epsilon_z M_z = \epsilon_z \vec{\epsilon}'^* \cdot \hat{q} A_9 + i \epsilon_z \vec{\epsilon}'^* \cdot \hat{q} \vec{\sigma} \cdot (\hat{q}' \times \hat{q}) A_{10} + i \epsilon_z \vec{\sigma} \cdot (\hat{q} \times \vec{\epsilon}'^*) A_{11} + i \epsilon_z \vec{\sigma} \cdot (\hat{q}' \times \vec{\epsilon}'^*) A_{12}. \quad (3.15)$$

The results for the functions A_i in the c.m. system expanded up to $O(2)$, i.e., $|\vec{q}'|^2$, $|\vec{q}'||\vec{q}|$, and $|\vec{q}|^2$, are shown in Tables I and II. The derivation of the corresponding expressions is outlined in the Appendix.

In the amplitudes A_5 and A_9 , terms of order $1/|\vec{q}'|$ appear. We have not further expanded them in $|\vec{q}|$ and kept, e.g., the on-shell form factors G_M and G_E . This was done mainly to stress that according to Low's theorem [9] these divergent terms are already entirely fixed. They are due to soft radiation off external lines, with the intermediate line approaching the mass shell in the pole terms. They can be obtained from, e.g., the Born terms that contain the same on-shell information (see however the caveats in Sec. IV). In order to uniquely identify these singular contributions we have expanded all relevant expressions in terms of $|\vec{q}|$ and $|\vec{q}'|$. This is the reason why the argument of the form factors is $Q^2 = q^2|_{|\vec{q}'|=0} = -2M(E_i - M)$, where $E_i = \sqrt{M^2 + \vec{q}^2}$.

Not only the irregular contribution, but all terms in the amplitudes up to terms linear in the photon momenta are uniquely determined through well-known properties of the free nucleon: its mass, charge, and magnetic moment. These are the terms that make up the LET for VCS. Up to $O(2)$, the coefficients in addition also involve the electric mean square radius, which we know from electron-proton scatter-

ing, as well as the electric and magnetic polarizabilities which also enter in real Compton scattering. The latter are the coefficients of the first model-dependent terms in the expansion of the scattering matrix element. Their specific form, and thus the definitions of the polarizabilities, depend on the particular representation one chooses for the effective Lagrangian; the $O(2)$ results in the tables are specific for the ‘‘canonical form,’’ which happens to be the standard choice for real Compton scattering. Note that to the order considered $A_7 = -A_8 = A_{10}$. The amplitudes A_1 and A_9 contain the electric polarizability $\bar{\alpha}$ of the proton whereas A_3 involves the magnetic polarizability $\bar{\beta}$ [4]. In terms of the functions $f_i(P^2)$ of Eq. (3.11) they are defined as

$$\bar{\alpha} = e^2 [f_1(4M^2) + 4M^2 f_2(4M^2)], \quad \bar{\beta} = -e^2 f_1(4M^2). \quad (3.16)$$

The function $f_3(4M^2)$ only contributes at order $O(3)$, since γ_5 connects upper and lower component of positive-energy spinors which effectively leads to an additional power of $|\vec{q}|$ or $|\vec{q}'|$ in the matrix element.

In Table III we also show the results for the transverse amplitudes in the c.m. frame for real Compton scattering. Since $A_5 = -A_6$ and $A_7 = -A_8$ the result effectively involves six independent amplitudes as required by time reversal in-

TABLE I. Transverse functions A_i of Eq. (3.14) in the c.m. frame. The functions are expanded in terms of $|\vec{q}'|$ and $|\vec{q}|$ of the final real and initial virtual photon, respectively. $N_i = \sqrt{(E_i + M)/2M}$ is the normalization factor of the initial spinor, where $E_i = \sqrt{M^2 + |\vec{q}|^2}$. $G_E(q^2) = F_1(q^2) + (q^2/4M^2)F_2(q^2)$ and $G_M(q^2) = F_1(q^2) - F_2(q^2)$ are the electric and magnetic Sachs form factors, respectively. $r_E^2 = 6G_E'(0) = (0.74 \pm 0.02) \text{ fm}^2$ is the electric mean square radius [31] and $\kappa = 1.79$ the anomalous magnetic moment of the proton. Q^μ is defined as $q^\mu|_{\vec{q}'=0} = (M - E_i, \vec{q})$, $Q^2 = -2M(E_i - M)$, and $z = \hat{q}' \cdot \hat{q}$. $\bar{\alpha}$ and $\bar{\beta}$ are the electric and magnetic Compton polarizabilities of the proton, respectively.

A_1	$-\frac{1}{M} + \frac{z}{M^2} \vec{q} - \left(\frac{1}{8M^3} + \frac{r_E^2}{6M} - \frac{\kappa}{4M^3} - \frac{\bar{\alpha}}{e^2}\right) \vec{q}' ^2 + \left(\frac{1}{8M^3} + \frac{r_E^2}{6M} - \frac{z^2}{M^3} + \frac{(1+\kappa)\kappa}{4M^3}\right) \vec{q} ^2$
A_2	$\frac{1+2\kappa}{2M^2} \vec{q}' - \frac{\kappa^2}{4M^3} \vec{q}' ^2 + \frac{z\kappa}{2M^3} \vec{q}' \vec{q} - \frac{(1+\kappa)^2}{4M^3} \vec{q} ^2$
A_3	$-\frac{1}{M^2} \vec{q} + \left(\frac{1}{4M^3} + \frac{\bar{\beta}}{e^2}\right) \vec{q}' \vec{q} + \frac{(3-2\kappa-\kappa^2)z}{4M^3} \vec{q} ^2$
A_4	$-\frac{(1+\kappa)^2}{2M^2} \vec{q} - \frac{(2+\kappa)\kappa}{4M^3} \vec{q}' \vec{q} + \frac{(1+\kappa)^2z}{4M^3} \vec{q} ^2$
A_5	$-\frac{N_i G_M(Q^2)}{(E_i + z \vec{q})(E_i + M)} \frac{ \vec{q} ^2}{ \vec{q}' } + \frac{(1+\kappa)\kappa}{4M^3} \vec{q} ^2$
A_6	$\frac{1+\kappa}{2M^2} \vec{q}' - \frac{(1+\kappa)\kappa}{4M^3} \vec{q}' ^2 - \frac{(1+\kappa)z}{2M^3} \vec{q}' \vec{q} $
A_7	$-\frac{1+3\kappa}{4M^3} \vec{q}' \vec{q} $
A_8	$\frac{1+3\kappa}{4M^3} \vec{q}' \vec{q} $

variance [4]. When comparing with other expressions in the literature [1,2,4,30], one has to keep in mind that they usually are given in the laboratory frame. This observation accounts, for example, for the difference of the LET for A_1 and A_3 of real Compton scattering in the laboratory and the c.m. frame. Note also that the $1/|\vec{q}'|$ singularity disappears in the real Compton scattering limit, since $\omega = |\vec{q}| = |\vec{q}'|$ in this case.

In conclusion, the low-energy behavior of the VCS matrix element for $e^- + p \rightarrow e^- + p + \gamma$, expanded up to order $O(2)$ in $|\vec{q}|$ and $|\vec{q}'|$, contains, in addition to the structure coefficients that enter into real Compton scattering, also the electric mean square radius and the electric and magnetic Sachs form factors in the spacelike region which all can be obtained from electron scattering off the proton. For the reaction $\gamma + p \rightarrow p + e^+ e^-$ [13], the analogous information for

timelike momentum transfers is needed. However, here one has to keep in mind that this information is not directly accessible for $0 < q^2 < 4M^2$.

IV. THE BORN TERMS

From the above discussion in a particular representation one might be tempted to conclude that the leading terms for VCS, on the operator level or for the amplitude, are in general simply given by ‘‘Born terms.’’ However, some caveats are in order. First, one has to keep in mind that the low-energy behavior was obtained from considerations involving the most general ansatz for the truncated four-point Green’s function. All terms one can think of are included into class A and class B. In, e.g., the derivation of [3] of the LET for Compton scattering, where no special representation is cho-

TABLE II. Longitudinal functions A_i of Eq. (3.15) in the c.m. frame. See caption of Table I.

A_9	$\frac{N_i G_E(Q^2)}{(E_i + z \vec{q})(E_i + M)} \frac{ \vec{q} ^2}{ \vec{q}' } - \frac{1}{M} + \frac{z}{M^2} \vec{q} - \left(\frac{1}{8M^3} + \frac{r_E^2}{6M} - \frac{\kappa}{4M^3} - \frac{\bar{\alpha}}{e^2}\right) \vec{q}' ^2 + \left(\frac{1}{8M^3} + \frac{r_E^2}{6M} - \frac{z^2}{M^3}\right) \vec{q} ^2$
A_{10}	$-\frac{1+3\kappa}{4M^3} \vec{q}' \vec{q} $
A_{11}	$-\frac{1+2\kappa}{2M^2} \vec{q}' + \frac{\kappa^2}{4M^3} \vec{q}' ^2 + \frac{(1+\kappa)z}{4M^3} \vec{q}' \vec{q} + \frac{1+2\kappa}{4M^3} \vec{q} ^2$
A_{12}	$\frac{(1+\kappa)z}{2M^2} \vec{q}' - \frac{(1+\kappa)\kappa z}{4M^3} \vec{q}' ^2 - \frac{(1+\kappa)(2z^2-1)}{4M^3} \vec{q}' \vec{q} - \frac{(1+\kappa)z}{4M^3} \vec{q} ^2$

TABLE III. Transverse functions A_i in the c.m. frame for both photons real: $q^2 = q'^2 = 0$, $|\vec{q}| = |\vec{q}'| = \omega$.

A_1	$-\frac{1}{M} + \frac{z}{M^2} \omega + \left(-\frac{z^2}{M^3} + \frac{(2+\kappa)\kappa}{4M^3} + \frac{\bar{\alpha}}{e^2} \right) \omega^2$
A_2	$\frac{1+2\kappa}{2M^2} \omega - \frac{1+2\kappa(1-z+\kappa)}{4M^3} \omega^2$
A_3	$-\frac{1}{M^2} \omega + \left(\frac{1+(3-\kappa(2+\kappa))z}{4M^3} + \frac{\bar{\beta}}{e^2} \right) \omega^2$
A_4	$-\frac{(1+\kappa)^2}{2M^2} \omega + \frac{z+\kappa(2+\kappa)(z-1)}{4M^3} \omega^2$
A_5	$-\frac{1+\kappa}{2M^2} \omega + \frac{(2z+\kappa)(1+\kappa)}{4M^3} \omega^2$
A_6	$\frac{1+\kappa}{2M^2} \omega - \frac{(2z+\kappa)(1+\kappa)}{4M^3} \omega^2$
A_7	$-\frac{1+3\kappa}{4M^3} \omega^2$
A_8	$\frac{1+3\kappa}{4M^3} \omega^2$

sen, it is clearly shown that *both* class A and class B terms are needed. Implicit in all derivations is of course the assumption that a description in terms of observable asymptotic hadronic degrees of freedom is sufficient and complete at low energies. Even though subnucleonic degrees of freedom, quarks and gluons, are ultimately the origin of the structure of the nucleon, it was shown [32,33] that an effective field-theory approach [34–37] in terms of hadrons is meaningful at low energies, thus allowing a classification into class A and class B.

Second, there is an ambiguity concerning what exactly is meant by “Born terms,” once phenomenological form factors are introduced and the result is not obtained from a microscopic Lagrangian. We will illustrate this ambiguity by considering different representations of the photon-nucleon vertex. All representations contain the same information concerning the electromagnetic structure of the on-shell nucleon, as obtained in electron-nucleon scattering, but differ in the

half-off-shell situation encountered in the s - and u -channel pole terms of Compton scattering. This difference is, of course, accompanied by different class B terms such that the total result is the same.

To explain the above points in more detail, we will first reconsider the most general expression for class A and, without going to a special representation, identify those terms which contain the irregular contribution for $q^\mu \rightarrow 0$ or $q'^\mu \rightarrow 0$. We find that these contributions can be expressed in terms of on-shell quantities, in our case the Dirac and Pauli form factors F_1 and F_2 . We then show that the use of on-shell equivalent electromagnetic vertices gives rise to the same results for the VCS matrix element as far as the *irregular* terms are concerned, but not every choice will result in “Born terms” which are gauge invariant.

A. Irregular contribution to the VCS matrix element

When calculating $V_{s_i s_f}^{\mu\nu}$, the irregular contribution originates from the singularities of the propagators $S(p_i + q)$ and $S(p_i - q')$ in the s and u channels, respectively. To be specific, below pion-production threshold the renormalized, full propagator can be written as

$$S(p + q) = S_F(p + q) + \text{regular terms},$$

$$(p + q)^2 < (M + m_\pi)^2, \quad (4.1)$$

where $S_F(p)$ denotes the free propagator of a nucleon with mass M . By “regular terms” we mean terms which have a well-defined, nonsingular limit for $q^\mu \rightarrow 0$. Thus, as long as we are only interested in the irregular terms, we can simply replace the full, renormalized propagator by the free Feynman propagator, S_F .

The most general form of the irreducible, electromagnetic vertex of the nucleon can be expressed in terms of 12 operators and associated form functions [38,39]. These functions depend on three scalar variables, e.g., the squared momentum transfer and the invariant masses of the initial and final nucleon lines. A convenient parametrization of $\Gamma^\mu(p_f, p_i)$ is given by

$$\Gamma^\mu(p_f, p_i) = \sum_{\alpha, \beta = +, -} \Lambda_\alpha(p_f) \left(\gamma^\mu F_1^{\alpha\beta} + i \frac{\sigma^{\mu\nu} q_\nu}{2M} F_2^{\alpha\beta} + \frac{q^\mu}{M} F_3^{\alpha\beta} \right) \Lambda_\beta(p_i), \quad F_i^{\alpha\beta} = F_i^{\alpha\beta}(q^2, p_f^2, p_i^2), \quad (4.2)$$

where $q = p_f - p_i$, and $\Lambda_\pm(p) = (M \pm \not{p})/2M$. We have chosen a form which differs slightly from the convention of [39] in the definition of the projection operators and the normalization of the F_3 form functions. We only need the following on-shell properties of the form functions:

$$F_1^{++}(q^2, M^2, M^2) = F_1(q^2), \quad F_2^{++}(q^2, M^2, M^2) = F_2(q^2), \quad F_3^{++}(q^2, M^2, M^2) = F_3(q^2) = 0, \quad (4.3)$$

where F_1 and F_2 are the standard Dirac and Pauli form factors and F_3 vanishes because of time-reversal invariance. One can also show that $F_3(q^2)$ vanishes due to current conservation.

We now systematically isolate the irregular part of, e.g., the s -channel pole diagram,²

$$V_{A,s}^{\mu\nu} = \bar{u}(p_f) \Gamma^\nu(p_f, p_f + q') i S(p_i + q) \Gamma^\mu(p_i + q, p_i) u(p_i) \approx \bar{u}(p_f) \Gamma^\nu(p_f, p_f + q') i S_F(p_i + q) \Gamma^\mu(p_i + q, p_i) u(p_i),$$

²In the following we omit the indices s_i and s_f .

where we made use of Eq. (4.1), and where the symbol $\approx$ denotes equality up to regular terms. We now insert Eq. (4.2) for $\Gamma^\mu(p_i+q, p_i)$ and use $\Lambda_-(p_i)u(p_i)=0$ and $\Lambda_+(p_i)u(p_i)=u(p_i)$ to obtain

$$V_{A,s}^{\mu\nu} \approx \bar{u}(p_f) \Gamma^\nu(p_f, p_f+q') iS_F(p_i+q) \{ \Lambda_+(p_i+q) [\gamma^\mu F_1^{++}(q^2, s, M^2) + \dots] + \Lambda_-(p_i+q) [\gamma^\mu F_1^{--}(q^2, s, M^2) + \dots] \} u(p_i),$$

where $s=(p_i+q)^2$. Since $S_F(p_i+q)\Lambda_-(p_i+q)$ results in a regular term, we have

$$V_{A,s}^{\mu\nu} \approx i\bar{u}(p_f) \Gamma^\nu(p_f, p_f+q') \frac{\not{p}_i + \not{q} + M}{s-M^2} \Lambda_+(p_i+q) [\gamma^\mu F_1^{++}(q^2, s, M^2) + \dots] u(p_i).$$

We now expand the form functions around $s_0=M^2$ and note that in the higher-order terms the powers of $(s-M^2)$ cancel the denominator, $s-M^2$, of the Feynman propagator and thus give rise to regular terms,

$$V_{A,s}^{\mu\nu} \approx \bar{u}(p_f) \Gamma^\nu(p_f, p_f+q') iS_F(p_i+q) \Lambda_+(p_i+q) \left(\gamma^\mu F_1(q^2) + i \frac{\sigma^{\mu\rho} q_\rho}{2M} F_2(q^2) \right) u(p_i),$$

where we made use of Eq. (4.3). Using $\Lambda_+(p_i+q)=1-\Lambda_-(p_i+q)$ and then repeating the same procedure for $\Gamma^\nu(p_f, p_f+q')$ we finally obtain

$$V_{A,s}^{\mu\nu} \approx \bar{u}(p_f, s_f) \Gamma_{F_1, F_2}^\nu(-q') iS_F(p_i+q) \Gamma_{F_1, F_2}^\mu(q) u(p_i, s_i), \quad (4.4)$$

where we introduced the abbreviation

$$\Gamma_{F_1, F_2}^\mu(q) = \gamma^\mu F_1(q^2) + i \frac{\sigma^{\mu\nu} q_\nu}{2M} F_2(q^2). \quad (4.5)$$

The procedure for the u -channel part, $V_{A,u}^{\mu\nu}$ is completely analogous and we obtain for the sum of s - and u -channel contributions

$$V_A^{\mu\nu} \approx \bar{u}(p_f) [\Gamma_{F_1, F_2}^\nu(-q') iS_F(p_i+q) \Gamma_{F_1, F_2}^\mu(q) + \Gamma_{F_1, F_2}^\mu(q) iS_F(p_i-q') \Gamma_{F_1, F_2}^\nu(-q')] u(p_i) \equiv V_{A'}^{\mu\nu}, \quad (4.6)$$

which only involves on-shell quantities, the Dirac and Pauli form factors and the nucleon mass. Explicit calculation, including the use of the Dirac equation, shows that $V_{A'}^{\mu\nu}$ of Eq. (4.6) is, in fact, identical with evaluating Eq. (3.4) between on-shell spinors. With the help of either Eq. (2.12) or by straightforward calculation it can easily be shown that Eq. (4.6) is gauge invariant. This is a special feature when working with these particular vertex operators and was essential for the derivation in [14]. It is quite unexpected, since the electromagnetic vertex, Eq. (4.5), and the nucleon propagator in Eq. (4.6) do not satisfy the Ward-Takahashi identity (except at the real photon point, $q^2=0$):

$$q_\mu \Gamma_{F_1, F_2}^\mu(p_f, p_i) = (\not{p}_f - \not{p}_i) F_1(q^2) \neq S_F^{-1}(p_f) - S_F^{-1}(p_i). \quad (4.7)$$

B. Different representations of the on-shell vertex

We now turn to ‘‘Born-term’’ calculations involving other representations of the nucleon current operator, i.e., other ways to introduce the model-independent information about the electromagnetic structure of the free nucleon into the calculation of the lowest-order terms. Two commonly used alternative ways to parametrize the nucleon current operator are [40]

$$\Gamma_{G_E, G_M}^\mu(p_f, p_i) = \left(1 - \frac{q^2}{4M^2} \right)^{-1} \left(\frac{P^\mu}{2M} G_E(q^2) + \frac{\gamma^\mu \not{P} \not{q} - \not{q} \not{P} \gamma^\mu}{8M^2} G_M(q^2) \right), \quad (4.8)$$

$$\Gamma_{H_1, H_2}^\mu(p_f, p_i) = \gamma^\mu H_1(q^2) - \frac{P^\mu}{2M} H_2(q^2), \quad (4.9)$$

where $P=p_i+p_f$ and

$$G_E = F_1 + \frac{q^2}{4M^2} F_2, \quad G_M = H_1 = F_1 + F_2, \quad H_2 = F_2. \quad (4.10)$$

Given Eqs. (4.10), it is straightforward to show the equivalence for the free proton current, the matrix elements of Eqs. (3.3), (4.5), (4.8), and (4.9) between free positive-energy spinors. On the other hand, the current operators in Eqs. (4.5), (4.8), and (4.9) do not satisfy the Ward-Takahashi identity when used in conjunction with free propagators, not even at the real-photon point [recall Eq. (4.7)]:

$$q_\mu \Gamma_{G_E, G_M}^\mu(p_f, p_i) = \left(1 - \frac{q^2}{4M^2} \right)^{-1} \frac{p_f^2 - p_i^2}{2M} G_E(q^2) \neq S_F^{-1}(p_f) - S_F^{-1}(p_i), \quad (4.11)$$

$$q_\mu \Gamma_{H_1, H_2}^\mu(p_f, p_i) = (\not{p}_f - \not{p}_i) H_1(q^2) - \frac{p_f^2 - p_i^2}{2M} H_2(q^2) \neq S_F^{-1}(p_f) - S_F^{-1}(p_i). \quad (4.12)$$

The ‘‘Born terms’’ calculated with these electromagnetic vertices and free-nucleon propagators are

$$V_X^{\mu\nu} = \bar{u}(p_f, s_f) [\Gamma_X^\nu(p_f, p_f + q') iS_F(p_i + q) \Gamma_X^\mu(p_i + q, p_i) + \Gamma_X^\mu(p_f, p_f - q) iS_F(p_i - q') \Gamma_X^\nu(p_i - q', p_i)] u(p_i, s_i), \quad (4.13)$$

where X denotes either the vertex of Eq. (4.8) in terms of G_E, G_M or the vertex of Eq. (4.9) involving H_1, H_2 , respectively. These ‘‘Born terms’’ are by construction crossing symmetric but in both cases not gauge invariant,

$$q_\mu V_{G_E, G_M}^{\mu\nu} = i \left(1 - \frac{q^2}{4M^2} \right)^{-1} G_E(q^2) \bar{u}(p_f, s_f) \left(\Gamma_{G_E, G_M}^\nu(p_f, p_i + q) - \Gamma_{G_E, G_M}^\nu(p_f - q, p_i) + \Gamma_{G_E, G_M}^\nu(p_f, p_i + q) \frac{q}{2M} \right. \\ \left. + \frac{q}{2M} \Gamma_{G_E, G_M}^\nu(p_f - q, p_i) \right) u(p_i, s_i), \quad (4.14)$$

$$q_\mu V_{H_1, H_2}^{\mu\nu} = i [H_1(q^2) - H_2(q^2)] \bar{u}(p_f, s_f) [\Gamma_{H_1, H_2}^\nu(p_f, p_i + q) - \Gamma_{H_1, H_2}^\nu(p_f - q, p_i)] u(p_i, s_i) \\ - i H_2(q^2) \bar{u}(p_f, s_f) \left(\Gamma_{H_1, H_2}^\nu(p_f, p_i + q) \frac{q}{2M} + \frac{q}{2M} \Gamma_{H_1, H_2}^\nu(p_f - q, p_i) \right) u(p_i, s_i). \quad (4.15)$$

With the help of Eqs. (4.8) and (4.9) it is easy to see that in each case the nonvanishing divergence is of order q . The fact that Eqs. (4.14) and (4.15) do not contain terms of the type $a \cdot q/b \cdot q$ suggests that in general the different ‘‘Born terms,’’ based on equivalent parametrizations of the on-shell currents, will differ by regular terms, only. It is of course conceivable that the amplitude could differ by irregular terms, which are separately gauge invariant. Thus the above argument is not stringent and can only serve as a motivation for the following claim which is essentially equivalent to Low’s theorem applied to the particular case of VCS: *Any ‘‘Born-term’’ calculation involving electromagnetic current operators which correctly reproduce the on-shell electromagnetic current of the nucleon will yield the same irregular contribution to the VCS matrix element.* The key to the proof of this statement is the fact that any current operator which transforms as a Lorentz four-vector can be brought into the form of Eq. (4.2). On-shell equivalence then amounts to the constraint that all operators have the same on-shell limit of the F_i^{++} form functions. In general, no statement can be made for either the other form functions or off-shell kinematics.³ However, as we have seen above, the irregular contribution of class A, and thus of the total VCS matrix element, only involves the on-shell information contained in

$F_1^{++}(q^2, M^2, M^2)$ and $F_2^{++}(q^2, M^2, M^2)$. Any information beyond this will give rise to regular terms. Thus the above statement is true.

To illustrate the above, we bring for example the current operators of Eqs. (4.5) and (4.9), involving F_1 and F_2 or H_1 and H_2 , respectively, into the general form of Eq. (4.2). By using $1 = \Lambda_+(p) + \Lambda_-(p)$, the result for the commonly used form of Eq. (4.5) is given by

$$F_1^{\alpha\beta}(q^2, p_f^2, p_i^2) = F_1(q^2), \\ F_2^{\alpha\beta}(q^2, p_f^2, p_i^2) = F_2(q^2), \\ F_3^{\alpha\beta}(q^2, p_f^2, p_i^2) = 0, \quad \alpha, \beta = +, -. \quad (4.16)$$

For the vertex given in Eq. (4.9), we use $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}$ and momentum conservation at the vertex to rewrite

$$\frac{(p_i + p_f)^\mu}{2M} = \frac{\not{p}_f \gamma^\mu + \gamma^\mu \not{p}_i}{2M} - i \frac{\sigma^{\mu\nu} q_\nu}{2M}. \quad (4.17)$$

By inserting appropriate projection operators in the form $\not{p} = M[\Lambda_+(p) - \Lambda_-(p)]$ and as above, the vertex of Eq. (4.9) can be expressed as

$$\Gamma_{H_1, H_2}^\mu(p_f, p_i) = \Lambda_+(p_f) \left(\gamma^\mu [H_1(q^2) - H_2(q^2)] + i \frac{\sigma^{\mu\nu} q_\nu}{2M} H_2(q^2) \right) \Lambda_+(p_i) + \Lambda_-(p_f) \left(\gamma^\mu H_1(q^2) \right. \\ \left. + i \frac{\sigma^{\mu\nu} q_\nu}{2M} H_2(q^2) \right) \Lambda_+(p_i) + \Lambda_+(p_f) \left(\gamma^\mu H_1(q^2) + i \frac{\sigma^{\mu\nu} q_\nu}{2M} H_2(q^2) \right) \Lambda_-(p_i) \\ \left. + \Lambda_-(p_f) \left(\gamma^\mu [H_1(q^2) + H_2(q^2)] + i \frac{\sigma^{\mu\nu} q_\nu}{2M} H_2(q^2) \right) \Lambda_-(p_i). \quad (4.18)$$

³Further conditions on the form functions can be derived from the Ward-Takahashi identity and discrete symmetry requirements [38,39].

Of course, Eq. (4.18) contains the same on-shell information as Eq. (4.5), since the form factors satisfy Eq. (4.10). On the other hand, the expressions for all the other $F_i^{\alpha\beta}$ form functions differ. This is why the two “Born-term” calculations based on these two vertices differ with respect to regular terms. It is straightforward to extend the same considerations to the vertex involving G_E and G_M .

In conclusion, we have shown that a calculation based on only the “Born terms,” built from any of the many possible on-shell equivalent vertices and free nucleon propagators, yields the same results for the *irregular* terms as the LET. Thus these “Born terms” will differ among each other through regular terms. Furthermore, such “Born terms” are, in general, not gauge invariant; an exception is the commonly used form involving the Dirac and Pauli form factors F_1 and F_2 . “Generalized Born terms” which are made gauge invariant by hand through an *ad hoc* prescription also differ by regular terms.

Important starting point for the derivation of the LET are the irregular terms. It is thus also possible to split the total VCS amplitude into “Born terms” plus “rest,” instead of class A and B amplitudes, to arrive at the same result for the LET, i.e., up to and including terms linear in the photon three-momenta. In general, this result will have contributions from “Born terms” and the “rest” amplitude. If one uses a “generalized Born amplitude,” all the terms appearing in the LET are due to the expansion of the Born amplitude. It is a well-known feature of soft-photon theorems that they cannot make statements about terms which are separately gauge invariant [41–43]. One has to keep this in mind when discussing the structure-dependent higher-order terms of VCS, i.e., one needs to specify which Born or class-A terms have been separated. For example, in [14] the “Born terms” involving F_1 and F_2 were separated since they provide without any further manipulation a gauge-invariant amplitude. Then the residual part with respect to these particular “Born terms” was parametrized in terms of generalized polarizabilities. A natural question to ask is what would have happened had one separated a different choice of “generalized Born terms” and defined generalized polarizabilities in an analogous fashion with respect to the corresponding residual amplitude. Obviously one would, in general, have found different numerical values for the new generalized polarizabilities in order to obtain the same total result.

V. CONCLUSIONS

In studying the structure of composite strongly interacting systems the electromagnetic interaction has been the traditional and precise tool of investigation. In scattering of electrons from a nucleon, our knowledge is restricted to two form factors that we can extract from experiments. Even though we have not yet been able to fully explain this information on the basis of QCD, it is important to look for other observables allowing us to test approximations to the exact QCD solution and effective, QCD inspired models. Such effective models are expected to work especially at low energies. The electron accelerators now make it possible to study virtual Compton scattering, which is clearly more powerful in probing the nucleon than the scattering of real photons. In analyzing Compton scattering it is important to know how

much of the prediction is not a true test of a model, but fixed due to general principles. These model-independent predictions for virtual Compton scattering were the main topic of our discussion.

The interest in virtual Compton scattering has also been due to another aspect: When studying reactions on a nucleus, such as $(e, e'p)$, the nucleon interacting with the electromagnetic probe is necessarily off its mass shell. We have no model-independent information for the behavior of such a nucleon and any conclusion about genuine medium modifications must be based on firm theoretical ground how to deal with a single nucleon under these kinematical circumstances. In fact, such a discussion depends very much on what one chooses as an interpolating field for the intermediate, not observed nucleon. This clearly makes the “off-shell behavior” of the nucleon representation dependent and unobservable. We discussed how certain features of the off-shell electromagnetic vertex of the nucleon can be shifted into irreducible, reaction-specific terms for the reaction amplitude. Two-step reactions on a free nucleon, like (virtual) Compton scattering, allow us to test many aspects of dealing with an intermediate, off-shell nucleon under simpler circumstances, without complications from, e.g., exchange currents or final state interactions. Understanding these aspects on the single-nucleon level would seem a prerequisite before any exotic claims can be made for nuclear reactions.

We have studied the virtual Compton scattering first on the *operator* level. Using the requirement of gauge invariance, as expressed by the Ward-Takahashi identity, we derived constraints for the operator that determine terms up to and including linear in the four-momenta q and q' . Also, we showed that on the operator level terms involving terms depending only on q or only on q' are determined model independently in terms of on-shell properties of the nucleon.

To obtain these results, we used the method of Gell-Mann and Goldberger, by splitting the contributions into general pole terms (class A) and the one-particle irreducible two-photon contributions (class B). We calculated class A below pion-production threshold in the framework of a specific representation for the most general effective Lagrangian compatible with Lorentz invariance, gauge invariance and discrete symmetries. This approach was introduced in [4] as a method for writing the general structure of the Compton scattering amplitude in a way that allows one best to discuss its low-energy behavior. In this connection, we also showed the origin for a commonly used form of the electromagnetic vertex of the nucleon and stated the consistency conditions for its use.

After discussing the leading terms of the VCS operator, we considered the *matrix element* for $\gamma^*p \rightarrow \gamma p$ in the photon-nucleon c.m. frame. We found that the VCS amplitude up to and including terms linear in the initial and final photon three-momentum can be expressed in terms of information one can obtain from electron-proton scattering. This is the result analogous to the LET for the real Compton scattering amplitude. As we also showed, the next order—terms involving $|\vec{q}'|^2$, $|\vec{q}'||\vec{q}|$, and $|\vec{q}|^2$ —is also completely specified but now requires in addition also the electromagnetic polarizabilities $\bar{\alpha}$ and $\bar{\beta}$ encountered in *real* Compton scattering. In other words, new structure-dependent information can only appear at order 3 or higher in the three-

momenta. Our results concern the expansion in terms of powers of both the initial and final photon momentum. This allowed us to determine more terms than in [14], where only the leading terms in the final momentum were concerned. On the other hand, by expanding in both momenta, the range of applicability is smaller since *both* kinematical variables should be small.

We then considered different commonly used methods to include the on-shell information contained in the electromagnetic form factors in a “Born-term” calculation of the VCS matrix element. The fact that the “Born terms” calculated with F_1 and F_2 are gauge invariant is not trivial, since the vertex and free propagator do not satisfy the Ward-Takahashi identity. We explained why different on-shell equivalent forms for the electromagnetic vertex operator lead to the same *irregular* contribution in the VCS matrix element. We emphasized the importance of stating with respect to which pole terms the structure-dependent terms are defined.

Using only gauge invariance, Lorentz invariance, crossing symmetry, and the discrete symmetries, we were able to make statements about the low-energy behavior up to $O(2)$. Further conclusions can be reached by also taking into account the constraints imposed by chiral symmetry. This would most naturally be done in the framework of chiral perturbation theory. In particular, predictions for the higher-order terms could be obtained.

ACKNOWLEDGMENTS

The work of J.H.K. is part of the research program of the Foundation for Fundamental Research of Matter (FOM) and the National Organization for Scientific Research (NWO). The collaboration between NIKHEF and the Institute for Nuclear Physics at Mainz is supported in part by a grant from NATO. A.Yu.K. thanks the Theory Group of NIKHEF-K for the kind hospitality and the Netherlands Organization for International Cooperation in Higher Education (NUFFIC) for financial support. A.Yu.K. also would like to thank SFB 201 of the Deutsche Forschungsgemeinschaft for its hospitality and financial support during his stay at Mainz. S.S. would like to thank H. W. Fearing for useful discussions on the soft-photon approximation. S.S. was supported by Deutsche Forschungsgemeinschaft.

APPENDIX

In this appendix we outline the calculation of the transverse and longitudinal functions A_i of Eqs. (3.14) and (3.15), respectively. For that purpose we split M^μ into its contributions from classes A and B, $M^\mu = M_A^\mu + M_B^\mu$. If we introduce

$$F(-q') = \epsilon'^* \left(1 + \frac{\kappa}{2M} \not{q}' \right) \quad (\text{A1})$$

for the vertex involving the final real photon, the contribution of class A [see Eq. (3.4)] reads

$$M_A^\mu = \bar{u}(p_f, s_f) (F(-q') S_F(p_i + q) \Gamma_{\text{eff}}^\mu(q) + \Gamma_{\text{eff}}^\mu(q) S_F(p_i - q') F(-q')) u(p_i, s_i), \quad (\text{A2})$$

where $\Gamma_{\text{eff}}^\mu(q)$ is defined in Eq. (3.3). Applying the Dirac equation, M_A^μ can be written as

$$M_A^\mu = \bar{u}(p_f, s_f) \left[2 \left(\frac{p_f \cdot \epsilon'^*}{s - M^2} + \frac{p_i \cdot \epsilon'^*}{u - M^2} \right) \Gamma_{\text{eff}}^\mu(q) + (1 + \kappa) \times \left(\frac{\epsilon'^* \not{q}' \Gamma_{\text{eff}}^\mu(q)}{s - M^2} + \frac{\Gamma_{\text{eff}}^\mu(q) \epsilon'^* \not{q}'}{u - M^2} \right) - \frac{\kappa}{M} \left(\frac{p_f \cdot \epsilon'^* \not{q}' \Gamma_{\text{eff}}^\mu(q)}{s - M^2} - \frac{p_i \cdot \epsilon'^* \Gamma_{\text{eff}}^\mu(q) \not{q}'}{u - M^2} \right) + \frac{\kappa}{2M} [\epsilon'^* \Gamma_{\text{eff}}^\mu(q) + \Gamma_{\text{eff}}^\mu(q) \epsilon'^*] \right] u(p_i, s_i), \quad (\text{A3})$$

where $s = (p_f + q')^2$ and $u = (p_i - q')^2$. Similarly, using Eqs. (3.10) and (3.11), M_B^μ can be written as

$$M_B^\mu = \bar{u}(p_f, s_f) ((\epsilon'^* \not{q} \cdot q' - q' \not{\epsilon}'^* \cdot q) f_1(P^2) + (\epsilon'^* \not{P} \cdot q P \cdot q' + P^\mu \epsilon'^* \cdot P q \cdot q' - P^\mu \epsilon'^* \cdot q P \cdot q' - q' \not{\epsilon}'^* \cdot P P \cdot q) f_2(P^2) - i \epsilon^{\mu\rho\nu\alpha} \epsilon'_\nu q_\rho q'_\alpha \gamma_5 f_3(P^2) + O(3)) u(p_i, s_i), \quad (\text{A4})$$

where $O(3)$ denotes terms of order 3 in q or q' .

The following considerations will be carried out in the center-of-mass (c.m.) frame, where the four-momenta are given by $q^\mu = (q_0, 0, 0, |\vec{q}|)$, $p_i^\mu = (E_q, -\vec{q})$, $q'^\mu = (|\vec{q}'|, \vec{q}')$, and $p_f^\mu = (E_{q'}, -\vec{q}')$. From energy conservation, $q_0 + E_q = |\vec{q}'| + E_{q'}$, we infer that we may choose $|\vec{q}|$, $|\vec{q}'|$, and $z = \hat{q} \cdot \hat{q}'$ as a set of independent variables. In terms of the c.m. variables the denominators of Eq. (A3) are proportional to $|\vec{q}'|$,

$$s - M^2 = 2|\vec{q}'|(E_{q'} + |\vec{q}|), \quad u - M^2 = -2|\vec{q}'|(E_q + z|\vec{q}|), \quad (\text{A5})$$

and thus M_A^μ can be written as

$$M_A^\mu = \frac{a^\mu(q, q')}{|\vec{q}'|} + b^\mu(q, q'). \quad (\text{A6})$$

The functions $a^\mu(q, q')$ and $b^\mu(q, q')$ are regular with respect to $|\vec{q}'|$ and $|\vec{q}|$, and are given by

$$a^\mu(q, q') = K(\vec{q}) \bar{u}(-\vec{q}', s_f) \Gamma_{\text{eff}}^\mu(q) u(-\vec{q}, s_i), \quad (\text{A7})$$

$$b^\mu(q, q') = \bar{u}(-\vec{q}', s_f) \left((1 + \kappa) \left(\frac{\epsilon'^* \not{h}' \Gamma_{\text{eff}}^\mu(q)}{2(E_{q'} + |\vec{q}'|)} - \frac{\Gamma_{\text{eff}}^\mu(q) \epsilon'^* \not{h}'}{2(E_q + z|\vec{q}|)} \right) - \kappa \frac{\vec{q} \cdot \vec{\epsilon}'^*}{M} \frac{\Gamma_{\text{eff}}^\mu(q) \not{h}'}{2(E_q + z|\vec{q}|)} + \frac{\kappa}{2M} [\epsilon'^* \Gamma_{\text{eff}}^\mu(q) + \Gamma_{\text{eff}}^\mu(q) \epsilon'^*] \right) u(-\vec{q}, s_i), \quad (\text{A8})$$

TABLE IV. Reduction of the coefficients $\vec{a}_{ij}$ of Eqs. (A17)–(A20) to Pauli space. For the definition of the corresponding Pauli spin operators see Eqs. (3.14) and (3.15). For further information see caption of Table I.

	$\vec{a}_{00}(Q)/ \vec{q}' $	$\vec{a}_{11} \vec{q} $	$\vec{a}_{21} \vec{q}' \vec{q} $	$\vec{a}_{12} \vec{q} ^2$
A_1	0	$\frac{z}{2M^2} \vec{q} $	$-\frac{z\kappa}{4M^3} \vec{q}' \vec{q} $	$-\frac{z^2}{2M^3} \vec{q} ^2$
A_2	0	0	0	0
A_3	0	$-\frac{1}{2M^2} \vec{q} $	$\frac{\kappa}{4M^3} \vec{q}' \vec{q} $	$\frac{z}{2M^3} \vec{q} ^2$
A_4	0	0	0	0
A_5	$-\frac{N_i G_M(Q^2)}{(E_i+z \vec{q})(E_i+M)}\frac{ \vec{q} ^2}{ \vec{q}' }$	0	0	$-\frac{\kappa}{4M^3} \vec{q} ^2$
A_6	0	0	0	0
A_7	0	$\frac{1}{2M^2} \vec{q} $	$-\frac{\kappa}{4M^3} \vec{q}' \vec{q} $	$-\frac{z}{2M^3} \vec{q} ^2$
A_8	0	0	0	0
A_9	$\frac{N_i G_E(Q^2)}{(E_i+z \vec{q})(E_i+M)}\frac{ \vec{q} ^2}{ \vec{q}' }$	$\frac{z}{2M^2} \vec{q} $	$-\frac{z\kappa}{4M^3} \vec{q}' \vec{q} $	$-\frac{z^2}{2M^3} \vec{q} ^2 + \frac{r_E^2}{6M} \vec{q} ^2$
A_{10}	0	$\frac{1}{2M^2} \vec{q} $	$-\frac{\kappa}{4M^3} \vec{q}' \vec{q} $	$-\frac{z}{2M^3} \vec{q} ^2$
A_{11}	0	0	0	0
A_{12}	0	0	0	0

where we introduced $n'^\mu = (1, \hat{q}')$, and

$$K(\vec{q}) = -\frac{\vec{q} \cdot \vec{\epsilon}'^*}{E_q + z|\vec{q}|}. \quad (\text{A9})$$

In order to obtain Eq. (A9) we explicitly made use of the Coulomb gauge for the final photon, namely, when using $p_f \cdot \epsilon'^* = 0$ and $p_i \cdot \epsilon'^* = \vec{q} \cdot \vec{\epsilon}'^*$.

We will now derive from Eqs. (A6)–(A9) the expansion of M_A^μ in powers of $|\vec{q}|$ and $|\vec{q}'|$. According to Eq. (3.13) it is sufficient to treat the space components of M^μ . Since the structure coefficients appear at $O(qq')$ and higher in the operator, we expand the contributions up to and including $|\vec{q}'|^2$, $|\vec{q}'||\vec{q}|$, and $|\vec{q}|^2$. When expanding the electromagnetic vertex we make use of the following relations:

$$\begin{aligned} q^\mu &= Q^\mu + |\vec{q}'|g^{\mu 0} + \dots, \\ \not{q} &= \not{Q} + |\vec{q}'|\gamma_0 + \dots, \\ q^2 &= Q^2 + 2Q_0|\vec{q}'| + \dots, \end{aligned} \quad (\text{A10})$$

where $Q^\mu = q^\mu|_{\vec{q}'=0} = (M - E_i, \vec{q})$. Furthermore, for $\vec{a}$ we need the expansion of the form factors around Q^2 :

$$F_{1,2}(q^2) = F_{1,2}(Q^2) + 2Q_0|\vec{q}'|F'_{1,2}(Q^2) + \dots, \quad (\text{A11})$$

with the notation $F'(x) = dF/dx$. Using the definition of $\Gamma_{\text{eff}}^\mu(q)$ of Eq. (3.3), it is straightforward to obtain

$$\vec{\Gamma}_{\text{eff}}(q) = \vec{\Gamma}_{00} + \vec{\Gamma}_{10}|\vec{q}'| + \vec{\Gamma}_{01}|\vec{q}| + \dots, \quad (\text{A12})$$

where

$$\vec{\Gamma}_{00} = \vec{\gamma}, \quad \vec{\Gamma}_{10} = -\frac{\kappa}{4M}[\vec{\gamma}, \gamma_0], \quad \vec{\Gamma}_{01} = \frac{\kappa}{4M}[\vec{\gamma}, \vec{\gamma} \cdot \hat{q}],$$

$$\vec{\Gamma}_{20} = F'_1(0)\vec{\gamma} - \frac{\kappa}{8M^2}[\vec{\gamma}, \gamma_0], \quad \vec{\Gamma}_{11} = -\hat{q}\gamma_0 F'_1(0),$$

$$\vec{\Gamma}_{02} = (-\vec{\gamma} + \vec{\gamma} \cdot \hat{q}\hat{q})F'_1(0) + \frac{\kappa}{8M^2}[\vec{\gamma}, \gamma_0], \dots \quad (\text{A13})$$

Since the dependence on the momenta $\vec{q}$ and $\vec{q}'$ is also contained in the initial and final nucleon spinors, respectively, we expand them to the required order:

$$\begin{aligned} u(-\vec{q}) &= \left(1 + \frac{\vec{\gamma} \cdot \vec{q}}{2M} + \frac{|\vec{q}|^2}{8M^2} + \dots\right) u(0), \\ \bar{u}(-\vec{q}') &= \bar{u}(0) \left(1 + \frac{\vec{\gamma} \cdot \vec{q}'}{2M} + \frac{|\vec{q}'|^2}{8M^2} + \dots\right), \end{aligned} \quad (\text{A14})$$

where from now on we suppress the spin indices. Finally, the expansion of the energy denominators reads

$$\frac{1}{E_{q'} + |\vec{q}'|} = \frac{1}{M} - \frac{|\vec{q}'|}{M^2} + \frac{|\vec{q}'|^2}{2M^3} + \dots,$$

$$\frac{1}{E_q + z|\vec{q}|} = \frac{1}{M} - \frac{z|\vec{q}|}{M^2} + \frac{(2z^2 - 1)|\vec{q}|^2}{2M^3} + \dots \quad (\text{A15})$$

TABLE V. Reduction of the coefficients $\vec{b}_{ij}$ of Eqs. (A23)–(A28) to Pauli space. See captions of Tables I and IV.

	$\vec{b}_{00}$	$\vec{b}_{10} \vec{q}' $	$\vec{b}_{01} \vec{q} $	$\vec{b}_{20} \vec{q}' ^2$	$\vec{b}_{11} \vec{q}' \vec{q} $	$\vec{b}_{02} \vec{q} ^2$
A_1	$-\frac{1}{M}$	0	$\frac{z}{2M^2} \vec{q} $	$-\left(\frac{1}{8M^3} + \frac{r_E^2}{6M} - \frac{\kappa}{4M^3}\right) \vec{q}' ^2$	$\frac{z\kappa}{4M^3} \vec{q}' \vec{q} $	$\left(\frac{1}{8M^3} + \frac{r_E^2}{6M} - \frac{z^2}{2M^3} + \frac{\kappa(1+\kappa)}{4M^3}\right) \vec{q} ^2$
A_2	0	$\frac{1+2\kappa}{2M^2} \vec{q}' $	0	$-\frac{\kappa^2}{4M^3} \vec{q}' ^2$	$\frac{z\kappa}{2M^3} \vec{q}' \vec{q} $	$-\frac{(1+\kappa)^2}{4M^3} \vec{q} ^2$
A_3	0	0	$-\frac{1}{2M^2} \vec{q} $	0	$\frac{1-\kappa}{4M^3} \vec{q}' \vec{q} $	$\frac{(1-2\kappa-\kappa^2)z}{4M^3} \vec{q} ^2$
A_4	0	0	$-\frac{(1+\kappa)^2}{2M^2} \vec{q} $	0	$-\frac{(2+\kappa)\kappa}{4M^3} \vec{q}' \vec{q} $	$\frac{(1+\kappa)^2z}{4M^3} \vec{q} ^2$
A_5	0	0	0	0	0	$\frac{(2+\kappa)\kappa}{4M^3} \vec{q} ^2$
A_6	0	$\frac{1+\kappa}{2M^2} \vec{q}' $	0	$-\frac{(1+\kappa)\kappa}{4M^3} \vec{q}' ^2$	$-\frac{(1+\kappa)z}{2M^3} \vec{q}' \vec{q} $	0
A_7	0	0	$-\frac{1}{2M^2} \vec{q} $	0	$-\frac{1+2\kappa}{4M^3} \vec{q}' \vec{q} $	$\frac{z}{2M^3} \vec{q} ^2$
A_8	0	0	0	0	$\frac{1+3\kappa}{4M^3} \vec{q}' \vec{q} $	0
A_9	$-\frac{1}{M}$	0	$\frac{z}{2M^2} \vec{q} $	$-\left(\frac{1}{8M^3} + \frac{r_E^2}{6M} - \frac{\kappa}{4M^3}\right) \vec{q}' ^2$	$\frac{z\kappa}{4M^3} \vec{q}' \vec{q} $	$\left(\frac{1}{8M^3} - \frac{z^2}{2M^3}\right) \vec{q} ^2$
A_{10}	0	0	$-\frac{1}{2M^2} \vec{q} $	0	$-\frac{1+2\kappa}{4M^3} \vec{q}' \vec{q} $	$\frac{z}{2M^3} \vec{q} ^2$
A_{11}	0	$-\frac{1+2\kappa}{2M^2} \vec{q}' $	0	$\frac{\kappa^2}{4M^3} \vec{q}' ^2$	$\frac{(1+\kappa)z}{4M^3} \vec{q}' \vec{q} $	$\frac{1+2\kappa}{4M^3} \vec{q} ^2$
A_{12}	0	$\frac{(1+\kappa)z}{2M^2} \vec{q}' $	0	$-\frac{(1+\kappa)\kappa z}{4M^3} \vec{q}' ^2$	$-\frac{(1+\kappa)(2z^2-1)}{4M^3} \vec{q}' \vec{q} $	$-\frac{(1+\kappa)z}{4M^3} \vec{q} ^2$

Let us first consider $\vec{a}(q, q')$ which we expand according to

$$\begin{aligned} \vec{a}(q, q') = & \vec{a}_{00}(Q) + |\vec{q}'|\vec{a}_{10} + |\vec{q}'|^2\vec{a}_{20} + |\vec{q}'||\vec{q}|\vec{a}_{11} \\ & + |\vec{q}'|^3\vec{a}_{30} + |\vec{q}'|^2|\vec{q}|\vec{a}_{21} + |\vec{q}'||\vec{q}|^2\vec{a}_{12} + \dots \end{aligned} \quad (\text{A16})$$

Using the relations of Eqs. (A10)–(A15) we obtain

$$\vec{a}_{00}(Q) = K(\vec{q})\vec{u}(0)\vec{\Gamma}_{\text{eff}}(Q)u(-\vec{q}), \quad (\text{A17})$$

$$\vec{a}_{11} = -\frac{\hat{q} \cdot \vec{\epsilon}'^*}{M}\vec{u}(0)\left(\frac{\vec{\gamma} \cdot \hat{q}'}{2M}\vec{\Gamma}_{00} + \vec{\Gamma}_{10}\right)u(0), \quad (\text{A18})$$

$$\vec{a}_{21} = -\frac{\hat{q} \cdot \vec{\epsilon}'^*}{M}\vec{u}(0)\left(\frac{1}{8M^2}\vec{\Gamma}_{00} + \frac{\vec{\gamma} \cdot \hat{q}'}{2M}\vec{\Gamma}_{10} + \vec{\Gamma}_{20}\right)u(0), \quad (\text{A19})$$

$$\vec{a}_{12} = -\frac{\hat{q} \cdot \vec{\epsilon}'^*}{M}\vec{u}(0)\left[-\frac{z}{M}\left(\frac{\vec{\gamma} \cdot \hat{q}'}{2M}\vec{\Gamma}_{00} + \vec{\Gamma}_{10}\right) + \frac{\vec{\gamma} \cdot \hat{q}'}{2M}\vec{\Gamma}_{01}\right] \quad (\text{A20})$$

$$+ \vec{\Gamma}_{11} + \left(\frac{\vec{\gamma} \cdot \hat{q}'}{2M}\vec{\Gamma}_{00} + \vec{\Gamma}_{10}\right)\frac{\vec{\gamma} \cdot \hat{q}}{2M}u(0),$$

$$\vec{a}_{j0} = 0, \quad j = 1, 2, \dots \quad (\text{A21})$$

The last equation follows from $K(0) = 0$. The function $\vec{a}_{00}(Q)$ is completely determined in terms of the electromagnetic form factors $F_{1,2}(Q^2)$ [or $G_{E,M}(Q^2)$]. Note that in Eq. (A17) we keep all powers in $|\vec{q}|$, since it will be multiplied with the $1/|\vec{q}'|$ singularity and there are no other terms which can generate such a singularity.

In order to determine $\vec{b}(q, q')$, we first expand $\vec{b}(q, q')$ in an analogous fashion to Eq. (A16)

$$\begin{aligned} \vec{b}(q, q') = & \vec{b}_{00} + |\vec{q}'|\vec{b}_{10} + |\vec{q}|\vec{b}_{01} + |\vec{q}'|^2\vec{b}_{20} \\ & + |\vec{q}'||\vec{q}|\vec{b}_{11} + |\vec{q}|^2\vec{b}_{02} + \dots \end{aligned} \quad (\text{A22})$$

Using the building blocks of Eqs. (A10)–(A15) we find for the coefficients $\vec{b}_{ij}$:

$$\vec{b}_{00} = \frac{1}{2M} \bar{u}(0) ((1 + \kappa) [\vec{\epsilon}'^* \vec{h}', \vec{\Gamma}_{00}] + \kappa \{ \vec{\epsilon}'^*, \vec{\Gamma}_{00} \}) u(0) = \frac{1}{2M} \bar{u}(0) X u(0), \quad (\text{A23})$$

$$\vec{b}_{10} = \frac{1}{2M} \bar{u}(0) \left[(1 + \kappa) \left([\vec{\epsilon}'^* \vec{h}', \vec{\Gamma}_{10}] - \frac{1}{M} \vec{\epsilon}'^* \vec{h}' \vec{\Gamma}_{00} \right) + \kappa \{ \vec{\epsilon}'^*, \vec{\Gamma}_{10} \} + \frac{\vec{\gamma} \cdot \hat{q}'}{2M} X \right] u(0) = \frac{1}{2M} \bar{u}(0) \left(Y + \frac{\vec{\gamma} \cdot \hat{q}'}{2M} X \right) u(0), \quad (\text{A24})$$

$$\begin{aligned} \vec{b}_{01} &= \frac{1}{2M} \bar{u}(0) \left[(1 + \kappa) \left([\vec{\epsilon}'^* \vec{h}', \vec{\Gamma}_{01}] + \frac{z}{M} \vec{\Gamma}_{00} \vec{\epsilon}'^* \vec{h}' \right) - \frac{\kappa}{M} \vec{\epsilon}'^* \cdot \hat{q} \vec{\Gamma}_{00} \vec{h}' + \kappa \{ \vec{\epsilon}'^*, \vec{\Gamma}_{01} \} + X \frac{\vec{\gamma} \cdot \hat{q}}{2M} \right] u(0) \\ &= \frac{1}{2M} \bar{u}(0) \left(Z + X \frac{\vec{\gamma} \cdot \hat{q}}{2M} \right) u(0), \end{aligned} \quad (\text{A25})$$

$$\vec{b}_{20} = \frac{1}{2M} \bar{u}(0) \left[(1 + \kappa) \left([\vec{\epsilon}'^* \vec{h}', \vec{\Gamma}_{20}] - \frac{1}{M} \vec{\epsilon}'^* \vec{h}' \vec{\Gamma}_{10} + \frac{1}{2M^2} \vec{\epsilon}'^* \vec{h}' \vec{\Gamma}_{00} \right) + \kappa \{ \vec{\epsilon}'^*, \vec{\Gamma}_{20} \} + \frac{\vec{\gamma} \cdot \hat{q}'}{2M} Y + \frac{1}{8M^2} X \right] u(0), \quad (\text{A26})$$

$$\begin{aligned} \vec{b}_{11} &= \frac{1}{2M} \bar{u}(0) \left[(1 + \kappa) \left([\vec{\epsilon}'^* \vec{h}', \vec{\Gamma}_{11}] - \frac{1}{M} \vec{\epsilon}'^* \vec{h}' \vec{\Gamma}_{01} + \frac{z}{M} \vec{\Gamma}_{10} \vec{\epsilon}'^* \vec{h}' \right) - \frac{\kappa}{M} \vec{\epsilon}'^* \cdot \hat{q} \vec{\Gamma}_{10} \vec{h}' + \kappa \{ \vec{\epsilon}'^*, \vec{\Gamma}_{11} \} \right. \\ &\quad \left. + Y \frac{\vec{\gamma} \cdot \hat{q}}{2M} + \frac{\vec{\gamma} \cdot \hat{q}'}{2M} Z + \frac{\vec{\gamma} \cdot \hat{q}'}{2M} X \frac{\vec{\gamma} \cdot \hat{q}}{2M} \right] u(0), \end{aligned} \quad (\text{A27})$$

$$\begin{aligned} \vec{b}_{02} &= \frac{1}{2M} \bar{u}(0) \left[(1 + \kappa) \left([\vec{\epsilon}'^* \vec{h}', \vec{\Gamma}_{02}] + \frac{z}{M} \vec{\Gamma}_{01} \vec{\epsilon}'^* \vec{h}' + \frac{1 - 2z^2}{2M^2} \vec{\Gamma}_{00} \vec{\epsilon}'^* \vec{h}' \right) - \frac{\kappa}{M} \vec{\epsilon}'^* \cdot \hat{q} \left(\vec{\Gamma}_{01} - \frac{z}{M} \vec{\Gamma}_{00} \right) \vec{h}' \right. \\ &\quad \left. + \kappa \{ \vec{\epsilon}'^*, \vec{\Gamma}_{02} \} + Z \frac{\vec{\gamma} \cdot \hat{q}}{2M} + \frac{1}{8M^2} X \right] u(0). \end{aligned} \quad (\text{A28})$$

The reduction of the above expression to Pauli space is straightforward but very tedious. The results are displayed in Tables IV and V.

Finally, it is straightforward to obtain the expansion of $\vec{M}_B$:

$$\vec{M}_B = \bar{u}(0) \left(|\vec{q}'|^2 \vec{\epsilon}'^* \frac{\vec{\alpha}}{e^2} + |\vec{q}'| |\vec{q}| (z \vec{\epsilon}'^* - \hat{q} \cdot \vec{\epsilon}'^* \hat{q}') \frac{\vec{\beta}}{e^2} + O(3) \right) u(0), \quad (\text{A29})$$

where we have defined the (real) Compton polarizabilities as

$$\vec{\alpha} = e^2 [f_1(4M^2) + 4M^2 f_2(4M^2)], \quad \vec{\beta} = -e^2 f_1(4M^2). \quad (\text{A30})$$

Due to the presence of the γ_5 matrix, the third function $f_3(4M^2)$ only contributes at $O(3)$ at the level of the matrix element.

-
<p>[1] F. E. Low, Phys. Rev. 96, 1428 (1954).
 [2] M. Gell-Mann and M. L. Goldberger, Phys. Rev. 96, 1433 (1954).
 [3] E. Kazes, Nuovo Cimento XIII, 1226 (1959).
 [4] A. I. L'vov, Int. J. Mod. Phys. A 8, 5267 (1993).
 [5] C. Audit <i>et al.</i>, CEBAF Report No. PR 93-050, 1993.
 [6] J. F. J. van den Brand <i>et al.</i>, CEBAF Report No. PR 94-011, 1994.
 [7] N. d'Hose, private communication.
 [8] P. Roman, <i>Introduction to Quantum Field Theory</i> (Wiley, New York, 1969), Chap. 10.
 [9] F. E. Low, Phys. Rev. 110, 974 (1958).
 [10] See, e.g., S. L. Adler and R. F. Dashen, <i>Current Algebras and</i></p> | <p><i>Applications to Particle Physics</i> (Benjamin, New York, 1968), Chap. 3.
 [11] R. A. Berg and C. N. Lindner, Nucl. Phys. 26, 259 (1961).
 [12] M. Perrottet, Lett. Nuovo Cimento 7, 915 (1973).
 [13] M. Schäfer, H. C. Dönges, and U. Mosel, Phys. Lett. B 342, 13 (1995).
 [14] P. A. M. Guichon, G. Q. Liu, and A. W. Thomas, Nucl. Phys. A591, 606 (1995).
 [15] M. Vanderhaeghen, Phys. Lett. B 368, 13 (1996).
 [16] J. C. Ward, Phys. Rev. 78, 182 (1950).
 [17] Y. Takahashi, Nuovo Cimento 6, 371 (1957).
 [18] See, e.g., C. Itzykson and J.-B. Zuber, <i>Quantum Field Theory</i> (McGraw-Hill, New York, 1985), Chaps. 6-1-4 and 11-3-1.</p> |
|---|--|

- [19] W. Melnitchouk, A. W. Schreiber, and A. W. Thomas, Phys. Rev. D **49**, 1183 (1994).
- [20] S. Scherer, J. H. Koch, and J. L. Friar, Nucl. Phys. **A552**, 515 (1993).
- [21] H. W. Fearing and S. Scherer (unpublished).
- [22] S. Scherer and H. W. Fearing, Phys. Rev. C **51**, 359 (1995).
- [23] S. Scherer and H. W. Fearing, Phys. Rev. D **52**, 6445 (1995).
- [24] J. S. R. Chisholm, Nucl. Phys. **26**, 469 (1961).
- [25] S. Kamefuchi, L. O’Raifeartaigh, and A. Salam, Nucl. Phys. **28**, 529 (1961).
- [26] C. Arzt, Phys. Lett. B **342**, 189 (1995).
- [27] F. Gross and D. O. Riska, Phys. Rev. C **36**, 1928 (1987).
- [28] J. D. Bjorken and S. D. Drell, *Relativistic Quantum Mechanics* (McGraw-Hill, New York, 1964).
- [29] See, e.g., E. Amaldi, S. Fubini, and G. Furlan, *Pion-Electroproduction* (Springer, Berlin, 1979), App. B.
- [30] S. Scherer, G. I. Poulis, and H. W. Fearing, Nucl. Phys. **A570**, 686 (1994).
- [31] G. G. Simon, Ch. Schmitt, F. Borkowski, and V. H. Walther, Nucl. Phys. **A333**, 381 (1980).
- [32] H. Leutwyler, Ann. Phys. (N.Y.) **235**, 165 (1994).
- [33] E. D’Hoker and S. Weinberg, Phys. Rev. D **50**, R6050 (1994).
- [34] S. Weinberg, Physica A **96**, 327 (1979).
- [35] J. Gasser and H. Leutwyler, Ann. Phys. (N.Y.) **158**, 142 (1984); Nucl. Phys. **B250**, 465 (1985).
- [36] J. Gasser, M. E. Sainio, and A. Svarc, Nucl. Phys. **B307**, 779 (1988).
- [37] S. Weinberg, Phys. Lett. B **251**, 288 (1990); Nucl. Phys. **B363**, 3 (1991).
- [38] A. M. Bincer, Phys. Rev. **118**, 855 (1960).
- [39] H. W. L. Naus and J. H. Koch, Phys. Rev. C **36**, 2459 (1987).
- [40] K. J. Barnes, Phys. Lett. **1**, 166 (1962).
- [41] R. Ferrari and M. Rosa-Clot, Nuovo Cimento A **56**, 582 (1968).
- [42] J. S. Bell and R. van Royen, Nuovo Cimento A **60**, 62 (1969).
- [43] H. W. Fearing, Phys. Rev. D **7**, 243 (1973).