



UvA-DARE (Digital Academic Repository)

Constipation in (early) infancy and childhood : pathogenesis and diagnostic procedures

de Lorijn, F.

[Link to publication](#)

Citation for published version (APA):

de Lorijn, F. (2005). Constipation in (early) infancy and childhood : pathogenesis and diagnostic procedures. Amsterdam.

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Chapter 9

Prognosis of constipation;
clinical factors and colonic transit time

Fleur de Lorijn, Michiel P. van Wijk, Johannes B. Reitsma,
Rijk van Ginkel, Jan A. Taminiau, Marc A. Benninga

Archives of Disease in Childhood 2004;89:723-727

Abstract

Background

Measurement of colonic transit time (CTT) is sometimes used in the evaluation of patients with chronic constipation.

Aim

To investigate the relation between symptoms and CTT, and to assess the importance of symptoms and CTT in predicting outcome.

Methods

Between 1995 and 2000, 169 consecutive patients (median age 8.4 years, 65% boys) fulfilling the criteria for constipation were enrolled. During the intervention and follow up period, all kept a diary to record symptoms. CTT was measured at entry to the study.

Results

At entry, defecation frequency was lower in girls than in boys, while the frequency of encopresis episodes was higher in boys. CTT values were significantly higher in those with a low defecation frequency (≤ 1 /week) and a high frequency of encopresis (≥ 2 /day). However, 50% had CTT values within the normal range. Successful outcome occurred more often in those with a rectal impaction. CTT results < 100 hours were not predictive of outcome. However, those with CTT > 100 hours were less likely to have had a successful outcome.

Conclusion

The presence of a rectal impaction at presentation is associated with a better outcome at one year. A CTT > 100 hours is associated with a poor outcome at one year.

Introduction

Constipation is a common problem in children, accounting for about 3% of consultations in an average paediatric practice and as much as 25% in a paediatric gastroenterology clinic¹⁻³. No specific organic cause can be found in approximately 90% of the children^{4,5}. The diagnosis is mainly based on clinical history and physical examination. Patients and/or their parents refer to the number of stools per week, to stool volume, to difficulty in defecation, and/or to sensation of abdominal fullness⁴. Several tests have been developed to objectify these complaints⁶⁻¹⁰.

A plain abdominal radiograph is frequently used to confirm the presence of retained stool or enlargement of the colon or rectum^{7,8}. However, the value of abdominal radiograph examination in this setting is a matter of debate¹¹. Assessment of total and segmental colonic transit time (CTT) using radio-opaque markers, is a non-invasive method which provides information about colorectal motor function^{6,9,10}. Furthermore, this technique has been used to localise a delayed transit in the colon and to evaluate the response to treatment^{9,11-13}. Some investigators have shown a good relation between symptoms of constipation and CTT in adults^{14,15}. However, others report a significant difference between previous reported symptoms and the CTT results^{9,16}. No previous studies have investigated the prognostic value of CTT measurement.

Our study had two main objectives. First, to investigate the relation between symptoms and CTT. Second, to evaluate the possible relation between symptoms and CTT, and the outcome after one year of follow up.

Patients and methods

Patients

All patients referred to our paediatric gastrointestinal outpatient clinic with constipation between 1995 and 2000 were eligible. They were referred by family practitioners, paediatricians, psychiatrists, and school doctors. At entry, patients had at least two of the following: (1) defecation <3/week; (2) encopresis episodes >1/week; (3) passing of very large stool every 7–30 days; and (4) a palpable abdominal or rectal faecal mass¹⁷. All were at least 5 years of age. Children with Hirschsprung's disease, spinal and anal anomalies, previous colon surgery, metabolic or renal abnormalities, mental retardation, or using drugs other than laxatives were excluded. All patients and parents gave written informed consent. The medical ethics committee of the hospital approved the protocol.

Colonic transit time

At entry, all children underwent a CTT study using the method described by Metcalf and colleagues¹⁰. Treatment with oral or rectal laxatives was discontinued for at least four days before the test; during this period they took one sachet of fibre (Volcolon, 6 g) each day. They then ingested a capsule with containing 20 radio-opaque markers on three consecutive mornings. Abdominal radiograph examinations were performed on days 4 and 7 in the morning. Additional abdominal radiograph examinations were performed on days 10, 13, and 16 if more than 20% of the markers remained on the previous film. Radiograph localisation of markers was based on the identification of bony landmarks and gaseous outlines as described by Arhan and colleagues⁶. Markers were counted in the right, left, and rectosigmoid regions, and mean segmental transit times were calculated according to a previously described formula^{6,10}. The normal ranges for total and segmental transit times were based on the upper limits (mean + 2 SD) from a study in healthy children⁶. Based on this study, a CTT of more than 62 hours was considered delayed. The upper limits of the normal range for right colon, left colon, and rectosigmoid transit time were 18, 20, and 34 hours, respectively⁶.

Medical history and physical examination

One week prior to entry and throughout the study the child and parents kept a diary in which defecation frequency, encopresis frequency, consistency and size of stool,

and pain during defecation were recorded. Associated symptoms such as abdominal pain, appetite suppression, absence of urge to defecate, and enuresis were also noted.

At entry, abdominal and rectal examinations were performed. Abdominal distension and palpable abdominal faecal masses were noted. Anal tone and the presence of a faecal impaction were recorded.

Treatment and follow up

The treatment regimens consisted of a high fibre diet, toilet advice (attempt defecation on the toilet for five minutes after meals), laxatives, and biofeedback training or anorectal manometry¹⁷. Oral laxative therapy consisted of an initial dosage of 6 g lactulose per day, increased stepwise to a maximum of 24 g if necessary. Enemas were given if severe rectal faecal impaction was present. A reward system was also employed.

The treatment intervention lasted for eight weeks. A detailed medical history was obtained during the intervention period, and at 6 and 12 months after entry. When necessary, children were also seen at other times at the outpatient clinic.

Definition of success

Treatment was considered to be "successful" if patients had three or more bowel movements weekly and less than one encopresis episode in a two week period, having discontinued laxatives for at least one month.

Statistical analysis

Baseline characteristics of the cohort were analysed in a descriptive way. Median values and 25th and 75th or 10th and 90th centiles were used if the distribution of continuous variables was skewed. Non-parametric (Mann-Whitney U and Kruskal-Wallis) and chi-square statistics were used to test for differences between groups. Logistic regression models were used to examine possible factors associated with success at one year. The following factors were considered: defecation frequency (in three groups: ≤ 1 /week, 1–3/week, and ≥ 3 /week), encopresis frequency (in four groups: no encopresis, < 1 /day, 1–2/day, ≥ 2 /day), the presence of night-time encopresis, the production of large stools, the presence of an abdominal or rectal mass, and CTT > 100 hours. Odds ratios with 95% confidence intervals were used to express the strength of the associations.

Results

Baseline characteristics

A total of 169 consecutive children (65% boys) were enrolled. Table 1 presents their baseline characteristics. The median age at intake was 8.4 years. The median defecation frequency at intake was lower in girls than in boys (1.0 versus 2.0 times per week; $p = 0.03$), while an encopresis frequency of more than twice weekly was reported more often in boys (94% versus 73%; $p = 0.0002$). Of the total group, 65% reported the passage of large stools every 7–30 days. Night-time encopresis occurred in 37%. A rectal impaction was present in 32%.

Table 2 shows the total and segmental colonic transit times at entry to the study. Approximately 50% had a total CTT in the normal range. No significant differences were found between boys and girls in the CTT or rectosigmoid transit time (RSTT). In 86% of those with an abnormal CTT, the RSTT was also prolonged. In children with a normal CTT, a delay in RSTT was present in only 14%. There was a significant correlation between CTT and RSTT (Pearson correlation coefficient 0.88, $p = 0.0002$). In the total group, a CTT >100 hours was found in 22%. Of these, 92% had a delayed RSTT and 39% had a delay in all segments.

Table 1 Baseline characteristics and clinical findings.

	Boys n=109	Girls n=60	Total n = 169	Missing
Age - yrs				
Median	8.5	8.0	8.4	
Percentiles 25 - 75	7.0 - 10.5	6.9 - 10.6	7.0 - 10.5	
Age of onset of symptoms - yrs				
Median	4.0	3.0	3.5	
Percentiles 25 - 75	1.0 - 4.0	0.0 - 4.0	1.0 - 4.0	
Defecation frequency / week				
Median	2.0	1.0 *	2.0	
Percentiles 25 - 75	1.0 - 3.0	1.0 - 2.0	1.0 - 2.0	
<3 times / week	74%	88% **	79%	
Encopresis frequency / week (daytime)				
Median	10.5	8.5	10.0	
Percentiles 25 - 75	7.0 - 21.0	1.0 - 21.0	5.5 - 21.0	
≥ 2 times per week	94%	73% **	86%	
no encopresis	6%	20% **	11%	
Nighttime Encopresis	38%	37%	37%	
Abdominal pain	57%	52%	56%	2
Pain during defecation	45%	47%	46%	2
No rectal sensation	22%	16%	20%	2
Large stools (every 7-30 days)	63%	68%	65%	
Palpable abdominal mass	26%	18%	23%	
Palpable rectal mass	32%	27%	30%	

* $p < 0.05$, Mann-Whitney U Test, ** $p < 0.05$, Chi-square

Table 2: Total and segmental transit times.

Transit time - hrs	Boys (n=109)	Girls (n=60)	Total group (n=169)
Total colon			
Median	60	53	58
Percentiles 25 - 75	38 - 103	37 - 74	37 - 92
Delayed > 62 hrs	49%	43%	47%
Ascending colon			
Median	10	11	10
Percentiles 25 - 75	5-16	5-15	5-16
Delayed > 18 hrs	23%	18%	21%
Descending colon			
Median	11	8	10
Percentiles 25 - 75	4-18	5-18	5-18
Delayed >20 hrs	21%	23%	22%
Rectosigmoid			
Median	37	31	32
Percentiles 25 - 75	19 - 68	17 - 47	18 - 63
Delayed > 34 hrs	53%	38%	48%

Correlation between CTT and symptom severity

At the first visit, children with a very low defecation frequency (≤ 1 /week) showed significantly prolonged median CTT (74 hours) compared to children with a defecation frequency of 1-3/week (50 hours) and a defecation frequency ≥ 3 /week (49 hours) ($p = 0.001$). In addition, in patients with 2 episodes of encopresis per day, median CTT values were significantly delayed (70 hours) compared to the group with 1-2 episodes of encopresis per day (50 hours), the group with < 1 episode of encopresis per day (52 hours), and those without encopresis (49 hours) ($p = 0.003$) (table 3). Furthermore, a significantly higher CTT was present in children with night-time encopresis and in those with a rectal mass (respectively $p < 0.0001$ and $p < 0.0001$) (table 3). As table 3 shows, similar associations were found between clinical symptoms and rectosigmoid transit time.

Clinical outcome and prognostic value

After one year 50% of the boys and 72% of the girls had a successful outcome (relative risk (RR) boys versus girls 0.70; 95% CI 0.55 to 0.90). These percentages increased to 61% in boys and to 80% in girls (RR 0.77; 95% CI 0.63 to 0.93) when patients were included who had a normal defecation frequency without encopresis but were still using laxatives.

The independent effect of various prognostic factors was analysed in a multivariate model predicting the probability of a successful outcome after one year. Boys were

less likely to be successfully treated at one year (OR 0.34; CI 0.16 to 0.70). The presence of a mass on rectal examination was a positive predictive sign for success (OR 3.39; CI 1.30 to 8.83). In those with a CTT <100 hours at entry, the transit time did not predict outcome at one year. However, those with a CTT >100 hours were less likely to have a successful outcome (OR 0.31; CI 0.12 to 0.85). Other possible prognostic factors such as a low defecation frequency, the presence of encopresis, and large stools at entry were not associated with outcome (table 4).

Table 3 Correlation between clinical parameters and transit time (hours).

	Number of patients	CTT (median)	p-value	RSTT (median)	p-value
Defecation frequency					
0-1/week	79	74	p = 0.001*	38	p = 0.009*
>1-3/week	55	50		30	
≥ 3/week	35	49		28	
Encopresis frequency (day and night)					
No encopresis	18	49	p = 0.003*	24	p = 0.03*
<1/day	24	52		31	
1-2/day	48	50		30	
≥ 2/day	79	70		38	
Nighttime encopresis					
Not present	106	47	p < 0.0001**	28	P < 0.0001**
Present	63	74		46	
Rectal mass					
Not present	118	48	p < 0.0001**	28	p < 0.0001**
Present	51	86		64	

*Kruskal-Wallis Test, **Mann-Whitney U Test

Table 4 Prognostic factors for cured at 1 year.

	Multivariate analysis Odds Ratio (95% CI)
Male	0.34 (0.16-0.70)
Defecation frequency	
≥ 3 / week	1.00 (reference)
>1-3/week	1.48 (0.54-4.08)
0-1/week	1.06 (0.40-2.80)
Encopresis frequency	
No encopresis	1.00 (reference)
< 1 / day	0.58 (0.14-2.43)
1-2/day	0.44 (0.11-1.68)
≥ 2 / day	0.44 (0.12-1.68)
Large stools	1.09 (0.51-2.30)
Nighttime encopresis	1.10 (0.51-2.41)
Palpable abdominal mass	1.23 (0.49-3.10)
Palpable rectal mass	3.39 (1.30-8.83)
CTT > 100 hrs	0.31 (0.12-0.85)

Discussion

This study shows that more severe symptoms, such as a relatively low defecation frequency, or a high encopresis frequency and the presence of a palpable rectal mass strongly correlate with a prolonged CTT and RSTT. Gender (girls) and a palpable rectal mass at entry were associated with a successful outcome at one year. An abdominal radiograph examination with the use of radio-opaque markers at intake had no predictive value. Measurement of CTT did not predict outcome if less than 100 hours. In contrast, a CTT above 100 hours predicted a poor outcome at one year.

The ingestion of radio-opaque markers and the use of one or more abdominal radiograph examinations allow the differentiation between children with normal or prolonged CTT¹⁸. Several techniques have been used to measure CTT in patients with gastrointestinal disorders and have proved to be reliable and reproducible¹⁹. In our study we used the three day Metcalf marker method. Radiograph examinations were repeated with a three day interval until more than 80% of the markers disappeared¹⁰. With this method, the radiation exposure is relatively high in children with severe constipation. In future studies, we will use the Bouchoucha method in which the patient ingests one capsule with 10 markers on six consecutive days. Subsequently, only one abdominal radiograph examination at day 7 is needed to calculate CTT²⁰.

In this study CTT was measured without first cleansing the colon. A recent study in adults with constipation showed no difference in distribution patterns of the markers before and after cleansing of the colon²¹.

In agreement with previous studies in adults and children we found that severe symptoms of constipation strongly correlated with prolonged CTT and RSTT^{12,15,22}. It has been suggested that constipation in children is mostly caused by the conscious or unconscious postponement of defecation (withholding behaviour). This could be learned behaviour due to pain with evacuation of a large faecal bolus. Prolonged stool retention in the rectum might result in increased anorectal sampling with failure of contraction of the external anal sphincter and consequent soiling. It is likely that the associated delay in RSTT leads to secondary prolongation of CTT.

In accordance with earlier studies in children with constipation a normal CTT was found in approximately 50% of the patients^{11,22-24}. It has been suggested that adults complaining of constipation with normal transit times have a high incidence of psychosocial disturbance and may not be truly constipated²⁵. In children, however, van der Plas et al. showed that reported symptoms correlate well with the actual

bowel habit²⁶. In a recent study we showed that colonic transit patterns, including normal colonic transit were not indicative of abnormal behaviour²⁷. Normal colonic transit time in constipated children is probably due to the fact that the measurement, even though it represents bowel pattern over a one week period, still remains a snapshot observation. Furthermore, the upper time limit for normal transit, as suggested by Arhan et al. is based on only 23 healthy children, and so may be misleading⁶. Papadopoulou et al. suggested that the markers themselves might accelerate transit, causing a false negative result¹³.

In our experience this marker test has proved useful in differentiating retentive from non-retentive defecation disorders, such as constipation and functional non-retentive faecal soiling²⁷. A normal CTT in combination with a normal defecation frequency and no rectal mass indicates the latter diagnosis²². Such children are best treated with a toilet training programme without laxatives²⁸. Furthermore, treatment of CTT is useful in cases in which information is unreliable, such as in eating disorders.

After one year, 58% of the patients in our study had been treated successfully. This was in accordance with success rates in other long term follow-up studies²⁹⁻³¹. The overall success rate was 68% when patients without symptoms of constipation were included. In a recent long term study, more than 50% of the children with constipation experienced at least one relapse within the five years of first treatment³².

To our surprise, the presence of a rectal mass at presentation was a positive predictor of success. After removal of the mass most children experience less pain with defecation. It is likely that rectal control of sensation and defecation improves after removal, and there is a decrease in encopresis and abdominal pain.

A CTT longer than 100 hours predicted a poor outcome at one year. Almost 40% of these children had a delay in transit in all segments. Some patients, mainly adolescent girls, might suffer from idiopathic slow transit constipation³³⁻³⁵. This severe form of constipation presents with a gradual reduction in bowel frequency and increasing abdominal pain.

In conclusion, the diagnostic and prognostic role of CTT measurements is limited. Only those with a CTT above 100 hours have a predictably poor outcome.

References

1. Loening-Baucke V. Chronic constipation in children. *Gastroenterology* 1993;105:1557-64.
2. Levine MD. Children with encopresis: A descriptive analysis. *Pediatrics* 1975;56:412-6.
3. Taitz LS, Wales JK, Urwin OM et al. Factors associated with outcome in management of defecation disorders. *Arch Dis Child* 1986;61:472-7.
4. Hinton JM and Lennard-Jones JE. Constipation: definition and classification. *Postgrad Med J* 1968;44:720-3.
5. Hinton JM, Lennard-Jones JE, and Young AC. A new method for studying gut transit times using radioopaque markers. *Gut* 1969;10:842-7.
6. Arhan P, Devroede G, Jehannin B et al. Segmental colonic transit time. *Dis Col Rect* 1981;24:625-9.
7. Barr RG, Levine MD, Wilkinson RH et al. Chronic and occult stool retention: a clinical tool for its evaluation in school-aged children. *Clin Pediatr* 1979;18:674-9.
8. Rockney RM, McQuade WH, and Days AL. The plain abdominal roentgenogram in the management of encopresis. *Arch Pediatr Adolesc Med* 1995;149:623-7.
9. Chaussade S, Khyari A, Roche H et al. Determination of total and segmental colonic transit time in constipated patients. *Dig Dis Sci* 1989;34:1168-72.
10. Metcalf AM, Phillips SF, Zinsmeister AR et al. Simplified assessment of segmental colonic transit. *Gastroenterology* 1987;92:40-47.
11. Benninga MA, Buller HA, Staalman CR et al. Defaecation disorders in children, colonic transit time versus the Barr-score. *Eur J Pediatr* 1995;154:277-84.
12. Benninga MA, Buller HA, Tytgat GN et al. Colonic transit time in constipated children: does pediatric slow-transit constipation exist? *J Pediatr Gastroenterol Nutr* 1996;23:241-51.
13. Papadopoulou A, Clayden GS, and Booth IW. The clinical value of solid marker transit studies in childhood constipation and soiling. *Eur J Pediatr* 1994;153:560-4.
14. Glia A, Lindberg G, Nilsson LH et al. Clinical value of symptom assessment in patients with constipation. *Dis Col Rect* 1999;42:1401-8.
15. Verduron A, Devroede G, Bouchoucha M et al. Megarectum. *Dig Dis Sci* 1988;33:1164-74.
16. Mollen RM, Claassen ATPM, and Kuijpers JHC. Measurement of colon transit time useful in the evaluation of functional constipation. *Ned Tijdschr Geneesk* 1998;142:357-61.

17. van der Plas RN, Benninga MA, Buller HA et al. Biofeedback training in treatment of childhood constipation: a randomised controlled study. *Lancet* 1996;348:776-80.
18. Aggarwal A, Cutts TF, Abell TL et al. Predominant symptoms in irritable bowel syndrome correlate with specific autonomic nervous system abnormalities. *Gastroenterology* 1994;106:945-50.
19. Nam YS, Pikarsky AJ, Wexner SD et al. Reproducibility of colonic transit study in patients with chronic constipation. *Dis Col Rect* 2001;44:86-92.
20. Bouchoucha M, Devroede G, Arhan P et al. What is the meaning of colorectal transit time measurement? *Dis Col Rect* 1992;35:773-82.
21. Sloots CE and Felt-Bersma RJ. Effect of bowel cleansing on colonic transit in constipation due to slow transit or evacuation disorder. *Neurogastroenterol Motil* 2002;14:55-61.
22. Benninga MA, Buller HA, Heymans HS et al. Is encopresis always the result of constipation? *Arch Dis Child* 1994;71:186-93.
23. Wald A. Colonic transit and anorectal manometry in chronic idiopathic constipation. *Arch Int Med* 1986;146:1713-6.
24. Gutierrez C, Marco A, Nogales A et al. Total and segmental colonic transit time and anorectal manometry in children with chronic idiopathic constipation. *J Pediatr Gastroenterol Nutr* 2002;35:31-8.
25. Wald A, Hinds JP, and Caruana BJ. Psychological and physiological characteristics of patients with severe idiopathic constipation. *Gastroenterology* 1989;97:932-7.
26. van der Plas RN, Benninga MA, Redekop WK et al. How accurate is the recall of bowel habits in children with defaecation disorders? *Eur J Pediatr* 1997;156:178-81.
27. Benninga MA, Voskuil WP, Akkerhuis GW et al. Colonic transit times and behaviour profiles in children with defecation disorders. *Arch Dis Child* 2004;89:13-6.
28. Louvel D, Delvaux M, Felez A et al. Oxytocin increases thresholds of colonic visceral perception in patients with irritable bowel syndrome. *Gut* 1996;39:741-7.
29. van Ginkel R, Benninga MA, Blommaart PJ et al. Lack of benefit of laxatives as adjunctive therapy for functional nonretentive fecal soiling in children. *J Pediatr* 2000;137:808-13.
30. Sutphen JL, Borowitz SM, Hutchison RL et al. Long-term follow-up of medically treated childhood constipation. *Clin Pediatr* 1995;34:576-80.
31. Loening-Baucke V. Constipation in early childhood: patient characteristics, treatment, and longterm follow up. *Gut* 1993;34:1400-4.
32. Keuzenkamp-Jansen CW, Fijnvandraat CJ, Kneepkens CM et al. Diagnostic dilemmas and results of treatment for chronic constipation. *Arch Dis Child* 1996;75:36-41.

33. van Ginkel R, Reitsma JB, Buller HA et al. Childhood constipation: longitudinal follow-up beyond puberty. *Gastroenterology* 2003;125:357-63.
34. Schouten WR, Ten Kate FJ, de Graaf EJ et al. Visceral neuropathy in slow transit constipation: an immunohistochemical investigation with monoclonal antibodies against neurofilament. *Dis Col Rect* 1993;36:1112-7.
35. Wedel T, Spiegler J, Soellner S et al. Enteric nerves and interstitial cells of Cajal are altered in patients with slow-transit constipation and megacolon. *Gastroenterology* 2002;123:1459-67.

