



UvA-DARE (Digital Academic Repository)

Suppression of plant resistance gene-based immunity by a fungal effector

Houterman, P.M.; Cornelissen, B.J.C.; Rep, M.

DOI

[10.1371/journal.ppat.1000061](https://doi.org/10.1371/journal.ppat.1000061)

Publication date

2008

Document Version

Final published version

Published in

PLoS Pathogens

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Houterman, P. M., Cornelissen, B. J. C., & Rep, M. (2008). Suppression of plant resistance gene-based immunity by a fungal effector. *PLoS Pathogens*, 4(5), Article e1000061. <https://doi.org/10.1371/journal.ppat.1000061>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Suppression of Plant Resistance Gene-Based Immunity by a Fungal Effector

Petra M. Houterman, Ben J. C. Cornelissen, Martijn Rep*

Plant Pathology, Swammerdam Institute for Life Sciences, University of Amsterdam, Amsterdam, The Netherlands

Abstract

The innate immune system of plants consists of two layers. The first layer, called basal resistance, governs recognition of conserved microbial molecules and fends off most attempted invasions. The second layer is based on Resistance (*R*) genes that mediate recognition of effectors, proteins secreted by pathogens to suppress or evade basal resistance. Here, we show that a plant-pathogenic fungus secretes an effector that can both trigger and suppress *R* gene-based immunity. This effector, Avr1, is secreted by the xylem-invading fungus *Fusarium oxysporum* f.sp. *lycopersici* (Fol) and triggers disease resistance when the host plant, tomato, carries a matching *R* gene (*I* or *I-1*). At the same time, Avr1 suppresses the protective effect of two other *R* genes, *I-2* and *I-3*. Based on these observations, we tentatively reconstruct the evolutionary arms race that has taken place between tomato *R* genes and effectors of Fol. This molecular analysis has revealed a hitherto unpredicted strategy for durable disease control based on resistance gene combinations.

Citation: Houterman PM, Cornelissen BJC, Rep M (2008) Suppression of Plant Resistance Gene-Based Immunity by a Fungal Effector. PLoS Pathog 4(5): e1000061. doi:10.1371/journal.ppat.1000061

Editor: Brendan P. Cormack, Johns Hopkins University School of Medicine, United States of America

Received: November 26, 2007; **Accepted:** April 9, 2008; **Published:** May 9, 2008

Copyright: © 2008 Houterman et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The study was funded by the University of Amsterdam and the Royal Academy of Sciences and Arts of the Netherlands.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: m.rep@uva.nl

Introduction

Long periods of co-evolution of plants and microorganisms have led to complex mechanisms of attack and defence, involving the innate immune system of plants and virulence factors of pathogens [1]. The first layer of plant defence, called basal immunity, is based on recognition of conserved microbial molecules but can be suppressed by microbial virulence factors known as “effectors”. Plants respond to this suppression by employing a second layer of defence, Resistance (*R*) gene-based immunity, which relies on recognition of effectors [2]. In turn, at least bacterial pathogens have found ways to manipulate or evade this second layer of defence [3]. It is unclear to what extent this capacity exists in eukaryotic plant pathogens like oomycetes and fungi.

Like bacteria, many plant-pathogenic fungi secrete proteins that are recognized by *R*-genes [4,5]. One of these fungi is *Fusarium oxysporum*, a common soil inhabitant. It propagates asexually and is mostly harmless. However, pathogenic and host-specific clonal lines have evolved that cause severe diseases in crops, such as banana, cotton, cucumber, melon and tomato [6,7]. Many of these diseases are caused by colonisation of the water-conducting xylem system of the roots followed by upward growth through xylem vessels, with wilting and death as a dramatic result. Strains of *F. oxysporum* that cause wilt of tomato plants are grouped in *forma specialis* (f.sp.) *lycopersici*. Several polymorphic resistance (*R*) genes have been identified in the tomato gene pool that each confer resistance against a subset of *F. oxysporum* f.sp. *lycopersici* (Fol) strains. These are *I* (for Immunity), *I-1*, *I-2* and *I-3* [8]. Races of Fol are named historically according to the *R* gene that is effective against them: the *I* gene and the (unlinked) *I-1* gene are effective against race 1, race 2 overcomes *I* and *I-1*, but is stopped by *I-2*, while race 3 overcomes *I*, *I-1* and *I-2* but is blocked by *I-3* [9].

Race 1 strains have been further divided into subgroups based on whether or not they are able to (partially) overcome *I-2* or *I-3* [9,10].

Based on the gene-for-gene hypothesis [11], it is assumed that disease resistance conferred by *R* genes in tomato requires ‘matching’ avirulence (*AVR*) genes in Fol. The *I* gene originates from *Solanum [Lycopersicon] pimpinellifolium* and resides on chromosome 11 [12,13], while the *I-1* gene is located on chromosome 7 in another wild relative of tomato, *Solanum [Lycopersicon] pennellii* [14]. The *I-2* gene has been cloned and encodes an R protein of the common NB-LRR class [15]. The *I-3* gene has not yet been cloned [16], but the matching *AVR* gene has: it encodes a small protein, Six1 (“Secreted in xylem 1”), which is secreted by Fol during colonization of the xylem system [17] and contributes to fungal virulence [9]. Six1 is now called Avr3 to indicate its gene-for-gene relationship with the *I-3* resistance gene.

We describe here the identification and analysis of a second avirulence factor of Fol, Avr1. Surprisingly, this protein does not only act as an avirulence factor in conjunction with the *I* gene, but also suppresses disease resistance mediated by *I-2* and *I-3*.

Results/Discussion

Identification of Avr1

In an initial analysis of the xylem sap proteome of tomato plants infected with Fol race 1 using 2-D gel electrophoresis and mass spectrometry, three small secreted proteins of Fol were identified in addition to Avr3 (Six1), named Six2, Six3 and Six4, and their genes cloned [18]. We now find that one of these, Six4, is not secreted by Fol race 2 (Fig. 1). For reasons detailed below, we now call this protein Avr1. Like the *AVR3 (SLX1)* gene, *AVR1* is surrounded by repetitive elements (Fig. 2A). In all of the race 1

Author Summary

In agriculture, the most environmentally friendly way to combat plant diseases is to make use of the innate immune system of plants, for instance by crossing into crop varieties polymorphic resistance genes that occur in natural populations of the crop plant or its close relatives. Plant pathogens, however, have co-evolved with their host plants and have developed ways to overcome the immune system. To effectively make use of components of the plant immune system, it is therefore important to understand the co-evolution of plants and their pathogens at the molecular level. For the interaction between a fungal pathogen and tomato, this paper presents a breakthrough in this respect. A small protein secreted by some strains of the fungus *Fusarium oxysporum* was found to suppress the activity of two disease resistance genes of tomato. However, a third resistance gene specifically targets this suppressor protein and renders the plant fully resistant against fungal strains that produce it. With this insight, together with knowledge of the genetic variation in the pathogen population, a combination of resistance genes is suggested that is expected to confer durable resistance in tomato against *Fusarium* wilt disease.

strains we examined, PCR experiments detected the presence of *AVR1* and no sequence polymorphism was detected in the coding regions of seven isolates from different clonal lines (see [9] for the list of strains; 17 of these are race 1, 23 are race 2 or 3). *AVR1* was not detected in race 2 or 3 strains by PCR nor is *AVR1* present in the genome sequence of the race 2 strain 4287 (*Fusarium oxysporum* Sequencing Project; Broad Institute of Harvard and MIT (<http://www.broad.mit.edu>)). Absence of *AVR1* or closely related genes in the race 2 and race 3 strains used in this study was confirmed by DNA gel blot analysis (Fig. 2B, lanes 4 and 7, respectively).

To test whether *AVR1* is indeed responsible for avirulence of Fol on plants carrying the *I* gene, we created an *AVR1* gene knock-out in a race 1 strain (Fol004) through *Agrobacterium*-mediated transformation (Fig. 2). For the *AVR1* gene, the frequency of homologous recombination leading to gene knock-out turned out to be extremely low, with only a single knock-out mutant obtained out of ~200 transformants (Fig. 2B, lane 2). A disease assay with this mutant (*avr1Δ*) confirmed that indeed deletion of *AVR1* leads to breaking of *I*-mediated disease resistance (Fig. 3A, panel A, quantified in Fig. 3B). Re-introduction of *AVR1* in the *avr1Δ* strain (Fig. 2B, lane 3) restored the original avirulence phenotype (results not shown). In addition, we found that disease resistance conferred by the unlinked *I-1* gene in tomato also depends on recognition of Avr1, since the *avr1Δ* strain (but not its parental strain) is virulent on a plant line carrying *I-1* (line 90E402F, results not shown). This suggests that *I* and *I-1* express the same resistance specificity.

To confirm that the *AVR1* gene is sufficient to trigger recognition by the *I* gene, we transformed *AVR1* to a race 2 strain (Fol007) and a race 3 strain (Fol029) that do not contain *AVR1* (Fig. 2B, lanes 4–9) and are virulent on *I*-containing tomato lines. Ten independent transformants (six of race 2 and four of race 3) containing *AVR1* were unable to cause disease on *I*-containing plants (Fig. 3A, panels B and C, quantified in Fig. 3B), confirming the avirulence character of *AVR1*. In contrast to Avr3 [9], Avr1 is dispensable for full virulence towards plants that do not contain *R* genes against Fol (results not shown).

Avr1 suppresses *I-2* and *I-3*-mediated disease resistance

Although all Fol strains possess an intact *AVR3* gene, most race 1 strains nevertheless cause disease on plants carrying only the *I-3* gene [9]. One explanation for this is that Avr1 itself is involved in suppression of *I-3* mediated disease resistance. To test this, we inoculated a plant line containing only the *I-3* gene with the set of Fol strains described above. The results clearly show that Avr1




Figure 1. Fol race 2 does not secrete Avr1/Six4. Proteins present in xylem sap of susceptible tomato plants infected with race 1 strain Fol004 (left panel) or race 2 strain Fol002 (right panel) were isolated and separated with 2-dimensional gel electrophoresis. Positions of isoelectric point markers are indicated at the top; positions of molecular weight markers are indicated on the left. The arrows in the left panel point to the two spots previously shown to contain Avr1 (Six4) [18]; the arrows in the right panel point to the corresponding (empty) positions. The right spot in the left panel likely represents a more extensively N-terminally processed form of Avr1 [18].
doi:10.1371/journal.ppat.1000061.g001




Figure 2. The AVR1 locus, gene deletion and complementation.

A) The AVR1 open reading frame (ORF; open arrow) is interrupted by a single intron (black box) [18] (accession AM234064). The ORF is flanked 714 bp upstream by a copy of the transposon *Tfo1* (striped arrow represents the end of the transposase ORF; triangle represents the inverted repeat), 485 bp upstream by a partial miniature impala repetitive element (*mimp-Δ*, grey box; triangle represents inverted repeat) and downstream by a *Fot5*-like repetitive element (the transposase ORF ends 541 bp downstream of the AVR1 ORF and is shown as a grey arrow). The small arrows denote the primers used to construct an AVR1 disruption construct and an AVR1 expression cassette for transformation to Fol (see Materials and methods). The insertion of a hygromycin resistance (*hygR*) cassette to create an AVR1 knock-out mutant is shown (not drawn to scale). The position of the probe and the restriction sites used for Southern blot analysis are indicated; H: *HindIII*, B: *BamHI*. B) Southern blot confirming AVR1 disruption and ectopic insertion of AVR1. A Southern blot of genomic DNA digested with *HindIII* and *BamHI* was probed with a 1.4 kb probe encompassing the AVR1 ORF and 3' sequences as indicated in Fig. 2A. The AVR1 locus in race 1 strain Fol004 (lane 1) is visible as a 1.25 kb *HindIII* band containing the ORF (AVR1) and a band of ~5 kb containing sequences 3' of the ORF (3'). In the race 1 *avr1Δ* strain (lane 2), replacement of the ORF with the disruption cassette through homologous recombination led to the expected replacement of the 1.25 *HindIII* band with a 1.1 kb *BamHI-HindIII* band containing part of the ORF and part of the disruption cassette (*avr1Δ*). Transformation of the AVR1 expression cassette to the *avr1Δ* strain (lane 3) led to reappearance of the AVR1 band. Race 2 strain Fol007 (lane 4) and race 3 strain Fol029 (lane 7) do not contain AVR1 (the AVR1 and 3' bands are absent). Transformation of the AVR1 expression cassette to these strains (lanes 5 and 6: race 2 transformants; lanes 8 and 9: race 3 transformants) leads to appearance of the 1.25 kb *HindIII* AVR1 band as well as a 0.56 kb *HindIII-BamHI* band (3' ectopic) that comprises sequences 3' of the AVR1 ORF until the *BamHI* site at the 3' end of the expression cassette (which is not present in the genomic locus but corresponds to the end of the probe shown in Fig. 2A). Note that in the *avr1Δ* strain (lane 2) the 0.56 kb band indicative of ectopic insertion is also present, indicating that this strain contains an additional copy of the disruption cassette. The additional, weaker bands are probably due to 104 bp of non-coding sequence of the *Fot5*-like transposon present at the 3' end of the probe (thick line

next to the grey arrow in Fig. 2A) – there are seven copies of this sequence in the latest release of the genome sequence of race 2 strain 4287 (*Fusarium oxysporum* Sequencing Project; Broad Institute of Harvard and MIT (<http://www.broad.mit.edu>). Molecular weight markers are indicated on the left (in kb).

doi:10.1371/journal.ppat.1000061.g002

indeed has this suppressive activity: deletion of AVR1 in race 1 leads to loss of virulence towards *I-3* plants (Fig. 3A, panel D, quantified in Fig. 3B), while introduction of AVR1 in race 2 or race 3 leads to gain of virulence towards *I-3* plants (Fig. 3A, panels E and F, quantified in Fig. 3B). Furthermore, we discovered that Avr1 also suppresses *I-2*-mediated disease resistance (Fig. 3A, panels D and E, quantified in Fig. 3B). This means that the ability of some race 1 strains to cause disease on *I-2* plants, as observed earlier [10], is likely to be caused by suppression of *I-2* rather than loss of AVR2. In accordance with earlier observations using *I-3* plants [9], we found that virulence due to suppression of *I-2* and *I-3* is partial compared to strains lacking the corresponding AVR gene (Fig. S1). It should be noted that not all race 1 strains are virulent on *I-2* and/or *I-3* plants [9,10], even though all contain AVR1 with identical sequences (results not shown). Apparently, suppression of R gene-based immunity by Avr1 is dependent on unknown factors in the genetic background of the fungus. Since suppression works in Fol007 (race 2) and Fol029 (race 3), the genetic background in which AVR1 is effective is not restricted to race 1 strains.

Possible function of Avr1

Our observation that Avr1 is not required for virulence to plants without *I* genes may be due to the existence of other effectors that are redundant for such an activity. Alternatively, the role of Avr1 is restricted to the suppression of *I-2* and *I-3*-mediated disease resistance. A mechanistic explanation for the latter role could be that Avr1 interferes directly with Avr2 and Avr3. However, at least Avr3 accumulates in xylem sap and remains unaltered in the presence of Avr1 [9,18]. A direct interaction between the two proteins could also not be demonstrated *in vitro* by pull down experiments (results not shown). Unlike bacteria, pathogenic fungi are not known to inject proteins directly into plant cells, but many are known to secrete small, frequently cysteine-rich, but otherwise unrelated proteins during colonization of plants [5]. Avr1, like Avr3, falls within this group, the predicted mature protein having 184 residues including 6 cysteines and lacking homology to other proteins [18]. The mode of action of most of these small secreted proteins has remained unclear. Molecular targets have been described for Avr2 and Avr4 from the leaf mold *Cladosporium fulvum*: Avr2 is a protease inhibitor [19] while Avr4 binds chitin in the fungal cell wall and protects it against attack by plant chitinases [20]. These two proteins act in the apoplast to enhance fungal virulence, but others act inside plant cells [4]. Uptake from the apoplast by plant cells has been shown directly for ToxA, a small secreted protein that acts as a host-selective toxin [21]. This may also occur with Avr2, since *I-2* is a cytoplasmic protein [15]. Avr1, then, may interfere with the uptake of Avr2 and Avr3. Alternatively, it may be taken up itself and interfere with *I-2* and *I-3* or with signal transduction processes downstream of these R proteins (Fig. 4).

Implications for the evolution of Avr-R gene interactions

Suppression of effector-triggered (R gene-mediated) immunity has been observed in bacteria [3,22,23]. In plant pathogenic fungi, suppression of avirulence by unlinked loci has been demonstrated by genetics in rust fungi [24]. In the flax rust fungus, two dominant




Figure 3. Avr1 suppresses I-2 and I-3 mediated resistance. Ten day old seedlings of tomato were inoculated with a fungal spore suspension and disease was scored after three weeks. Tomato lines carrying only a single resistance gene or no resistance gene were used to determine the effect of Avr1 on the activity of each resistance gene (see Materials and methods for description of plant lines). All lines were inoculated with the following Fol strains: race 1 (strain Fol004), race 2 (strain Fol007), race 3 (strain Fol029), race 1 *avr1Δ* (Fol004 with *AVR1* deleted by gene replacement), race 2+*AVR1* (Fol007 transformed with *AVR1*; similar virulence patterns were obtained with six independent transformants) and race 3+*AVR1* (Fol029 transformed with *AVR1*; similar virulence patterns were obtained with four independent transformants). A) Representative plants are shown three weeks after infection. Panel A shows that loss of *AVR1* leads to breaking of I-mediated resistance. Panel B and C show that gain of *AVR1* triggers I-mediated resistance. Panel D shows that loss of *AVR1* leads to loss of virulence on I-2 and I-3-containing plant lines. Panels E and F show that gain of *AVR1* by race 2 or race 3 leads to virulence on I-2 and I-3-containing plant lines. B): Quantification of disease assays. The outcomes of the disease assays depicted in (A) were quantified in two ways: 1) average plant weight above the cotyledons and 2) phenotype scoring according to a disease index ranging from zero (no disease) to four (heavily diseased or

dead). Error bars indicate the 95% confidence interval of the mean. Interactions where Avr1 induces I-mediated resistance are indicated with a circle. Interactions where Avr1 suppresses I-2 or I-3 are indicated with an asterisk. N.I.: not infected. doi:10.1371/journal.ppat.1000061.g003

alleles or tightly linked genes at the I (“inhibitor”) locus suppress – sometimes partially – either one (*MI*) or several (*MI*, *LI*, *7*, *8*, *10*) *R* genes out of 30 against flax rust [24,25]. The flax rust inhibitor locus is not itself linked to avirulence. Here, we report the identification of a fungal avirulence factor that suppresses disease resistance conferred by two *R* genes.

Interpreting this phenomenon in terms of molecular arms races between plants and their pathogens [1], we envisage the following scenario. During evolution of the tomato-Fol pathosystem, I-2 and I-3 have evolved to recognize, respectively, Avr2 and Avr3. Since Avr3 is required for full virulence of Fol, evasion of I-3 recognition through loss of the *AVR3* gene would entail a serious fitness penalty. This explains why all Fol strains analysed so far retained *AVR3* [9,26]. Point mutations in *AVR3* preventing recognition have not been found either [9]. A possible explanation for this is that the I-3 protein operates in accordance with the guard model, in which not the Avr3 protein itself but the effect it has on its virulence target is recognized [27]. In any case, Fol has (partially) regained virulence towards I-3-containing plants by acquisition of *AVR1*, which, as shown here, suppresses the function of I-3. Subsequently, tomato responded to this ‘invention’ with the employment of the I gene, or the unlinked I-1 gene, to specifically recognize and respond to Avr1. Apparently, I and I-1 are themselves insensitive to the suppressive effect of Avr1 (Fig. 4).

The agricultural ‘arms race’ between Fol and tomato is different from the natural one because it is dictated by successive *R* gene deployment in commercial cultivars [8]. The I gene from the wild tomato relative *Solanum [Lycopersicon] pimpinellifolium* was the first *R* gene to be introgressed into tomato cultivars to resist Fusarium wilt in the 1940s [12]. At that time, Fol strains without Avr1 may already have been present in some locations, since I-breaking race 2 strains were quickly discovered [28] even though major outbreaks did not occur before 1960 [29]. The I-2 gene, also from *S. pimpinellifolium* and directed against Avr2, was introduced in commercial cultivars in the 1960s to protect tomato against Fol race 2 [29,30]. The combination of I and I-2 was effective for about two decades until the appearance of race 3 in both Australia




Figure 4. Schematic summary of the interactions between Fol Avr proteins and tomato resistance (I) proteins. Arrows signify activation, lines ending in a cross bar signify suppression. Avr1 is synonymous to Six4, Avr3 is synonymous to Six1. doi:10.1371/journal.ppat.1000061.g004

and North America [31], which probably emerged from a race 2 background through selection for loss or mutation of *AVR2*. To combat race 3, the *I-3* gene was introgressed from *S. pennellii* [31]. From the results presented here, we deduce that the combination of *I* (or *I-1*) and *I-3* may yield durable resistance of tomato to Fusarium wilt disease of tomato, since *I-3* is directed against a virulence factor (*Avr3*) and *I* (and *I-1*) against the suppressor of *I-3* (*Avr1*).

The molecular toolbox that is now gradually filling up (*Avr1*, *Avr3*, *I-2*) will help us to define host targets and evolutionary bottlenecks that govern the arms race in the Fol-tomato pathosystem. It also may allow development of new strategies for breeding plants with durable resistance against fungal pathogens.

Materials and Methods

Plant lines and fungal strains

The following tomato lines were used (Fol resistance genes between brackets): GCR161 (*I*) [32], 90E402F (*I-1*) [31,33]; 90E341F (*I-2*) [29] and E779 (*I-3*) [31], C32 (no *I* gene) [32]. The following Fol strains were used: Fol004 (race 1), Fol002 (race 2), Fol007 (race 2), Fol029 (race 3), Fol004avr1Δ (Fol004 with *AVR1* deleted by gene replacement), Fol004avr1Δ+*AVR1* (Fol004avr1Δ transformed with *AVR1*), Fol007+*AVR1* (Fol007 transformed with *AVR1*), Fol029+*AVR1* (Fol029 transformed with *AVR1*). See Rep *et al.* (2005) [9] for a more detailed description of the wild type Fol strains.

Xylem sap proteome analysis

Proteins present in xylem sap of tomato plants infected with Fol were isolated and separated with 2-dimensional gel electrophoresis as described earlier [18], using for the first dimension an Immobiline DryStrip of 13 cm, pH 6–11 NL (Amersham Biosciences).

Disease assays

Ten day old seedlings of tomato were inoculated with a fungal spore suspension and disease was scored after three weeks as described earlier [17]. The outcome of the disease assays was quantified in two ways: 1) average plant weight above the cotyledons and 2) phenotype scoring according to a disease index ranging from zero (no disease) to four (heavily diseased or dead) [17].

AVR1 disruption and complementation constructs

The *AVR1* disruption construct was made by PCR amplification of *AVR1* upstream and downstream sequences for homologous recombination, and their insertion in front of and behind the hygromycin resistance gene in the vector pRW2h (see below): an upstream fragment, from 714 bp to 1 bp upstream of the start codon, was cloned into pRW2h between the *PacI* and *KpnI* sites, and a downstream fragment, from 375 bp after the start codon to 537 bp downstream of the stop codon, was cloned into pRW2h between the *XbaI* and *BssHII* sites (see Fig. 2A for location of the primers). The construct for complementation was made by amplification of a *AVR1* expression cassette from 714 bp upstream of the start codon to 537 bp downstream of the stop codon (Fig. 2A), which was inserted between the *XbaI* and *StuI* sites of pRW1p (see below). Transformation of these constructs to Fol was done with *Agrobacterium* as described earlier [34].

pRW2h is a binary vector for *Agrobacterium*-mediated transformation of fungi. It was made through insertion of a *NheI-XbaI*

fragment from pAN7.1, carrying the hygromycin resistance gene *hph* under control of the *Aspergillus (Emericella) nidulans gpd* promoter and *trpC* terminator [35], into the unique *XbaI* site of pPZP-201BK [36]. Similarly, pRW1p was derived from pPZP-201BK through insertion of a *NheI-XbaI* fragment from pAN8.1 [35] carrying the phleomycin resistance gene *ble* under control of the same *gpd* promoter and *trpC* terminator.

Southern blotting

Genomic DNA of *F. oxysporum* was isolated according to Raeder and Broda [37], digested with *HindIII* and *BamHI*, separated in a 1% agarose gel and blotted to Hybond N+ according to Sambrook *et al.* [38]. The probe containing the *AVR1* ORF and 3' sequences (1402 bp, Fig. 2A) was generated by PCR and contains sequences from 72 bp upstream to 537 bp downstream of the ORF. The probe was radioactively labelled with $\alpha^{32}\text{P}$ dATP using the DecaLabel™ DNA labeling kit from MBI Fermentas (Vilnius, Lithuania). Hybridization was done overnight at 65°C in 0.5M phosphate buffer pH 7.2 containing 7% SDS and 1 mM EDTA. Blots were washed at 65°C with 0.2 X SSC, 0.1% SDS. The position of sequences hybridizing to the probe were visualized by phosphoimaging (Molecular Dynamics).

Accession numbers

The *AVR1* (*SIX4*) locus: AM234064

The *Avr1* (*Six4*) protein: CAJ84000

Supporting Information

Figure S1 Suppression of *I-2* and *I-3* is partial. Ten day old seedlings of tomato were inoculated with a fungal spore suspension and disease was scored after three weeks as described earlier. Tomato lines carrying only *I-2* (90E341F) or *I-3* (E779) were either mock-inoculated (A,B) or inoculated with race 1 strain Fol004 that suppress *I-2* and *I-3* (C, D) or with strains that avoid recognition by *I-2* or *I-3* through absence of the corresponding *AVR* gene (E, F). In (E), race 3 strain Fol029 (no *AVR2*) was used. In (F), Fol004 *avr3*Δ (race 1 strain Fol004 with *AVR3* (*SIX1*) deleted by gene replacement) was used. Representative plants are shown three weeks after infection. Note that although *AVR3* is required for full virulence towards susceptible plants of three weeks and older, *AVR3* is not required for virulence in the seedling assay used here, allowing assessment of the effectiveness of individual *R* genes [9].

Found at: doi:10.1371/journal.ppat.1000061.s001 (5.94 MB TIF)

Acknowledgments

We thank Frank Takken, Charlotte van der Does and Michel Haring for critical review of the manuscript. We thank Ringo van Wijk for construction of pRW2h and pRW1p, and Michiel Meijer for producing tagged versions of *Avr1* and *Avr3* and assessing their possible interaction *in vitro*.

Author Contributions

Conceived and designed the experiments: MR. Performed the experiments: PH. Analyzed the data: PH MR. Wrote the paper: MR. Originated the research leading up to this paper and provided guidance and review: BC.

References

1. Stahl EA, Bishop JG (2000) Plant-pathogen arms races at the molecular level. *Curr Opin Plant Biol* 3: 299–304.
2. Jones JDG, Dangl JL (2006) The plant immune system. *Nature* 444: 323–329.
3. Rosebrock TR, Zeng L, Brady JJ, Abramovitch RB, Xiao F, et al. (2007) A bacterial E3 ubiquitin ligase targets a host protein kinase to disrupt plant immunity. *Nature* 448: 370–374.

4. Catanzariti AM, Dodds PN, Ellis JG (2007) Avirulence proteins from haustoria-forming pathogens. *FEMS Microbiol Lett* 269: 181–188.
5. Rep M (2005) Small proteins of plant-pathogenic fungi secreted during host colonization. *FEMS Microbiol Lett* 253: 19–27.
6. Di Pietro A, Madrid MP, Caracul Z, Delgado-Jarana J, Roncero MIG (2003) *Fusarium oxysporum*: exploring the molecular arsenal of a vascular wilt fungus. *Mol Plant Pathol* 4: 315–325.
7. Gordon TR, Martyn RD (1997) The evolutionary biology of *Fusarium oxysporum*. *Annu Rev Phytopathol* 35: 111–128.
8. Huang CC, Lindhout P (1997) Screening for resistance in wild *Lycopersicon* species to *Fusarium oxysporum* f. sp. *lycopersici* race 1 and race 2. *Euphytica* 93: 145–153.
9. Rep M, Meijer M, Houterman PM, van der Does HC, Cornelissen BJC (2005) *Fusarium oxysporum* evades I-3-mediated resistance without altering the matching avirulence gene. *Mol Plant-Microbe Interact* 18: 15–23.
10. Mes JJ, Weststeijn EA, Herlaar F, Lambalk JJM, Wijbrandi J, et al. (1999) Biological and molecular characterization of *Fusarium oxysporum* f. sp. *lycopersici* divides race 1 isolates into separate virulence groups. *Phytopathology* 89: 156–160.
11. Flor HH (1971) Current status of the gene-for-gene concept. *Annu Rev Phytopathol* 9: 275–296.
12. Bohn GW, Tucker CM (1939) Immunity to *Fusarium* wilt of tomato. *Science* 89: 603–604.
13. Sela-Buurlage MB, Budai-Hadrian O, Pan Q, Carmel-Goren L, Vunsch R, et al. (2001) Genome-wide dissection of *Fusarium* resistance in tomato reveals multiple complex loci. *Mol Genet Genomics* 265: 1104–1111.
14. Sarfatti M, Abu-Abied M, Katan J, Zamir D (1991) RFLP mapping of *I1*, a new locus in tomato conferring resistance against *Fusarium oxysporum* f. sp. *lycopersici* race 1. *Theor Appl Genet* 82: 22–26.
15. Simons G, Groenendijk J, Wijbrandi J, Reijers MGJ, Diergaarde P, et al. (1998) Dissection of the *Fusarium I2* gene cluster in Tomato reveals six homologs and one active gene copy. *Plant Cell* 10: 1055–1068.
16. Hemming MN, Basuki S, McGrath DJ, Carroll BJ, Jones DA (2004) Fine mapping of the tomato I-3 gene for fusarium wilt resistance and elimination of a co-segregating resistance gene analogue as a candidate for I-3. *Theor Appl Genet* 109: 409–418.
17. Rep M, van der Does HC, Meijer M, van Wijk R, Houterman PM, et al. (2004) A small, cysteine-rich protein secreted by *Fusarium oxysporum* during colonization of xylem vessels is required for I-3-mediated resistance in tomato. *Mol Microbiol* 53: 1373–1383.
18. Houterman PM, Speijer D, Dekker HL, de Koster CG, Cornelissen BJC, et al. (2007) The mixed xylem sap proteome of *Fusarium oxysporum*-infected tomato plants. *Molecular Plant Pathology* 8: 215–221.
19. Rooney HC, van't Klooster JW, van der Hoorn RA, Joosten MH, Jones JD, et al. (2005) Cladosporium Avr2 Inhibits Tomato Rcr3 Protease Required for Cf-2-Dependent Disease Resistance. *Science* 308: 1783–1786.
20. van den Burg HA, Harrison SJ, Joosten MH, Vervoort J, de Wit PJ (2006) *Cladosporium fulvum* Avr4 protects fungal cell walls against hydrolysis by plant chitinases accumulating during infection. *Mol Plant-Microbe Interact* 19: 1420–1430.
21. Manning VA, Ciuffetti LM (2005) Localization of Ptr ToxA Produced by *Pyrenophora tritici-repentis* Reveals Protein Import into Wheat Mesophyll Cells. *Plant Cell* 17: 3203–3212.
22. Chisholm ST, Coaker G, Day B, Staskawicz BJ (2006) Host-microbe interactions: shaping the evolution of the plant immune response. *Cell* 124: 803–814.
23. Abramovitch RB, Anderson JC, Martin GB (2006) Bacterial elicitation and evasion of plant innate immunity. *Nat Rev Mol Cell Biol* 7: 601–611.
24. Lawrence GJ, Mayo GME, Shepherd KW (1981) Interactions between genes controlling pathogenicity in the flax rust fungus. *Phytopathology* 71: 12–19.
25. Jones DA (1988) Genetic properties of inhibitor genes in flax rust that alter avirulence to virulence on flax. *Phytopathology* 78: 342–344.
26. van der Does HC, Lievens B, Claes L, Houterman PM, Cornelissen BJC, et al. (2008) The presence of a virulence locus discriminates *Fusarium oxysporum* isolates causing tomato wilt from other isolates. *Environmental Microbiology*; in press.
27. Van der Hoorn RA, De Wit PJGM, Joosten MHAJ (2002) Balancing selection favors guarding resistance proteins. *Trends Plant Sci* 7: 67–71.
28. Alexander LJ, Tucker CM (1945) Physiological specialization in the tomato wilt fungus *Fusarium oxysporum* f.sp. *lycopersici*. *J Agric Res* 70: 303–313.
29. Stall RE, Walter JM (1965) Selection and inheritance of resistance in tomato to isolates of races 1 and 2 of the *Fusarium* wilt organism. *Phytopathology* 55: 1213–1215.
30. Cirulli M, Alexander LJ (1966) A comparison of pathogenic isolates of *Fusarium oxysporum* f.sp. *lycopersici* and different sources of resistance in tomato. *Phytopathology* 56: 1301–1304.
31. Scott JW, Jones JP (1989) Monogenic resistance in tomato to *Fusarium oxysporum* f. sp. *lycopersici* race 3. *Euphytica* 40: 49–53.
32. Kroon BAM, Elgersma DM (1993) Interactions between race 2 of *Fusarium oxysporum* f. sp. *lycopersici* and near-isogenic resistant and susceptible lines of intact plants or callus of tomato. *J Phytopathology* 137: 1–9.
33. Mes JJ, Wit R, Testerink CS, de Groot F, Haring MA, et al. (1999) Loss of avirulence and reduced pathogenicity of a gamma-irradiated mutant of *Fusarium oxysporum* f. sp. *lycopersici*. *Phytopathology* 89: 1131–1137.
34. Takken FLW, Van Wijk R, Michielse CB, Houterman PM, Ram AF, et al. (2004) A one-step method to convert vectors into binary vectors suited for *Agrobacterium*-mediated transformation. *Curr Genet* 45: 242–248.
35. Punt PJ, van den Hondel CA (1992) Transformation of filamentous fungi based on hygromycin B and phleomycin resistance markers. *Methods Enzymol* 216: 447–457.
36. Covert SF, Kapoor P, Lee MH, Briley A, Nairn CJ (2001) *Agrobacterium tumefaciens*-mediated transformation of *Fusarium circinatum*. *Mycological Research* 105: 259–264.
37. Raeder U, Broda P (1985) Rapid preparation of DNA from filamentous fungi. *Lett Appl Microb* 1: 17–20.
38. Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning. A Laboratory Manual*. Cold Spring Harbor New York: Cold Spring Harbor Laboratory Press.