



## UvA-DARE (Digital Academic Repository)

### The effect of altered expression of transcriptional regulators of catabolism on the transcription profile and physiology of *Saccharomyces cerevisiae*

Schuurmans, J.M.

**Publication date**  
2008

[Link to publication](#)

#### **Citation for published version (APA):**

Schuurmans, J. M. (2008). *The effect of altered expression of transcriptional regulators of catabolism on the transcription profile and physiology of Saccharomyces cerevisiae*. [Thesis, fully internal, Universiteit van Amsterdam].

#### **General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

#### **Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

## References

## References

- Ahuatzi, D., Herrero, P., de la Cera, T. and Moreno, F.** (2004). The glucose-regulated nuclear localization of hexokinase 2 in *Saccharomyces cerevisiae* is Mig1-dependent. *J Biol Chem* **279**, 14440-6.
- Ahuatzi, D., Riera, A., Pelaez, R., Herrero, P. and Moreno, F.** (2007). Hxk2 regulates the phosphorylation state of Mig1 and therefore its nucleocytoplasmic distribution. *J Biol Chem* **282**, 4485-93.
- Bailey, R. B. and Woodward, A.** (1984). Isolation and characterization of a pleiotropic glucose repression resistant mutant of *Saccharomyces cerevisiae*. *Mol Gen Genet* **193**, 507-12.
- Balzi, E. and Goffeau, A.** (1995). Yeast multidrug resistance: the PDR network. *J Bioenerg Biomembr* **27**, 71-6.
- Barbara, K. E., Haley, T. M., Willis, K. A. and Santangelo, G. M.** (2007a). The transcription factor Gcr1 stimulates cell growth by participating in nutrient-responsive gene expression on a global level. *Mol Genet Genomics* **277**, 171-88.
- Barbara, K. E., Willis, K. A., Haley, T. M., Deminoff, S. J. and Santangelo, G. M.** (2007b). Coiled coil structures and transcription: an analysis of the *S. cerevisiae* coilome. *Mol Genet Genomics* **278**, 135-47.
- Barnett, J. A.** (1976). The utilization of sugars by yeasts. *Adv Carbohydr Chem Biochem* **32**, 125-234.
- Barnett, J. A.** (1998). A history of research on yeasts. 1: Work by chemists and biologists 1789-1850. *Yeast* **14**, 1439-51.
- Barnett, J. A.** (2000). A history of research on yeasts 2: Louis Pasteur and his contemporaries, 1850-1880. *Yeast* **16**, 755-71.
- Battle, M., Lu, A., Green, D. A., Xue, Y. and Hirsch, J. P.** (2003). Krh1p and Krh2p act downstream of the Gpa2p G(alpha) subunit to negatively regulate haploid invasive growth. *J Cell Sci* **116**, 701-10.
- Berkey, C. D., Vyas, V. K. and Carlson, M.** (2004). Nrg1 and nrg2 transcriptional repressors are differently regulated in response to carbon source. *Eukaryot Cell* **3**, 311-7.
- Bird, A. J., Gordon, M., Eide, D. J. and Winge, D. R.** (2006). Repression of *ADH1* and *ADH3* during zinc deficiency by Zap1-induced intergenic RNA transcripts. *Embo J* **25**, 5726-34.
- Bisson, L. F. and Fraenkel, D. G.** (1984). Expression of kinase-dependent glucose uptake in *Saccharomyces cerevisiae*. *J Bacteriol* **159**, 1013-7.
- Blackman, F. F.** (1905). Optima and limiting factors. *Ann Bot* **19**, 281-295.
- Bloecher, A. and Tatchell, K.** (2000). Dynamic localization of protein phosphatase type 1 in the mitotic cell cycle of *Saccharomyces cerevisiae*. *J Cell Biol* **149**, 125-40.
- Blom, J., De Mattos, M. J. and Grivell, L. A.** (2000). Redirection of the respiratory flux distribution in *Saccharomyces cerevisiae* by overexpression of the transcription factor Hap4p. *Appl Environ Microbiol* **66**, 1970-3.
- Boer, V. M., de Winde, J. H., Pronk, J. T. and Piper, M. D.** (2003). The genome-wide transcriptional responses of *Saccharomyces cerevisiae* grown on glucose in aerobic chemostat cultures limited for carbon, nitrogen, phosphorus, or sulfur. *J Biol Chem* **278**, 3265-74.

## References

- Bojunga, N. and Entian, K. D.** (1999). Cat8p, the activator of gluconeogenic genes in *Saccharomyces cerevisiae*, regulates carbon source-dependent expression of NADP-dependent cytosolic isocitrate dehydrogenase (Idp2p) and lactate permease (Jen1p). *Mol Gen Genet* **262**, 869-75.
- Bojunga, N., Kotter, P. and Entian, K. D.** (1998). The succinate/fumarate transporter Acr1p of *Saccharomyces cerevisiae* is part of the gluconeogenic pathway and its expression is regulated by Cat8p. *Mol Gen Genet* **260**, 453-61.
- Boles, E. and Hollenberg, C. P.** (1997). The molecular genetics of hexose transport in yeasts. *FEMS Microbiol Rev* **21**, 85-111.
- Boorsma, A., Foat, B. C., Vis, D., Klis, F. and Bussemaker, H. J.** (2005). T-profiler: scoring the activity of predefined groups of genes using gene expression data. *Nucleic Acids Res* **33**, W592-5.
- Bruinenberg, P. M.** (1986). The NADP(H) redox couple in yeast metabolism. *Antonie Van Leeuwenhoek* **52**, 411-29.
- Button, D. K.** (1993). Nutrient-limited microbial growth kinetics: overview and recent advances. *Antonie Van Leeuwenhoek* **63**, 225-35.
- Cameron, S., Levin, L., Zoller, M. and Wigler, M.** (1988). cAMP-independent control of sporulation, glycogen metabolism, and heat shock resistance in *S. cerevisiae*. *Cell* **53**, 555-66.
- Carlson, M., Osmond, B. C. and Botstein, D.** (1981). Mutants of yeast defective in sucrose utilization. *Genetics* **98**, 25-40.
- Celenza, J. L. and Carlson, M.** (1986). A yeast gene that is essential for release from glucose repression encodes a protein kinase. *Science* **233**, 1175-80.
- Celenza, J. L., Eng, F. J. and Carlson, M.** (1989). Molecular analysis of the SNF4 gene of *Saccharomyces cerevisiae*: evidence for physical association of the SNF4 protein with the SNF1 protein kinase. *Mol Cell Biol* **9**, 5045-54.
- Chang, J. S., Henry, K., Wolf, B. L., Geli, M. and Lemmon, S. K.** (2002). Protein phosphatase-1 binding to scd5p is important for regulation of actin organization and endocytosis in yeast. *J Biol Chem* **277**, 48002-8.
- Chaves, R. S., Herrero, P. and Moreno, F.** (1999). Med8, a subunit of the mediator CTD complex of RNA polymerase II, directly binds to regulatory elements of *SUC2* and *HXK2* genes. *Biochem Biophys Res Commun* **254**, 345-50.
- Cheng, C., Huang, D. and Roach, P. J.** (1997). Yeast *PIG* genes: *PIG1* encodes a putative type 1 phosphatase subunit that interacts with the yeast glycogen synthase Gsy2p. *Yeast* **13**, 1-8.
- Ciriacy, M.** (1977). Isolation and characterization of yeast mutants defective in intermediary carbon metabolism and in carbon catabolite derepression. *Mol Gen Genet* **154**, 213-20.
- Colombo, S., Ronchetti, D., Thevelein, J. M., Winderickx, J. and Martegani, E.** (2004). Activation state of the Ras2 protein and glucose-induced signaling in *Saccharomyces cerevisiae*. *J Biol Chem* **279**, 46715-22.
- Contois, D. E.** (1959). Kinetics of bacterial growth: relationship between population density and specific growth rate of continuous cultures. *J Gen Microbiol* **21**, 40-50.

## References

- Coons, D. M., Boulton, R. B. and Bisson, L. F.** (1995). Computer-assisted nonlinear regression analysis of the multicomponent glucose uptake kinetics of *Saccharomyces cerevisiae*. *J Bacteriol* **177**, 3251-8.
- Cui, D. Y., Brown, C. R. and Chiang, H. L.** (2004). The type 1 phosphatase Reg1p-Glc7p is required for the glucose-induced degradation of fructose-1,6-bisphosphatase in the vacuole. *J Biol Chem* **279**, 9713-24.
- Dang, V. D., Valens, M., Bolotin-Fukuhara, M. and Daignan-Fornier, B.** (1994). A genetic screen to isolate genes regulated by the yeast CCAAT-box binding protein Hap2p. *Yeast* **10**, 1273-83.
- Daran-Lapujade, P., Rossell, S., van Gulik, W. M., Luttk, M. A., de Groot, M. J., Slijper, M., Heck, A. J., Daran, J. M., de Winde, J. H., Westerhoff, H. V., Pronk, J. T. and Bakker, B. M.** (2007). The fluxes through glycolytic enzymes in *Saccharomyces cerevisiae* are predominantly regulated at posttranscriptional levels. *Proc Natl Acad Sci U S A* **104**, 15753-8.
- De Deken, R. H.** (1966). The crabtree effect: a regulatory system in yeast. *J. Gen. Microbiol.* **44**, 149-156.
- de Groot, M. J., Daran-Lapujade, P., van Breukelen, B., Knijnenburg, T. A., de Hulster, E. A., Reinders, M. J., Pronk, J. T., Heck, A. J. and Slijper, M.** (2007). Quantitative proteomics and transcriptomics of anaerobic and aerobic yeast cultures reveals post-transcriptional regulation of key cellular processes. *Microbiology* **153**, 3864-78.
- de la Cera, T., Herrero, P., Moreno-Herrero, F., Chaves, R. S. and Moreno, F.** (2002). Mediator factor Med8p interacts with the hexokinase 2: implication in the glucose signalling pathway of *Saccharomyces cerevisiae*. *J Mol Biol* **319**, 703-14.
- De Nicola, R., Hazelwood, L. A., De Hulster, E. A., Walsh, M. C., Knijnenburg, T. A., Reinders, M. J., Walker, G. M., Pronk, J. T., Daran, J. M. and Daran-Lapujade, P.** (2007). Physiological and transcriptional responses of *Saccharomyces cerevisiae* to zinc limitation in chemostat cultures. *Appl Environ Microbiol.*
- de Winde, J. H. and Grivell, L. A.** (1993). Global regulation of mitochondrial biogenesis in *Saccharomyces cerevisiae*. *Prog Nucleic Acid Res Mol Biol* **46**, 51-91.
- Diderich, J. A., Raamsdonk, L. M., Kruckeberg, A. L., Berden, J. A. and Van Dam, K.** (2001a). Physiological properties of *Saccharomyces cerevisiae* from which hexokinase II has been deleted. *Appl Environ Microbiol* **67**, 1587-93.
- Diderich, J. A., Raamsdonk, L. M., Kuiper, A., Kruckeberg, A. L., Berden, J. A., Teixeira de Mattos, M. J. and van Dam, K.** (2002). Effects of a hexokinase II deletion on the dynamics of glycolysis in continuous cultures of *Saccharomyces cerevisiae*. *FEMS Yeast Res* **2**, 165-72.
- Diderich, J. A., Schepper, M., van Hoek, P., Luttk, M. A., van Dijken, J. P., Pronk, J. T., Klaassen, P., Boelens, H. F., de Mattos, M. J., van Dam, K. and Kruckeberg, A. L.** (1999). Glucose uptake kinetics and transcription of HXT genes in chemostat cultures of *Saccharomyces cerevisiae*. *J Biol Chem* **274**, 15350-9.

## References

- Diderich, J. A., Schuurmans, J. M., Van Gaalen, M. C., Kruckeberg, A. L. and Van Dam, K.** (2001b). Functional analysis of the hexose transporter homologue HXT5 in *Saccharomyces cerevisiae*. *Yeast* **18**, 1515-24.
- Dombek, K. M., Kacherovsky, N. and Young, E. T.** (2004). The Reg1-interacting proteins, Bmh1, Bmh2, Ssb1, and Ssb2, have roles in maintaining glucose repression in *Saccharomyces cerevisiae*. *J Biol Chem* **279**, 39165-74.
- Dombek, K. M., Voronkova, V., Raney, A. and Young, E. T.** (1999). Functional analysis of the yeast Glc7-binding protein Reg1 identifies a protein phosphatase type 1-binding motif as essential for repression of *ADH2* expression. *Mol Cell Biol* **19**, 6029-40.
- Dubacq, C., Chevalier, A. and Mann, C.** (2004). The protein kinase Snf1 is required for tolerance to the ribonucleotide reductase inhibitor hydroxyurea. *Mol Cell Biol* **24**, 2560-72.
- Elbing, K., Larsson, C., Bill, R. M., Albers, E., Snoep, J. L., Boles, E., Hohmann, S. and Gustafsson, L.** (2004). Role of hexose transport in control of glycolytic flux in *Saccharomyces cerevisiae*. *Appl Environ Microbiol* **70**, 5323-30.
- Entian, K. D. and Zimmermann, F. K.** (1982). New genes involved in carbon catabolite repression and derepression in the yeast *Saccharomyces cerevisiae*. *J Bacteriol* **151**, 1123-8.
- Estruch, F., Treitel, M. A., Yang, X. and Carlson, M.** (1992). N-terminal mutations modulate yeast SNF1 protein kinase function. *Genetics* **132**, 639-50.
- Fell, D.** (1996). Understanding the control of metabolism. *Portland press*.
- Fell, D. A.** (1992). Metabolic control analysis: a survey of its theoretical and experimental development. *Biochem J* **286 (Pt 2)**, 313-30.
- Feng, Z. H., Wilson, S. E., Peng, Z. Y., Schlender, K. K., Reimann, E. M. and Trumbly, R. J.** (1991). The yeast *GLC7* gene required for glycogen accumulation encodes a type 1 protein phosphatase. *J Biol Chem* **266**, 23796-801.
- Flick, J. S. and Johnston, M.** (1991). *GRR1* of *Saccharomyces cerevisiae* is required for glucose repression and encodes a protein with leucine-rich repeats. *Mol Cell Biol* **11**, 5101-12.
- Flikweert, M. T., de Swaaf, M., van Dijken, J. P. and Pronk, J. T.** (1999). Growth requirements of pyruvate-decarboxylase-negative *Saccharomyces cerevisiae*. *FEMS Microbiol Lett* **174**, 73-9.
- Forsburg, S. L. and Guarente, L.** (1989). Identification and characterization of HAP4: a third component of the CCAAT-bound HAP2/HAP3 heteromer. *Genes Dev* **3**, 1166-78.
- Frederick, D. L. and Tatchell, K.** (1996). The *REG2* gene of *Saccharomyces cerevisiae* encodes a type 1 protein phosphatase-binding protein that functions with Reg1p and the Snf1 protein kinase to regulate growth. *Mol Cell Biol* **16**, 2922-31.
- Gamo, F. J., Moreno, E. and Lagunas, R.** (1995). The low-affinity component of the glucose transport system in *Saccharomyces cerevisiae* is not due to passive diffusion. *Yeast* **11**, 1393-8.
- Gancedo, J. M.** (1998). Yeast carbon catabolite repression. *Microbiol Mol Biol Rev* **62**, 334-61.
- Griffioen, G. and Thevelein, J. M.** (2002). Molecular mechanisms controlling the localisation of protein kinase A. *Curr Genet* **41**, 199-207.

## References

- Harashima, T. and Heitman, J.** (2002). The Galpha protein Gpa2 controls yeast differentiation by interacting with kelch repeat proteins that mimic Gbeta subunits. *Mol Cell* **10**, 163-73.
- Hartwell, L. H., Mortimer, R. K., Culotti, J. and Culotti, M.** (1973). Genetic Control of the Cell Division Cycle in Yeast: V. Genetic Analysis of cdc Mutants. *Genetics* **74**, 267-286.
- Haurie, V., Perrot, M., Mini, T., Jeno, P., Sagliocco, F. and Boucherie, H.** (2001). The transcriptional activator Cat8p provides a major contribution to the reprogramming of carbon metabolism during the diauxic shift in *Saccharomyces cerevisiae*. *J Biol Chem* **276**, 76-85.
- Hedges, D., Proft, M. and Entian, K. D.** (1995). CAT8, a new zinc cluster-encoding gene necessary for derepression of gluconeogenic enzymes in the yeast *Saccharomyces cerevisiae*. *Mol Cell Biol* **15**, 1915-22.
- Heinrich, R. and Rapoport, T. A.** (1974). A linear steady-state treatment of enzymatic chains. General properties, control and effector strength. *Eur J Biochem* **42**, 89-95.
- Heinrich, R., Schuster, S.** (1996). The regulation of cellular systems. *Chapman and Hall*.
- Herrero, P., Martínez-Campa, C. and Moreno, F.** (1998). The hexokinase 2 protein participates in regulatory DNA-protein complexes necessary for glucose repression of the SUC2 gene in *Saccharomyces cerevisiae*. *FEBS Lett* **434**, 71-6.
- Hirayama, T., Maeda, T., Saito, H. and Shinozaki, K.** (1995). Cloning and characterization of seven cDNAs for hyperosmolarity-responsive (HOR) genes of *Saccharomyces cerevisiae*. *Mol Gen Genet* **249**, 127-38.
- Hohmann, S., Winderickx, J., de Winde, J. H., Valckx, D., Cobbaert, P., Luyten, K., de Meirsmann, C., Ramos, J. and Thevelein, J. M.** (1999). Novel alleles of yeast hexokinase PII with distinct effects on catalytic activity and catabolite repression of SUC2. *Microbiology* **145** ( Pt 3), 703-14.
- Horak, J. and Wolf, D. H.** (1997). Catabolite inactivation of the galactose transporter in the yeast *Saccharomyces cerevisiae*: ubiquitination, endocytosis, and degradation in the vacuole. *J Bacteriol* **179**, 1541-9.
- Ichimura, T., Kubota, H., Goma, T., Mizushima, N., Ohsumi, Y., Iwago, M., Kakiuchi, K., Shekhar, H. U., Shinkawa, T., Taoka, M., Ito, T. and Isobe, T.** (2004). Transcriptomic and proteomic analysis of a 14-3-3 gene-deficient yeast. *Biochemistry* **43**, 6149-58.
- Inc, W.** (2007). Mathematica Edition: Version 6, Wolfram Research, Inc.
- Jackson, M.** (1977). *The world guide to beer*. Ballantine books.
- Jannasch, H. W. and Egli, T.** (1993). Microbial growth kinetics: a historical perspective. *Antonie Van Leeuwenhoek* **63**, 213-24.
- Jansen, M. L., De Winde, J. H. and Pronk, J. T.** (2002). Hxt-carrier-mediated glucose efflux upon exposure of *Saccharomyces cerevisiae* to excess maltose. *Appl Environ Microbiol* **68**, 4259-65.
- Jiang, R. and Carlson, M.** (1996). Glucose regulates protein interactions within the yeast SNF1 protein kinase complex. *Genes Dev* **10**, 3105-15.

## References

- Jiang, R. and Carlson, M.** (1997). The Snf1 protein kinase and its activating subunit, Snf4, interact with distinct domains of the Sip1/Sip2/Gal83 component in the kinase complex. *Mol Cell Biol* **17**, 2099-106.
- Kacser, H. and Burns, J. A.** (1973). The control of flux. *Symp Soc Exp Biol* **27**, 65-104.
- Klein, C. J., Olsson, L. and Nielsen, J.** (1998). Glucose control in *Saccharomyces cerevisiae*: the role of Mig1 in metabolic functions. *Microbiology* **144** ( Pt 1), 13-24.
- Klein, C. J., Rasmussen, J. J., Ronnow, B., Olsson, L. and Nielsen, J.** (1999). Investigation of the impact of MIG1 and MIG2 on the physiology of *Saccharomyces cerevisiae*. *J Biotechnol* **68**, 197-212.
- Koerkamp, M. G., Rep, M., Bussemaker, H. J., Hardy, G. P., Mul, A., Piekarska, K., Szigyarto, C. A., De Mattos, J. M. and Tabak, H. F.** (2002). Dissection of transient oxidative stress response in *Saccharomyces cerevisiae* by using DNA microarrays. *Mol Biol Cell* **13**, 2783-94.
- Kolkman, A., Daran-Lapujade, P., Fullaondo, A., Olsthoorn, M. M., Pronk, J. T., Slijper, M. and Heck, A. J.** (2006). Proteome analysis of yeast response to various nutrient limitations. *Mol Syst Biol* **2**, 2006 0026.
- Kraakman, L., Lemaire, K., Ma, P., Teunissen, A. W., Donaton, M. C., Van Dijck, P., Winderickx, J., de Winde, J. H. and Thevelein, J. M.** (1999). A *Saccharomyces cerevisiae* G-protein coupled receptor, Gpr1, is specifically required for glucose activation of the cAMP pathway during the transition to growth on glucose. *Mol Microbiol* **32**, 1002-12.
- Kruckeberg, A. L.** (1996). The hexose transporter family of *Saccharomyces cerevisiae*. *Arch Microbiol* **166**, 283-92.
- Lagunas, R.** (1993). Sugar transport in *Saccharomyces cerevisiae*. *FEMS Microbiol Rev* **10**, 229-42.
- Lascaris, R., Piwowarski, J., van der Spek, H., Teixeira de Mattos, J., Grivell, L. and Blom, J.** (2004). Overexpression of HAP4 in glucose-derepressed yeast cells reveals respiratory control of glucose-regulated genes. *Microbiology* **150**, 929-34.
- Lesage, P., Yang, X. and Carlson, M.** (1996). Yeast SNF1 protein kinase interacts with SIP4, a C6 zinc cluster transcriptional activator: a new role for SNF1 in the glucose response. *Mol Cell Biol* **16**, 1921-8.
- Li, C., Hung Wong, W.** (2001). Model-based analysis of oligonucleotide arrays: Expression index computation and outlier detection. *Proc. Natl. Acad. Sci.* **98**, 31-36.
- Lin, S. J., Kaerberlein, M., Andalis, A. A., Sturtz, L. A., Defosse, P. A., Culotta, V. C., Fink, G. R. and Guarente, L.** (2002). Calorie restriction extends *Saccharomyces cerevisiae* lifespan by increasing respiration. *Nature* **418**, 344-8.
- Liu, Y.** (2007). Overview of some theoretical approaches for derivation of the Monod equation. *Appl Microbiol Biotechnol* **73**, 1241-50.
- Maaheimo, H., Fiaux, J., Cakar, Z. P., Bailey, J. E., Sauer, U. and Szyperski, T.** (2001). Central carbon metabolism of *Saccharomyces cerevisiae* explored by biosynthetic fractional (<sup>13</sup>C) labeling of common amino acids. *Eur J Biochem* **268**, 2464-79.



## References

- Marger, M. D. and Saier, M. H., Jr.** (1993). A major superfamily of transmembrane facilitators that catalyse uniport, symport and antiport. *Trends Biochem Sci* **18**, 13-20.
- Marshall-Carlson, L., Celenza, J. L., Laurent, B. C. and Carlson, M.** (1990). Mutational analysis of the SNF3 glucose transporter of *Saccharomyces cerevisiae*. *Mol Cell Biol* **10**, 1105-15.
- Marshall-Carlson, L., Neigeborn, L., Coons, D., Bisson, L. and Carlson, M.** (1991). Dominant and recessive suppressors that restore glucose transport in a yeast *snf3* mutant. *Genetics* **128**, 505-12.
- Matsumoto, K., Uno, I., Oshima, Y. and Ishikawa, T.** (1982). Isolation and characterization of yeast mutants deficient in adenylate cyclase and cAMP-dependent protein kinase. *Proc Natl Acad Sci U S A* **79**, 2355-9.
- Mayordomo, I. and Sanz, P.** (2001). Hexokinase PII: structural analysis and glucose signalling in the yeast *Saccharomyces cerevisiae*. *Yeast* **18**, 923-30.
- Mensonides, F. I., Schuurmans, J. M., Teixeira de Mattos, M. J., Hellingwerf, K. J. and Brul, S.** (2002). The metabolic response of *Saccharomyces cerevisiae* to continuous heat stress. *Mol Biol Rep* **29**, 103-6.
- Monod, J.** (1949). The growth of bacterial cultures. *Ann Rev Microbiol* **3**, 371-349.
- Moreno, F. and Herrero, P.** (2002). The hexokinase 2-dependent glucose signal transduction pathway of *Saccharomyces cerevisiae*. *FEMS Microbiol Rev* **26**, 83-90.
- Moser, A.** (1958). The dynamics of bacterial populations maintained in the chemostat. *volume Publication 614. Washington D.C.: The Carnegie Institute.*
- Nehlin, J. O. and Ronne, H.** (1990). Yeast *MIG1* repressor is related to the mammalian early growth response and Wilms' tumour finger proteins. *Embo J* **9**, 2891-8.
- Neigeborn, L. and Carlson, M.** (1987). Mutations causing constitutive invertase synthesis in yeast: genetic interactions with *snf* mutations. *Genetics* **115**, 247-53.
- Nikawa, J., Sass, P. and Wigler, M.** (1987). Cloning and characterization of the low-affinity cyclic AMP phosphodiesterase gene of *Saccharomyces cerevisiae*. *Mol Cell Biol* **7**, 3629-36.
- Nourani, A., Wesolowski-Louvel, M., Delaveau, T., Jacq, C. and Delahodde, A.** (1997). Multiple-drug-resistance phenomenon in the yeast *Saccharomyces cerevisiae*: involvement of two hexose transporters. *Mol Cell Biol* **17**, 5453-60.
- Oehlen, L. J., Scholte, M. E., de Koning, W. and van Dam, K.** (1994). Decrease in glycolytic flux in *Saccharomyces cerevisiae* *cdc35-1* cells at restrictive temperature correlates with a decrease in glucose transport. *Microbiology* **140** ( Pt 8), 1891-8.
- Otterstedt, K., Larsson, C., Bill, R. M., Stahlberg, A., Boles, E., Hohmann, S. and Gustafsson, L.** (2004). Switching the mode of metabolism in the yeast *Saccharomyces cerevisiae*. *EMBO Rep* **5**, 532-7.
- Ozcan, S., Dover, J. and Johnston, M.** (1998). Glucose sensing and signaling by two glucose receptors in the yeast *Saccharomyces cerevisiae*. *Embo J* **17**, 2566-73.
- Ozcan, S., Dover, J., Rosenwald, A. G., Wolff, S. and Johnston, M.** (1996a). Two glucose transporters in *Saccharomyces cerevisiae* are glucose sensors that generate a signal for induction of gene expression. *Proc Natl Acad Sci U S A* **93**, 12428-32.

## References

- Ozcan, S. and Johnston, M.** (1995). Three different regulatory mechanisms enable yeast hexose transporter (*HXT*) genes to be induced by different levels of glucose. *Mol Cell Biol* **15**, 1564-72.
- Ozcan, S. and Johnston, M.** (1996). Two different repressors collaborate to restrict expression of the yeast glucose transporter genes *HXT2* and *HXT4* to low levels of glucose. *Mol Cell Biol* **16**, 5536-45.
- Ozcan, S. and Johnston, M.** (1999). Function and regulation of yeast hexose transporters. *Microbiol Mol Biol Rev* **63**, 554-69.
- Ozcan, S., Leong, T. and Johnston, M.** (1996b). Rgt1p of *Saccharomyces cerevisiae*, a key regulator of glucose-induced genes, is both an activator and a repressor of transcription. *Mol Cell Biol* **16**, 6419-26.
- Ozcan, S., Schulte, F., Freidel, K., Weber, A. and Ciriacy, M.** (1994). Glucose uptake and metabolism in *grr1/cat80* mutants of *Saccharomyces cerevisiae*. *Eur J Biochem* **224**, 605-11.
- Park, S. H., Koh, S. S., Chun, J. H., Hwang, H. J. and Kang, H. S.** (1999). Nrg1 is a transcriptional repressor for glucose repression of *STA1* gene expression in *Saccharomyces cerevisiae*. *Mol Cell Biol* **19**, 2044-50.
- Peeters, T., Louwet, W., Gelade, R., Nauwelaers, D., Thevelein, J. M. and Versele, M.** (2006). Kelch-repeat proteins interacting with the Galpha protein Gpa2 bypass adenylate cyclase for direct regulation of protein kinase A in yeast. *Proc Natl Acad Sci U S A* **103**, 13034-9.
- Peggie, M. W., MacKelvie, S. H., Bloecher, A., Knatko, E. V., Tatchell, K. and Stark, M. J.** (2002). Essential functions of Sds22p in chromosome stability and nuclear localization of PP1. *J Cell Sci* **115**, 195-206.
- Petit, T., Diderich, J. A., Kruckeberg, A. L., Gancedo, C. and Van Dam, K.** (2000). Hexokinase regulates kinetics of glucose transport and expression of genes encoding hexose transporters in *Saccharomyces cerevisiae*. *J Bacteriol* **182**, 6815-8.
- Postma, E., Scheffers, W. A. and van Dijken, J. P.** (1989). Kinetics of growth and glucose transport in glucose-limited chemostat cultures of *Saccharomyces cerevisiae* CBS 8066. *Yeast* **5**, 159-65.
- Raamsdonk, L. M., Diderich, J. A., Kuiper, A., van Gaalen, M., Kruckeberg, A. L., Berden, J. A., Van Dam, K. and Kruckberg, A. L.** (2001). Co-consumption of sugars or ethanol and glucose in a *Saccharomyces cerevisiae* strain deleted in the *HXK2* gene. *Yeast* **18**, 1023-33.
- Raghevedran, V., Gombert, A. K., Christensen, B., Kotter, P. and Nielsen, J.** (2004). Phenotypic characterization of glucose repression mutants of *Saccharomyces cerevisiae* using experiments with <sup>13</sup>C-labelled glucose. *Yeast* **21**, 769-79.
- Ramaswamy, N. T., Li, L., Khalil, M. and Cannon, J. F.** (1998). Regulation of yeast glycogen metabolism and sporulation by Glc7p protein phosphatase. *Genetics* **149**, 57-72.
- Randez-Gil, F., Bojunga, N., Proft, M. and Entian, K. D.** (1997). Glucose derepression of gluconeogenic enzymes in *Saccharomyces cerevisiae* correlates with phosphorylation of the gene activator Cat8p. *Mol Cell Biol* **17**, 2502-10.

## References

- Reifenberger, E., Boles, E. and Ciriacy, M.** (1997). Kinetic characterization of individual hexose transporters of *Saccharomyces cerevisiae* and their relation to the triggering mechanisms of glucose repression. *Eur J Biochem* **245**, 324-33.
- Reifenberger, E., Freidel, K. and Ciriacy, M.** (1995). Identification of novel *HXT* genes in *Saccharomyces cerevisiae* reveals the impact of individual hexose transporters on glycolytic flux. *Mol Microbiol* **16**, 157-67.
- Rodriguez, A., De La Cera, T., Herrero, P. and Moreno, F.** (2001). The hexokinase 2 protein regulates the expression of the *GLK1*, *HXK1* and *HXK2* genes of *Saccharomyces cerevisiae*. *Biochem J* **355**, 625-31.
- Rohwer, J. M., Meadow, N. D., Roseman, S., Westerhoff, H. V. and Postma, P. W.** (2000). Understanding glucose transport by the bacterial phosphoenolpyruvate:glycose phosphotransferase system on the basis of kinetic measurements in vitro. *J Biol Chem* **275**, 34909-21.
- Rolland, F., Winderickx, J. and Thevelein, J. M.** (2002). Glucose-sensing and -signalling mechanisms in yeast. *FEMS Yeast Res* **2**, 183-201.
- Rossell, S., Lindenbergh, A., van der Weijden, C. C., Kruckeberg, A. L., van Eunen, K., Westerhoff, H. V. and Bakker, B. M.** (2007). Mixed and diverse metabolic and gene-expression regulation of the glycolytic and fermentative pathways in response to a *HXK2* deletion in *Saccharomyces cerevisiae*. *FEMS Yeast Res*.
- Rossell, S., van der Weijden, C. C., Kruckeberg, A. L., Bakker, B. M. and Westerhoff, H. V.** (2005). Hierarchical and metabolic regulation of glucose influx in starved *Saccharomyces cerevisiae*. *FEMS Yeast Res* **5**, 611-9.
- Santangelo, G. M.** (2006). Glucose signaling in *S.cerevisiae*. *Microbiol Mol Biol Rev* **70**, 253-282.
- Sanz, P., Alms, G. R., Haystead, T. A. and Carlson, M.** (2000). Regulatory interactions between the *Reg1-Glc7* protein phosphatase and the *Snf1* protein kinase. *Mol Cell Biol* **20**, 1321-8.
- Sasaki, H. and Uemura, H.** (2005). Influence of low glycolytic activities in *gcr1* and *gcr2* mutants on the expression of other metabolic pathway genes in *Saccharomyces cerevisiae*. *Yeast* **22**, 111-27.
- Schaaff, I., Heinisch, J. and Zimmermann, F. K.** (1989). Overproduction of glycolytic enzymes in yeast. *Yeast* **5**, 285-90.
- Schierbeek, A.** (1963). *Antoni van Leeuwenhoek en zijn voornaamste ontdekkingen*. Kruseman.
- Schuurmans, J. M., Boorsma, A., Lascaris, R., Hellingwerf, K. J. and Teixeira de Mattos, M. J.** (2008). Physiological and transcriptional characterization of *Saccharomyces cerevisiae* strains with modified expression of catabolic regulators. *FEMS Yeast Res* **8**, 26-34.
- Senn, H., Lendenmann, U., Snozzi, M., Hamer, G. and Egli, T.** (1994). The growth of *Escherichia coli* in glucose-limited chemostat cultures: a re-examination of the kinetics. *Biochim Biophys Acta* **1201**, 424-36.
- Small, J. R.** (1994). Design and analysis of chemostat experiments using metabolic control analysis: a top down approach. *Microbiol* **140**, 2439-2449.
- Smith, A., Ward, M. P. and Garrett, S.** (1998). Yeast PKA represses *Msn2p/Msn4p*-dependent gene expression to regulate growth, stress response and glycogen accumulation. *Embo J* **17**, 3556-64.

## References

- Smits, H. P., Hauf, J., Muller, S., Hobley, T. J., Zimmermann, F. K., Hahn-Hagerdal, B., Nielsen, J. and Olsson, L. (2000). Simultaneous overexpression of enzymes of the lower part of glycolysis can enhance the fermentative capacity of *Saccharomyces cerevisiae*. *Yeast* **16**, 1325-34.
- Snoep, J. L., Jensen, P. R., Groeneveld, P., Molenaar, D., Kholodenko, B. N. and Westerhoff, H. V. (1994). How to determine control of growth rate in a chemostat. Using metabolic control analysis to resolve the paradox. *Biochem Mol Biol Int* **33**, 1023-32.
- Snoep, J. L., Westerhoff, H. (2005). *From isolation to integration, a systems biology approach for building the silicon cell*. Springer-Verlag.
- Stock, J. B., Waygood, E. B., Meadow, N. D., Postma, P. W. and Roseman, S. (1982). Sugar transport by the bacterial phosphotransferase system. The glucose receptors of the *Salmonella typhimurium* phosphotransferase system. *J Biol Chem* **257**, 14543-52.
- Stuart, J. S., Frederick, D. L., Varner, C. M. and Tatchell, K. (1994). The mutant type 1 protein phosphatase encoded by *glc7-1* from *Saccharomyces cerevisiae* fails to interact productively with the GAC1-encoded regulatory subunit. *Mol Cell Biol* **14**, 896-905.
- Tanaka, K., Lin, B. K., Wood, D. R. and Tamanoi, F. (1991). IRA2, an upstream negative regulator of RAS in yeast, is a RAS GTPase-activating protein. *Proc Natl Acad Sci U S A* **88**, 468-72.
- Tanaka, K., Matsumoto, K. and Toh, E. A. (1989). IRA1, an inhibitory regulator of the RAS-cyclic AMP pathway in *Saccharomyces cerevisiae*. *Mol Cell Biol* **9**, 757-68.
- Tatchell, K., Chaleff, D. T., DeFeo-Jones, D. and Scolnick, E. M. (1984). Requirement of either of a pair of ras-related genes of *Saccharomyces cerevisiae* for spore viability. *Nature* **309**, 523-7.
- Teissier, G. (1936). Les lois quantitatives de la croissance. *Ann Physiol Physicochim Biol* **12**, 527-586.
- Teusink, B., Diderich, J. A., Westerhoff, H. V., van Dam, K. and Walsh, M. C. (1998). Intracellular glucose concentration in derepressed yeast cells consuming glucose is high enough to reduce the glucose transport rate by 50%. *J Bacteriol* **180**, 556-62.
- Teusink, B., Passarge, J., Reijenga, C. A., Esgalhado, E., van der Weijden, C. C., Schepper, M., Walsh, M. C., Bakker, B. M., van Dam, K., Westerhoff, H. V. and Snoep, J. L. (2000). Can yeast glycolysis be understood in terms of in vitro kinetics of the constituent enzymes? Testing biochemistry. *Eur J Biochem* **267**, 5313-29.
- Thevelein, J. M., Beullens, M., Mbonyi, K. and Van Aelst, L. (1989). The glucose-induced CDC25- and RAS-mediated cAMP signal in the yeast *Saccharomyces cerevisiae*. *Yeast* **5 Spec No**, S421-5.
- Toda, T., Cameron, S., Sass, P., Zoller, M. and Wigler, M. (1987). Three different genes in *S. cerevisiae* encode the catalytic subunits of the cAMP-dependent protein kinase. *Cell* **50**, 277-87.
- Toda, T., Uno, I., Ishikawa, T., Powers, S., Kataoka, T., Broek, D., Cameron, S., Broach, J., Matsumoto, K. and Wigler, M. (1985). In yeast, RAS proteins are controlling elements of adenylate cyclase. *Cell* **40**, 27-36.

## References

- Treitel, M. A., Kuchin, S. and Carlson, M.** (1998). Snf1 protein kinase regulates phosphorylation of the Mig1 repressor in *Saccharomyces cerevisiae*. *Mol Cell Biol* **18**, 6273-80.
- Tu, J. and Carlson, M.** (1995). REG1 binds to protein phosphatase type 1 and regulates glucose repression in *Saccharomyces cerevisiae*. *Embo J* **14**, 5939-46.
- Tu, J., Song, W. and Carlson, M.** (1996). Protein phosphatase type 1 interacts with proteins required for meiosis and other cellular processes in *Saccharomyces cerevisiae*. *Mol Cell Biol* **16**, 4199-206.
- Vallier, L. G. and Carlson, M.** (1994). Synergistic release from glucose repression by mig1 and ssn mutations in *Saccharomyces cerevisiae*. *Genetics* **137**, 49-54.
- Vallier, L. G., Coons, D., Bisson, L. F. and Carlson, M.** (1994). Altered regulatory responses to glucose are associated with a glucose transport defect in grr1 mutants of *Saccharomyces cerevisiae*. *Genetics* **136**, 1279-85.
- van der Werf, M. J., Overkamp, K. M., Mulwijk, B., Coulier, L. and Hankemeier, T.** (2007). Microbial metabolomics: toward a platform with full metabolome coverage. *Anal Biochem* **370**, 17-25.
- Van Hoek, P., Van Dijken, J. P. and Pronk, J. T.** (1998). Effect of specific growth rate on fermentative capacity of baker's yeast. *Appl Environ Microbiol* **64**, 4226-33.
- van Maris, A. J., Bakker, B. M., Brandt, M., Boorsma, A., Teixeira de Mattos, M. J., Grivell, L. A., Pronk, J. T. and Blom, J.** (2001). Modulating the distribution of fluxes among respiration and fermentation by overexpression of *HAP4* in *Saccharomyces cerevisiae*. *FEMS Yeast Res* **1**, 139-49.
- Van Urk, H., Bruinenberg, P. M., Veenhuis, M., Scheffers, W. A. and Van Dijken, J. P.** (1989a). Respiratory capacities of mitochondria of *Saccharomyces cerevisiae* CBS 8066 and *Candida utilis* CBS 621 grown under glucose limitation. *Antonie Van Leeuwenhoek* **56**, 211-20.
- van Urk, H., Postma, E., Scheffers, W. A. and van Dijken, J. P.** (1989b). Glucose transport in crabtree-positive and crabtree-negative yeasts. *J Gen Microbiol* **135**, 2399-406.
- Verduyn, C., Postma, E., Scheffers, W. A. and Van Dijken, J. P.** (1992). Effect of benzoic acid on metabolic fluxes in yeasts: a continuous-culture study on the regulation of respiration and alcoholic fermentation. *Yeast* **8**, 501-17.
- Verduyn, C., Stouthamer, A. H., Scheffers, W. A. and van Dijken, J. P.** (1991). A theoretical evaluation of growth yields of yeasts. *Antonie Van Leeuwenhoek* **59**, 49-63.
- Versele, M., de Winde, J. H. and Thevelein, J. M.** (1999). A novel regulator of G protein signalling in yeast, Rgs2, downregulates glucose-activation of the cAMP pathway through direct inhibition of Gpa2. *Embo J* **18**, 5577-91.
- Vincent, O. and Carlson, M.** (1998). Sip4, a Snf1 kinase-dependent transcriptional activator, binds to the carbon source-responsive element of gluconeogenic genes. *Embo J* **17**, 7002-8.
- Vyas, V. K., Kuchin, S. and Carlson, M.** (2001). Interaction of the repressors Nrg1 and Nrg2 with the Snf1 protein kinase in *Saccharomyces cerevisiae*. *Genetics* **158**, 563-72.

## References

- Walsh, M. C., Smits, H. P., Scholte, M. and van Dam, K.** (1994). Affinity of glucose transport in *Saccharomyces cerevisiae* is modulated during growth on glucose. *J Bacteriol* **176**, 953-8.
- Wang, Y., Pierce, M., Schnepfer, L., Guldal, C. G., Zhang, X., Tavazoie, S. and Broach, J. R.** (2004). Ras and Gpa2 mediate one branch of a redundant glucose signaling pathway in yeast. *PLoS Biol* **2**, E128.
- Wendell, D. L. and Bisson, L. F.** (1994). Expression of high-affinity glucose transport protein Hxt2p of *Saccharomyces cerevisiae* is both repressed and induced by glucose and appears to be regulated posttranslationally. *J Bacteriol* **176**, 3730-7.
- Westerhoff, H. V., Lolkema, J. S., Otto, R. and Hellingwerf, K. J.** (1982). Thermodynamics of growth. Non-equilibrium thermodynamics of bacterial growth. The phenomenological and the mosaic approach. *Biochim Biophys Acta* **683**, 181-220.
- Wu, L., Mashego, M. R., van Dam, J. C., Proell, A. M., Vinke, J. L., Ras, C., van Winden, W. A., van Gulik, W. M. and Heijnen, J. J.** (2005). Quantitative analysis of the microbial metabolome by isotope dilution mass spectrometry using uniformly <sup>13</sup>C-labeled cell extracts as internal standards. *Anal Biochem* **336**, 164-71.
- Xue, Y., Battle, M. and Hirsch, J. P.** (1998). GPR1 encodes a putative G protein-coupled receptor that associates with the Gpa2p Galpha subunit and functions in a Ras-independent pathway. *Embo J* **17**, 1996-2007.
- Young, E. T., Dombek, K. M., Tachibana, C. and Ideker, T.** (2003). Multiple pathways are co-regulated by the protein kinase Snf1 and the transcription factors Adr1 and Cat8. *J Biol Chem* **278**, 26146-58.
- Zakrzewska, A., Boorsma, A., Brul, S., Hellingwerf, K. J. and Klis, F. M.** (2005). Transcriptional response of *Saccharomyces cerevisiae* to the plasma membrane-perturbing compound chitosan. *Eukaryot Cell* **4**, 703-15.
- Zenke, F. T., Engles, R., Vollenbroich, V., Meyer, J., Hollenberg, C. P. and Breunig, K. D.** (1996). Activation of Gal4p by galactose-dependent interaction of galactokinase and Gal80p. *Science* **272**, 1662-5.
- Zhu, X., Demolis, N., Jacquet, M. and Michaeli, T.** (2000). MSI1 suppresses hyperactive RAS via the cAMP-dependent protein kinase and independently of chromatin assembly factor-1. *Curr Genet* **38**, 60-70.
- Zimmermann, F. K. and Scheel, I.** (1977). Mutants of *Saccharomyces cerevisiae* resistant to carbon catabolite repression. *Mol Gen Genet* **154**, 75-82.