UvA-DARE (Digital Academic Repository)

The authors' reply

Verouden, N.J.; de Winter, R.J.

DOI

[10.1136/hrt.2009.181222](https://doi.org/10.1136/hrt.2009.181222)

Publication date

2009

Document Version

Final published version

Published in

Heart

[Link to publication](#)

Citation for published version (APA):

Verouden, N. J., & de Winter, R. J. (2009). The authors' reply. *Heart*, 95(23), 1952-1953. <https://doi.org/10.1136/hrt.2009.181222>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Featured correspondence

side branch) treatable with percutaneous coronary intervention with a favourable outcome.^{4,5} The limited area of ischaemia has been proposed as the basis for the discordance between the ST and T vectors.

Particular subtlety is recommended when introducing new observations in the scientific community.

M J Eskola,¹ K C Nikus,¹ S Sclarovsky²

¹ Heart Center, Tampere University Hospital, Tampere, Finland; ² Procardia Medical Center, Tel Aviv, Israel

Correspondence to: Dr K C Nikus, Heart Center, Tampere University Hospital, Tampere, Finland; kjell.nikus@pshp.fi

Competing interests: None.

Provenance and peer review: Not commissioned; not externally peer reviewed.

Heart 2009;**95**:1951–1952. doi:10.1136/hrt.2009.181214

REFERENCES

1. Verouden NJ, Koch KT, Peters RJ, *et al*. Persistent precordial “hyperacute” T-waves signify proximal LAD artery occlusion. *Heart* 2009;**95**:1701–6.
2. Dressler W, Roesler H. High T waves in the earliest stage of myocardial infarction. *Am Heart J* 1947;**34**:627–46.
3. Sclarovsky S, ed. Angina at rest and acute myocardial ischemia. In: *Electrocardiography of acute myocardial ischaemic syndromes*. London, UK: Martin Dunitz Ltd, 1999:1–29.
4. Sclarovsky S, Rechavia E, Strasberg B, *et al*. Unstable angina: ST segment depression with positive versus negative T wave deflections – clinical course, ECG evolution, and angiographic correlation. *Am Heart J* 1988;**116**:933–41.
5. Nikus KC, Eskola MJ, Virtanen VK, *et al*. ST-Depression with negative T waves in leads V4–V5 – a marker of severe coronary artery disease in non-ST elevation acute coronary syndrome: a prospective study of angina at rest, with troponin, clinical, electrocardiographic, and angiographic correlation. *Ann Noninvasive Electrocardiol* 2004;**9**:1–8.

The authors’ reply: With interest we read the comment by Eskola and colleagues¹ on our systematic observation of a characteristic ECG pattern associated with acute and persistent occlusion of the proximal left anterior descending (LAD) artery.² The pattern of “hyperacute T waves” has been described before, but has invariably been described previously as a transient phenomenon associated with anterior ischaemia, but not with persistent LAD occlusion. With careful serial observation and simultaneous coronary angiography, we have shown, in contrast to previous reports, that these ECG changes may be associated with persistent LAD occlusion.

We are grateful for the reference by Dressler and Roesler,³ which was overlooked by us. Indeed, in the report by Dressler and Roesler,³ the ECG pattern of case number 4 recorded 3 h after the onset of symptoms is very similar to our description, and the clinical history and subsequent follow-up ECG strongly suggest that this patient had persistent LAD occlusion instead of transient LAD occlusion (see fig 1). In the second report, Sclarovsky *et al*⁴ published clinical and angiographic data on 21 patients

Figure 1 ST-segment depression and peaked, positive T waves followed by loss of R wave and inverted T waves one day later. With permission from the *American Heart Journal*.

presenting with precordial ST-segment depression and peaked, positive T waves. Importantly, the ECG pattern was described as a transient feature in that study and did not result in myocardial necrosis in over 60% of the patients. In patients with elevated cardiac enzymes, the investigators observed a transition to overt ST-segment elevation on evolving ECG. In all but one patient, a non-occlusive stenosis

of the LAD artery was observed during coronary angiography.

We would like to emphasise three pivotal observations described in the current issue of this journal,² which sharply contrast with the above-mentioned reports. We described an ECG pattern of precordial J-point depression and tall, positive T waves that remained unchanged from first medical contact until arrival at the catheterisa-

tion laboratory and did not convert to overt ST-segment elevation. Furthermore, the majority of our patients showed a total occlusion of the proximal LAD artery and, despite successful primary percutaneous coronary intervention in all cases, had a considerable loss of myocardium (median myocardial type creatine kinase 290 µg/l). This seems in contradiction to previous suggestions that ST-segment depression and tall positive T waves in the precordial leads are associated with regional subendocardial ischaemia⁵ and a favourable outcome.⁴ Therefore, we think it is important to emphasise that in some patients, this novel ECG pattern may be associated with persistent proximal LAD artery occlusion and transmural ischaemia of the anterior myocardium. These patients must be distinguished from patients with regional subendocardial ischaemia and be referred for immediate reperfusion therapy.

N J Verouden, R J de Winter

Department of Cardiology of the Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

Correspondence to: Dr R J de Winter, Department of Cardiology, B2-137, Academic Medical Center, Meibergdreef 9, PO box 22660, 1100 DD Amsterdam, The Netherlands; r.j.dewinter@amc.uva.nl

Competing interests: None.

Provenance and peer review: Not commissioned; not externally peer reviewed.

Heart 2009;**95**:1952–1953. doi:10.1136/hrt.2009.181222

REFERENCES

1. **Eskola MJ**, Nikus KC, Sclarovsky S. Persistent precordial “hyper acute” T waves signify proximal left anterior descending artery occlusion. *Heart* 2009;**95**:1701–6.
2. **Verouden NJ**, Koch KT, Peters RJ, *et al.* Persistent precordial “hyper acute” T-waves signify proximal LAD artery occlusion. *Heart* 2009;**95**:1701–6.
3. **Dressler W**, Roesler H. High T waves in the earliest stage of myocardial infarction. *Am Heart J* 1947;**34**:627–46.

4. **Sclarovsky S**, Rechavia E, Strasberg B, *et al.* Unstable angina: ST segment depression with positive versus negative T wave deflections—clinical course, ECG evolution, and angiographic correlation. *Am Heart J* 1988;**116**:933–41.
5. **Sclarovsky S**, ed. Angina at rest and acute myocardial infarction. In: *Electrocardiography of acute myocardial ischaemic syndromes*. London: Martin Dunitz Ltd, 1999:1–29.

CORRECTION

doi:10.1136/hrt.2009.179390corr1

Wackers J. Chest pain in the emergency department: role of cardiac imaging. The author's reply. *Heart* 2009;**95**:1802. In the third paragraph, the first sentence should read “I believe though that there is a place for exercise ECG in the evaluation of patients in an ED chest pain centre (CPC).” The journal apologises for the error which has been corrected online.

The authors' reply:

N J Verouden and R J de Winter

Heart 2009 95: 1952-1953

doi: 10.1136/hrt.2009.181222

Updated information and services can be found at:

<http://heart.bmj.com/content/95/23/1952.full.html>

These include:

References

This article cites 4 articles, 2 of which can be accessed free at:

<http://heart.bmj.com/content/95/23/1952.full.html#ref-list-1>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Interventional cardiology](#) (7282 articles)

[Percutaneous intervention](#) (495 articles)

[Clinical diagnostic tests](#) (19854 articles)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>