



UvA-DARE (Digital Academic Repository)

Three-manifold quantum invariants and mock theta functions

Cheng, M.C.N.; Ferrari, F.; Sgroi, G.

DOI

[10.1098/rsta.2018.0439](https://doi.org/10.1098/rsta.2018.0439)

Publication date

2019

Document Version

Submitted manuscript

Published in

Philosophical transactions. Series A, Mathematical, physical, and engineering sciences

[Link to publication](#)

Citation for published version (APA):

Cheng, M. C. N., Ferrari, F., & Sgroi, G. (2019). Three-manifold quantum invariants and mock theta functions. *Philosophical transactions. Series A, Mathematical, physical, and engineering sciences*, 378(2163), [0439]. <https://doi.org/10.1098/rsta.2018.0439>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

THREE-MANIFOLD QUANTUM INVARIANTS AND MOCK THETA FUNCTIONS

Miranda C. N. Cheng^{*1,2}, Francesca Ferrari^{3,4}, and Gabriele Sgroi^{†2}

¹Korteweg-de Vries Institute for Mathematics, Amsterdam, the Netherlands

²Institute of Physics, University of Amsterdam, Amsterdam, the Netherlands

³International School for Advanced Studies (SISSA), Trieste, Italy

⁴INFN, Sezione di Trieste, Trieste, Italy

Abstract

Mock modular forms have found applications in numerous branches of mathematical sciences since they were first introduced by Ramanujan nearly a century ago. In this proceeding we highlight a new area where mock modular forms start to play an important role, namely the study of three-manifold invariants. For a certain class of Seifert three-manifolds, we describe a conjecture on the mock modular properties of a recently proposed quantum invariant. As an illustration, we include concrete computations for a specific three-manifold, the Brieskorn sphere $\Sigma(2, 3, 7)$. This note is partially based on the talk by the first author in the conference “Srinivasa Ramanujan: in celebration of the centenary of his election as FRS” held at the Royal Society in 2018.

*On leave from CNRS, France.

†g.sgroi@uva.nl

Contents

1	Introduction	2
2	False, Mock, and Three-Manifolds	5
2.1	False Theta Functions and Negative Three-Star Graphs	5
2.2	False, Mock, and Quantum	8
2.3	A Mock Conjecture	11
3	Explicit Calculations	13
3.1	Quantum Invariants via Knots	13
3.2	Relation to indefinite theta functions	14

1 Introduction

Mock modular forms have been a source of fascination since Ramanujan first introduced them about a century ago [Ram88], [Ram00]. In 2002, Zwegner’s doctoral thesis provided the crucial starting point for a structural theory of mock modular forms. These functions have found applications in various branches of mathematical sciences, including combinatorics, moonshine, conformal field theory, string theory and more. We refer to [Zag10a], [Ono09], [Fol10], [Duk14], [CDH14a], [CDH14b], [BFOR17] and other contributions to this volume for a partial overview of these developments. The purpose of this article is to highlight the appearance of mock modular forms in a different context: the topology of three-dimensional manifolds. This appearance was first anticipated in [CCFGH18] and later further developed via concrete computations in [GM19] and [CS].

To see how mock modular forms appear in the study of three-manifolds, we first introduce a set of topological invariants, noted by \widehat{Z}_a , defined in [GPPV17] for *weakly negative plumbed manifolds*. Roughly speaking, these are three-manifolds obtained through surgeries along links that are in turn determined by weighted graphs (cf. Figure 1), which moreover satisfy a certain negativity condition [CCFGH18].

More precisely, the data we need is a plumbing graph, which is a a weighted simple graph (V, E, λ) specified by the set V of vertices, the set E of edges, and an integral weight function $\lambda : V \rightarrow \mathbb{Z}$. Equivalently, the data can be captured by an adjacency matrix M , which is a square matrix of size $|V|$ with entries given by $M_{vv'} = \lambda(v)$ if $v = v'$, 1 if $(v, v') \in E$ and 0 otherwise. The data determines a three-manifold¹. We say that M_3 is a weakly negative plumbed manifold if M^{-1} is negative-definite when restricted to the subspace generated by all vertices with degree larger than 2.

Definition 1.1. For M_3 a weakly negative plumbed three-manifold and using the above notation, we define the quantum invariants $\widehat{Z}_a(M_3; \tau)$ via the following principal value $|V|$ -dimensional integral:

$$\widehat{Z}_a(M_3; \tau) := (-1)^\pi q^{\frac{3\sigma - \sum_{v \in V} \lambda(v)}{4}} \sum_{\mathbf{n} \in 2M\mathbb{Z}^{|V|+\mathbf{a}}} \text{vp} \prod_{v \in V} \oint_{|w_v|=1} \frac{dw_v}{2\pi i w_v} \left(w_v - \frac{1}{w_v} \right)^{2-\deg(v)} q^{-\frac{\mathbf{n}^T M^{-1} \mathbf{n}}{4}} e^{2\pi i \mathbf{z}^T \mathbf{n}} \tag{1.1}$$

¹Different weighted graph related by the so-called Kirby moves can lead to the same topological three-manifold M_3 . See [GM19] for a proof the Kirby-invariance of the quantum invariants $\widehat{Z}_a(M_3)$ defined in Definition 1.1.




Figure 1: Weighted graphs, adjacency matrices, links and plumbed three-manifolds.

where we write $q := e^{2\pi i\tau}$ and $w_v := e^{2\pi iz_v}$ as usual, and use the bold-faced letters to denote elements in $\mathbb{Z}^{|V|}$. When M^{-1} is moreover negative definite, the above can be rewritten as

$$\widehat{Z}_a(M_3; \tau) := (-1)^\pi q^{\frac{3\sigma - \sum_{v \in V} \lambda(v)}{4}} \text{vp} \prod_{v \in V} \oint_{|w_v|=1} \frac{dw_v}{2\pi i w_v} \left(w_v - \frac{1}{w_v} \right)^{2 - \deg(v)} \Theta_a^{-M}(\tau, \mathbf{z}) \quad (1.2)$$

In the above, π denotes the number of positive eigenvalues, and σ is the signature of M^{-1} . The label a of the quantum invariants $\widehat{Z}_a(M_3)$ can be identified with elements of the set $\text{Spin}^c(Y) \cong \pi_0 \mathcal{M}_{\text{ab}}(M_3) \cong (2\mathbb{Z}^{|V|} + \delta)/(2M\mathbb{Z}^{|V|})$, where $\delta \in \mathbb{Z}^{|V|}/2\mathbb{Z}^{|V|}$ is defined by $\delta_v = \deg(v) \bmod 2$, and $\mathcal{M}_{\text{ab}}(M_3)$ denotes the moduli space of Abelian flat connections. Denote by \mathbf{a} the corresponding element of $(2\mathbb{Z}^{|V|} + \delta)/(2M\mathbb{Z}^{|V|})$, the theta function reads

$$\Theta_a^{-M}(\tau, \mathbf{z}) = \sum_{\mathbf{n} \in 2M\mathbb{Z}^{|V|} + \mathbf{a}} q^{-\frac{\mathbf{n}^T M^{-1} \mathbf{n}}{4}} e^{2\pi i \mathbf{z}^T \mathbf{n}}. \quad (1.3)$$

A well-known topological invariant for three-manifolds is the Witten-Reshetikhin-Turaev (WRT) invariant, defined for all three-manifolds. Physically speaking, it is (up to certain well-understood prefactors) the partition function of Chern-Simons theory on the three-manifold M_3 which we denote by $Z_{\text{CS}}(M_3)$. For a given three-manifold M_3 (and a simple Lie group G which we will take to be $G = SU(2)$ for the sake of concreteness), we obtain a function $Z_{\text{CS}}(M_3) : \mathbb{Z} \rightarrow \mathbb{C}$ defined on all integers, namely the (shifted) Chern-Simons levels. Analogous to knot theory, it would be desirable to have a q -series version of the invariants defined on a continuous domain, such as the upper-half plane. This would then be the first step towards a categorification of three-manifold invariants, analogous to the categorification programme of knot invariants. It was shown for weakly negative plumbed manifolds that the quantum invariants \widehat{Z}_a provide exactly such a q -series generalisation of

the WRT invariants. To be more precise, for weakly negative plumbed manifolds one has [GPPV17]

$$(i\sqrt{2k})Z_{\text{CS}}(M_3; k) = \sum_{a,b} X_{ab} e^{2\pi i \text{CS}(a)k} \lim_{\tau \rightarrow \frac{1}{k}} \widehat{Z}_b(M_3; \tau). \quad (1.4)$$

In the above equation, the sum over a is over the set $((2\mathbb{Z}^{|V|} + \delta)/2M\mathbb{Z}^{|V|})/\mathbb{Z}_2$, which can be identified with the space of gauge-inequivalent $SU(2)$ Abelian flat connections on M_3 , and $\text{CS}(a)$ denotes the corresponding Chern-Simons invariant when we regard a as a label for Abelian $SU(2)$ flat connections. The sum over b is over the set $(\mathbb{Z}^{|V|}/M\mathbb{Z}^{|V|})/\mathbb{Z}_2$, and the matrix X has as elements

$$X_{ab} = \frac{\sum_{(a',b') \in \{\mathbb{Z}_2 \times \mathbb{Z}_2 \text{ orbit of } (a,b)\}} e^{2\pi i(a', M^{-1}b')}}{2\sqrt{|\text{Det}M|}}. \quad (1.5)$$

To summarise, two steps need to be taken in order to retrieve Z_{CS} from \widehat{Z}_a . First, \widehat{Z}_a has an extra label a indexing the $SU(2)$ Abelian flat connections while Z_{CS} does not, and this label needs therefore to be summed over. Second, a so-called radial limit $\tau \rightarrow \frac{1}{k}$ taking $\tau \in \mathbb{H}$ to the boundary $\mathbb{Q} \cup \{i\infty\}$ of the upper-half plane needs to be taken in order to relate the continuous variable τ and the (shifted) Chern-Simons level.

The modular-like properties of the quantum invariants \widehat{Z}_a is a rich subject that has been in development since [CCFGH18]. So far it develops in parallel to the study of modular-like properties of knot invariants (see for instance [Zag10b], [GZ18], [DG18], [HL15] for a sample of work in this direction), although it is expected that the two topics are related both in their physical and mathematical contexts.

For concreteness and in order to make direct contact with Ramanujan's mock theta functions, here we restrict our attention to the simplest non-trivial plumbing graphs: the so-called three-star weighted graphs. These are, as the name suggests, weighted simple graphs with one vertex of degree three, three vertices of degree one, while the rest of the vertices (if any) have degree two. See Figure 2. We will denote the unique vertex with degree three by v_0 . Such graphs are either weakly negative or not, depending on the sign of $(M^{-1})_{00}$. When $(M^{-1})_{00} < 0$, Definition 1.1 is readily applicable and it is not hard to show that the quantum invariants \widehat{Z}_a are always holomorphic functions on the upper-half plane with well-defined q -expansions and moreover have integral coefficients. In fact, a lot more is true: up to a possible addition of a polynomial, the quantum invariants \widehat{Z}_a are linear combinations of false theta functions multiplied by a rational q -power (cf. §2.1 and [BMM18]).

A puzzle immediately arises given the simple result for weakly negative three-star graphs: what happens when one flips the orientation of the three-manifold? While this might sound like an innocuous operation, it can in fact have rather dramatic consequences due to the *pseudo-chiral symmetry* (or CP symmetry in physical terms)

$$Z_{\text{CS}}(M_3; k) = Z_{\text{CS}}(-M_3; -k) \quad (1.6)$$

of Chern-Simons theory. From the relation (1.4) between the quantum invariants $\widehat{Z}_a(M_3; \tau)$ and

$Z_{\text{CS}}(M_3; k)$, and in particular the relation “ $\tau \rightarrow \frac{1}{k}$ ” between the two variables, one is led to the guess

$$\widehat{Z}_a(-M_3; \tau) \text{ “=” } \widehat{Z}_a(M_3; -\tau). \quad (1.7)$$

There are a few immediate problems with this guess. Recall that for a weakly negative plumbed manifold M_3 , Definition 1.1 defines a function $\widehat{Z}_a(M_3; \tau)$ on the upper-half plane \mathbb{H} , which is not preserved by the action $\tau \mapsto -\tau$. As a result it is not clear what the right-hand side of the equation (1.7) even means. More concretely, it is clear from (1.1) that for plumbed manifolds one has $\tau \mapsto -\tau \Leftrightarrow M \mapsto -M$, which flips the sign of the adjacency matrix and hence flips the signature of the lattice for which the theta function Θ_a^{-M} should be defined, and as a result does not render a function on \mathbb{H} when one tries to literally apply Definition 1.1.

This is when the question starts to become interesting from the perspective of mock modular forms. To be concrete, we let M_3 be a weakly negative three-star plumbed three-manifold. As mentioned before, for such cases $\widehat{Z}_a(M_3; \tau)$ are basically false theta functions, which are known to furnish (rather simple) examples of the so-called *quantum modular forms*, as will be explained in §2. The quantum modular properties of the quantum invariants \widehat{Z}_a are essentially what makes their relation (1.4) to Z_{CS} possible. At the same time, it can be shown that a mock theta function and the corresponding false theta function lead to a pair of quantum modular forms that are in fact basically equivalent (cf. Lemma 2.6), in a way that precisely leads to the radial limit relation (1.6). This leads to the natural guess that the quantum invariants $\widehat{Z}_a(-M_3; \tau)$ for the orientation-reversed three-star plumbed manifold are given by mock theta functions. This conjecture, proposed in [CCFGH18], will be discussed in §2.7.

In §3, we will review some recent results supporting the conjecture. The first involves building the relevant orientation-reversed three-manifold via Dehn surgeries on knot complements [GM19], and the second involves employing the indefinite theta series to extend the definition of $\widehat{Z}_a(M_3)$ to general plumbed manifolds [CS]. To illustrate the various ideas discussed in this proceeding, we will discuss in details the specific example of the Brieskorn sphere $M_3 = \Sigma(2, 3, 7)$.

2 False, Mock, and Three-Manifolds

In this section we argue that mock modular forms play a role in three-manifold quantum invariants. In §2.1 we introduce the relevant class of quantum invariants and review their relation to false theta functions. In §2.2 we review the quantum modular properties of false and mock theta functions and explain their relevance for three-manifold topology. In §2.3 we discuss a mock conjecture for \widehat{Z}_a and its motivation and consequences.

2.1 False Theta Functions and Negative Three-Star Graphs

For concreteness, we focus on the simplest type of non-trivial graph: the three-star graphs (see Figure 2). These type of graphs correspond via plumbing (cf. Figure 1) to Seifert manifolds with three singular fibers. The relation between false theta functions and the WRT invariants for this family of three-manifolds was first pointed out in [LZ99] and later extensively studied in [Hik05a], [Hik05b],

[Hik11]. Here we are interested in their quantum invariants $\widehat{Z}_a(M_3)$. It is easy to see [CCFGH18] that Definition 1.1 leads to a function well-defined on \mathbb{H} if and only if $(M^{-1})_{00} < 0$, namely when the resulting plumbed three-manifold M_3 is weakly negative.




Figure 2: A three-star graph.

In order to describe the quantum modular properties of $\widehat{Z}_a(M_3)$, we will need the following definitions.

Definition 2.1. Let $m \in \mathbb{Z}_{>0}$ and $r \in \mathbb{Z}/2m$. Define false theta functions

$$\widetilde{\theta}_{m,r}^1(\tau) := \sum_{\substack{k \in \mathbb{Z} \\ k \equiv r \pmod{2m}}} \operatorname{sgn}(k) q^{\frac{k^2}{4m}}. \quad (2.1)$$

Note that this is nothing but the usual theta function for one-dimensional lattice $\sqrt{2m}\mathbb{Z}$ when the sign factor in the summand is removed. This leads to the nomenclature [Rog17, Fin88].

It will also be convenient to define, after [BMM18], the following functions for $m \in \mathbb{Z}_{>0}$ and $j \in \mathbb{Z}$

$$F_{j,m}(\tau) := \sum_{k \in \mathbb{Z}} \operatorname{sgn}\left(k + \frac{1}{2}\right) q^{\left(k + \frac{j}{2m}\right)^2}. \quad (2.2)$$

Furthermore, we have

$$F_{j,m}(m\tau) = \widetilde{\theta}_{m,j}^1(\tau) + p_{m,j}(\tau), \quad (2.3)$$

where $p_{m,j}(\tau)$ is the polynomial in q given by

$$p_{m,j}(\tau) = \begin{cases} -2 \sum_{k=1}^{\lfloor \frac{j}{2m} \rfloor} q^{\frac{(j-2mk)^2}{4m}} & , \text{ if } j \geq 2m \\ 0 & , \text{ if } 0 \leq j < 2m \\ 2 \sum_{k=0}^{-\lfloor \frac{j}{2m} \rfloor - 1} q^{\frac{(j+2mk)^2}{4m}} & , \text{ if } j < 0. \end{cases} \quad (2.4)$$

Note that the definition (2.2) can be extended to $m, j \in \mathbb{Q}^*$ since the right hand side only depends on their ratio $\frac{j}{m}$.

In terms of the above building blocks, it can be shown that given a 3-star weighted graph, the corresponding $\widehat{Z}_a(M_3)$ can be written in terms of $F_{j,p}$ for some p and j . See Theorem 4.2 in [BMM18] for the result on a closely related quantity, denoted $Z(q)$ in [BMM18], and [CCFGH18] for numerous examples.

In particular, in what follows we will further restrict our attention to weakly negative plumbed manifolds with three-star plumbing graphs with four nodes. Denote by M the corresponding adjacency matrix, let $A := -\frac{1}{2}M^{-1}$ and let v_0 be the unique vertex with degree three. Moreover, assume that the corresponding adjacency matrix M is unimodular. As a result there is only one quantum invariant $\widehat{Z}_0(M_3; \tau) := \widehat{Z}_a(M_3; \tau)$ with $a = \delta \pmod{2M\mathbb{Z}^{|V|}}$, as defined in 1.1. Write also

$$\begin{aligned} m &= 2A_{00} \\ b_0 &= 2 \sum_{j=1}^3 A_{j0}, \quad b_i = 4A_{i0} - 2 \sum_{j=1}^3 A_{j0} \\ c_0 &= A_{12} + A_{23} + A_{31} + \frac{1}{2} \sum_{j=1}^3 A_{jj}, \quad c_i = c_0 - 2 \sum_{\substack{j \in \{1,2,3\} \\ j \neq i}} A_{ij} \end{aligned} \tag{2.5}$$

for $i = 1, 2, 3$. Note that $d_i := -\frac{b_i^2}{4m} + c_j$ satisfy $d_i = d_j =: d$ for all $i, j \in \{0, 1, 2, 3\}$. In the above notation we have the following Proposition.

Proposition 2.2. [BMM18] *Consider a weakly negative three-star plumbing graph with four nodes and unimodular adjacency matrix, denote by M_3 the corresponding plumbed three-manifold. Its unique quantum invariant satisfies*

$$(-1)^\pi q^{-c} \widehat{Z}_0(M_3; \tau) = \sum_{j=0}^3 F_{m-b_j, m}(m\tau) \tag{2.6}$$

with $c = d + \frac{3\sigma - \sum_v m(v)}{4}$, where m, b_j and d are defined as above and where σ and π as defined as in Definition 1.1.

Note that, using (2.3) this immediately shows

$$q^{-c} \widehat{Z}_0(M_3; \tau) = \sum_{j=1}^4 \theta_{m, m-b_j}^1(\tau) + p(\tau) \tag{2.7}$$

where $p(\tau)$ is a polynomial which one can work out explicitly using (2.3). Often times, one has $-m < b_j \leq m$ for all $j \in \{0, 1, 2, 3\}$ and $p(\tau) = 0$. In other words, up to an overall rational power of q and possibly the addition of a polynomial, the quantum invariants \widehat{Z}_0 is given by a false theta function.

Example. In this section, we will illustrate the computation of the quantum invariant and in particular Proposition 2.2, with the example of the Brieskorn sphere $M_3 = \Sigma(2, 3, 7)$, which can be described as the intersection between the algebraic surface $\{x^2 + y^3 + z^7 = 0\}$ and the five sphere $\{|x|^2 + |y|^2 + |z|^2 = 1\}$. It can be obtained as a plumbed manifold with the plumbing graph shown in Figure 3. Note that M is indeed unimodular, consistent with the fact that Brieskorn sphere is a integral homology sphere with trivial $H_1(M_3; \mathbb{Z}) \cong \mathbb{Z}^4/M\mathbb{Z}^4$ and there is hence just one quantum invariant $\widehat{Z}_0(M_3; \tau)$.

Plugging the adjacency matrix in (2.5) one obtains

$$m = 42, \quad (b_j, 4c_j) = (1, 1), (-13, 5), (-29, 21), (41, 41) \text{ for } j = 0, 1, 2, 3. \tag{2.8}$$




Figure 3: Plumbing graph and adjacency matrix for $\Sigma(2, 3, 7)$

Using Definition 1.1 (or (2.6)), one obtains that

$$\begin{aligned} \widehat{Z}_0(\Sigma(2, 3, 7); \tau) &= q^{\frac{83}{168}} \sum_{\substack{k \geq 0 \\ k^2 \equiv 1 \pmod{42}}} \binom{k}{21} q^{\frac{k^2}{168}} \\ &= q^{\frac{83}{168}} \left(\widetilde{\theta_{42,1}^1} - \widetilde{\theta_{42,13}^1} - \widetilde{\theta_{42,29}^1} + \widetilde{\theta_{42,41}^1} \right) (\tau) \\ &= q^{\frac{1}{2}} (1 - q - q^5 + q^{10} - q^{11} + \dots) \end{aligned} \tag{2.9}$$

which is indeed a false theta function of weight $1/2$. The fact that $Z_{\text{CS}}(\Sigma(2, 3, 7))$ is given by the above function by taking the limit

$$Z_{\text{CS}}(\Sigma(2, 3, 7); k) = \frac{1}{i\sqrt{2k}} \lim_{t \rightarrow 0^+} (\widehat{Z}_0(\Sigma(2, 3, 7); \frac{1}{k} + t)) \tag{2.10}$$

was first established by [LZ99].

2.2 False, Mock, and Quantum

As we have seen, a pre-requisite for a q -series to play the role of the quantum invariants $\widehat{Z}_a(M_3)$ is to have a specific behaviour when taking the radial limit, so that it gives the WRT invariants via (1.4). This is demonstrated in the $\Sigma(2, 3, 7)$ example in (2.10). This leads us to the concept of *quantum modular forms* (QMF), first introduced by D. Zagier [Zag10b]. Roughly speaking, a quantum modular form is a function defined on \mathbb{Q} with a certain modular-like property: the deviation from modularity, measured by a modular difference function denoted by p_γ , has nice analytic properties that are not a priori manifest or expected. In this proceeding we work with a specific version of the definition that is often referred to as defining *strong quantum modular forms*. We refer to §7.3 of [CCFGH18] for details about modular forms in the current context.

To state the definition, let us recall the usual definition of the slash operator acting on the space of holomorphic functions on \mathbb{H} for weight w and multiplier χ on Γ , which we take to be a subgroup of the modular group $SL_2(\mathbb{Z})$:

$$f(\tau)|_{w, \chi \gamma} := f\left(\frac{a\tau + b}{c\tau + d}\right) \chi(\gamma) (c\tau + d)^{-w}, \quad \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma. \tag{2.11}$$

Definition 2.3. [Zag10b] Consider a function $Q : \mathbb{Q} \rightarrow \mathbb{C}$. It is called a *strong quantum modular*

form of weight w and multiplier χ for Γ if for every $\gamma \in \Gamma$ the modular difference function $p_\gamma(x) : \mathbb{Q} \setminus \{\gamma^{-1}(\infty)\} \rightarrow \mathbb{C}$, defined by

$$p_\gamma(x) := Q(x) - Q|_{w,\chi}\gamma(x) \quad (2.12)$$

is a real-analytic function of \mathbb{R} minus finitely many points.

The false theta functions we encountered in §2.1 are examples of quantum modular forms. The simplest way to see this is to note that false theta functions defined in (2.1) are examples of Eichler integrals. Given a cusp form $g = \sum_{n>0} a_g(n)q^n$ of weight $w \in \frac{1}{2}\mathbb{Z}$, its Eichler integral is defined as

$$\tilde{g}(\tau) := \sum_{n>0} n^{1-w} a_g(n) q^n. \quad (2.13)$$

It is easy to see that the false theta function $\widetilde{\theta_{m,r}^1}$ is the Eichler integral of the weight $3/2$ unary theta function

$$\theta_{m,r}^1(\tau) := \frac{1}{\sqrt{4m}} \sum_{\substack{\ell \in \mathbb{Z} \\ \ell \equiv r \pmod{2m}}} \ell q^{\ell^2/4m}, \quad (2.14)$$

as the notation suggests.

Note that this is equal to the following integral²

$$\tilde{g}(\tau) = C \int_{\tau}^{i\infty} g(z')(z' - \tau)^{w-2} dz', \quad (2.15)$$

where $C = \frac{(2\pi i)^{w-1}}{\Gamma(w-1)}$. Letting $Q(x) := \lim_{t \rightarrow 0^+} \tilde{g}(x + it)$ in Definition 2.3, one immediately sees that the modular difference function $p_\gamma(x)$ admits an expression as a period integral

$$p_\gamma(x) = C \int_{\gamma^{-1}(i\infty)}^{i\infty} g(z')(z' - x)^{w-2} dz' \quad (2.16)$$

and is hence equipped with the desired analytic properties.

An analogous argument demonstrates that mock modular forms also lead to quantum modular forms. To see that, we first recall the definition of mock modular forms, adapted to the classes of functions that are relevant for our context. In particular, we assume that the shadow g is a cusp form with real Fourier coefficients, namely $\overline{g(-\bar{\tau})} = g(\tau)$.

Definition 2.4. We say that a holomorphic function f on \mathbb{H} is a *mock modular form* of weight k and multiplier χ on Γ if there exists a weight $2 - k$ cusp form g on Γ such that the non-holomorphic completion of f , defined as

$$\hat{f}(\tau) = f(\tau) - g^*(\tau)$$

satisfies $\hat{f} = \hat{f}|_{k,\chi}\gamma$ for every $\gamma \in \Gamma$. In the above, g^* denotes the non-holomorphic Eichler integral

$$g^*(\tau) := C \int_{-\bar{\tau}}^{i\infty} (\tau' + \tau)^{-k} g(\tau') d\tau', \quad (2.17)$$

defined for $\tau \in \mathbb{H}$.

²We choose the branch to be the principal branch $-\pi < \arg x \leq \pi$.

Note that there is no canonical normalization for the shadow and we choose ours to simplify the comparison between mock modular forms and Eichler integrals (2.13). Assuming that the limit $\lim_{t \rightarrow 0^+} f(x + it)$ exists for a given $x \in \mathbb{Q}$, let $Q(x) := \lim_{t \rightarrow 0^+} f(x + it)$, and consider $\gamma \in \Gamma$ in the notation of Definition 2.3. The γ -invariance of the completion \hat{f} leads to an expression for the corresponding modular difference function $p_\gamma(x)$ given by the modular difference function associated to $g^*(\tau)$. Through the related function (given by $\tau \mapsto -\tau$)

$$\tilde{g}^*(\tau) = C \int_{\bar{\tau}}^{i\infty} (\tau' - \tau)^{-k} g(\tau') d\tau' \quad (2.18)$$

and the fact that the modular difference function associated with \tilde{g}^* is again a period integral completely analogous to (2.16), it follows that mock modular forms indeed lead to quantum modular forms. Moreover, the quantum modular forms arising from a mock modular form and the Eichler integral of its shadow are clearly closely related.

However, just as Ramanujan already pointed out in his original work [Ram88], mock modular forms inevitably encounter infinities when approaching certain rational numbers from within the upper-half plane. Nonetheless, there exists a finite collection of weakly holomorphic modular forms that can be used to “cut out” these infinities and render a well-defined radial limit. More explicitly, we have the following theorem.

Theorem 2.5. [GOR13, CLR16] *Let f be a mock modular form of weight k and multiplier system χ for $\Gamma_0(N)$ with non-vanishing shadow g , and let $\{x_1, \dots, x_t\} \subset \mathbb{Q} \cup \{i\infty\}$ be a set of representatives of $\Gamma_0(N)$ -inequivalent cusps, then*

1. *the function $f(\tau)$ has exponential singularities at infinitely many rational numbers,*
2. *for every weakly holomorphic modular form G of weight k and multiplier system χ for $\Gamma_0(N)$, $f - G$ has exponential singularities at infinitely many rational numbers,*
3. *there exists a collection $\{G_j\}_{j=1}^t$ of weakly holomorphic modular forms with the following property.*

Given any cusp x , let x_j be the cusp representative that is $\Gamma_0(N)$ -equivalent to x and write $G_x = G_j$. Then $f - G_x$ is bounded towards x .

Moreover, following the arguments sketched above, the mock modular form and the Eichler integral of its shadow leads to a pair of closely related strong quantum modular forms.

Lemma 2.6. [CCFGH18] *With the notation of Theorem 2.5, let g be the shadow of f , the asymptotic expansions of the Eichler integral \tilde{g} and the mock modular form $f - G_x$ near x take the form*

$$(f - G_x)(-x + it) \sim \sum_{n \geq 0} \alpha_x(n) (-t)^n \quad \text{and} \quad \tilde{g}(x + it) \sim \sum_{n \geq 0} \alpha_x(n) t^n. \quad (2.19)$$

In particular, when the shadow g is a weight $3/2$ unary theta function, the mock modular forms are (up to an overall rational power of q) called mock theta functions in the terminology of [Zag10a], and the Eichler integral are the false theta functions encountered in §2.1. The false-mock pair satisfies

$$\lim_{t \rightarrow 0^+} (f - G_x)(x + it) = \lim_{t \rightarrow 0^+} \tilde{g}(-x + it), \quad (2.20)$$

reminiscent of the relation (1.6) between $Z_{\text{CS}}(M_3)$ and $Z_{\text{CS}}(-M_3)$ when taking $x = \frac{1}{k}$. Focusing on the cusp $x = 0$, we can see that the false and mock forms have the “same” asymptotic series,

approaching from the upper- and lower-half plane, in the sense that the asymptotic expansions in the limit $t \rightarrow 0^+$ satisfy

$$(f - G_0)(it) \sim \sum_{n \geq 0} \alpha_0(n)(-t)^n \text{ and } \tilde{g}(it) \sim \sum_{n \geq 0} \alpha_0(n)t^n. \quad (2.21)$$

On the three-manifold side, the cusp $x = 0$ is relevant for the perturbative invariants (the so-called Ohtsuki series), capturing the expansion around the semi-classical $k \rightarrow \infty$ limit.

To end this subsection, we provide an explicit example of such a false-mock pair.

Example. Consider the order seven mock theta function $F_0(q)$ by Ramanujan [Ram00]. It is, up to an overall power of $q^{-\frac{1}{168}}$, a mock modular form of weight $1/2$

$$f(\tau) = q^{-\frac{1}{168}} F_0(q) = q^{-\frac{1}{168}} (1 + q + q^3 + q^4 + q^5 + 2q^7 + O(q^8)), \quad (2.22)$$

whose shadow is given by the unary theta function

$$g(\tau) = (\theta_{42,1}^1 - \theta_{42,13}^1 - \theta_{42,29}^1 + \theta_{42,41}^1)(\tau). \quad (2.23)$$

Compared to (2.9), we see that the Eichler integral is (up to a factor $q^{\frac{83}{168}}$) precisely the quantum invariant of the Brieskorn sphere $\Sigma(2, 3, 7)$:

$$\widehat{Z}_0(\Sigma(2, 3, 7); \tau) = q^{\frac{83}{168}} \tilde{g}(\tau). \quad (2.24)$$

2.3 A Mock Conjecture

In [CCFGH18], the following relation between mock modular forms and three-manifold quantum invariants is proposed³.

Conjecture 2.7. *Let M_3 be a three-manifold whose quantum invariants take the form*

$$\widehat{Z}_a(M_3; \tau) = q^c (\tilde{g}(\tau) + p(\tau)) \quad (2.25)$$

where $c \in \mathbb{Q}$, $\tilde{g}(\tau)$ is the Eichler integral of a theta function $g(\tau)$ of weight $w = \frac{3}{2}$ and $p(\tau)$ is a polynomial in q , then

$$\widehat{Z}_a(-M_3; \tau) = q^{-c} (f(\tau) + p(-\tau)), \quad (2.26)$$

where $f(\tau)$ is a weight $1/2$ mock modular form whose shadow is given by $g(\tau)$.

The relevance of the above conjecture can be seen in Proposition 2.2, which guarantees the existence of M_3 satisfying the condition of the conjecture. More generally, we also expect mixed weight and higher-depth mock modular forms to play a role in three-manifolds quantum invariants.

See [CCFGH18] and [CCFFGH]. In what follows we briefly describe the three general motivations for

³Note that when $-M_3$ is not a weakly negative plumbed manifold, the mathematical definition 1.1 does not apply and this conjecture can be seen as rather a definition. However, recall that a physical definition of $\widehat{Z}_a(M_3)$ does exist for all closed three-manifolds [GPPV17]. As a result, independent computations can in principle be carried out for $-M_3$, as we will demonstrate in §3.1 for certain classes of $-M_3$. With this in mind we regard (2.26) as a conjecture.

the above conjecture, first discussed in [CCFGH18]. In §3 we will present explicit calculations which render results predicted by Conjecture 2.7, and hence constitute further evidence for it.

- As mentioned in the previous subsection, the asymptotic values (2.20) and expansions (2.21) of a false-mock pair are analogous to the relation (1.6) among the WRT invariants of a pair of three-manifolds related by a flip in orientation.
- Some false theta functions have known expressions as q -hypergeometric series, which converge not only inside but also outside the unit circle (when considered as a function of q). In some cases the expression on the other side is given by a mock theta function. See §7.4 of [CCFGH18] for details.
- When a weight $1/2$ mock modular form can be expressed as a so-called Rademacher sum, one can prove in general that the same Rademacher sum, now performed in the lower rather than the upper half-plane, yields precisely the corresponding Eichler integral. In other words, the Rademacher sum yields a function defined on both \mathbb{H} and \mathbb{H}^- , where they coincide with the mock resp. false theta function in question.

We refer to §7.4 of [CCFGH18] for a detailed discussion of the third point above. To illustrate the second point, let us consider an example that is again relevant for the Brieskorn sphere $M_3 = \Sigma(2, 3, 7)$.

Example. Let us define a function $\psi : \mathbb{H} \cup \mathbb{H}^-$ in terms of the q -hypergeometric series:

$$\psi(\tau) := \sum_{n \geq 0} \frac{q^{n^2}}{(q^{n+1}; q)_n}. \quad (2.27)$$

Note that the q -hypergeometric series converges both for $|q| < 1$ and $|q| > 1$. It can be shown that [Hik05b]

$$\psi(\tau) = \begin{cases} q^{-\frac{1}{168}} \tilde{g}(\tau) & , \tau \in \mathbb{H} \\ F_0(q^{-1}) & , \tau \in \mathbb{H}^- \end{cases}. \quad (2.28)$$

See also [BFR12] for a more general discussion. As a result, since $\widehat{Z}_0(\Sigma(2, 3, 7), \tau) = q^{\frac{1}{2}} \psi(\tau)$ for $\tau \in \mathbb{H}$, we can try to extend the definition of LHS to \mathbb{H}^- using the RHS. It is hence natural to guess that (cf. (1.7))

$$\widehat{Z}_0(-\Sigma(2, 3, 7), \tau) \text{ “ = ” } \widehat{Z}_0(\Sigma(2, 3, 7), -\tau) \text{ “ = ” } q^{-\frac{1}{2}} F_0(q). \quad (2.29)$$

We now end this section with a discussion on certain important open questions. First, note that Conjecture 2.7 does not specify, given a shadow, *which* mock modular form f should be the correct quantum invariant for the orientation-reversed manifold M_3 . Recall that two mock modular forms differing by a (weakly holomorphic) modular form have the same shadow. This question is of crucial importance since, as proposed in [GPPV17], the Fourier coefficients of the quantum invariants \widehat{Z}_a are (up to a possible factor of 2) integers which have the physical interpretation of counting supersymmetric quantum states in the underlying quantum physical theory. This said, we do expect the leading term of \widehat{Z}_a in the $\tau \rightarrow i\infty$ expansion to obey the naive $q \leftrightarrow q^{-1}$ relation and this puts meaningful constraints on the mock modular forms. Second, as we have seen in §2.2, mock and false theta functions relate to the WRT invariants in a slightly different way. While the radial limit of false theta functions are well-defined, for many cusps x one has to subtract the singular terms (by

subtracting a modular form G_x which cuts out the singularity (for instance) of the mock form in order to have a well-defined limit when approaching x from within the upper-half plane (cf. Lemma 2.6). The asymmetry might not be so surprising from the physical point; the $M5$ -brane theory is known to be a chiral theory. It would be extremely interesting to understand the physical or topological interpretation of the singular terms when taking radial limit of mock theta functions.

3 Explicit Calculations

In this section we summarise recent developments which make it possible to define and to compute the quantum invariants $\widehat{Z}_a(-M_3)$ for certain three-manifolds $-M_3$ that are relevant for the mock conjecture discussed in §2.3. We illustrate these methods with explicit computations for the Brieskorn sphere $M_3 = \Sigma(2, 3, 7)$.

3.1 Quantum Invariants via Knots

In this subsection, we review a (conjectural) way, introduced in [GM19], to compute the quantum invariants \widehat{Z}_a for some of the three-manifolds that are relevant for the mock conjecture 2.7, by constructing them via Dehn surgeries of knot complements.

Consider a knot K . Let $Y(K)$ be the knot complement of K in an integral homology sphere \hat{Y} . A closed manifold $Y_{p/r}(K)$ can be obtained by $Y(K)$ via Dehn surgery with coefficient $p/r \in \mathbb{Q}^*$. Roughly speaking, p/r specifies the diffeomorphism of $\partial Y(K)$, dictating the way a solid torus is glued along $\partial Y(K)$ to obtain $Y_{p/r}(K)$.

Now consider the special case when $\hat{Y} = S^3$. Given this choice, one associates to a knot K a two variable series

$$F_K(x, q) \in 2^{-c} q^\Delta \mathbb{Z}[x^{1/2}, x^{-1/2}][q^{-1}, q] \quad (3.1)$$

where $c \in \mathbb{Z}_+$ and $\Delta \in \mathbb{Q}$. For instance, for K a positive torus knot, an explicit expression for $F_K(x, q)$ has been given in [GM19]. Define a ‘‘Laplace transform’’ $\mathcal{L}_{p/r}^{(a)}$, given by (see also [GMP16])

$$\mathcal{L}_{p/r}^{(a)} : x^u q^v \rightarrow \begin{cases} q^{-u^2 r/p} \cdot q^v & \text{if } ru - a \in p\mathbb{Z}, \\ 0 & \text{otherwise.} \end{cases} \quad (3.2)$$

It has been shown for positive torus knots K (Theorem 1.2 of [GM19]) and conjectured for general knots (Conjecture 1.7 of [GM19]) that, for values of p/r such that the right hand side is well defined and for some $d \in \mathbb{Q}$ and $\varepsilon \in \{\pm 1\}$, one has

$$\widehat{Z}_a(\tau, S_{p/r}^3(K)) = \varepsilon q^d \cdot \mathcal{L}_{p/r}^{(a)}[(x^{\frac{1}{2r}} - x^{-\frac{1}{2r}})F_K(x, q)] \quad (3.3)$$

where we canonically identify the Spin^c -structure of $S_{p/r}^3(K)$ with

$$a \in \mathbb{Z} + \frac{r+1}{2} \text{ mod } p\mathbb{Z}. \quad (3.4)$$

Now it remains to compute F_K for general knots. It is convenient to define a rescaled version of

$F_K(q, x)$:

$$f_K(x, q) := \frac{F_K(x, q)}{x^{\frac{1}{2}} - x^{-\frac{1}{2}}}. \quad (3.5)$$

Based on physical expectations, a relation between the Borel resummation of the colored Jones polynomial $J_n(e^{\hbar})$ and $f_K(x, q)$, where $q = e^{\hbar}$ and $x = e^{n\hbar}$, is conjectured in (Conjecture 1.5 of) [GM19]. This, together with the AJ conjecture for the Jones polynomials [Gar04], in turn leads to the following conjecture

Conjecture 3.1. [GM19] *For any knot $K \subset S^3$ the quantum polynomial \widehat{A} of K annihilates the series $f_K(x, q)$*

$$\widehat{A}f_K(x, q) = 0 \quad (3.6)$$

and

$$\lim_{q \rightarrow 1} f_K(x, q) = s.e. \frac{1}{\Delta_K(x)} \quad (3.7)$$

where the symmetric expansion *s.e.* denotes the average of the expansions of the given rational function as $x \rightarrow 0$ (as a Laurent power series in x) and as $x \rightarrow \infty$ (as a Laurent power series in x^{-1}).

Note that (3.6) sets up a recursion relation for the coefficients $f_m(q)$ in $f_K(q, x) = \sum_m f_m(q)x^m$, while the relation (3.7) to the Alexander polynomial $\Delta_K(x)$ provides a boundary condition for the recursion equation. This is often sufficient to determine F_K to any desired order.

Example. For the figure-eight knot $K = \mathbf{4}_1$, the above-mentioned procedure leads to the leading order expansion [GM19]

$$F_{\mathbf{4}_1}(x, q) = \frac{1}{2} (\Xi(x, q) - \Xi(x^{-1}, q)) \quad (3.8)$$

where

$$\Xi(x, q) = x^{1/2} + 2x^{3/2} + (q^{-1} + 3 + q)x^{5/2} + (2q^{-2} + 2q^{-1} + 5 + 2q^1 + 2q^2) + \dots \quad (3.9)$$

The orientation-flipped Brieskorn sphere $-\Sigma(2, 3, 7)$ can be constructed through surgery on the complement in S^3 of the figure-eight knot $\mathbf{4}_1$, namely $-\Sigma(2, 3, 7) = S^3_{-1}(\mathbf{4}_1)$. Exploiting the conjecture (3.3) and plugging in (3.8)-(3.9), we obtain the result:

$$\widehat{Z}_0(-\Sigma(2, 3, 7)) = -q^{-\frac{1}{2}}(1 + q + q^3 + q^4 + q^5 + 2q^7 + q^8 + 2q^9 + q^{10} + 2q^{11} + \dots). \quad (3.10)$$

Note that the above leading terms in the q -expansion coincide (up to a sign) with the guess (2.29) based on quantum modular properties and on q -hypergeometric identities. However, the procedure outlined in this subsection does not immediately lead to a way to prove the modularity of (3.10). We will see yet another way to compute $\widehat{Z}_0(-\Sigma(2, 3, 7))$ in the following subsection.

3.2 Relation to indefinite theta functions

As mentioned earlier, one immediate problem with the proposal $\widehat{Z}_a(-M_3; \tau) \stackrel{?}{=} \widehat{Z}_a(M_3; -\tau)$ (1.7) is the fact that in Definition 1.1 one has $\tau \leftrightarrow -\tau \Leftrightarrow M \leftrightarrow -M$, and after this flipping of signature one no longer obtains a theta function Θ_a^{-M} (1.3) (and an integral (1.1)) that makes sense on the upper-half plane.

While it seems to be the end of the road as far as Definition 1.1 is concerned, a natural possibility is to replace the naive theta series with a regularised theta function. Indeed, building on previous work by Vignéras [Vig77], Zwegers [Zwe02] has devised a way to define a regularisation for theta functions of signature $(1, n)$ which retains its holomorphicity, and moreover established the relation to mock theta functions. The regularisation of general indefinite theta functions and the relation to higher-depth mixed mock modular forms has recently been developed in [ABMP16], [Naz16], [Wes16], [FK17], [ZZ]. In [CS], we apply these results to define and to compute quantum invariants for plumbed three-manifolds that are not weakly negative.

For the sake of concreteness and in order to establish a direct relation to Ramanujan's mock theta function, we focus on the class of three-manifolds discussed in Proposition 2.2. In the notation of Proposition 2.2 and of Figure 2, after performing the integration over w_{v_i} for $i \in \{1, 2, 3\}$ and write the $w_{v_0} = w$, we obtain

$$\widehat{Z}_0(M_3; \tau) = (-1)^\pi q^{\frac{3\sigma - \sum_v m(v)}{4}} \text{vp} \oint_{|w|=1} \frac{dw}{2\pi i w(w - w^{-1})} h(\tau, z) \quad (3.11)$$

where

$$h(\tau, z) = \sum_{j=0}^3 \sum_{\varepsilon \in \{\pm 1\}} \varepsilon \sum_{k \in 1+2\mathbb{Z}} q^{\frac{m}{4}k^2 - \frac{\varepsilon b_j}{2}k + c_j} w^k. \quad (3.12)$$

Note that naively taking $\tau \mapsto -\tau$ in $h(\tau, z)$ gives

$$\begin{aligned} & \frac{q^{\frac{1}{24}}}{\eta(\tau)} \sum_{j=0}^3 \sum_{\varepsilon \in \{\pm 1\}} \varepsilon \sum_{k \in 1+2\mathbb{Z}} \sum_{n \in \mathbb{Z}} (-1)^n q^{-\frac{m}{4}k^2 + \frac{\varepsilon b_j}{2}k - c_j + \frac{3n^2 - n}{2}} w^k \\ &= \frac{q^{-d} e^{\frac{\pi i}{6}}}{\eta(\tau)} \sum_{j=0}^3 \sum_{\varepsilon \in \{\pm 1\}} \varepsilon w^{\frac{\varepsilon b_j}{m}} \sum_{\mathbf{v} \in \Lambda_{j,\varepsilon}} q^{\frac{(\mathbf{v}, \mathbf{v})}{2}} e^{2\pi i(z \frac{1}{2}) \cdot \mathbf{v}} \end{aligned} \quad (3.13)$$

where we have inserted $1 = \frac{\eta(\tau)}{\eta(\tau)} = \frac{q^{\frac{1}{24}} \sum_{n \in \mathbb{Z}} (-1)^n q^{\frac{3n^2 - n}{2}}}{\eta(\tau)}$, and the bilinear form in the second line is given by

$$(\mathbf{v}', \mathbf{v}) := \mathbf{v}'^T K \mathbf{v}, \quad K := \begin{pmatrix} -\frac{m}{2} & 0 \\ 0 & 3 \end{pmatrix}, \quad (3.14)$$

and the set of summation is given by

$$\Lambda_{j,\varepsilon} = \left\{ \mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \mid v_1 \in 2\mathbb{Z} + 1 - \frac{\varepsilon b_j}{m}, v_2 \in \mathbb{Z} - \frac{1}{6} \right\}. \quad (3.15)$$

In other words, the key ingredient $h(\tau, z)$ of the integrand becomes, after taking $\tau \mapsto -\tau$ and multiplying by $\eta(\tau)$, a sum of theta functions of signature $(1, 1)$ that we would like to make sense of. As a result, we propose the following definition for three-manifolds M_3 that satisfy the conditions specified in Proposition 2.2 (*i.e.*, when M_3 is a weakly negative plumbed manifold which can be obtained from a three-star plumbing graph with four nodes and unimodular adjacency matrix):

$$\widehat{Z}_0(-M_3; \tau) := \frac{(-1)^\pi q^{\frac{3\sigma - \sum_v m(v)}{4}}}{\eta(\tau)} \text{vp} \oint_{|w|=1} \frac{dw}{2\pi i w(w - w^{-1})} \vartheta_a^M(\tau, z) \quad (3.16)$$

where

$$\vartheta_a^M(\tau, z) := q^{-d} e^{\frac{\pi i}{6}} \sum_{j=0}^3 \sum_{\varepsilon \in \{\pm 1\}} \varepsilon w^{\frac{\varepsilon b_j}{m}} \sum_{\mathbf{v} \in \Lambda_{j,\varepsilon}} \rho(\mathbf{v}) q^{\frac{(\mathbf{v}, \mathbf{v})}{2}} e^{2\pi i(z \frac{1}{2}) \cdot \mathbf{v}} \quad (3.17)$$

for an appropriately chosen ‘‘regularisation factor’’ $\rho(\mathbf{v})$ which will be discussed in details in [CS] and will be described explicitly in the example below.

Example. We will again take the example of $M_3 = \Sigma(2, 3, 7)$, with the plumbing graph and the adjacency matrix given in Figure 3. The relevant parameters m , b_j , and c_j are given in (2.8). Adapting [Zwe02] to preserve the symmetry⁴ $\vartheta_a^M(\tau, z) = -\vartheta_a^M(\tau, -z)$, we choose the regularising factor

$$\rho(\mathbf{v}) = \rho^{c,c'}(\mathbf{v}) := \frac{1}{2} (\operatorname{sgn}(\bar{\mathbf{v}}, c) - \operatorname{sgn}(\bar{\mathbf{v}}, c')) \quad (3.18)$$

where $\bar{\mathbf{v}} = (\frac{|v_1|}{v_2})$ for $\mathbf{v} = (\frac{v_1}{v_2})$ and the two timelike vectors are chosen to be

$$c = (1, 0), \quad c' = (8, 21). \quad (3.19)$$

See [CS] for a discussion on the choices of the timelike vectors c and c' . Putting things together, we have the following result (see [CS] for a proof).

Proposition 3.2. *When the regularisation factor $\rho(\mathbf{v})$ is given as in (3.18), (3.19), the definition (3.16) leads to*

$$\begin{aligned} q^{\frac{1}{2}} \widehat{Z}_0(-\Sigma(2, 3, 7); \tau) &= F_0(q) \\ &= 1 + q + q^3 + q^4 + q^5 + 2q^7 + O(q^8) \end{aligned} \quad (3.20)$$

where $F_0(q)$ is the order 7 mock theta function of Ramanujan.

Note that this result, given by the order 7 mock theta function, was precisely what was expected in [CCFGH18] (2.29). Moreover, at least the leading orders of q -expansion also, up to a sign, coincides with the result (3.10) which was obtained via a logically totally independent computation. These results constitute supporting evidence for Conjecture 2.7.

References

- [ABMP16] Sergei Alexandrov, Sibasish Banerjee, Jan Manschot, and Boris Pioline *Indefinite theta series and generalized error functions*. arXiv:1606.05495, 2016.
- [AP18] Sergei Alexandrov, and Boris Pioline. *Black holes and higher depth mock modular forms*. arXiv:1808.08479, 2018.
- [BFOR17] Kathrin Bringmann, Amanda Folsom, Ken Ono, and Larry Rolin. *Harmonic Maass Forms and Mock Modular Forms: Theory and Applications*. AMS, Colloquium Publications, Volume: 64; 2017.

⁴In this case, as noted in [BMM18], the principal value contour integral renders the same result as integrating over a contour lying inside (or outside) the unit disk.

- [BFR12] K. Bringmann, A. Folsom, R. C. Rhoades. *Partial theta functions and mock modular forms as q -hypergeometric series* R.C. Ramanujan J (2012) 29: 295.
- [BMM18] Kathrin Bringmann, Karl Mahlburg, and Antun Milas. *Quantum modular forms and plumbing graphs of three-manifolds*. arXiv:1810.05612 (v2), 2018.
- [CCFGH18] Miranda C. N. Cheng, Sungbong Chun, Francesca Ferrari, Sergei Gukov, and Sarah M. Harrison. *3d modularity*. arXiv:1809.10148, 2018.
- [CCFFGH] Miranda C. N. Cheng, Sungbong Chun, Boris Feigin, Francesca Ferrari, Sergei Gukov, and Sarah M. Harrison. *3d modularity and log VOAs*. to appear.
- [CDH14a] Miranda C. N. Cheng, John F. R. Duncan, and Jeffrey A. Harvey. *Umbral Moonshine*. Commun. Number Theory Phys., 8(2):101–242, 2014.
- [CDH14b] Miranda C. N. Cheng, John F. R. Duncan, and Jeffrey A. Harvey. *Umbral Moonshine and the Niemeier Lattices*. Research in the Mathematical Sciences, 1(3):1–81, 2014.
- [CLR16] D. Choi, S. Lim, and R. C. Rhoades, *Mock modular forms and quantum modular forms*, *Proc. Amer. Math. Soc.* **144** no. 6, (2016) 2337–2349. <https://doi.org/10.1090/proc/12907>.
- [CS] Miranda C. N. Cheng, and Gabriele Sgroi. *Three-manifolds and Indefinite Theta Functions*. To appear.
- [Duk14] W. Duke. *Almost a century of answering the question: what is a mock theta function?* Notices Amer. Math. Soc., 61(11):1314–1320, 2014.
- [DG18] Tudor Dimofte, and Stavros Garoufalidis. *Quantum modularity and complex Chern-Simons theory*. Commun. Number Theory Phys. 12 (2018), no. 1, 152, 2018.
- [DMZ14] Atish Dabholkar, Sameer Murthy, and Don Zagier. *Quantum Black Holes, Wall Crossing, and Mock Modular Forms*. arXiv:1208.4074, 2014.
- [Fin88] N. J., Fine. *Basic hypergeometric series and applications*. Math. Surveys and Monographs. vol. 27, Amer. Math. Soc., Providence, 1988.
- [Fol10] Amanda Folsom. *What is ... a mock modular form?* Notices Amer. Math. Soc., 57(11):1441–1443, 2010.
- [FH17] Francesca Ferrari, and Sarah M. Harrison. *Properties of extremal CFTs with small central charge*. arXiv:1710.10563, 2017
- [FK17] Jens Funke, and Stephen S. Kudla. *Mock modular forms and geometric theta functions for indefinite quadratic forms*. J. Phys. A 50 (2017), no. 40, 404001.
- [Gar04] S. Garoufalidis. *On the characteristic and deformation varieties of a knot*. in Proceedings of the Casson Fest, Geom. Topol. Publ., Coventry, volume 7 of Geom. Topol. Monogr., pp. 291?309, 2004.
- [GM19] Sergei Gukov and Ciprian Manolescu. *A two-variable series for knot complements*. arXiv:1904.06057, 2019.
- [GMP16] S. Gukov, M. Marino and P. Putrov. *Resurgence in complex Chern-Simons theory*. arXiv:1605.07615.

- [GOR13] M. Griffin, K. Ono, and L. Rolin, *Ramanujan’s mock theta functions*, *Proc. Natl. Acad. Sci. USA* **110** no. 15, (2013) 5765–5768.
- [GPPV17] Sergei Gukov, Du Pei, Pavel Putrov, and Cumrun Vafa. *BPS spectra and three-manifold invariants*. arXiv:1701.06567, 2017.
- [GZ18] Stavros Garoufalidis, and Don Zagier. *Asymptotics of Nahm sums at roots of unity*. arXiv:1812.07690, 2018, and to appear.
- [Hik05a] Kazuhiro Hikami *On the quantum invariants for the spherical Seifert manifolds* Commun. Math. Phys. , Vol. 268, 285-319. arXiv:0504082.
- [Hik05b] Kazuhiro Hikami *Mock (False) Theta functions as quantum invariants* Regular and Chaotic Dynamics **10**, pp. 509–530 (2005), arXiv:0506073.
- [Hik11] Kazuhiro Hikami *Decomposition of Witten-Reshetikhin-Turaev invariant: linking pairing and modular forms* AMS/IP Stud. Adv. Math. **50** (2011) 131
- [HL15] Kazuhiro Hikami, Jeremy Lovejoy *Torus knots and quantum modular forms*. Res. Math. Sci. 2 (2015).
- [LZ99] Ruth Lawrence, and Don Zagier *Modular forms and quantum invariants of 3- manifolds* Asian Journal of Mathematics **3**, (1999) 93–108.
- [Naz16] Caner Nazaroglu *r-Tuple Error Functions and Indefinite Theta Series of Higher-Depth*. arXiv:1609.01224, 2016.
- [Ono09] Ken Ono. *Unearthing the visions of a master: harmonic Maass forms and number theory*. In Current developments in mathematics, 2008, pages 347–454. Int. Press, Somerville, MA, 2009.
- [Ram88] Srinivasa Ramanujan. *The lost notebook and other unpublished papers*. Springer-Verlag, Berlin, 1988. With an introduction by George E. Andrews.
- [Ram00] Srinivasa Ramanujan. *Collected papers of Srinivasa Ramanujan*. AMS Chelsea Publishing, Providence, RI, 2000. Edited by G. H. Hardy, P. V. Seshu Aiyar and B. M. Wilson, Third printing of the 1927 original, With a new preface and commentary by Bruce C. Berndt.
- [Rog17] L. J. Rogers. *On two theorems of combinatory analysis and some allied identities*. Proc. London Math. Soc. (2) 16 (1917), pp. 316-336.
- [Vig77] Marie-France Vignéras. *Séries theta des formes quadratiques indéfinies*. Modular functions of one variable VI, Springer lecture notes 627 (1977), 227-239..
- [Wes16] Martin Westerholt-Raum *Indefinite Theta Series on Cones*. arXiv:1608.08874, 2016.
- [Zag10a] Don Zagier. *Ramanujan’s mock theta functions and their applications (after Zwegers and Ono-Bringmann)*. Astérisque, (326):Exp. No. 986, vii–viii, 143–164 (2010). Séminaire Bourbaki. Vol. 2007/2008.
- [Zag10b] Don Zagier *Quantum modular forms*. Quanta of maths, 659 - 675, Clay Math. Proc., 11, Amer. Math. Soc., Providence, RI, 2010.
- [Zwe02] Sander Zwegers. *Mock Theta Functions*. PhD thesis, Utrecht University, 2002.
- [ZZ] Don Zagier and Sander Zwegers. To appear.