UvA-DARE (Digital Academic Repository)

Parasite-host specificity: A cross-infection study of the parasite *Ophryocystis elektroscirrha*

Gao, K.; Muijderman, D.; Nichols, S.; Heckel, D.G.; Wang, P.; Zalucki, M.P.; Groot, A.T.

DOI

[10.1016/j.jip.2020.107328](https://doi.org/10.1016/j.jip.2020.107328)

Publication date

2020

Document Version

Final published version

Published in

Journal of Invertebrate Pathology

License

Article 25fa Dutch Copyright Act

[Link to publication](#)

Citation for published version (APA):

Gao, K., Muijderman, D., Nichols, S., Heckel, D. G., Wang, P., Zalucki, M. P., & Groot, A. T. (2020). Parasite-host specificity: A cross-infection study of the parasite *Ophryocystis elektroscirrha*. *Journal of Invertebrate Pathology*, 170, [107328].
<https://doi.org/10.1016/j.jip.2020.107328>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Parasite-host specificity: A cross-infection study of the parasite *Ophryocystis elektroscirrha*

Ke Gao^{a,*}, Daphne Muijderland^a, Sarah Nichols^a, David G. Heckel^b, Peng Wang^c,
Myron P. Zalucki^c, Astrid T. Groot^{a,b}

^a Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Science Park 904, Amsterdam, the Netherlands

^b Max Planck Institute for Chemical Ecology, Department of Entomology, Jena, Germany

^c School of Biological Science, The University of Queensland, 4072 Brisbane, Australia

ARTICLE INFO

Keywords:

Host specificity

Ophryocystis elektroscirrha

Parasites

Cross-infection

Helicoverpa

Parthenos sylvia

ABSTRACT

Many parasites are constrained to only one or a few hosts, showing host specificity. It remains unclear why some parasites are specialists and other parasites are generalists. The parasite *Ophryocystis elektroscirrha* (OE) is a neogregarine protozoan thought to be restricted to monarch butterflies, *Danaus plexippus* (Nymphalidae) and *D. gilippus*. Recently, we found OE-like spores in other Lepidoptera, specifically in three noctuid moths: *Helicoverpa armigera*, *H. assulta* and *H. punctigera*, as well as another nymphalid, *Parthenos sylvia*. To our knowledge, this is the first report of OE-like parasite infections in species other than the genus *Danaus*. In sequencing 558 bp of 18S rRNA, we found the genetic similarity between OE from *D. plexippus* and OE-like parasite from the moths *H. armigera* and *H. punctigera* to be 95.2%. When we conducted cross-species infection experiments, we could not infect the moths with OE from *D. plexippus*, but OE-like parasite from *H. armigera* did infect *D. plexippus* and a closely related moth species *Heliothis virescens*. Interestingly, we did not find the OE-like parasite in the *H. armigera* population from Spain. Inter-population infection experiments with *H. armigera* demonstrated a higher sensitivity to OE-like infection in the population from Spain compared to the populations from Australia and China. These results suggest geographic variation in OE-like susceptibility and coevolution between parasite and host. Our findings give important new insights into the prevalence and host specificity of OE and OE-like parasites, and provide opportunities to study parasite transmission over spatial and temporal scales.

1. Introduction

Parasites are present in all kinds of organisms and affect host fitness in variable ways (Schmid Hempel, 2011). Parasitic infections can result in the decline of populations, especially under changing environmental conditions including global warming (Altizer et al., 2003; Wood et al., 2007). Most parasites of both plants and animals show host specificity (Little et al., 2006; Norton and Carpenter, 1998). With global warming, many parasites and their hosts can expand their geographic ranges, which likely affects host-parasite interactions (Hellgren et al., 2007; Pickles et al., 2013; Polley and Thompson, 2009). For example, parasites may infect closely related host species when introduced into new habitats, as they share similar immune defenses and physiological traits as the original host (Norton & Carpenter, 1998; Davies & Pedersen, 2008; Poulin et al., 2011). When host shifts occur, the virulence of parasites may change (Little et al., 2006). Interestingly, it remains unclear why some parasites are highly specialized on only one or few hosts

and other parasites are shared across a wide range of host species (Cooper et al., 2012; Mácová et al., 2018). Understanding the specific range of parasite infections is crucial to predicting future infectious diseases in plants and animals, including humans (Cooper et al., 2012; Keesing et al., 2010; Li et al., 2016).

The parasite *Ophryocystis elektroscirrha* (OE) is a neogregarine protozoan and has been thought to be an obligate parasite restricted to the monarch butterfly, *Danaus plexippus*, and closely related queen butterfly, *D. gilippus*. This host specificity is deduced from the fact that OE was first observed on these host species in Florida (McLaughlin and Myers, 1970) and the life-cycle of OE is closely linked to its host (Altizer and Oberhauser, 1999). After early-instar larvae consume OE spores, they undergo asexual, vegetative replication in the larval gut (Altizer et al., 2000). Upon lysis, the spores form micronuclear schizonts which penetrate the gut wall, after which they migrate to the hypoderm where a combination of sexual and asexual replication commences (Vickerman et al., 1999). Sporulation occurs during host pupation and

* Corresponding author.

E-mail address: K.Gao@uva.nl (K. Gao).

<https://doi.org/10.1016/j.jip.2020.107328>

Received 8 October 2019; Received in revised form 18 December 2019; Accepted 11 January 2020

Available online 14 January 2020

0022-2011/ © 2020 Published by Elsevier Inc.

dormant spores emerge on the abdominal scales of adult butterflies (McLaughlin and Myers, 1970). OE infections have been found to be detrimental to host fitness in *D. plexippus*, reducing longevity, eclosion, mating success, fecundity and flight ability (Altizer and Oberhauser, 1999; Bradley and Altizer, 2005; De Roode et al., 2009).

Transmission of this parasite is predominantly vertical, as females scatter spores on their eggs or host plants during oviposition and egg shells are ingested by hatching larvae (Leong et al., 1992). However, paternal or horizontal transmission routes are possible, e.g., through mating or spores consumed by unrelated larvae (Altizer et al., 2004; De Roode et al., 2009). Recently, Barriga et al. (2016) firstly reported OE-like infections in the lesser wanderer butterfly, *D. petilia*, suggesting that the OE-like parasites could have a wider host range than previously assumed.

In checking field-collected samples of the cotton bollworm, *Helicoverpa armigera* (Hübner) (Lepidoptera, Noctuidae) in Australia to start laboratory rearing, we noticed that some adults did not fully emerge from their pupal case or could not expand their wings, a symptom indicative of OE infection (Leong et al., 1992). A check of adults indicated OE-like spores present. These observations led us to investigate the identity of the parasite infecting *H. armigera* by sequencing a subset of the 18S rRNA from parasites collected from field-collected *H. armigera* populations in China, Australia and Spain, as well as from the closely related species *H. assulta* in China and *H. punctigera* in Australia, and comparing these sequences to the *D. plexippus* OE. We further explored the OE or OE-like parasite host range by investigating its presence in butterfly species in the butterfly garden of Artis Zoo in Amsterdam, which are flown in biweekly as pupae from so-called butterfly farms in South-East Asia, Africa and South America. In addition to sequencing, we determined the host-specificity of OE and OE-like parasites by conducting cross-infection experiments with *D. plexippus*, *H. armigera* and *Heliothis virescens*, and we tested whether the susceptibility of OE-like infection varied between geographic origins of *H. armigera* populations. As *H. armigera* is highly polyphagous, feeding on a wide variety of host plants from at least 172 species within 68 families (Cunningham and Zalucki, 2014; Fitt, 1989), and is widely distributed throughout Africa, Europe, Asia, Oceania (Fitt, 1989) and recently South America (Czapak and Albernaz, 2013), these experiments give a first indication of the potential risk of OE-like parasite to spread to other closely related *Helicoverpa* and *Heliothis* species. (Pogue (2013) has recently resurrected the name *Chloridea* for a monophyletic genus containing *virescens* and *subflexa*, but for consistency with the older literature we continue to refer to *Heliothis virescens* here.)

2. Materials and methods

2.1. Occurrence of OE-like parasite in Lepidoptera other than *Danaus*

Three *Helicoverpa* moth species were collected from the field and checked for OE parasites: *H. armigera* from China, Australia and Spain, and *H. assulta* and *H. punctigera* in China and in Australia, respectively (see Table 1). The moths were collected by using a hand net or caught in pheromone traps, after which the moths were killed, placed individually in Eppendorf tubes and brought back to the laboratory at the University of Amsterdam for further diagnosis. To diagnose OE-like parasite infection, the abdomens of the moths were sampled using

2.5 cm diameter transparent tape and checked for parasite spores, as described in detail by Altizer et al. (2000). OE spores are approx. 14 µm long, 9 µm wide, dark when seen in scales or on the egg, but amber in transmitted light (McLaughlin and Myers, 1970). OE-like spores on *H. armigera* from Australia are morphologically similar as OE spores, with dimension of approx. 12 µm length and 9 µm width.

To survey OE-like parasite occurrence in other butterfly species, specimens were collected from the butterfly garden at Artis Zoo, in Amsterdam, Netherlands in May-June 2018. In total 32 butterfly species, originating from Asia, Africa and South America, were sampled (see Supplementary Table 1). Butterfly specimens were placed individually in paper bags with the species identification and collected location. All specimens were brought back to the laboratory and checked OE-like parasite under the microscope, as described above.

2.2. Genetic similarity between OE and OE-like parasites

2.2.1. DNA extraction and PCR

To identify OE-like parasite that we found in the different species in the field and the Amsterdam Zoo, we extracted DNA from the collected spores from five different host species (Table 2). Specifically, the abdomen of infected adults was placed in a 1.5 ml Eppendorf tube with 100% ethanol. The tube was agitated for 1 min using a vortex mixer followed by a 5 min rest interval, and repeated three times. The abdomen was removed, after which the spore suspension was centrifuged for 10 min at 13,000 rpm and the supernatant discarded and 200 µl of 2% CTAB buffer with 5 µl proteinase-K were added to the tube with parasite spores. The spores were then ground with a clean pestle, and incubated for 1.5 h at 55 °C. After incubation, the supernatant was transferred into a sterile tube with chloroform: IAA (24:1) after centrifugation at 12,000 rpm for 5 min 300 µl of 100% ice-cold ethanol was added to precipitate the parasite DNA. The DNA sample was centrifuged at 13,000 for 10 min and the supernatant discarded. The DNA pellet was washed with 70% ethanol and dried in a speedvac for 10 min at 30 °C. The pellet was dissolved in 25 µl distilled H₂O and stored at –20 °C.

To sequence the 18S rRNA, PCR reactions were prepared in 10 µl that included 3.5 µl distilled H₂O, 2 µl 5x PCR buffer, 2 µl 5 mM dNTPs, 0.2 µl 10 mM primers, 0.1 µl Phire hotstart Taq and 2 µl DNA template. The PCR amplification was performed by 35 cycles of 30 s at 98 °C, 10 s at 56 °C, 2 min at 72 °C. The forward primer 5'-CCCGTGTGAGTCAA ATTAAG-3' and reverse primer 5'-AGGGCAAGTCTGGTGCCAG-3' were used to amplify a subset of the 18S rRNA gene in this study. The PCR products were examined on a 1.5% Agarose gel.

2.2.2. Phylogenetic analysis

The subset of the 18S rRNA sequences were checked manually using CodonCode Aligner (CodonCode Corp., USA), and were aligned to the sequence of *O. elektroscirra* (AF129883) from GenBank, using MEGA version 10.0.5. The sequences in this study were deposited in GenBank (accession no. MK720123-MK720132). Outgroup species were selected based on previous phylogenetic studies by Carreno et al. (1999) and Bekircan et al. (2017), based on which the representative species *Ascogregarina culicis*, *Aranciocystis muskarensis* and *Acarus siro* were chosen.

To assess interspecific genetic distances of the OE and OE-like

Table 1
Field collections of *Helicoverpa* species sampled for OE-like infection.

Species	Locations	Coordinates	Years	Number of Individuals	Infection rate (%)
<i>H. armigera</i>	Guadajira, Badajoz, Spain	38°51'08.8"N, 6°40'48.9"W	2016, 2018	174	0
<i>H. armigera</i>	Gatton, Brisbane, Australia	27°32'11.6"S, 152°20'16.3"E	2017	48	19
<i>H. armigera</i>	Dali county, Shaanxi, China	34°45'01.9"N, 110°09'56.1"E	2017, 2018	66	2
<i>H. assulta</i>	Dali county, Shaanxi, China	34°45'01.9"N, 110°09'56.1"E	2017	20	10
<i>H. punctigera</i>	Gatton, Brisbane, Australia	27°32'11.6"S, 152°20'16.3"E	2017	45	7

Table 2
Samples used for phylogenetic analyses.

Species	Voucher	Host	Collection Location	Collection Date	GenBank Accession No.
<i>Ophryocystis elektroscirrha</i>	OEDP1	<i>Danaus plexippus</i>	Costa Rica, South America ¹	2018.06	MK720123
	OEDP2	<i>Danaus plexippus</i>	Costa Rica, South America ¹	2018.06	MK720124
	OEDP3	<i>Danaus plexippus</i>	Netherlands	2018.08	MK720125
	OEPC4	<i>Parthenos sylvia</i>	Asia ¹	2018.06	MK720126
	OEPC5	<i>Parthenos sylvia</i>	Asia ¹	2018.06	MK720127
	OEPC6	<i>Parthenos sylvia</i>	Asia ¹	2018.06	MK720128
	OEHP7	<i>Helicoverpa punctigera</i>	Australia	2017.02	MK720129
	OEHA8	<i>Helicoverpa armigera</i>	Australia	2017.03	MK720130
	OEHA9	<i>Helicoverpa armigera</i>	China	2017.08	MK720131
	OEHA10	<i>Helicoverpa assulta</i>	China	2017.08	MK720132
<i>Ascogregarina culicis</i>		<i>Danaus plexippus</i>			AF129883*
<i>Aranciocystis muskarensis</i>					DQ462456*
<i>Acarus siro</i>					KU926299*
					AY490099*

* Sequences download from GenBank.

¹ Collections from Artis Zoo, Amsterdam.

parasites, Kimura-2-parameter distances (K2P distance) were calculated in MEGA. For phylogenetic analysis, a maximum likelihood (ML) tree was constructed using MEGA. The substitution model for ML tree was based on Tamura 3-parameter method with uniform rates among sites. Reliability of ML tree was tested with 1000 bootstrap replicates.

2.3. Host-specificity of OE

2.3.1. Cross-species infection experiment

To determine whether the OE or OE-like parasites can infect different host species, cross-infection experiments were conducted. For these cross-infection experiments, larvae of *H. armigera* were collected from Australia, Spain and China during 2017 and 2018, and reared individually on artificial pinto bean diet in the same climate chamber (60% relative humidity (RH); 25 °C; 14 h light: 10 h dark with lights off at 11.00 am) in the laboratory at the Institute for Biodiversity and Ecosystem and Dynamics, University of Amsterdam. Larvae of *H. virescens* used were also reared individually on artificial pinto bean diet. Larvae of monarch butterfly *D. plexippus* were originally from five mated females collected at a butterfly garden in Harskamp, Netherlands in July 2018. Monarch larvae were reared on the milkweeds *Asclepias incarnata*. These plants were grown and supplied in the greenhouse at the University of Amsterdam.

First, we used the OE-like parasite spores collected from *H. armigera* to infect three species: *H. armigera* from Australia, a closely related species *H. virescens* and *D. plexippus*. The OE-like parasite spores used in the experiments were obtained from the lab-reared *H. armigera* from CSIRO, Black Mountain Laboratory, ACT, Australia. The spore inoculum was obtained using a method modified from Leong et al. (1992). Specifically, the abdomen of an infected *H. armigera* adult was placed in a 2 ml Eppendorf tube with a washing solution of 0.5 ml 0.05% wetting agent (Tween 20) and 1 ml distilled water. The tube was agitated for 1 min using a vortex mixer, followed by a 5 min rest interval, and repeated three times. The abdomen was removed, and the spore suspension was then centrifuged for 5 min at 10,000 rpm. The supernatant was discarded and the remaining pellet was re-suspended in 1 ml distilled water to ensure the spores were evenly dispersed. To estimate the spore load, 10 droplets of 1 µl spore suspension were randomly chosen and the number of spores in each droplet counted under the microscope. From these counts we calculated the average number of spores with standard error (SE) in 1 µl droplet of spore suspension.

To infect the larvae of *H. armigera* and *H. virescens*, 1 µl of spore suspension (spores load: 122 ± 10 spores/µl) was pipetted onto a piece of 1 cm² pinto bean-based artificial diet in a 37 ml cup. To prevent the diet from drying out before consumption, the cups were lined with cotton soaked in distilled water. Third instar larvae of *H. armigera* from

six families and *H. virescens* from four families were used in the experiments. All the larvae placed into the cups individually and observed to consume the diet completely, after which the larvae were transferred to the normal diet until pupation.

To infect the larvae of the monarch butterfly with OE-like spores from *H. armigera*, second instar caterpillars were infected with 1 µl of spore suspension (spores load: 76 ± 2 spores/µl) on a piece of 1 cm² *A. incarnata*. After consumption of the spores, the caterpillars were placed in 0.6 m³ mesh cages supplied with clean *A. incarnata* plants that were replaced every day until the caterpillars pupated. Pupae were individually transferred into clear plastic cups until eclosion.

Second, we used the OE parasite spores collected from *D. plexippus* to infect *D. plexippus*, *H. armigera* from Australia, and *H. virescens*. The OE spores were collected from newly hatched heavily infected *D. plexippus* at the butterfly garden in Harskamp, Netherlands. All the larvae were infected with 1 µl of spore suspension (spores load: 55 ± 3 spores/µl). As none of the moths were infected from *D. plexippus* OE spores, we repeated the experiment and infected *H. armigera* and *H. virescens* larvae with 1 µl of spore suspension (spores load: 135 ± 2 spores/µl) that we collected from *D. plexippus*. These two spore suspensions were prepared as described above, and all the larvae were infected in the same way. Third instar larvae of *H. armigera* from eight families and *H. virescens* from five families were used in the experiments. Upon eclosion, the adults of all species in the study were frozen in individual Eppendorf tubes or in paper bags, after which their abdomens were examined for infection of parasite spores as described above.

2.3.2. Inter-population infection experiment

To determine the occurrence and specificity of OE-like parasite among *H. armigera* populations, the OE-like parasite spores collected on *H. armigera* from Australia were used to infect the three populations of *H. armigera* sourced from Australia, China and Spain. The larvae were infected with a low concentration (16 ± 1 spores/µl) and a high concentration of spore suspension (99 ± 3 spores/µl), using the same methods as described above.

2.4. Effect of OE-like parasite on moth emergence success

To assess the effect of OE-like spore concentration on infection probability, third instar larvae of *H. armigera* from Australia were infected with nine different concentrations of spores (from 1 spores/µl to 221 ± 3 spores/µl). For these infections, the OE-like parasite spores collected from *H. armigera* from Australia were used, and multiple spore suspensions were prepared as described above. At least 30 larvae were exposed to each concentration or distilled water as control. All the

larvae were reared individually on pinto bean diet until eclosion. The adults were examined for infection. We also recorded the number of adults that were stuck in their pupal case, and the number of adults with crinkled wings that did not fully expand. To assess effects of OE parasite on the emergence success of *H. armigera*, we compared that the number of stuck in pupae and adults with crinkled wings between infected individuals from the larvae exposed to OE-like spores and uninfected individuals from larvae treated with distilled water.

2.5. Statistical analyses

To determine the difference of infection rate in host species in the cross-species infection experiment, we performed a chi-square test. The same test was used to determine the difference of OE-like infection rate among *H. armigera* populations from Australia, China and Spain, when treated with a low or high concentration of spores. To test the effect of OE-like parasite spore concentration on infection probability, a regression analysis was carried out. We used a chi-square test to determine the difference between infected and uninfected individuals of the number of pupae stuck and adults with crinkled wings in *H. armigera*. All statistical analyses were performed in R software, version 3.4.1 (R_Core_Team 2018).

3. Results

3.1. Occurrence of OE or OE-like parasites

In the field, the OE-like parasite was detected in all three *Helicoverpa* moths. We found OE-like infection in *H. armigera* populations in China and Australia, but not in Spain. We also found the OE-like infection in other closely related species, such as in *H. assulta* in China and *H. punctigera* in Australia (Table 1). The prevalence of OE-like parasite infection varied among the locations, with highest infection rate of 19% (9 out of 48 individuals) on *H. armigera* in Australia.

From the specimens collected in the Amsterdam Zoo, two out of 32 butterfly species were found to carry OE or OE-like parasites. Specifically, 86% of *D. plexippus* (12 out of 14 individuals) that were originally from South America were infected and 58% of *Parthenos sylvia* (Nymphalidae) (15 out of 26 individuals) originating from Asia were infected (Table S1).

3.2. Genetic similarity between OE and OE-like parasites

The subset of the 18S rRNA sequences were 558 bp in length. Aligning all sequences, we found the interspecific K2P distance between *D. plexippus* and *P. sylvia* to be 1.5%, and the K2P distances between *D. plexippus* and *H. armigera* (also *H. punctigera*) was 4.8%. The mean K2P distance between OE-like spores collected within the *Helicoverpa* genus was < 0.2%.

All parasites collected from *D. plexippus* individuals, as well as all parasites collected from *P. sylvia* individuals, formed a monophyletic group with strong (99%) bootstrap support (Fig. 1). The remaining three *Helicoverpa* moths formed another monophyletic clade, with 100% bootstrap support.

3.3. Host specificity of OE or OE-like parasites

3.3.1. Cross-species infection experiment

When larvae from different species were treated with OE spores collected from *D. plexippus*, 97% (37 out of 38 adults) of *D. plexippus* were infected, while none of the adult *H. armigera* ($n = 63$) or *H. virescens* ($n = 95$) were infected ($\chi^2 = 194.8$, $df = 2$, $P < 0.0001$). In contrast, when larvae from the different species were treated with OE-like spores collected from *H. armigera* in Australia, the infection rate was 71% in *D. plexippus* (20 out of 28 adults) and 85% in *H. armigera* (93 out of 110 adults), but significantly lower in *H. virescens*, i.e., 48%

(13 out of 27 adults) ($\chi^2 = 10.0$, $df = 2$, $P = 0.0067$) (Fig. 2).

3.3.2. Inter-population infection experiment

When we treated *H. armigera* larvae from different geographic locations with a low concentration (16 ± 1 spores/ μ l) of OE-like spores collected on *H. armigera* from Australia, 53% (9 out of 17 adults) of *H. armigera* from Spain were infected, 15% (3 out of 20 adults) from China were infected, while no ($n = 10$) adults from Australia were infected. The infection rate in individuals from Spain was significantly higher than that from China and Australia ($\chi^2 = 65.68$, $df = 2$, $P < 0.0001$).

When treated with a high concentration of spores (99 ± 3 spores/ μ l), 75% (18 out of 24 adults) of *H. armigera* from Spain were infected, while 73% (24 out of 33 adults) from China and 57% (13 out of 23 adults) from Australia were infected. There was no significant difference in the infection rate among the three *H. armigera* populations at this high dose ($\chi^2 = 2.99$, $df = 2$, $P = 0.22$) (Fig. 3).

3.4. Effect of OE-like parasite on moth emergence success

When larvae of *H. armigera* from Australia were infected with different concentrations of OE-like spores, the infection rates in adult moths best fitted a logarithmic regression line ($R^2 = 0.71$, $P = 0.0045$) (Fig. 4a). This logarithmic relationship suggests that the proportion of infected individuals increased with concentration of OE-like spores, reaching an asymptote. The number of pupae stuck ($\chi^2 = 4.63$, $df = 1$, $P = 0.031$) or adults with wings crinkled ($\chi^2 = 6.01$, $df = 1$, $P = 0.014$) from infected individuals were significantly higher than that from uninfected individuals (Fig. 4b).

4. Discussion

To our knowledge, this is the first report of OE-like parasite infections in species of Lepidoptera other than *Danaus* species. In the past, the protozoan OE and OE-like parasites were thought to be specialized on monarchs and closely related butterflies (Barriga et al., 2016; McLaughlin and Myers, 1970). In our survey, OE-like parasites were found in three noctuid moth species *H. armigera*, *H. assulta* and *H. punctigera* collected in China and Australia, as well as in the butterfly *P. sylvia* (Nymphalidae) collected in Asia. This indicates that OE and OE-like parasites are more widespread and has more potential hosts in nature.

On the basis of partial 18S rRNA sequences in the different OE and OE-like spore collections, we found only 1.5% genetic distance between the butterflies *P. sylvia* and *D. plexippus*, indicating a high similarity. The OE-like parasite that we found in noctuid moth species (especially *H. armigera* and *H. punctigera*) exhibited a 4.8% genetic distance from *D. plexippus* OE. Our phylogenetic analysis did show an OE and OE-like parasites divergence into two groups, representing butterflies and moths. It strongly suggests that OE from *D. plexippus* and OE-like parasite from moths may be different species, which displays host specificity or coevolutionary processes of OE and OE-like parasites among different hosts.

As we found that OE from *D. plexippus* did not infect *H. armigera* or *H. virescens*, it suggests that OE populations present on monarch butterflies are more specialized than OE-like parasite that we found on *H. armigera*. Barriga et al. (2016) also found that monarch butterflies were more susceptible to their own natal OE parasite, while queen butterfly *D. gilippus* was barely infected by monarch OE parasite. In contrast, both *D. plexippus* and *D. gilippus* could be infected by OE-like parasite collected from lesser wanderer butterfly *D. petilia*. Similarly, OE-like parasite collected from *H. armigera* could not only infect the closely related host *H. virescens*, but were capable of infecting monarch butterflies. Such an asymmetrical distribution of a parasite between two or more hosts has been found in other parasites, e.g., helminths and copepods and indicates that local specialization of parasites to their hosts (Poulin et al., 2011).

Fig. 1. Phylogenetic relationship of OE and OE-like spores collected from different hosts. Maximum likelihood (ML) tree based on 558 bp 18S RNA gene sequences. Values above the branches indicate clade bootstrap support.

Fig. 2. Infection rate with OE spores collected from *D. plexippus* (black bar) or OE-like spores collected from *H. armigera* (gray bars). Significant differences are indicated by asterisks (***: $P < 0.001$) and different letters ($P < 0.05$).

Interestingly, there seems to be geographic variation in host susceptibility to OE-like parasite in *H. armigera*, as we found that *H. armigera* larvae from Spain were more susceptible to OE-like infections than *H. armigera* larvae from China and Australia. As the genetic variation of OE-like parasite was less than 0.2% within the *Helicoverpa* genus, the geographic variation is likely to be due to local environmental conditions. Previous studies have found that environmental factors, such as temperature and humidity, could have a major impact on prevalence, transmission and intensity of parasites (Duncan and Little,

2007; Laine and Tellier, 2008; Poulin, 2006; Stromberg, 1997). Furthermore, the resistance to parasites could vary among the host populations from different geographic locations (Barriga et al., 2016; Kaltz et al., 1999; Sternberg et al., 2013). Since OE-like parasite was not found on any *H. armigera* individual collected from Spain, this population may be more susceptible to infection, while *H. armigera* populations in China and Australia could have developed some level of resistance or tolerance to OE-like infections.

Host migration could have a strong influence on the interaction

Fig. 3. Infection rate of OE-like spores collected from *H. armigera* in Australia in different *H. armigera* populations. Small letters indicate significant differences between different *H. armigera* populations with low concentration (16 ± 1 spores/ μ l) ($P < 0.05$), and capital letters indicate no significant differences with high concentration (99 ± 3 spores/ μ l) ($P > 0.05$).

between host and parasite (Altizer et al., 2011; Gandon et al., 1996; Kaltz and Shykoff, 1998). For example, long-distance migration can help to reduce parasite transmission and spread host resistance genes (Bartel et al., 2011; Bradley and Altizer, 2005; Loehle, 1995). In *D. plexippus*, the prevalence of OE parasite has been found to be higher in resident than in migratory populations (Bartel et al., 2011). *Helicoverpa armigera* also has the ability to undergo seasonal migration covering long distances (Zalucki & Furlong, 2005; Feng et al., 2009), which could affect OE-like parasite transmission over spatial and temporal scales. Taking these factors into account, we can expect that the prevalence of OE-like parasite and host susceptibility or resistance varies among *H. armigera* populations.

Climate change can alter the dynamics of parasite transmission and increase the potential for host switching (Brooks and Hoberg, 2007; Cooper et al., 2012). Correspondingly, the OE or OE-like parasites could pose a potential risk to other Lepidoptera and expand their distributions into new areas, or even new hosts. As *H. armigera* has recently invaded in South America (Czepak and Albernaz, 2013), and we already found that OE-like parasites can infect closely related host species, non-pest moth species and butterflies in all continents may be at risk. Once the parasites are introduced into new areas or new hosts, a number of questions arise, such as: What are the major factors that drive host shifts

in these parasites? How does OE or OE-like parasites affect the fitness of different host species and how do different host species respond to parasitic infections? As temperature and relative humidity affect many life history traits in plants and animals in general (Alcázar and Parker, 2011; Peng et al., 2019), therefore climate change likely also has profound consequences in coevolutionary process between hosts and parasites.

5. Conclusion

In summary, our study gives the first evidence of OE-like infections in Lepidoptera other than the genus *Danaus*. The genetic similarity between OE from *D. plexippus* and OE-like parasite from the moths *H. armigera* and *H. punctigera* was 95.2%. Cross-infection experiments showed a higher host specificity of OE collected from monarchs and lower host specificity from OE-like parasite collected from *H. armigera*. Interestingly, the *H. armigera* population in Spain, in which we did not find OE-like infections, showed higher sensitivity to OE-like infection than the *H. armigera* populations in Australia and China, indicating geographic variation in the level of susceptibility, resistance or tolerance to OE-like in *H. armigera*. Further studies should give insights into how parasitic OE or OE-like differs among lepidopteran hosts and the level and extent of parasite transmission over spatial and temporal scales.

Acknowledgements

We thank Tom Walsh for supplying OE-like parasites in Australia, and Kevin Wakeman for his help to identify OE by giving the primer sequences and suggesting DNA extraction protocols for OE. We also thank Peter Kuperus and Hans Breeuwer for their help with parasite DNA isolation and phylogenetic analysis, and Dennis van Veldhuizen for his help with rearing insects. Two anonymous reviewers provided comments that helped improve this manuscript. This research was supported by the China Scholarship Council (CSC) (award no. 201506300162), the National Science Foundation (award IOS-1456973), the Netherlands Organisation for Scientific Research (NOW-ALW, awards 822.01.012 and ALWOP 2015.075) and the Max Planck Society.

Declarations of Competing Interest

The authors declared that there is no conflict of interest.

Fig. 4. (a) OE-like infection rates in *H. armigera* adults when third instar larvae were treated with different spore concentrations. (b) Percentage of stuck pupae or adults with crinkled wings in infected individuals and uninfected individuals. n = total number of individuals checked. Significant differences are indicated by asterisks (*: $P < 0.05$).

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jip.2020.107328>.

References

- Alcázar, R., Parker, J.E., 2011. The impact of temperature on balancing immune responsiveness and growth in *Arabidopsis*. *Trends Plant Sci.* 16, 666–675.
- Altizer, S., Bartel, R., Han, B.A., 2011. Animal migration and infectious. *Science* 331, 296–302. <https://doi.org/10.1126/science.1194694>.
- Altizer, S., Harvell, D., Friedle, E., 2003. Rapid evolutionary dynamics and disease threats to biodiversity. *Trends Ecol. Evol.* 18, 589–596. <https://doi.org/10.1016/j.tree.2003.08.013>.
- Altizer, S.M., Oberhauser, K.S., 1999. Effects of the protozoan parasite *Ophryocystis elektroscirrha* on the fitness of monarch butterflies (*Danaus plexippus*). *J. Invertebr. Pathol.* 74, 76–88. <https://doi.org/10.1006/jipa.1999.4853>.
- Altizer, S.M., Oberhauser, K.S., Brower, L.P., 2000. Associations between host migration and the prevalence of a protozoan parasite in natural populations of adult monarch butterflies. *Ecol. Entomol.* 25, 125–139.
- Altizer, S.M., Oberhauser, K.S., Geurts, K.A., 2004. Transmission of the protozoan parasite, *Ophryocystis elektroscirrha*, in monarch butterfly populations: implications for prevalence and population-level impacts. In: Oberhauser, K.S., Solensky, M.J. (Eds.), *The Monarch Butterfly: Biology and Conservation*. Cornell University Press, pp. 203–218.
- Barriga, P.A., Sternberg, E.D., Lefèvre, T., de Roode, J.C., Altizer, S., 2016. Occurrence and host specificity of a neogregarine protozoan in four milkweed butterfly hosts (*Danaus* spp.). *J. Invertebr. Pathol.* 140, 75–82. <https://doi.org/10.1016/j.jip.2016.09.003>.
- Bartel, R.A., Oberhauser, K.S., De Roode, J.C., Altizer, S.M., 2011. Monarch butterfly migration and parasite transmission in eastern North America. *Ecology* 92, 342–351. <https://doi.org/10.1890/10-0489.1>.
- Bekircan, Ç., Güce, M., Baki, H., Tosun, O., 2017. *Arancicystis muskarensis* n. gen., n. sp., a neogregarine pathogen of the *Anisoplia segetum* Herbst (Coleoptera: Scarabaeidae). *J. Invertebr. Pathol.* 144, 58–64. <https://doi.org/10.1016/j.jip.2017.01.014>.
- Bradley, C.A., Altizer, S., 2005. Parasites hinder monarch butterfly flight: Implications for disease spread in migratory hosts. *Ecol. Lett.* 8, 290–300. <https://doi.org/10.1111/j.1461-0248.2005.00722.x>.
- Brooks, D.R., Hoberg, E.P., 2007. How will global climate change affect parasite-host assemblages? *Trends Parasitol.* 23, 571–574. <https://doi.org/10.1016/j.pt.2007.08.016>.
- Carreno, R.A., Matrin, D.S., Barta, J.R., 1999. *Cryptosporidium* is more closely related to the gregarines than to coccidia as shown by phylogenetic analysis of apicomplexan parasites inferred using small-subunit ribosomal RNA gene sequences. *Parasitol. Res.* 85, 899–904. <https://doi.org/10.1007/s004360050655>.
- Cooper, N., Griffin, R., Franz, M., Omotayo, M., Nunn, C.L., 2012. Phylogenetic host specificity and understanding parasite sharing in primates. *Ecol. Lett.* 15, 1370–1377. <https://doi.org/10.1111/j.1461-0248.2012.01858.x>.
- Cunningham, J.P., Zalucki, M.P., 2014. Understanding heliothine (Lepidoptera: Heliothinae) pests: what is a host plant? *J. Econ. Entomol.* 107, 881–896. <https://doi.org/10.1603/ec14036>.
- Czapak, C., Albernaz, K.C., 2013. First reported occurrence of *Helicoverpa armigera* in Brazil. *Pesqui. Agropecuária Trop.* 43, 110–113. <https://doi.org/10.1590/S1983-40632013000100015>.
- Davies, T.J., Pedersen, A.B., 2008. Phylogeny and geography predict pathogen community similarity in wild primates and humans. *Proc. R. Soc. B Biol. Sci.* 275, 1695–1701. <https://doi.org/10.1098/rspb.2008.0284>.
- De Roode, J.C., Chi, J., Rarick, R.M., Altizer, S., 2009. Strength in numbers: High parasite burdens increase transmission of a protozoan parasite of monarch butterflies (*Danaus plexippus*). *Oecologia* 161, 67–75. <https://doi.org/10.1007/s00442-009-1361-6>.
- Duncan, A.B., Little, T.J., 2007. Parasite-driven genetic change in a natural population of *Daphnia*. *Evolution* 61, 796–803. <https://doi.org/10.1111/j.1558-5646.2007.00072.x>.
- Feng, H.Q., Wu, K.M., Cheng, D.F., Guo, Y.Y., 2009. Northward migration of *Helicoverpa armigera* (Lepidoptera: Noctuidae) and other moths in early summer observed with radar in northern China. *J. Econ. Entomol.* 97, 1874–1883. <https://doi.org/10.1603/0022-0493.97.6.1874>.
- Fitt, G.P., 1989. The ecology of *Heliothis* species in relation to agroecosystems. *Annu. Rev. Entomol.* 34, 17–52.
- Gandon, S., Capowiez, Y., Dubois, Y., Michalakakis, Y., Olivieri, I., 1996. Local adaptation and gene-for-gene coevolution in a metapopulation model. *Proc. R. Soc. London. Ser. B Biol. Sci.* 263, 1003–1009.
- Hellgren, O., Waldenström, J., Pérez-Tris, J., Szöllősi, E., Hasselquist, D., Krizanauskienė, A., Ottosson, U., Bensch, S., 2007. Detecting shifts of transmission areas in avian blood parasites – a phylogenetic approach. *Mol. Ecol.* 16, 1281–1290. <https://doi.org/10.1111/j.1365-294X.2007.03227.x>.
- Kaltz, O., Gandon, S., Michalakakis, Y., Shykoff, J.A., 1999. Local maladaptation in the anther-smut fungus *Microbotryum violaceum* to its host plant *Silene latifolia*: evidence from a cross-inoculation experiment. *Evolution* 53, 395–407. <https://doi.org/10.2307/2640776>.
- Kaltz, O., Shykoff, J.A., 1998. Local adaptation in host – parasite systems. *Heredity* 81, 361–370.
- Keesing, F., Belden, L.K., Daszak, P., Dobson, A., Harvell, C.D., Holt, R.D., Hudson, P., Jolles, A., Jones, K.E., Mitchell, C.E., Myers, S.S., Bogich, T., Ostfeld, R.S., 2010. Impacts of biodiversity on the emergence and transmission of infectious diseases. *Nature* 468, 647–652. <https://doi.org/10.1038/nature09575>.
- Laine, A.L., Tellier, A., 2008. Heterogeneous selection promotes maintenance of polymorphism in host-parasite interactions. *Oikos* 117, 1281–1288. <https://doi.org/10.1111/j.0030-1299.2008.16563.x>.
- Leong, K.L., Kaya, H.K., Yoshimura, M.A., Frey, D.F., 1992. The occurrence and effect of a protozoan parasite, *Ophryocystis elektroscirrha* (Neogregarinida: Ophryocystidae) on overwintering monarch butterflies, *Danaus plexippus* (Lepidoptera: Danaidae) from two California winter sites. *Ecol. Entomol.* 17, 338–342. <https://doi.org/10.1111/j.1365-2311.1992.tb01067.x>.
- Li, Q., Li, L., Tao, W., Jiang, Y., Wan, Q., Lin, Y., Li, W., 2016. Molecular investigation of *Cryptosporidium* in small caged pets in northeast China: host specificity and zoonotic implications. *Parasitol. Res.* 115, 2905–2911. <https://doi.org/10.1007/s00436-016-5076-4>.
- Little, T.J., Watt, K., Ebert, D., 2006. Parasite-host specificity: experimental studies on the basis of parasite adaptation. *Evolution* 60, 31–38. <https://doi.org/10.1554/05-316.1>.
- Loehle, C., 1995. Social barriers to pathogen transmission in wild animal populations. *Ecology* 76, 326–335.
- Máková, A., Hoblíková, A., Hypša, V., Stanko, M., Martinů, J., Kvičárová, J., 2018. Mysteries of host switching: diversification and host specificity in rodent-coccidia associations. *Mol. Phylogenet. Evol.* 127, 179–189. <https://doi.org/10.1016/j.ympev.2018.05.009>.
- McLaughlin, R.E., Myers, 1970. *Ophryocystis elektroscirrha* sp. n., a neogregarine pathogen of the monarch butterfly *Danaus plexippus* (L.) and the Florida queen butterfly *D. glipippus berenice* Cramer. *J. Protozool.* 17, 300–305.
- Norton, D.A., Carpenter, M.A., 1998. Mistletoes as parasites: host specificity and speciation. *Trends Ecol. Evol.* 13, 101–105. <https://doi.org/10.1007/978-94-011-5868-8.3>.
- Peng, X., Zhao, Q., Guo, X., Su, S., Liu, L., Li, Y., Song, C., Chen, M., 2019. Effects of variable maternal temperature on offspring development and reproduction of *Rhopalosiphum padi*, a serious global pest of wheat. *Ecol. Entomol.* <https://doi.org/10.1111/een.12796>.
- Pickles, R.O.B.S.A., Thornton, D., Feldman, R., Marques, A., 2013. Predicting shifts in parasite distribution with climate change: a multitrophic level approach. *Glob. Chang. Biol.* 19, 2645–2654. <https://doi.org/10.1111/gcb.12255>.
- Pogue, M.G., 2013. Revised status of *Chloridea* Duncan and (Westwood), 1841, for the *Heliothis virescens* species group (Lepidoptera: Noctuidae: Heliothinae) based on morphology and three genes. *Syst. Entomol.* 38, 523–542. <https://doi.org/10.1111/syen.12010>.
- Polley, L., Thompson, R.C.A., 2009. Parasite zoonoses and climate change: molecular tools for tracking shifting boundaries. *Trends Parasitol.* 25, 285–291. <https://doi.org/10.1016/j.pt.2009.03.007>.
- Poulin, R., 2006. Variation in infection parameters among populations within parasite species: intrinsic properties versus local factors. *Int. J. Parasitol.* 36, 877–885. <https://doi.org/10.1016/j.ijpara.2006.02.021>.
- Poulin, R., Krasnov, B.R., Mouillot, D., 2011. Host specificity in phylogenetic and geographic space. *Trends Parasitol.* 27, 355–361. <https://doi.org/10.1016/j.pt.2011.05.003>.
- Schmid Hempel, P., 2011. *Evolutionary Parasitology: The Integrated Study of Infections, Immunology, Ecology, and Genetics*. Oxford University Press, Oxford.
- Sternberg, E.D., Li, H., Wang, R., Gowler, C., de Roode, J.C., 2013. Patterns of host-parasite adaptation in three populations of monarch butterflies infected with a naturally occurring protozoan disease: virulence, resistance, and tolerance. *Am. Nat.* 182. <https://doi.org/10.1086/673442>.
- Stromberg, B.E., 1997. Environmental factors influencing transmission. *Vet. Parasitol.* 72, 247–264. [https://doi.org/10.1016/S0304-4017\(97\)00100-3](https://doi.org/10.1016/S0304-4017(97)00100-3).
- Vickerman, D., Michels, A., Burrows, P.A., 1999. Levels of infection of migrating monarch butterflies, *Danaus plexippus* (Lepidoptera: Nymphalidae) by the parasite *Ophryocystis elektroscirrha* (Neogregarinida: Ophryocystidae), and evidence of a new mode of spore transmission between adults. *J. Kansas Entomol. Soc.* 124–128.
- Wood, C.L., Byers, J.E., Cottingham, K.L., Altman, I., Donahue, M.J., Blakeslee, A.M.H., 2007. Parasites alter community structure. *Proc. Natl. Acad. Sci.* 104, 9335–9339. <https://doi.org/10.1073/pnas.0700062104>.
- Zalucki, M.P., Furlong, M.J., 2005. Forecasting *Helicoverpa* populations in Australia: a comparison of regression based models and a bioclimatic based modelling approach. *Insect Sci.* 12, 45–56. <https://doi.org/10.1111/j.1672-9609.2005.00007.x>.