



UvA-DARE (Digital Academic Repository)

Waldo Heliodoor Zagwijn (1928-2018): the instigator and architect of European chronostratigraphy

Hooghiemstra, H.; Hoek, W.Z.

Published in:
Geologie en Mijnbouw/Netherlands Journal of Geosciences

DOI:
[10.1017/njg.2019.9](https://doi.org/10.1017/njg.2019.9)

[Link to publication](#)

License
CC BY

Citation for published version (APA):
Hooghiemstra, H., & Hoek, W. Z. (2019). Waldo Heliodoor Zagwijn (1928-2018): the instigator and architect of European chronostratigraphy. *Geologie en Mijnbouw/Netherlands Journal of Geosciences*, 98, [e7].
<https://doi.org/10.1017/njg.2019.9>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<http://dare.uva.nl>)

Review

Cite this article: Hooghiemstra H and Hoek WZ. Waldo Heliendoor Zagwijn (1928–2018): the instigator and architect of European chronostratigraphy. *Netherlands Journal of Geosciences*, Volume 98, e7. <https://doi.org/10.1017/njg.2019.9>

Received: 7 November 2019

Accepted: 7 November 2019


Keywords:

quaternary; stratigraphy; palynology; interglacial; palaeogeography

Author for correspondence: Henry

Hooghiemstra, Email: h.hooghiemstra@uva.nl

Waldo Heliendoor Zagwijn (1928–2018): the instigator and architect of European chronostratigraphy

Henry Hooghiemstra¹ and Wim Z. Hoek² 

¹University of Amsterdam, Science Park 904, 1098XH Amsterdam, the Netherlands and ²Faculty of Geosciences, Utrecht University, Utrecht, the Netherlands

Abstract

On 26 June 2018 Waldo Heliendoor Zagwijn died at the age of 89. He was an Emeritus Professor of the Faculty of Earth Sciences, Vrije Universiteit in Amsterdam. As a geologist, palynologist and palaeobotanist he focused on developing a stratigraphy of the Netherlands based on changes in vegetation and climate. The Dutch setting of a subsiding basin, and the clear signal of a sequence of glacial–interglacial cycles, was promising. As early as the late 1950s it became clear that the Quaternary Period included more than the previously assumed four ice ages in the Netherlands. In his PhD thesis Zagwijn defined the start of the Quaternary around 2.5 million years before the present (2.5 Ma). The international community accepted Zagwijn's arguments after he retired. He showed how the rivers Meuse, Scheldt and Rhine had built the Netherlands in four dimensions. He is the instigator and architect of the climate- and chronostratigraphy of the Quaternary Period of Western Europe.

Early education

Waldo Zagwijn was born on 16 October 1928 in The Hague. His father, a composer, had married when he was 50. Waldo's uncle was a music conductor but Waldo did not inherit the family's musical genes. His mother was much younger and had travelled a lot in her 20s, to Africa and elsewhere. Later, she worked as a telephone operator. She had a collection of minerals, and perhaps this interest explains Waldo's second name 'Heliendoor' (Al₂Be₃, a light-green mineral). Is it surprising that Waldo became a geologist?

Waldo attended the Haganum Gymnasium in The Hague where the well-known Dr Schierbeek taught biology. During the war he persuaded Waldo to spend a summer in the province of Limburg, where he collected fossils in the famous 'Heimansgroeve'. Waldo collected fossil seeds and fruits, and his interest in Earth sciences mushroomed. When Waldo left the gymnasium in 1947, studying geology was an obvious choice.

Academic education in Leiden

Waldo studied geology at Leiden University from 1947 to 1952. He was taught by professors Escher, Niggli, de Sitter, Van der Vlerk (geology), Florschütz (palaeobotany) and Lam (botany). He chose to study the Quaternary, a field of research in which his teachers Van der Vlerk and Florschütz had already achieved significant recognition. Florschütz did pioneering work in palynology in the 1930s and is considered the founding father of Dutch palynology. Later, Waldo Zagwijn played an important part in the application of palynology in applied geology, particularly in the 1950s and 1960s. Within five years, while also having a job teaching students the basics of palynology, Waldo finished his studies (July 1952). Florschütz and Van der Vlerk must have been inspiring teachers. Thomas van der Hammen writes in his obituary on Florschütz, 'Nearly three months before his death I received a letter from Van der Vlerk full of the same spirit as 25 years before. His mental capacity had in no way diminished and it seemed as if he had not aged' (modified after Van der Hammen, 1974).

Van der Vlerk and Florschütz published a very inspiring book *Nederland in het ijstijdvak; de geschiedenis van flora, fauna en klimaat, toen aap en mammoet ons land bewoonden* ('The Netherlands during the Ice Ages; history of flora, fauna and climate, when ape and mammoth lived in our country') (Van der Vlerk & Florschütz, 1950). The concepts explained in this book had a substantial influence on the development of Quaternary research in the Netherlands and on the education of a new generation of geo-ecologists, including Thomas van der Hammen and Waldo Zagwijn. Zagwijn took no interest in plants a priori, but he realised how this type of information could supplement regular geological information. Zagwijn became an all-round man early in his career. In 1952 he obtained his doctorate *summa cum laude* with geology, palaeontology and botany as the main subjects. Then, he joined the army.

© The Author(s) 2019. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

Development of pollen analysis

Between the two world wars, palynological research developed rapidly (Birks & Berglund, 2017), becoming an essential instrument in stratigraphical correlation and relative age determination. Based on a geological–palaeobotanical–palaeoecological context, Waldo was able to reconstruct the regional climate history. Subsequently he compared the climate history of the Netherlands to that of Western and Mediterranean Europe. He was interested in relatively warm intervals (interglacials and interstadials) for practical reasons. During warm intervals the Netherlands was covered by vegetation, leaving evidence (seeds, fruits, pollen) in the sediments (Fig. 1). The lack of this palaeobotanical evidence in glacial deposits leads to the conclusion that during the coldest phases of glacial periods, the Netherlands had hardly any vegetation cover. Zagwijn placed the changing patterns of vegetation in a four-dimensional time–space image and reconstructed the palaeogeographical evolution of Western Europe. In the Netherlands pollen analysis was applied in e.g. archaeology, palaeoclimatology and industry, and in many government-funded projects varying from geological mapping to infrastructural works.

A career at the Geological Survey

In April 1954 Waldo was employed at the Geological Survey. One of his first duties was to contribute several chapters of the *Geological History of the Netherlands*, edited by Pannekoek (1956), which served as an explanatory volume to the *Geological Map of the Netherlands* on the scale of 1:200,000 published by the Royal Netherlands Geological and Mining Society (KNGMG) and the Geological Survey (RGD). The contents of this volume gave an early glimpse of his future career, with his contributions on: continental Miocene, continental Pliocene, Praetiglian, Tiglian, Needian, stratigraphy and periglacial phenomena of the Tubantian, aeolian sands of the Tubantian, Older Holocene deeper peat, peat, and recent blown sand and inland dunes. Together with Jan Zonneveld he wrote one of his first papers on the complexity of the Cromerian (Zagwijn & Zonneveld, 1956). At the Geological Survey, he became head of the Department of Palaeobotany and Stratigraphy and was encouraged to study for a PhD. Professor Van der Vlerk, studying the Pleistocene of the Netherlands, was his supervisor. In contrast to older geological periods, defined on the basis of biostratigraphic evidence, Pleistocene stratigraphy was based on geomorphological evidence. The four Pleistocene ice ages of Western Europe (Günz, Mindel, Riss and Würm) had been defined on the evidence of moraines and river terraces using the terminology of Penck and Brückner (1901–1909).

While working on his thesis Zagwijn published papers showing that the classical subdivision of the Pleistocene was too simplistic (Zagwijn, 1956, 1957). In 1960 Zagwijn was awarded a PhD *summa cum laude* for his thesis ‘Aspects of the Pliocene and early Pleistocene vegetation of the Netherlands’. He was the first to show that the Pleistocene included more than four ice ages (Fig. 2). In Waldo’s thesis (Zagwijn, 1960) the warm period of the Tiglian was not addressed, as Florschütz had already reserved this period for his own studies. Later, Waldo also published a paper about the Tiglian (Zagwijn, 1963a) and a well-illustrated synthesis of his thesis for a wider audience (Zagwijn, 1963b).

Waldo married Reinske Sjoerds, a petrological laboratory technician at the Geological Survey. They had two sons, Peter (1965) and Erik (1968). Waldo was a well-organised man who worked full days at the Geological Survey but brought little work home.



Fig. 1. Waldo Zagwijn in his favourite surroundings, studying plant remains in the Belfeld Clay in a pit at Maalbeek, 1992 (after de Jong, 1995).

He had friendly contact with many colleagues, but maintained some distance as he liked his privacy.

Pollen analysis and palaeobotany were not the main objectives, but he used vegetation and climate history as a way to develop Quaternary stratigraphy. After Libby had discovered the radiocarbon-dating method Waldo immediately applied it and was able to date his earlier-published ‘Amersfoort’ interstadial at 64,000 years BP (uncalibrated) (Zagwijn, 1961). The Geological Survey focused on geological mapping and applied geology but the relevance of using palaeobotany and palynology as a scientific instrument was well-recognised. Zagwijn was often out in the field, particularly in the area around Reuver and Tegelen in the Southeastern Netherlands, near the Dutch–German border (see also Westerhoff, 2009). He was in close contact with survey geologists in Krefeld, Nordrhein-Westphalia. At this time, the opencast lignite mines in the Lower Rhine Embayment were opened and steadily excavated to a depth of 400 m. Well-developed lowland sequences of Miocene, Pliocene and Pleistocene deposits were exposed there and the stratigraphy could be studied to new levels of detail. Later in his career Zagwijn also included the Holocene period in his studies.

Conceptual contributions to the start of the Quaternary

Zagwijn was much involved in the definition of the base of the Pleistocene. In 1957 he had already argued that the Pleistocene should start at the moment when thermophilous trees began to disappear from the European flora: this was during the cold period of the Pretiglian (Zagwijn, 1957). He showed that climate cycles had also occurred in previous periods of the Pliocene but with less impact on Western European vegetation.

Zagwijn had expanded his terrestrial chronostratigraphy with sequences in Belgium (Zagwijn & Paepe, 1968; Paepe & Zagwijn, 1972), Germany (Zagwijn & Menke, 1971) and France




Fig. 2. The early and middle Pleistocene division nomenclature developed by Zagwijn (after Cohen & Gibbard 2019). Courtesy of K. Cohen.

(Suc & Zagwijn, 1983; Zagwijn & Suc, 1984) and found the climate cycles largely corresponded with oscillations in the marine oxygen isotope record. While more and more $\delta^{18}\text{O}$ records had been constructed from long marine sediment cores, the marine $\delta^{18}\text{O}$ record such as the V28-239 record of Shackleton & Opdyke (1976) had grown into a detailed record of global significance. In contrast, Zagwijn had to study short sediment cores and exposed profiles from many different locations, and his job was to complete a difficult jigsaw puzzle. His locations were close to the expansions of the glacial ice sheets, and the sedimentary archive had either partially been eroded or was no longer *in situ*. All the parts of the puzzle showed short and incomplete intervals of the Quaternary. In Western Europe the first long and continuous sedimentary archive covering a full glacial–interglacial cycle is found in northern France (Woillard & Mook, 1982). Zagwijn's ability to deal with a four-dimensional dataset allowed him to build a solid chronostratigraphical framework from many short intervals and from a suite of locations. His encyclopaedic knowledge of the details of many published pollen records was very helpful. There is an interesting correspondence with Nick Shackleton who also remembered the details of many $\delta^{18}\text{O}$ records. In discussions after congress presentations Zagwijn showed his erudition by knowing the evidence; discussions were always sharp, with much content, but sympathetic.

In the 1970s, discussions focused on the number and intensity of Pleistocene glaciations affecting Northwest Europe (de Jong, 1988). A paper in collaboration with English colleagues Phil Gibbard and Richard West well reflects the state of knowledge at the end of the 1980s (Gibbard et al., 1991). Marine geologists preferred to place the start of the Pleistocene at c.1.77 Ma, close to the start of the Olduvai palaeomagnetic subchron. However, considering climate, landscape and biotic evolution on the continents, changes in conditions around 1.77 Ma had little significance. Zagwijn saw the earlier slotted Pretiglian interval as the time when thermophilous trees disappeared from the Western European flora – more strongly than in Pliocene climate fluctuations before. The east–west orientation of European mountains (Alps, Pyrenees), and the southward expansion of the Scandinavian ice sheet that squeezed the latitudinally organised European vegetation zones, was leaving powerful evidence in the sedimentary archive in this part of the world, whereas in other parts of continents conditions and outcomes were different. Therefore, using 'a first ice age' to explain changing biogeographical patterns may be misleading. This process of migrating vegetation zones is well illustrated in the chapter 'The floral record of the late Cenozoic of Europe' (Van der Hammen et al., 1971), a classic paper still frequently cited.

Zagwijn used palaeobotanic evidence and its implications to build a chronostratigraphic framework. Every time he found the Pretiglian to be the logical start of the Quaternary period (Zagwijn, 1963a, 1974a, 1975a, 1982, 1985, 1992). Zagwijn defined several warm interglacials and interstadials and named them. On the basis of regional vegetation reconstructions he showed that up to one million years ago, glacial refugia of thermophilous trees were relatively close to the Netherlands (Central France and Southern Germany), whereas during the last million years these were further away (near the Alps and Mediterranean shores). In terms of palynologically inferred temperature fluctuations, there was good correlation with the ocean-averaged marine $\delta^{18}\text{O}$ record (e.g. the LR04 stack of Lisiecky & Raymo, 2005) that showed similar differences in intensity before and after c.1 Ma, a change dubbed 'the mid-Pleistocene revolution' (e.g. Head & Gibbard, 2005). The early and middle Pleistocene division nomenclature developed by Zagwijn is still used today (Fig. 2 after Cohen & Gibbard, 2019).

Serving geological mapping of the Netherlands, Zagwijn was involved in the lithostratigraphical division and nomenclature of the Dutch Quaternary (Zagwijn & Van Staaldin, 1975). In the pre-Quaternary period Zagwijn had recognised several less significant cool intervals, such as the Reuverian. Based on the presence of thermophilous trees such as *Carya*, *Pterocarya* and *Tsuga*, Zagwijn defined the Eburonian glacial, the Waalian interglacial and the Menapian glacial periods (Zagwijn, 1957, 1960). Based on the same arboreal taxa, he argued that the early–middle Pleistocene transition should be placed between the Menapian and the Cromerian Complex. He studied the interglacial Bavelian Complex, chronostratigraphically positioned between the Menapian and the Cromerian Complex (Zagwijn et al., 1971), and marking the end of the early Pleistocene, with Jan de Jong (Zagwijn & de Jong, 1984). He worked on the Holsteinian and Saalian organic beds in the Middle Pleistocene (de Ridder & Zagwijn, 1962; Zagwijn, 1973, 1978). For the Eemian of the Netherlands, he published a sea-level record (Zagwijn, 1983, 1996). In the late Pleistocene Zagwijn studied the Amersfoort, Hengelo and Moershoofd interstadials (Zagwijn, 1961, 1974b, 1989a; van der Hammen et al., 1967). Towards the end of his career he studied the Eemian and Holocene interglacials, using marker taxa to build consistent climate reconstructions (Zagwijn, 1994, 1996). Together with Saskia Jelgersma, Jan de Jong and the physical geographer Jean François van Regteren Altena, Zagwijn studied the coastal dunes in profiles several kilometres long near IJmuiden, along the Noordzeekanaal and in the Waterleidingduinen (dune area with infiltration basins to produce drinking water for Amsterdam). They identified the so-called Old Dunes and Young Dunes between the villages of Bergen and Monster. Radiocarbon-dated soil horizons reflected alternating dry and wet periods. The authors show an increasing human impact on the dune area since the late Middle Ages. These studies contributed to understanding of the geology, archaeology and vegetation history of the coastal dune landscape (Zagwijn, 1965, 1969, 1971, 1984; Jelgersma et al., 1970; van der Maarel & Zagwijn, 1971). This understanding aggregated in the monumental series of books *Wilde Planten; flora en vegetatie in onze natuurgebieden* ('Wild plants, flora and vegetation of our nature reserves') (Westhoff et al., 1971) in which Zagwijn showed how the present-day surface of the Netherlands is related to the complex structure of the subsurface.

The Geological Survey was mapping the North Sea basin, mainly for aggregates and other mineral resources; however, for Zagwijn the North Sea basin sediments were of importance in developing palaeomagnetic records. After the start of the Pleistocene had been defined (Zagwijn, 1960), palaeomagnetism showed 2.3 million years to be the base of the Quaternary (Van Montfrans, 1971). Where Zagwijn placed the start of the Pleistocene in the North Sea basin, sediment accumulation increased tenfold. Tectonic uplift of the hinterland was a possible explanation; however, according to Zagwijn, the increasing amplitude of climate fluctuations from the Pretiglian onwards (2.56–2.44 Ma) meant that Western Europe was mostly north of the forest line during much of a glacial period, and forest was gradually replaced by steppe, tundra and polar deserts. With the help of palaeomagnetic records Zagwijn and Suc (Montpellier) found a reliable correlation between the chronostratigraphies of Western Europe and the Mediterranean area (Suc & Zagwijn, 1983; Zagwijn & Suc, 1984).

The start of the Quaternary period has long been discussed. Arguments from marine geology (a start at 1.77 Ma) long prevailed over compelling evidence from the continents (a start at c. 2.6 Ma). In a summary at the end of his career, Zagwijn again argued for a

start around 2.3 Ma (Zagwijn, 1992); as he was nearing retirement he could make a definite statement: *‘Despite recent attempts, in particular by marine palaeontologists, to minimize the status of the Quaternary and even to deny it a place of its own in the stratigraphic scale, the present author is of the opinion that the exceptional climatic evolution that the planet Earth has witnessed in its latest history and which resulted in repeated large-scale glaciation in the northern hemisphere, is a sufficiently valid criterion to assign this interval a status of its own on the Period level.’* When he was 82 the International Commission on Stratigraphy of the International Union of Geological Sciences (IUGS-ICS) formalised the redefinition of the start of the Pleistocene at *c.*2.58 Ma (Gibbard et al., 2010). This was not only a recognition of the concept published in Zagwijn’s 1960 thesis, but also a plea for a more prominent role of climate swings, also reflected in long palynological records, in the stratigraphy of the Quaternary (Suc et al., 2018). Zagwijn did not see this acceptance as a personal victory; he had his own reasons for placing the start of the Pleistocene around 2.3 Ma, and he knew that other criteria are used in other regions. Zagwijn was too much of a scientist to be excited by other colleagues accepting or not his arguments.

Palaeogeographical maps

Probably during two glacial periods of the Pleistocene only, the northern part of the Netherlands was covered by ice. This caused the courses of the rivers Meuse, Scheldt and Rhine to change repeatedly. Zagwijn drew maps and showed in a spatial–temporal framework how the Netherlands were shaped during successive stages of the Quaternary (Zagwijn, 1974c, 1975b, 1979, 1989b; Van Staaldunin et al., 1979) and he became well known for his popular scientific palaeogeographical maps of the Netherlands, showing how the Low Countries got their shape over Pleistocene and Holocene times (Zagwijn, 1986). At the turn of the twenty-first century, palaeogeographical understanding of the Netherlands continued to expand, partly inspired by the work of Waldo Zagwijn (e.g. Berendsen & Stouthamer, 2001; Busschers, 2008; Westerhoff, 2009; Peeters et al., 2016). The set of 12 palaeogeographical maps in the *Atlas van Nederland in het Holoceen* (Vos et al., 2011) and updated versions (Vos & De Vries, 2013) show the impressive detail currently reached for different parts of the Netherlands – owing to continued data collection and computerised data combination (Hoek, 1997b; Van der Meulen et al., 2013; Pierik et al., 2016; Cohen et al., 2017a,b; Woolderink et al., 2019) as well as to the inspiration of Zagwijn’s palaeogeographical legacy (Berendsen & Stouthamer, 2001; Westerhoff, 2009; Vos, 2015).

Zagwijn’s expertise was frequently consulted

Zagwijn’s research was key to palaeobotany, geology, climatology, physics and soil science coming together. Florschütz (1887–1965) had developed the foundations and contacts between academia, industry, museums and governmental organisations. Zagwijn expanded this field using new methods (C14-dating, palaeomagnetism) and new proxies. His personal archive contains much correspondence with colleagues from institutes representing many disciplines such as geology, oceanography, anthropology, archaeology, botany, zoology and palynology. Correspondence with governmental offices also increased substantially: foreign geological surveys and biostratigraphical companies. In the Netherlands Zagwijn was consulted by governmental offices about drinking-water, the establishment of polders in the IJsselmeer, the

construction of canals in Twente, and by the Directorate-General for Public Works and Water Management (Rijkswaterstaat) for coastal protection. Many of his colleagues were well trained in thinking long-term. Robert C. Bright from the University of Minnesota wrote to Saskia Jelgersma – in a letter forwarded to Zagwijn because he was a member of the Netherlands’ Radiocarbon Dating Commission – on results of the samples he had submitted: his letter starts ‘It seems like a million years ago when we had a short meeting in Boulder . . .’

Zagwijn’s archive is an impressive source of information and it shows he was frequently consulted by colleagues about the interpretation of new data. Although time-consuming, it contributed to an ever-improving understanding of the climate stratigraphy of Western Europe. It is hardly surprising that he was often asked to contribute to national and international commissions; to name but a few: the IUGS-ICS Subcommittee for Stratigraphic Classification; the INQUA (International Union for Quaternary Research) Subcommittee on the Pleistocene of Europe (1967–1993); the Scientific Committee UNESCO–IUGS International Geological Correlation Programme (1976–1981); the Commission for Biosociology and Palaeobotany of the Royal Netherlands Botanical Society (1958–1961); the Foundation for Isotope Geological Research (1970–1985); the ministerial Advisory Commission of Natural History Museums (1981–1993); and the section Earth Sciences of the Royal Netherlands Academy of Sciences (1986–1993). His contributions to the INQUA were fundamental. He was a member, later chairman, of the INQUA-Netherlands Commission (1963–1993), and a member of the INQUA Commission for the Palaeogeographic Atlas of the Quaternary.

Zagwijn was a member of the editorial boards of *Review of Palaeobotany and Palynology* (1966–1971), *Géologie Méditerranéenne* (1975–1993), *Bulletin de l’Association Française pour l’Etude du Quaternaire* (1980–1993), *Acta Palynologica* (1989–1993) and *Quaternary International* (1989–1993); he was also an external examiner of PhD committees in the Netherlands and France.

Both Zagwijn and Van der Hammen were involved in the famous Climate/Long Range Investigation Mapping and Predictions Project (CLIMAP) of the US National Science Foundation. In 1974, in Amsterdam, they organised a workshop entitled ‘The climate record of the last million years’, in which top international scientists were involved. The results of the CLIMAP project were cited for decades, showing that the understanding of this group of scientists was far ahead of general societal perception of the impact of climate change.

International recognition

For a long time Zagwijn was active in the German Quaternary Association (DEUQUA) and he received the Albrecht Penck Medal in 1972. In 1974, Zagwijn and Van der Hammen received the Van Waterschoot van der Gracht Medal of the Royal Geological and Mining Society of the Netherlands (KNGMG). In 1980, Zagwijn was appointed a member of the Royal Netherlands Academy of Sciences (KNAW) and in 1983 he was appointed *Officier in de Orde van Oranje Nassau* by the queen. He was elected an Honorary Life Fellow of INQUA in 1987, and an Honorary Member of the Quaternary Research Association in 1994.

Professor at Vrije Universiteit Amsterdam

In 1989 Zagwijn was appointed Professor of ‘Quaternary Palynology’ at Vrije Universiteit in Amsterdam (Fig. 3). He was



Fig. 3. Waldo Zagwijn interviewed on the occasion of his appointment at Vrije Universiteit Amsterdam in 1989. Courtesy of Mrs. R. Zagwijn-Sjoerds and collection photo press agency De Boer, Haarlem.

proud of this move so near the end of his career and held his inaugural lecture on Interglacials (Zagwijn, 1991a). His teaching as professor included a fascinating series of lectures on the Quaternary vegetation history of Europe (Zagwijn, 1991b). His unprecedented knowledge of the evidence was obvious and it became clear why he was the person who had completed the climate stratigraphical jigsaw puzzle. As there was only a short time before his retirement Zagwijn supervised only one PhD student, Wim Hoek, who got his PhD with the thesis 'Palaeogeography of Lateglacial vegetations' (Hoek, 1997a,b). This subject had developed in the 1970s but Zagwijn himself was unable to compile that much data; he had to wait for this unique opportunity. The Lateglacial, though, was a period which he investigated for his very first scientific publication (Zagwijn, 1952).

As well as teaching at Vrije Universiteit for one day a week, Zagwijn was involved in developing the new building for the Geological Survey. Hindered by delays in the building process, Zagwijn presented his inaugural lecture, entitled *Interglacieren* ('Interglacials') on 20 March 1991 (Zagwijn, 1991a). The new building was opened in 1992, coinciding with Zagwijn's retirement from the Geological Survey, but he worked a further year at the university to finish several projects. During his four years of professorship Zagwijn developed climate reconstructions of the last two interglacial periods, the Eemian and Holocene (Zagwijn 1994, 1996).

Professor Zagwijn as an Emeritus

In 1993, on the occasion of Zagwijn's retirement from Vrije Universiteit in Amsterdam, an international symposium was organised in the Royal Academy of Sciences building. In a special issue of *Mededelingen van de Rijks Geologische Dienst*, entitled *Neogene and Quaternary Geology of North-West Europe; Contribution on the occasion of Waldo H. Zagwijn's retirement*, 34 papers written by 67 of his colleagues were published

(Herngreen & Van der Valk, 1995). This special issue also includes a list of the almost 100 publications by Zagwijn (De Jong, 1995).

Zagwijn had decided to end his scientific work after retirement. He devoted his time to his family, to travelling and to his hobbies. In 2004 he was hospitalised and needed a period of convalescence. About ten years after this his mental health deteriorated and he spent his last year in a nursing home, where he died. Zagwijn was cremated after a funeral with only family members present.

We remember Waldo Zagwijn as an eminent, erudite and modest scientist with a fabulous memory for detail. He had a remarkable ability to develop an image in time and space of the Netherlands below the surface. He developed early understanding of the Pliocene and Quaternary palaeogeographical, geological and climatological developments of Western Europe, and the Netherlands in particular. He is the instigator and architect of the chrono- and climate stratigraphy of Western Europe.

Acknowledgements. This paper is an abridged version of an obituary in Dutch published at the Royal Netherlands Academy of Sciences (KNAW). H.H. thanks Noord-Hollands Archief (Haarlem) for permission to consult Zagwijn's personal archive, and also thanks Mrs Zagwijn and Jan de Jong for interviews, Phil Gibbard and Thijs van Kolfschoten for helpful correspondence, and Jean-Pierre Suc and co-authors for their informative paper about Zagwijn's work in southern Europe. Kim Cohen is thanked for constructing the stratigraphic chart and his useful comments. Jeroen Schokker is thanked for improving a final draft. We thank Nicolas Broadbridge for improving the English text.

References

- Berendsen, H.J.A. & Stouthamer, E.**, 2001. Palaeogeographic development of the Rhine-Meuse delta, the Netherlands. Van Gorcum (Assen).
- Birks, H.J.B. & Berglund, B.E.**, 2017. One hundred years of Quaternary pollen analysis 1916-2016. *Vegetation History Archaeobotany* 27(2): 271-309.
- Busschers, F.S.**, 2008. Unravelling the Rhine. PhD Thesis. Vrije Universiteit Amsterdam (Amsterdam), Geology of the Netherlands 1, TNO & Geological Survey of the Netherlands.
- Cohen, K.M. & Gibbard, P.L.**, 2019. Global chronostratigraphical correlation table for the last 2.7 million years, version 2018 QI-500. *Quaternary International* 500: 20-31.
- Cohen, K.M., Dambrink, R., De Bruijn, R., Marges, V.C., Erkens, G., Pierik, H.J., Koster, K., Stafleu, J., Schokker, J. & Hijma, M.P.**, 2017a. Mapping buried Holocene landscapes: past lowland environments, palaeoDEMs and preservation in GIS. In: Lauwerier *et al.* (eds): Knowledge for informed choices. Netherlands Archeological Reports 55. Rijksdienst Cultureel Erfgoed (Amersfoort): 73-95.
- Cohen, K.M., Dambrink, R., De Bruijn, R., Schokker, J. & Hijma, M.P.**, 2017b. Vervaardiging van hoogtemodellen en landschapskaarten naar periode en diepte voor archeologisch gebruik in Holoceen-afgedekte delen van Nederland. *Deltares Reports* 1210450-000-BGS-0012.
- De Jong, J.**, 1988. Climatic variability during the past three million years, as indicated by vegetational evolution in northwest Europe and with emphasis on data from The Netherlands. *Philosophical Transactions of the Royal Society of London* B318: 603-617.
- De Jong, J.**, 1995. At the retirement of Prof. Dr. W.H. Zagwijn: review of his work and bibliography. *Mededelingen Rijks Geologische Dienst* 92: 13-19.
- De Ridder, N.A. & Zagwijn, W.H.**, 1962. A mixed Rhine-Meuse deposit of Holsteinian age from the south-eastern part of The Netherlands. *Geologie en Mijnbouw* 41: 125-130.
- Gibbard, P.L., West, R.G., Zagwijn, W.H., Balson, P.S., Burger, A.W., Funnell, B.M., Jeffrey, D.H., De Jong, J., Van Kolfschoten, T., Lister, A.M., Meijer, T., Norton, P.E.P., Preece, R.C., Rose, J., Stuart, A.J., Whiteman, C.A. & Zalasiewicz, J.A.**, 1991. Early and Early Middle Pleistocene correlations in the southern North Sea basin. *Quaternary Science Reviews* 10: 23-52.
- Gibbard P.L., Head M.J. & Walker, M.J.C.**, 2010. Formal ratification of the Quaternary System/Period and the Pleistocene Series/Epoch with a base at 2.58 Ma. *Journal of Quaternary Science* 25: 96-102.

- Head, M.J. & Gibbard, P.L.** (eds), 2005. Early-Middle Pleistocene transitions: an overview and recommendation for the defining boundary. Geological Society of London, Special Publication 247(1): 1–18.
- Herngreen, G.F.W. & Van der Valk, L.** (eds), 1995. Neogene and Quaternary geology of North-West Europe. Mededelingen Rijks Geologische Dienst 52: 1–510.
- Hoek, W.Z.**, 1997a. Palaeogeography of Lateglacial vegetations: aspects of Lateglacial and early Holocene vegetation, abiotic landscape, and climate in the Netherlands. PhD Thesis, Volume I, Netherlands Geographical Studies (Utrecht–Amsterdam) 230: 160 pp.
- Hoek, W.Z.**, 1997b. Atlas to Palaeogeography of Lateglacial vegetations: maps of Lateglacial and early Holocene landscape and vegetation in the Netherlands, with an extensive review of available palynological data. PhD Thesis, Volume II, Netherlands Geographical Studies (Utrecht–Amsterdam) 231: 176 pp.
- Jelgersma, S., De Jong, J., Zagwijn, W.H. & Van Regteren Altena, J.F.**, 1970. The coastal dunes of the western Netherlands: geology, vegetational history and archaeology. Mededelingen Rijks Geologische Dienst (Nieuwe Serie) 21: 93–167.
- Lisiecky, L.E. & Raymo, M.E.**, 2005. A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records. *Paleoceanography* 20: PA1003, 1–17.
- Paepé, R. & Zagwijn, W.H.**, 1972. Possibilités de corrélation des dépôts Weichseliens de la Belgique et des Pays-Bas. Bulletin Association Française pour l'étude du Quaternaire 1: 59–69.
- Pannekoek, A.J.** (ed.), 1956. Geological history of the Netherlands: explanation to the General Geological Map of The Netherlands on the scale of 1:200,000. Government Printing and Publishing Office (The Hague): 147 pp.
- Peeters, J., Busschers, F.S., Stouthamer, E., Bosch, J.H.A., Van den Berg, M.W., Wallinga, J., Versendaal, A.J., Bunnik, F.P.M. & Middelkoop, H.**, 2016. Sedimentary architecture and chronostratigraphy of a late Quaternary incised-valley fill: a case study of the late Middle and Late Pleistocene Rhine system in the Netherlands. *Quaternary Science Reviews* 131: 211–236.
- Penck, A. & Brückner, E.**, 1901–1909. Die Alpen im Eiszeitalter. Tauchnitz (Leipzig): 3 volumes.
- Pierik, H.J., Cohen, K.M. & Stouthamer, E.**, 2016. A new GIS approach for reconstructing and mapping dynamic late Holocene coastal plain palaeogeography. *Geomorphology* 270: 55–70.
- Shackleton, N.J. & Opdyke, N.D.**, 1976. Oxygen isotope and paleomagnetic stratigraphy of Pacific core V28-239, Late Pliocene to Latest Pleistocene. *Geological Society of America Memoir* 145: 449–464.
- Suc, J.-P. & Zagwijn, W.H.**, 1983. Plio-Pleistocene correlations between the northwestern Mediterranean region and northwestern Europe according to recent biostratigraphic and palaeoclimatic data. *Boreas* 12: 153–166.
- Suc, J.-P., Popescu, S.-M., Bertini, A., Head, M.J., Gibbard, P.L. & Diniz, F.**, 2018. The pioneering contributions and legacy of Waldo H. Zagwijn (1928–2018). *Alpine and Mediterranean Quaternary* 31: iii–viii.
- Van der Hammen, T.**, 1974. In memoriam prof. dr. I.M. van der Vlerk. *Geologie en Mijnbouw* 53: 241–243.
- Van der Hammen, T., Wijmstra, T.A. & Zagwijn, W.H.**, 1971. The floral record of the Late Cenozoic of Europe. In: Turekian, K.K. (ed.): Late Cenozoic glacial ages. Yale University Press (New Haven–London): 391–424.
- Van der Hammen, T., Maarleveld, G.C., Vogel, J.C. & Zagwijn, W.H.**, 1967. Stratigraphy, climatic succession radiocarbon dating of the Last Glacial in The Netherlands. *Geologie en Mijnbouw* 46: 79–95.
- Van der Maarel, E. & Zagwijn, W.H.**, 1971. Historic and dynamic aspects of coastal dune vegetations in the Netherlands. *Acta Botanica Neerlandica* 20: 173–182.
- Van der Meulen, M.J., Doornenbal, J.C., Gunnink, J.L., Stafleu, J., Schokker, J., Vernes, R.W., Van Geer, F.C., Van Gessel, S.F., Van Heteren, S., Van Leeuwen, R.J.W., Bakker, M.A.J., Bogaard, P.J.F., Busschers, F.S., Griffioen, J., Gruijters, S.H.L.L., Kiden, P., Schroot, B.M., Simmelink, H.J., Van Berkel, W.O., Van der Krogt, R.A.A., Westerhoff, W.E. & Van Daalen, T.M.**, 2013. 3D geology in a 2D country: perspectives for geological surveying in the Netherlands. *Journal Geosciences* 92: 217–241.
- Van der Vlerk, I.M. & Florschütz, F.**, 1950. Nederland in het ijstijdvak. De geschiedenis van flora, fauna en klimaat toen aap en mammoet ons land bewoonden. De Haan (Utrecht): 287 pp.
- Van Montfrans, H.M.**, 1971. Paleomagnetic dating in the North Sea basin. PhD Thesis. University of Amsterdam (Amsterdam).
- Van Staalduinen, C.J., Van Adrichem-Bogaert, H.A., Bless, M.J.M., Doppert, J.W.C., Harsveldt, H.M., Oele, E., Van Montfrans, H.M., Wermuth, R.A. & Zagwijn, W.H.**, 1979. The geology of the Netherlands. Mededelingen Rijks Geologische Dienst 31: 9–49.
- Vos, P.C.**, 2015. Origin of the Dutch coastal landscape: long-term landscape evolution of the Netherlands during the Holocene described and visualized in national, regional and local palaeogeographical map series. PhD Thesis. University of Utrecht & Barkhuis (Groningen).
- Vos, P.C., Bazelmans, L., Weerts, H.J.T. & Van der Meulen, M.J.** (eds), 2011. Atlas van Nederland in het Holoceen. Bert Bakker (Amsterdam): 94 pp.
- Vos, P.C. & De Vries, S.**, 2013. 2e generatie palaeogeografische kaarten van Nederland (versie 2.0). Deltares, Utrecht & Rijksdienst Cultureel Erfgoed (Amersfoort).
- Westerhoff, W.E.**, 2009. Stratigraphy and sedimentary evolution: the lower Rhine-Meuse system during the Late Pliocene and Early Pleistocene (southern North Sea basin). PhD Thesis. Vrije Universiteit Amsterdam (Amsterdam), Geology of the Netherlands 2, TNO & Geological Survey of the Netherlands.
- Westhoff, V., Bakker, P.A., Van Leeuwen, C.G. & Van der Voo, E.E.**, 1971. Wilde planten; flora en vegetatie in onze natuurgebieden. Vereniging tot Behoud van Natuurmonumenten in Nederland (Amsterdam): 3 volumes.
- Woillard, G.M. & Mook, W.G.**, 1982. Carbon-14 dates at Grande Pile: correlation of land and sea chronologies. *Science* 215: 159–161.
- Woolderink, H.A.G., Kasse, C., Cohen, K.M., Hoek, W.Z. & Van Balen, R.T.**, 2019. Spatial and temporal variations in river terrace formation, preservation and morphology in the Lower Meuse Valley, the Netherlands. *Quaternary Research* 91: 548–569.
- Zagwijn, W.H.**, 1952. Pollenanalytische Untersuchungen einer spätglacialen Seeablagerung aus Tirol. *Geologie en Mijnbouw* 14: 235–239.
- Zagwijn, W.H.**, 1956. Zum heutigen Stand der Pollenanalytischen Untersuchungen des Pleistozäns in den Niederlanden. *Geologie en Mijnbouw (Nieuwe Serie)* 18: 426–427.
- Zagwijn, W.H.**, 1957. Vegetation, climate and time-correlations in the Early Pleistocene of Europe. *Geologie en Mijnbouw (Nieuwe Serie)* 19: 233–244.
- Zagwijn, W.H.**, 1960. Aspects of the Pliocene and early Pleistocene vegetation of the Netherlands. PhD Thesis. Leiden University (Leiden) & Mededelingen van de Geologische Stichting, Serie C-III-1(5): 1–78.
- Zagwijn, W.H.**, 1961. Vegetation, climate and radiocarbon datings in the Late Pleistocene of the Netherlands. Part 1: Eemian and Early Weichselian. Mededelingen van de Geologische Stichting, Nieuwe Serie 14: 15–45.
- Zagwijn, W.H.**, 1963a. Pollen-analytic investigations in the Tiglian of the Netherlands. Mededelingen van de Geologische Stichting, Nieuwe Serie 16: 49–72.
- Zagwijn, W.H.**, 1963b. Pleistocene stratigraphy in the Netherlands based on changes in vegetation and climate. Verhandelingen Koninklijk Nederlands Geologisch Mijnbouwkundig Genootschap, Geologische Serie 21: 173–196.
- Zagwijn, W.H.**, 1965. Pollen-analytic correlations in the coastal barrier deposits near The Hague (The Netherlands). Mededelingen van de Geologische Stichting, Nieuwe Serie 17: 83–88.
- Zagwijn, W.H.**, 1969. Geologie en vegetatiegeschiedenis van de Nederlandse kustduinen. Koninklijke Maatschappij voor Natuurkunde 'Diligentia', Den Haag, Jaarboek 1968–69: 167–176.
- Zagwijn, W.H.**, 1971. Vegetational history of the coastal dunes in the western Netherlands. *Acta Botanica Neerlandica* 20: 174–182.
- Zagwijn, W.H.**, 1973. Pollenanalytic studies of Holsteinian and Saalian beds in the northern Netherlands. Mededelingen Rijks Geologische Dienst, Nieuwe Serie 24: 139–156.
- Zagwijn, W.H.**, 1974a. The Pliocene–Pleistocene boundary in western and southern Europe. *Boreas* 3: 75–97.
- Zagwijn, W.H.**, 1974b. Vegetation, climate and radiocarbon datings in the Late Pleistocene of the Netherlands. Part II: Middle Weichselian. Mededelingen Rijks Geologische Dienst, Nieuwe Serie 25: 101–110.
- Zagwijn, W.H.**, 1974c. The palaeogeographic evolution of the Netherlands during the Quaternary. *Geologie en Mijnbouw* 53: 369–385.
- Zagwijn, W.H.**, 1975a. A model-theory for the Pliocene-Pleistocene boundary determination, based on past climate changes. In: Saito LT, Burkle LH (eds): Late Neogene epoch boundaries. *Micropalaeontology*, Special Publication 1: 71–74.

- Zagwijn, W.H.**, 1975b. De paleogeografische ontwikkeling van Nederland in de laatste drie miljoen jaar. *Geografisch Tijdschrift* **9**: 181–201 (with 12 palaeogeographical maps).
- Zagwijn, W.H.**, 1978. A macroflora of Holsteinian age from the northern part of the Netherlands. *Review of Palaeobotany and Palynology* **26**: 243–248.
- Zagwijn, W.H.**, 1979. Early and middle Pleistocene coast lines in the southern North Sea basin. *In*: Oele, E., Schüttenhelm, R.T.E. & Wiggers, A.J. (eds): *The Quaternary history of the North Sea. Symposia Universitatis Upsaliensis annum quingentesimum celebrantis, Book 2* (Uppsala): 31–42.
- Zagwijn, W.H.**, 1982. Het begin van het IJstijdvak. *Verslag Afdeling Natuurkunde* **91**: 54–58.
- Zagwijn, W.H.**, 1983. Sea-level changes in the Netherlands during the Eemian. *Geologie en Mijnbouw* **62**: 437–450.
- Zagwijn, W.H.**, 1984. The formation of the Younger Dunes of the west coast of the Netherlands (AD 1000–1600). *Geologie en Mijnbouw* **63**: 259–268.
- Zagwijn, W.H.**, 1985. An outline of the Quaternary stratigraphy of the Netherlands. *Geologie en Mijnbouw* **64**: 17–24.
- Zagwijn, W.H.**, 1986. Nedearland in het Holoceen. *Geologie van Nederland, deel 1*, Haarlem/Den Haag: 46 pp.
- Zagwijn, W.H.**, 1989a. Vegetation and climate during warmer intervals in the Late Pleistocene of Western and Central Europe. *Quaternary International* **3–4**: 57–67.
- Zagwijn, W.H.**, 1989b. The Netherlands during the Tertiary and the Quaternary: a case history of Coastal Lowland evolution. *Geologie en Mijnbouw* **68**: 107–120.
- Zagwijn, W.H.**, 1991a. Interglacialen. Oratie, Faculteit der Aardwetenschappen, 20 maart 1991. Vrije Universiteit Amsterdam (Amsterdam).
- Zagwijn, W.H.**, 1991b. Lectures 'Vegetatiegeschiedenis van het Europese Kwartaair'. Vrije Universiteit Amsterdam (Amsterdam): **6** volumes.
- Zagwijn, W.H.**, 1992. The beginning of the ice age in Europe and its major subdivisions. *Quaternary Science Reviews* **11**: 583–591.
- Zagwijn, W.H.**, 1994. Reconstruction of climate change during the Holocene in western and central Europe based on pollen records of indicator species. *Vegetation History and Archaeobotany* **3**: 65–88.
- Zagwijn, W.H.**, 1996. An analysis of Eemian climate in Western and Central Europe. *Quaternary Science Reviews* **15**: 451–469.
- Zagwijn, W.H. & De Jong, J.**, 1984. Die Interglaziale von Bavel und Leerdam und ihre stratigraphische Stellung im niederländischen Früh-Pleistozän. *Mededelingen Rijks Geologische Dienst* **37**: 155–169.
- Zagwijn, W.H. & Menke, B.**, 1971. Arbeitsgruppe für Biostratigraphie des Pleistozäns im nordeuropäischen Vereisungsgebiet. Zusammenkunft vom 25–27.5.1971 in Haarlem. *Eiszeitalter und Gegenwart* **22**: 192–195.
- Zagwijn, W.H. & Paepe, R.**, 1968. Die Stratigraphie der weichselzeitlichen Ablagerungen der Niederlande und Belgiens. *Eiszeitalter und Gegenwart* **19**: 129–146.
- Zagwijn, W.H. & Suc, J.-P.**, 1984. Palynostratigraphie du Plio-Pléistocène d'Europe et de Méditerranée nord-occidentales: corrélations chronostratigraphiques. *Histoire de la végétation et du climat. Paléobiologie Continentale* **14**: 475–483.
- Zagwijn, W.H., Van Montfrans, H.M. & Zandstra, J.G.**, 1971. Subdivision of the 'Cromerian' in the Netherlands: pollen-analysis, paleomagnetism and sedimentary petrology. *Geologie en Mijnbouw* **50**: 41–58.
- Zagwijn, W.H. & Van Staalduinen, C.J.**, 1975. Toelichting bij Geologische overzichtskaarten van Nederland. Rijks Geologische Dienst (Haarlem): 134 pp.
- Zagwijn, W.H. & Zonneveld, J.I.S.**, 1956. The interglacial of Westerhoven. *Geologie en Mijnbouw (Nieuwe Serie)* **18**: 37–46.