

Multimedia Appendix 3. Questionnaire results on test-retest reliability and practice effects (significant *t* values in italics). HADS: Hospital Anxiety and Depression Scale; ICC: intraclass correlation coefficient; MFI: Multidimensional Fatigue Inventory; SDC: smallest detectable change; SEM: standard error of measurement.

Questionnaire	n	Mean 1 (SD)	Mean 2 (SD)	<i>t</i> value (degrees of freedom), <i>P</i> value	SEM	SDC group ^a	ICC ^c	Pearson <i>r</i> ^c	Spearman ρ ^c
HADS anxiety	248	5.09 (3.24)	4.75 (3.28)	<i>2.55</i> (247), <i>P</i> =.01	1.49	0.60	.79	.80	.78
HADS depression	248	3.11 (2.78)	3.02 (2.84)	0.69(247), <i>P</i> =.49	1.45	0.26	.74	.74	.72
MFI total	248	40.49 (15.27)	40.25 (14.99)	0.38(247), <i>P</i> =.70	7.02	1.24	.79	.79	.77

^aSDC group: SDC / \sqrt{n} .

^bFor measures with a significant practice effect (*t* value in italics) Mean1 - Mean2 difference scores were added.

^cAll correlation coefficients were significant at *P*<.001.