



UvA-DARE (Digital Academic Repository)

VLA radio detection of the very-faint X-ray transient IGR J17285-2922

van den Eijnden, J.; Degenaar, N.; Russell, T.; Wijnands, R.; Hernandez Santisteban, J.V.; Russell, D.M.; Maitra, D.; Heinke, C.O.; Sivakoff, G.R.; Shaw, A.W.; Maccarone, T.J.; Miller-Jones, J.C.A.; Bahramian, A.; Armas Padilla, M.

Publication date

2019

Document Version

Final published version

Published in

The astronomer's telegram

License

Unspecified

[Link to publication](#)

Citation for published version (APA):

van den Eijnden, J., Degenaar, N., Russell, T., Wijnands, R., Hernandez Santisteban, J. V., Russell, D. M., Maitra, D., Heinke, C. O., Sivakoff, G. R., Shaw, A. W., Maccarone, T. J., Miller-Jones, J. C. A., Bahramian, A., & Armas Padilla, M. (2019). VLA radio detection of the very-faint X-ray transient IGR J17285-2922. *The astronomer's telegram*, 12669. <http://www.astronomerstelegam.org/?read=12669>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

2 Apr 2020; 14:23 UT

This space for free for your conference.



Outside

GCN
IAUCs

Other

ATel on [Twitter](#) and [Facebook](#)
ATELstream
ATel Community Site

CCV Nederland

[[Previous](#) | [Next](#) | [ADS](#)]

VLA radio detection of the very-faint X-ray transient IGR J17285-2922

ATel #12669; *J. van den Eijnden, N. Degenaar, T. Russell, R. Wijnands (University of Amsterdam), J. V. Hernandez Santisteban (University of St. Andrews), D. M. Russell (NYU Abu Dhabi), D. Maitra (Wheaton College), C. O. Heinke, G. R. Sivakoff, A. W. Shaw (University of Alberta), T. J. Maccarone (Texas Tech University), J. C. A. Miller-Jones, A. Bahramian (ICRAR-Curtin), M. Armas Padilla (Instituto de Astrofísica de Canarias)*

on 18 Apr 2019; 15:54 UT

Credential Certification: *Jakob Van den Eijnden (a.j.vandeneijnden@uva.nl)*

Subjects: Radio, X-ray, Black Hole, Neutron Star, Transient

IGR J17285-2922 is a very-faint X-ray transient that was reported to be in outburst by INTEGRAL/IBIS-ISGRI (ATel #12646). We triggered a joint X-ray and radio monitoring campaign of this X-ray binary with the Neil Gehrels Swift Observatory (Swift) and the Karl G. Jansky Very Large Array (VLA).

We observed IGR J17285-2922 on 13 April 2019 from 09:44 to 11:43 UTC (MJD 58586.45 +/- 0.04) while the VLA was in B configuration. These observations were taken simultaneously at central frequencies of 4.5 and 7.5 GHz, each with a bandwidth of 1 GHz. 3C286 and J1743-3058 were used for primary and secondary calibration, respectively. We used the Common Astronomy Software Applications package (CASA v4.7.2; McMullin et al. 2007, ASPC, 376, 127) to calibrate and image the data, following standard procedures. Imaging was carried out using Briggs weighting with a robustness of zero to balance sensitivity and resolution.

To measure the flux density, we fitted a point source in the image plane by forcing an elliptical Gaussian with a FWHM equal to the synthesized beam (2.99" x 0.97" with a -4.5 degree position angle east of north at 4.5 GHz; 1.67" x 0.59" with a -4.75 degree position angle at 7.5 GHz). We measure flux densities of 109 +/- 7 μ Jy at 4.5 GHz and 126 +/- 7 μ Jy at 7.5 GHz. These values yield a spectral index measurement of $\alpha = 0.3 +/- 0.2$ (where the flux density $S_\nu \propto \nu^\alpha$), consistent with a self-absorbed compact jet.

The best fit source position, determined at 7.5 GHz, is:

RA (J2000) = 17h 28m 38.853s +/- 0.01s

Dec (J2000) = -29d 21' 43.181" +/- 0.06",

Related

- | | |
|-------|---|
| 12669 | VLA radio detection of the very-faint X-ray transient IGR J17285-2922 |
| 12651 | Swift/XRT detection of the very faint X-ray binary transient IGR J17285-2922 |
| 12646 | INTEGRAL detection of hard X-ray emission from the Very Faint X-ray Transient IGR J17285-2922 |
| 3011 | Pan-STARRS1 observations of the very faint X-ray transient XTE J1728-295 during quiescence |
| 2997 | Monitoring of the likely optical counterpart of XTE J1728-295 |
| 2991 | Swift/XRT follow-up of the X-ray transient XTE J1728-295 during its decaying phase |
| 2870 | Optical observations of the candidate optical counterpart to XTE J1728-295 (=IGR J17285-2922) |
| 2869 | Chandra Localization of the X-ray Transient XTE J1728-295 |
| 2827 | Candidate optical counterparts of XTE J1728-295 = IGR J17285-2922 |
| 2825 | INTEGRAL confirms that XTE J1728-295 = IGR J17285-2922 |
| 2824 | Swift/XRT follow-up and a refined position of the X-ray transient XTE J1728-295 |
| 2823 | RXTE PCA Detection of a New Outburst of XTE J1728-295 (probably IGR J17285-2922) |
| 904 | Announcement of the Swift/BAT Hard X-ray Transient Monitor |

where the errors are calculated using the standard VLA astrometric accuracy of 10% of the synthesized beam. This position is separated by 0.5" from the Chandra position reported in ATel #2869 during a 2010 outburst of this transient, which has a 0.6" 90%-confidence radius.

While no simultaneous X-ray observations were performed, the closest Swift/XRT observation taken on 10 April 2019 was reported in ATel #12651. A preliminary analysis of later Swift observations shows that the source remains at relatively constant X-ray flux. Therefore, we assume that the X-ray luminosity did not change significantly between the closest X-ray observation and our radio epoch, to place the source on the X-ray/radio luminosity plane. Adopting the assumed 8 kpc source distance from ATel #12651, we estimate a $4.2\text{E}+28 \text{ erg s}^{-1}$ radio luminosity at 5 GHz. Combined with the $1.7\text{E}+36 (D/8 \text{ kpc})^2 \text{ erg s}^{-1}$ X-ray luminosity (ATel #12651), this makes IGR J17285-2922 consistent with both the sample of neutron star X-ray binaries and the radio-faint black hole X-ray binaries (see e.g., Bahramian et al. 2018). For significantly smaller distances, the source appears most similar to the accreting neutron stars instead.

Continued Swift and VLA monitoring of IGR J17285-2922 is planned. Further multiwavelength observations are encouraged. We thank the Swift and VLA teams for rapidly performing the observations.

[[Telegram Index](#)]

R. E. Rutledge, Editor-in-Chief

Derek Fox, Editor

Mansi M. Kasliwal, Co-Editor

rrutledge@astronomerstelegam.org

dfox@astronomerstelegam.org

mansi@astronomerstelegam.org