



UvA-DARE (Digital Academic Repository)

ATCA detects the radio brightening of the X-ray transient MAXI J1348-630

Russell, T.; Anderson, G.; Miller-Jones, J.; Degenaar, N.; Eijnden, J. van den; Sivakoff, G.R.; Tetarenko, A.

Publication date

2019

Document Version

Final published version

Published in

The astronomer's telegram

License

Unspecified

[Link to publication](#)

Citation for published version (APA):

Russell, T., Anderson, G., Miller-Jones, J., Degenaar, N., Eijnden, J. V. D., Sivakoff, G. R., & Tetarenko, A. (2019). ATCA detects the radio brightening of the X-ray transient MAXI J1348-630. *The astronomer's telegram*, 12456. <http://www.astronomerstelegram.org/?read=12456>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

2 Apr 2020; 15:40 UT

This space for free for your conference.



Outside

GCN
IAUCs

Other

ATel on [Twitter](#) and [Facebook](#)
ATELstream
ATel Community Site[[Previous](#) | [Next](#) | [ADS](#)]

ATCA detects the radio brightening of the X-ray transient MAXI J1348-630

ATel #12456; *T. Russell (UvA), G. Anderson, J. Miller-Jones (ICRAR/Curtin), N. Degenaar, J. van den Eijnden (UvA), Gregory R. Sivakoff (UAlberta), and A. Tetarenko (EAO)*
on 30 Jan 2019; 20:28 UT

Credential Certification: [Thomas Russell \(t.d.russell@uva.nl\)](mailto:Thomas.Russell@uva.nl)

Subjects: Radio, X-ray, Black Hole, Transient

Referred to by ATel #: [12470](#), [12477](#), [12480](#), [12491](#), [12497](#), [12520](#), [13459](#), [13465](#), [13539](#)

Following the discovery of the X-ray transient MAXI J1348-630 (ATels #[12425](#), #[12430](#), #[12434](#), #[12439](#), #[12441](#), #[12447](#), #[12448](#)), we conducted radio observations with the Australia Telescope Compact Array (ATCA) from 2019-01-26 19:55 UT to 2019-01-27 00:33 UT (MJD 58509.9 +/- 0.1), and from 2019-01-27 21:30 UT to 2019-01-28 03:55 UT (MJD 58511.03 +/- 0.15). For both observations, the telescope was in its most compact H75 configuration, with the core of the array having a longest baseline of 75 m, and a single fixed antenna located 6 km from the array core. Observations were taken simultaneously at 5.5 and 9.0 GHz, with a bandwidth of 2 GHz at both frequencies. We used PKS 1934-638 for bandpass and flux calibration, while 1352-63 was used for phase calibration. Data were calibrated and imaged following standard standard procedures within CASA (version 5.1.1; McMullin et al. 2007), where imaging (with the inclusion of the isolated antenna) was carried out with a Briggs robust parameter of -1 at both frequencies.

We detect a radio source consistent with the X-ray position (ATel #[12434](#)), with a radio position (at 9 GHz) of:

R.A. (J2000): 13:48:12.79 +/- 0.03

Dec (J2000): -63:16:28.48 +/- 0.04,

where the R.A. errors are from beam centroiding and Declination errors are statistical.

To determine the source flux density, we fit for a point source in the image plane. Due to the compact configuration and single isolated antenna, the flux densities were also checked by fitting a delta function in the uv-plane with UVMULTIFIT (Marti-Vidal et al. 2014) within CASA to ensure the results were consistent. Our first observation (on MJD 58509), detected the source at a flux density of 3.4 +/- 0.2 mJy at 5.5 GHz and 3.5 +/- 0.2 mJy at 9 GHz, implying a radio spectral

α

Related

- [13539](#) X-ray and near-infrared observation of rebrightening of MAXI J1348-630
- [13467](#) MeerKAT and Swift/XRT detection of MAXI J1348-630
- [13465](#) Re-brightening and decaying of MAXI J1348-630 as observed with NICER
- [13459](#) MAXI J1348-630: MAXI/GSC detection
- [13454](#) XB-NEWS detects a new optical rise during the current outburst of MAXI J0637-430
- [13451](#) XB-NEWS detection of a new outburst of MAXI J1348-630
- [13188](#) Rebrightening of MAXI J1348-630
- [12838](#) MAXI/GSC detection of X-ray rebrightening of MAXI J1348-630
- [12829](#) Optical re-brightening of MAXI J1348-630
- [12520](#) Low-frequency radio detection of MAXI J1348-630 with the MWA
- [12505](#) Preliminary Result on MAXI J1348-630 using Swift data: Detection of 0.56 Hz QPO and Mass Estimation
- [12497](#) MeerKAT follow-up observations of MAXI J1348-630 reveal bright radio flare at state transition
- [12491](#) Optical fade in MAXI J1348-630 during transition towards the soft state
- [12480](#) MAXI J1348-630: SALT optical spectroscopy during outburst
- [12477](#) Monitoring the transient MAXI J1348-630 with the Neil Gehrels Swift observatory
- [12471](#) INTEGRAL sees the ongoing spectral transition of the black hole candidate MAXI J1348-630
- [12470](#) Insight-HXMT observations of MAXI J1348-630

index of $\alpha = 0.0 \pm 0.2$ (where $S_\nu \propto \nu^{-\alpha}$), consistent with a flat radio spectrum from a compact jet. Our second radio observation (on MJD 58511) shows the radio counterpart brightening to 6.2 ± 0.4 mJy at 5.5 GHz and 6.5 ± 0.5 at 9 GHz, where $\alpha = 0.1 \pm 0.3$, also indicating a flat or slightly inverted radio spectrum from a compact jet.

Our initial radio observation translates to a 5 GHz radio luminosity of $\sim 1.3 \times 10^{30} (D/8\text{kpc})^2$ erg/s. Swift-XRT observations taken close in time to our first radio epoch (ATel #12434), show a 1-10 keV X-ray luminosity of $\sim 3 \times 10^{37} (D/8\text{kpc})^2$ erg/s. Combining the radio and X-ray luminosities from this date and placing them on the radio/X-ray luminosity plane supports the classification as a black hole X-ray binary from the X-ray timing and spectral properties (ATel #12447).

Radio monitoring will continue.

We thank Jamie Stevens and ATNF staff for scheduling these radio observations.

- 12469 MAXI/GSC detects an X-ray spectral softening of the X-ray transient MAXI J1348-630
- 12457 Reduction of the rate of flux increase during recent INTEGRAL hard state observations of the new transient black hole candidate MAXI J1348-630
- 12456 ATCA detects the radio brightening of the X-ray transient MAXI J1348-630
- 12448 Optical colors of MAXI J1348-630
- 12447 NICER identification of MAXI J1348-630 as a probable black hole X-ray binary
- 12441 INTEGRAL's detection of evolution in MAXI J1348-630
- 12439 Optical brightening of MAXI J1348-630
- 12434 MAXI J1348-630: Swift XRT localization, possible periodicity
- 12430 Optical counterpart of MAXI J1348-630
- 12425 MAXI/GSC discovery of a new X-ray transient MAXI J1348-630
- 12128 Optical brightening of MAXI J1820+070 over the soft to hard transition observed with LCO and the AI Sadeem Observatory

[Telegram Index]

R. E. Rutledge, Editor-in-Chief

Derek Fox, Editor

Mansi M. Kasliwal, Co-Editor

rrutledge@astronomerstelegam.org

dfox@astronomerstelegam.org

mansii@astronomerstelegam.org