



## UvA-DARE (Digital Academic Repository)

### NICER follow-up observations of the transient system Swift J1728.9-3613

Enoto, T.; Altamirano, D.; Bult, P.M.; Gendreau, K.C.; Arzoumanian, Z.; Strohmayer, T.E.; Ray, P.S.; Guillot, S.; Jaisawal, G.K.; Uttley, P.; Homan, J.; NICER team

**Publication date**

2019

**Document Version**

Final published version

**Published in**

The astronomer's telegram

**License**

Unspecified

[Link to publication](#)

**Citation for published version (APA):**

Enoto, T., Altamirano, D., Bult, P. M., Gendreau, K. C., Arzoumanian, Z., Strohmayer, T. E., Ray, P. S., Guillot, S., Jaisawal, G. K., Uttley, P., Homan, J., & NICER team (2019). NICER follow-up observations of the transient system Swift J1728.9-3613. *The astronomer's telegram*, 12455. <http://www.astronomerstelegam.org/?read=12455>

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

3 Apr 2020; 13:49 UT

**Outside**GCN  
IAUCs**Other**ATel on [Twitter](#) and [Facebook](#)  
ATELstream  
ATel Community Site

This space for free for your conference.

Growing Black Holes:  
Accretion & Mergers[\[ Previous](#) | [Next](#) | [ADS](#) ]

## NICER follow-up observations of the transient system Swift J1728.9-3613

ATel #12455; *T. Enoto (Kyoto Univ.), D. Altamirano (Univ. of Southampton), P. M. Bult, K. C. Gendreau, Z. Arzoumanian, T. E. Strohmayer (NASA/GSFC), P. S. Ray (NRL), S. Guillot (IRAP/CNRS/CNES), G. K. Jaisawal (DTU Space), P. Uttley (Univ. of Amsterdam), J. Homan (Eureka Scientific and SRON) and the NICER team*

on 30 Jan 2019; 04:17 UT

Credential Certification: [Teruaki Enoto \(teruaki.enoto@gmail.com\)](mailto:teruaki.enoto@gmail.com)

Subjects: X-ray, Black Hole, Transient

Referred to by ATel #: [12502](#)

Following reports of the new Galactic transient Swift J1728.9-3613 (MAXI J1728-360) reported by Swift and MAXI (ATel #[12436](#), #[12437](#), #[12445](#), GCN #[23800](#)), NICER started follow-up observations from 15:20 UT 2019 January 29. The source count rate is ~600 count/s in the 1-10 keV energy band during the first three orbits (~840 s exposure). The average source count rate is nearly constant over the three orbits.

The 1-10 keV X-ray spectrum is approximated by an absorbed power-law model after applying a Crab correction to deal with instrumental uncertainties (see, e.g. Ludlam et al. 2018, ApJL 858, L5). A column density of  $N_{\text{H}}=3.86(1)e+22$  cm<sup>-2</sup> and photon index of 2.69(1) are obtained from preliminary fitting (reduced chi-square of 1.70 with 615 d.o.f.). The fit is improved when adding a blackbody component in soft X-rays (reduced chi-square of 1.16 with 613 d.o.f.). The best fitting parameters are  $N_{\text{H}}=3.42(3)e+22$  cm<sup>-2</sup>, Photon index=2.40(3), and  $kT=0.67(1)$  keV. The absorbed X-ray flux is estimated to be  $4.4(1)e-9$  ergs/cm<sup>2</sup> in the 1-10 keV. The quoted errors are statistical 1-sigma range.

A preliminary timing analysis showed band-limited variability below 1 Hz and a narrow QPO at 5.5 Hz. Fitting the power spectrum with a superposition of Lorentzian profiles, we measured the amplitude of the band-limited noise at 11.7(2)% rms, with a break frequency of 0.034(2) Hz. The QPO was detected at 6 sigma significance, with an amplitude of 4.1(6)% rms, width 0.5(2) Hz and centroid frequency 5.46(5) Hz. No excess power was observed at higher frequencies.

The shape of the average power spectrum is reminiscent of that observed when black hole X-ray binaries display Type-B QPO. If this interpretation is correct, then Swift J1728.9-3613 is a black

**Related**

- 12522** MeerKAT discovery of a variable radio source associated with Swift J1728.9-3613
- 12502** INTEGRAL observations of the on-going outburst from the new X-ray transient Swift J1728.9-3613
- 12455** NICER follow-up observations of the transient system Swift J1728.9-3613
- 12445** Swift J1728.9-3613: Swift/XRT localization
- 12443** Swift J1728.9-3613: BOOTES-3/YA optical counterpart candidate observation
- 12437** MAXI/GSC detection of the newly discovered X-ray transient Swift J1728.9-3613 (= MAXI J1728-360)
- 12436** Swift detection of a new transient Swift J1728.9-3613

hole candidate currently in the soft-intermediate state. This interpretation implies that we missed the rise of the outburst. Further observations both in X-rays and at other wavelengths are needed to test this interpretation.

NICER is a 0.2-12 keV X-ray telescope operating on the International Space Station. The NICER mission and portions of the NICER science team activities are funded by NASA.

---

---

[ [Telegram Index](#) ]

R. E. Rutledge, Editor-in-Chief

[rrutledge@astronomerstelegam.org](mailto:rrutledge@astronomerstelegam.org)

Derek Fox, Editor

[dfox@astronomerstelegam.org](mailto:dfox@astronomerstelegam.org)

Mansi M. Kasliwal, Co-Editor

[mansi@astronomerstelegam.org](mailto:mansi@astronomerstelegam.org)