



UvA-DARE (Digital Academic Repository)

Clocking Stars with Radio Telescopes

Timing Four Pulsars from the GBNCC Survey

Aloisi, R.; Cruz, A.; Daniels, L.; Meyers, N.; Roekle, R.; Schuett, A.; Swiggum, J.; DeCesar, M.E.; Kaplan, D.; Lynch, R.S.; Stovall, K.; Levin, L.S.; Archibald, A.; Banaszak, S.; Chawla, P.; Cui, B.; Fonseca, E.; Kaspi, V.; Kondratiev, V.; McLaughlin, M.; Al Noori, H.; Ransom, S.M.; Spiewak, R.; Stairs, I.; van Leeuwen, J.; Boyles, J.; Hessels, J.; Roberts, M.; Karako-Argaman, C.; Siemens, X.

Publication date

2019

Document Version

Final published version

Published in

American Astronomical Society Meeting

[Link to publication](#)

Citation for published version (APA):

Aloisi, R., Cruz, A., Daniels, L., Meyers, N., Roekle, R., Schuett, A., Swiggum, J., DeCesar, M. E., Kaplan, D., Lynch, R. S., Stovall, K., Levin, L. S., Archibald, A., Banaszak, S., Chawla, P., Cui, B., Fonseca, E., Kaspi, V., Kondratiev, V., ... Siemens, X. (2019). Clocking Stars with Radio Telescopes: Timing Four Pulsars from the GBNCC Survey. *American Astronomical Society Meeting*, 233, 156. [153.07].

<https://assets.pubpub.org/qmxqtups/61582749069276.pdf>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library@uva.nl, or by mail to: Library@uva.nl, P.O. Box 1012, 3000 AA Amsterdam, The Netherlands. You will be contacted as soon as possible.

153.06 — Finding Compact Sources in the VLA Sky Survey

Cameron Riley¹; Shami Chatterjee¹; James Cordes¹

¹ Cornell University (Ithaca, New York, United States)

The VLA Sky Survey (VLASS) is a three epoch radio wavelength survey which covers over 33,000 deg² of the sky as visible from the Very Large Array in New Mexico, USA. At a resolution of 2.5 arcseconds and 120 microJansky RMS sensitivity per epoch, the VLASS is expected to locate around five million sources. We have initiated a project to identify candidate radio pulsars from the VLASS images for targeted follow up observations. To do so, we require a compact source catalog with minimal false positive detections that simultaneously does not miss out on real sources. We are testing the performance of various source finding tools to optimize such a catalog. In particular, Aegean Source Finder and PyBDSF were used to create catalogs for sample VLASS Quicklook images, which were analyzed for accuracy and completeness. Although more testing is required, our preliminary analysis shows that PyBDSF has a lower false positive detection rate and excels in differentiating sources within clusters, while Aegean has better performance in locating point sources. After further testing of other tools, we will construct a compact source catalog and use multiwavelength matches to rank pulsar candidates for follow up.

153.07 — Clocking Stars with Radio Telescopes: Timing Four Pulsars from the GBNCC Survey

Robert Aloisi¹; Aristeo Cruz¹; Luke Daniels¹; Natalie Meyers¹; Ryan Roekle¹; Abigail Schuett¹; Joe Swiggum¹; Megan E. DeCesar²; David Kaplan¹; Ryan S. Lynch^{3,4}; Kevin Stovall⁴; Lina S. Levin⁵; Anne Archibald⁶; Shawn Banaszak¹; Pragya Chawla¹⁰; Bingyi Cui³; Emmanuel Fonseca¹⁰; Victoria Kaspi¹⁰; Vlad Kondratiev⁶; Maura McLaughlin⁷; Hind Al Noori¹¹; Scott M. Ransom⁴; Renee Spiewak¹; Ingrid Stairs⁸; Joeri van Leeuwen⁹; Jason Boyles¹²; Jason Hessels⁹; Mallory Roberts¹¹; Chen Karako-Argaman¹⁰; Xavier Siemens¹

¹ Center for Gravitation, Cosmology, and Astrophysics, Department of Physics, University of Wisconsin - Milwaukee (Milwaukee, Wisconsin, United States)

¹⁰ Department of Physics, McGill Space Institute for Radio Astronomy (Dwingeloo, Netherlands)

¹¹ New York University (Abu Dhabi, United Arab Emirates)

¹² Department of Physics and Astronomy, Western Kentucky University (Bowling Green, Kentucky, United States)

² Department of Physics, Lafayette College (Easton, Pennsylvania, United States)

³ Center for Gravitational Waves and Cosmology, Department of Physics and Astronomy, West Virginia University (Morgantown, West Virginia, United States)

⁴ National Radio Astronomy Observatory (Socorro, New Mexico, United States)

⁵ Jodrell Bank Centre for Astrophysics, School of Physics and Astronomy, The University of Manchester, (Manchester, United Kingdom)

⁶ U. Amsterdam, the Netherlands Institute for Radio Astronomy (Dwingeloo, Netherlands)

⁷ Department of Physics and Astronomy, West Virginia University (Morgantown, West Virginia, United States)

⁸ Department of Physics and Astronomy, University of British Columbia (Vancouver, British Columbia, Canada)

⁹ Anton Pannekoek Institute for Astronomy, University of Amsterdam (Amsterdam, Netherlands)

We present the timing solutions for four pulsars discovered in the Green Bank Northern Celestial Cap (GBNCC) survey. Timing observations were processed and timing solutions were obtained by undergraduate students participating in course-based research at the University of Wisconsin - Milwaukee. Both discovery and timing observations were conducted at a center frequency of 350 MHz using the 100-m Robert C. Byrd Green Bank Telescope. All four pulsars are isolated with spin periods between 0.26 s and 1.84 s. PSR J0038-2501 has a 0.26 s period and a period derivative of $7.6 \times 10^{-19} \text{ s s}^{-1}$, which is unusually low for isolated, longer period pulsars. This low period derivative may be simply an extreme value for an isolated pulsar or it could indicate an unusual evolution path for PSR J0038-2501, such as a disrupted recycled pulsar (DRP) from a binary system or an orphaned central compact object (CCO). Correcting the observed spin-down rate for the Shklovkii effect suggests that this pulsar may have an unusually low space velocity, which is consistent with expectations for DRPs since they come from disrupted binaries. There is no X-ray emission detected from PSR J0038-2501 in an archival *Neil Gehrels Swift Observatory* observation, which suggests that it is not a young orphaned CCO. A second pulsar, PSR J1949+3426 has a high dispersion measure suggesting that it is one of the most distant pulsars discovered in the GBNCC survey at an estimated distance of 12.3 kpc. Among the pulsars discovered in the GBNCC survey that makes it one of the brightest pulsars discovered in the GBNCC survey with a pseudoluminosity of 570 mJy kpc².