



UvA-DARE (Digital Academic Repository)

STROBE-X

X-ray Timing & Spectroscopy on Dynamical Timescales from Microseconds to Years

Wilson-Hodge, C.A.; Ray, P.S.; Maccarone, T.J.; Chakrabarty, D.; Gendreau, K.C.; Arzoumanian, Z.; Jenke, P.; Ballantyne, D.; Bozzo, E.; Brandt, S.; Brenneman, L.; Christophersen, M.; DeRosa, A.; Feroci, M.; Goldstein, A.; Hartmann, D.; Hernanz, M.; McDonald, M.; Nowak, M.; Phlips, B.F.; Remillard, R.; Stevens, A.L.; Tomsick, J.; Watts, A.; Wood, K.S.; Zane, S.

Publication date

2019

Document Version

Final published version

Published in

American Astronomical Society Meeting

[Link to publication](#)

Citation for published version (APA):

Wilson-Hodge, C. A., Ray, P. S., Maccarone, T. J., Chakrabarty, D., Gendreau, K. C., Arzoumanian, Z., Jenke, P., Ballantyne, D., Bozzo, E., Brandt, S., Brenneman, L., Christophersen, M., DeRosa, A., Feroci, M., Goldstein, A., Hartmann, D., Hernanz, M., McDonald, M., Nowak, M., ... Zane, S. (2019). STROBE-X: X-ray Timing & Spectroscopy on Dynamical Timescales from Microseconds to Years. *American Astronomical Society Meeting*, 233, 186. Article 158.11. <https://assets.pubpub.org/qmxqtups/61582749069276.pdf>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library@the-university-of-amsterdam.nl, Secretariat, Singel 425, 1012 CA Amsterdam, The Netherlands. You will be contacted as soon as possible.

by relaxing tolerances and stability requirements; by use of a smaller launch vehicle. The starshade would provide excellent exoplanet science performance, but for a smaller number of detected exoplanets of all types, including exoEarth candidates, and a smaller fraction of exoplanets with measured orbits. The full suite of HabEx observatory science is supported. Our approach uses a non-deployed segmented primary mirror, whose manufacture is within current capabilities.

158.11 — STROBE-X: X-ray Timing & Spectroscopy on Dynamical Timescales from Microseconds to Years

Colleen A. Wilson-Hodge¹; Paul S. Ray²; Thomas J. Maccarone³; Deepto Chakrabarty⁴; Keith C. Gendreau⁵; Zaven Arzoumanian⁵; Peter Jenke⁶; David Ballantyne⁷; Enrico Bozzo⁸; Soren Brandt⁹; Laura Brenneman¹⁰; Marc Christophersen²; Alessandra DeRosa¹¹; Marco Feroci¹¹; Adam Goldstein¹²; Dieter Hartmann¹³; Margarita Hernanz¹⁴; Michael McDonald⁴; Michael Nowak⁴; Bernard F. Phillips²; Ronald Remillard⁴; Abigail L. Stevens¹⁵; John Tomsick¹⁶; Anna Watts¹⁷; Kent S. Wood¹⁸; Silvia Zane¹⁹

¹ NASA MSFC (Huntsville, Alabama, United States)

¹⁰ SAO (Cambridge, Massachusetts, United States)

¹¹ IAPS/INAF (Rome, Italy)

¹² USRA (Huntsville, Alabama, United States)

¹³ Clemson University (Clemson, South Carolina, United States)

¹⁴ Institute of Space Sciences, CSIC-IEEC (Barcelona, Spain)

¹⁵ Michigan State University (East Lansing, Michigan, United States)

States)

¹⁶ UC Berkeley (Berkeley, California, United States)

¹⁷ University of Amsterdam (Amsterdam, Netherlands)

¹⁸ Praxis Inc. (resident at NRL) (Washington, District of Columbia, United States)

Columbia, United States)

¹⁹ MSSL/UCL (London, United Kingdom)

² NRL (Washington, District of Columbia, United States)

³ Texas Tech (Lubbock, Texas, United States)

⁴ MIT (Cambridge, Massachusetts, United States)

⁵ NASA GSFC (Greenbelt, Maryland, United States)

⁶ UAH (Huntsville, Alabama, United States)

⁷ Georgia Tech (Atlanta, Georgia, United States)

⁸ University of Geneva (Geneva, Switzerland)

⁹ DTU Space (Copenhagen, Denmark)

We describe a probe-class mission concept that provides an unprecedented view of the X-ray sky, performing timing and 0.2-30 keV spectroscopy over timescales from microseconds to years. The Spectroscopic Time-Resolving Observatory for Broadband Energy X-rays (STROBE-X) has three key science drivers: (1) measuring the spin distribution of accreting black holes, (2) understanding the equation

of state of dense matter, and (3) exploring the properties of the precursors and electromagnetic counterparts of gravitational wave sources. To perform these science investigations, STROBE-X comprises three primary instruments. The first uses an array of lightweight optics (3-m focal length) that concentrate incident photons onto solid state detectors with CCD-level (85-130 eV) energy resolution, 100 ns time resolution, and low background rates to cover the 0.2-12 keV band. This technology is scaled up from NICER, with enhanced optics to take advantage of the longer focal length of STROBE-X. The second uses large-area collimated silicon drift detectors, developed for ESA's LOFT, to cover the 2-30 keV band. These two instruments, with effective areas of 2 m² at 1.5 keV and 5 m² at 10 keV, respectively, each provide an order of magnitude improvement in effective area compared with its predecessor (NICER and RXTE, respectively). Finally, a sensitive sky monitor triggers pointed observations, provides high duty cycle, high time resolution, high spectral resolution monitoring of the X-ray sky with 20 times the sensitivity of the RXTE ASM, and enables multi-wavelength and multi-messenger studies on a continuous, rather than scanning basis. The STROBE-X mission concept is a rapidly re-pointable observatory in low-Earth orbit, similar to RXTE or Swift, and will be presented to the 2020 Astrophysics Decadal Survey for consideration as a probe-class mission.

158.12 — Reducing the Athena WFI Background with the Science Products Module: Results from Geant4 Simulations

Eric D. Miller¹; Catherine Grant¹; Marshall Bautz¹; Jonathan Keelan²; David Hall²; Andrew Holland²; Esra Bulbul³; Ralph Kraft³; Paul Nulsen³; David Burrows⁴; Steven Allen⁵

¹ Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology (Cambridge, Massachusetts, United States)

² Open University (Milton Keynes, United Kingdom)

³ Harvard-Smithsonian Center for Astrophysics (Cambridge, Massachusetts, United States)

⁴ The Pennsylvania State University (University Park, Pennsylvania, United States)

⁵ Stanford University (Stanford, California, United States)

The Wide Field Imager (WFI) on ESA's Athena X-ray observatory will include the Science Products Module, a secondary CPU that can perform special processing on the science data stream. Our goal is to identify on-board processing algorithms that can reduce WFI charged particle background and improve knowledge of the background to reduce systematics.