UvA-DARE (Digital Academic Repository)

Facile Phenylphosphinidene Transfer Reactions from Carbene-Phosphinidene Zinc Complexes

Krachko, T.; Bispinghoff, M.; Tondreau, A.M.; Stein, D.; Baker, M.; Ehlers, A.W.; Slootweg, J.C.; Grutzmacher, H.

DOI

[10.1002/ange.201703672](https://doi.org/10.1002/ange.201703672)

[10.1002/anie.201703672](https://doi.org/10.1002/anie.201703672)

Publication date

2017

Document Version

Final published version

Published in

Angewandte Chemie

License

Article 25fa Dutch Copyright Act

[Link to publication](#)

Citation for published version (APA):

Krachko, T., Bispinghoff, M., Tondreau, A. M., Stein, D., Baker, M., Ehlers, A. W., Slootweg, J. C., & Grutzmacher, H. (2017). Facile Phenylphosphinidene Transfer Reactions from Carbene-Phosphinidene Zinc Complexes. *Angewandte Chemie*, 129(27), 8056-8059. <https://doi.org/10.1002/ange.201703672>, <https://doi.org/10.1002/anie.201703672>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Phosphorus Heterocycles

Deutsche Ausgabe: DOI: 10.1002/ange.201703672
Internationale Ausgabe: DOI: 10.1002/anie.201703672

Facile Phenylphosphinidene Transfer Reactions from Carbene–Phosphinidene Zinc Complexes

Tetiana Krachko, Mark Bispinchoff, Aaron M. Tondreau, Daniel Stein, Matthew Baker, Andreas W. Ehlers, J. Chris Slootweg,* and Hansjörg Grützmacher*

Dedicated to Professor Evamarie Hey-Hawkins on the occasion of her 60th birthday

Abstract: Phosphinidenes [R-P] are convenient P_1 building blocks for the synthesis of a plethora of organophosphorus compounds. Thus far, transition-metal-complexed phosphinidenes have been used for their singlet ground-state reactivity to promote selective addition and insertion reactions. One disadvantage of this approach is that after transfer of the P_1 moiety to the substrate, a challenging demetallation step is required to provide the free phosphine. We report a simple method that enables the Lewis acid promoted transfer of phenylphosphinidene, [PhP], from $NHC=PPh$ adducts ($NHC=N$ -heterocyclic carbene) to various substrates to produce directly uncoordinated phosphorus heterocycles that are difficult to obtain otherwise.

Considerable effort has been devoted to the development of phosphinidene [R-P] transfer agents^[1,2] that do not require a metal.^[3] Cummins and co-workers developed the unprotected dibenzo-7 λ^3 -phosphanorbornadiene **A** as a source of the transient [iPr₂NP], which undergoes a [1+4]-cycloaddition with 1,3-cyclohexadiene.^[4] Driess and co-workers successfully transferred the parent phosphinidene [HP] from phosphasilene **B** (Dipp = 2,6-diisopropylphenyl) to an N-heterocyclic carbene,^[5] and Weber et al. demonstrated the acyclic carbene–phosphinidene adducts **C** (R = *t*Bu, Cy, 1-Ad, Ph, Mes) to be viable [RP] transfer agents to diphenylketene.^[6] Interestingly, Arduengo and co-workers showed that the Lewis acid BPh₃ induces formation of cyclopolyphosphines (PPh)_{*n*} (*n* = 3–5) from ^{Mes}NHC=PPh **D**,^[7] but no transfer reactions of the extruded phenylphosphinidene

moiety have been reported to date. This inspired us to target the sterically little encumbered ^{Mes}NHC=PPh (**1**; Scheme 1) and study its ability to transfer [PhP] to suitable substrates in the presence of an appropriate Lewis acid.

Scheme 1. Synthesis of ^{Mes}NHC=PPh (**1**; left) and molecular structure (right; hydrogen atoms and C₆H₆ solvent molecule are omitted for clarity, one crystallographic independent molecule is shown). Selected bond lengths [Å] and angles [°] (values for the second molecule in square brackets): P1–C1 1.7917(14) [1.7911(15)], P1–C6 1.8157(16) [1.8132(14)], N1–C1 1.359(2) [1.3546(19)], N2–C1 1.3562(18) [1.356(2)]; C1–P1–C6 101.30(6) [99.98(7)], N1–C1–N2 104.82(11) [105.15(13)]; N2–C1–P1–C6 48.01(16) [50.42(15)].

First, we developed a scalable, efficient one-pot synthesis of carbene–phosphinidene adduct ^{Mes}NHC=PPh (**1**) that avoids the need for free carbenes. Reaction of 1,3-dimethylimidazolium iodide with (PPh)₅ and sodium *tert*-butoxide in THF afforded **1** after a simple work-up (removal of all volatiles under reduced pressure, extraction into toluene, and crystallization) as an orange solid in 95 % yield. The molecular structure of **1** ($\delta^{31}P$ (C₆D₆) = –49.1 ppm; Scheme 1, right),^[8] when compared to the bulkier ^{Mes}NHC=PMes reported by Hey-Hawkins et al. ($\delta^{31}P$ (C₆D₆) = –73.8 ppm),^[9] displays a slightly elongated P1–C1 bond [1.7917(14)/1.7911(15) vs. 1.768(4) Å], a more acute C1–P1–C6 angle [101.30(6)/99.98(7) vs. 104.6(2)°], and a larger dihedral angle between the imidazole plane and the P1–C6 bond [48.01(16)/50.42(15) vs. 32.5(4)°], thus indicating considerable ylide character.^[9] Beside the nature of the carbene moiety,^[10] the P substituent also has a marked influence on the

[*] T. Krachko, Dr. A. W. Ehlers, Prof. Dr. J. C. Slootweg
Van't Hoff Institute for Molecular Sciences
University of Amsterdam, Science Park 904
PO Box 94157, 1090 GD Amsterdam (The Netherlands)
E-mail: j.c.slootweg@uva.nl

M. Bispinchoff, Dr. A. M. Tondreau, Dr. D. Stein, M. Baker,
Prof. Dr. H. Grützmacher
Laboratory of Inorganic Chemistry, ETH Zürich
Vladimir-Prelog-Weg 1, 8093 Zürich (Switzerland)
E-mail: hgruetzmacher@ethz.ch

Dr. A. W. Ehlers
Department of Chemistry, Science Faculty
University of Johannesburg
PO Box 254, Auckland Park, Johannesburg (South Africa)

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.201703672>.

structure^[11] and thus the $\delta^{31}\text{P}$ chemical shift of the carbene–phosphinidene adduct. $\text{Me}^{\text{NHC}}\text{=PPh}$ **D**^[12] (93 %) and $\text{Dipp}^{\text{NHC}}\text{=PPh}$ ^[10] (87 %) were also conveniently prepared by this method, thus complementing the known procedures for making carbene–phosphinidene adducts.^[13]

Next, we targeted the synthesis of a Lewis acid adduct of **1** that can controllably release phenylphosphinidene and, ideally, simultaneously capture the free carbene. Treatment of **1** with BPh_3 , AlCl_3 , MgCl_2 , or $\text{Zn}(\text{OAc})_2$ merely resulted in the direct formation of cyclopolyphosphines, which is consistent with Arduengo's observations.^[7,14] However, slow addition of 0.5 equiv of ZnCl_2 to a THF solution of **1** afforded zinc adduct **2** (1:ZnCl_2) as a pale yellow precipitate (76%; Scheme 2, top), which is poorly soluble in common organic

Scheme 2. Synthesis of ZnCl_2 complexes **2** and **3** (top) and molecular structures (bottom) of **2** (DME solvent molecule omitted) and **3** (THF solvent molecule omitted). Selected bond lengths [Å] and angles [°] for **2**: Zn1–P1 2.3959(11), Zn1–P2 2.3867(11), P1–C1 1.818(4), P1–C6 1.826(4), P2–C12 1.827(4), P2–C17 1.841(5); C1–P1–C6 100.2(2), C12–P2–C17 100.8(2); N2–C1–P1–C6 31.1(4), N4–C12–P2–C17 42.9(4). **3**: Zn1–P1 2.3793(14), Zn1–P2 2.4386(13), Zn2–P2 2.4110(14), P1–C1 1.819(4), P1–C6 1.832(4), P2–C12 1.818(4), P2–C17 1.833(4); N2–C1–P1–C6 33.3(4). DME = dimethoxyethane, THF = tetrahydrofuran.

solvents. Single crystals suitable for X-ray diffraction analysis were obtained from DME, which unequivocally established the formation of a 2:1 complex of a carbene–phosphinidene adduct (Scheme 2, bottom).^[8,15] Upon complexation, significant structural changes occur: the central P–C bonds [1.818(4), 1.827(4) Å] become longer, the C–P–C angles more acute [100.2(2), 100.8(2)°], and the dihedral angles smaller [31.1(4), 42.9(4)°; Scheme 2, bottom].^[8] Reaction of **1** with 1 equiv of ZnCl_2 in THF afforded the soluble 1:1 adduct **3**, which was isolated as colorless crystals in 87% yield (Scheme 2, top). The structure of **3** in the solid state shows a di-zinc complex that can be related to the one of **2** by adding $\text{ZnCl}_2(\text{THF})$ to P2 such that one $\text{Me}^{\text{NHC}}\text{=PPh}$ ligand bridges two Zn centers via P2, while the other $\text{Me}^{\text{NHC}}\text{=PPh}$ unit takes a terminal position (Scheme 2, bottom).^[8] Consequently, the phosphorus atoms are inequivalent, yet in solution, even at -80°C , only one $\delta^{31}\text{P}$ resonance was observed in $[\text{D}_8]\text{THF}$ at -88.1 ppm, thus indicating that **3** has a different structure in solution

(either monomeric or a dimer with symmetric cyclic Zn_2P_2 core).

The soluble Lewis adduct **3** was tested as a phosphinidene-transfer agent with phenanthrene-9,10-quinone, diphenylketene, and *trans*-chalcone as suitable acceptors (Scheme 3).^[16]

Scheme 3. Reactivity of **3** towards phenanthrene-9,10-quinone, diphenylketene, and *trans*-chalcone.

Treatment of **3** with phenanthrene-9,10-quinone in THF afforded phosphonite **4** as a pale green solid (46%; $\delta^{31}\text{P}$ (C_6D_6) = 183.3 ppm), which was characterized by single-crystal X-ray diffraction analysis (Figure 1).^[8] Previously, **4** has only been accessible through thermal fragmentation of the corresponding phosphorane.^[17] Heterocycle **4** was devoid of zinc chloride, which was transferred to the carbene to

Figure 1. Molecular structures of **4**, **5** (only a fragment is shown), **6**, and **8**. Selected bond lengths [Å] and angles [°] for **4**: P1–O1 1.6859(19), P1–O2 1.675(2), P1–C1 1.827(3), C7–C8 1.347(4). **5**: Zn1–Cl1 2.2179(18), Zn1–Cl2 2.3670(11), Zn1–Cl2a 2.3670(11), Zn2a–Cl2a 2.3597(12), Zn1–C1 2.021(6), N1–C1 1.329(8), N2–C1 1.377(8); **6**: P1–C1 1.826(2), P1–C7 1.902(2), P1–C9 1.824(2), O1–C8 1.365(2), O1–C9 1.415(2), O2–C8 1.197(2), C7–C8 1.526(3), C9–C22 1.340(3); **8**: P1–O1 1.6636(13), P1–C1 1.876(2), P1–C16 1.8241(18), O1–C3 1.400(2), C1–C2 1.503(3), C2–C3 1.325(3); C2–C3–C10–C11 $-3.2(3)$.

afford NHC complex **5** as an insoluble, colorless, crystalline solid.^[18] The molecular structure of **5** displays a one-dimensional coordination polymer with unusual zigzag zinc chloride chains with almost identical Zn–Cl distances [Zn1–Cl2 2.3670(11), Zn2–Cl2 2.3597(12) Å] and the NHC moiety positioned orthogonally to the main chain [Zn1–C1 2.021(6) Å; Figure 1].^[8,19]

Reaction of di-zinc complex **3** with 4 equiv of diphenylketene afforded phosphorus heterocycles **6** and **7** in a 2:3 ratio ($\delta^{31}\text{P}$ ([D₈]THF): –2.9 and 90.9 ppm, respectively; Scheme 3) together with NHC complex **5**. Whereas 1,4,2-dioxaphospholane **7** was reported previously,^[6] 1,3-oxaphospholan-5-one **6** is new, and only derivatives thereof lacking the exocyclic C=C double bond are known.^[20,21] The molecular structure of **6** was unambiguously established by single-crystal X-ray analysis (Figure 1).^[8,16] It shows a five-membered heterocycle resulting from a formal [1+2+2] cycloaddition, where the endocyclic P1–C9 [1.824(2) Å] and exocyclic P1–C1 bonds [1.826(2) Å] are of similar length, while the endocyclic P1–C7 bond [1.902(2) Å] is significantly elongated, most likely due to steric hindrance from neighboring phenyl rings. Treatment of **3** with only 2 equiv of diphenylketene yielded **10** after work-up as an off-white solid (82 %, $\delta^{31}\text{P}$ ([D₈]THF) = –16.6 ppm; Scheme 4), which contained traces of **6** and **7**. Single crystals

Treatment of **3** with 2 equiv of *trans*-chalcone afforded the rare tricoordinate oxo-3-phospholene **8** as a single diastereomer in 80 % yield ($\delta^{31}\text{P}$ (C₆D₆) = 133.9 ppm; Scheme 3), simply after filtering off [^MNHC·ZnCl₂]_n **5** (83 %), extraction into toluene, and crystallization. Such an unprotected five-membered heterocycle has only been prepared once before, through a two-step procedure using an electrophilic phosphinidene complex followed by demetallation.^[22] The molecular structure of **8** firmly establishes the phenyl rings to be in the *trans* position and shows that the third phenyl ring (on C3) is in conjugation with the C1=C2 double bond in the ring [C2–C3–C10–C11 –3.2(3)°; Figure 1].^[8] We turned to DFT calculations to provide insight into the formation of **8**.^[14] In contrast to the ketene, where the carbonyl carbon is attacked first, now the reactions start with coordination of *trans*-chalcone to **3** by Zn–O bond formation, which selectively affords **8** after P–C bond formation and subsequent ring closure by P–O bond formation and elimination of (NHC)Zn complex **5** (see the Supporting Information).

In summary, the sterically little hindered carbene–phosphinidene adduct ^MNHC=PPh allows the synthesis of new zinc complexes, of which the soluble Lewis adduct **3** selectively transfers a phenylphosphinidene fragment, thereby providing access to uncoordinated phosphorus heterocycles. The driving force for these reactions is likely the formation of the insoluble coordination polymer [^MNHC·ZnCl₂]_n **5**, which explains why only ZnCl₂ proved to be efficient to date. Highly reactive or unstable main-group fragments can be stabilized by NHCs,^[23] however, their transfer to other substrates has very rarely been observed.^[24] The Lewis acid promoted transfer reaction may help to develop this chemistry further.

Scheme 4. Synthesis of intermediate **10** (left) and molecular structure of **10** (right; hydrogen atoms, one disordered DME molecule, and toluene solvent are omitted for clarity). Selected bond lengths [Å] and angles [°] for **10**: P1–C1 1.835(3), P1–C6 1.838(4), P1–C12 1.854(3), O1–C12 1.330(4), C12–C13 1.362(4); O1–C12–C13 126.3(3).

suitable for X-ray diffraction analysis were obtained from DME, which established the formation of a 1:1 adduct (Scheme 4, right).^[8] The structural parameters are consistent with the formula shown in Scheme 4 [P1–C1 1.835(3) Å, P–C12 1.854(3) Å, C12–O1 1.330(4) Å are single bonds, C12–C13 1.362(4) Å is a double bond]. Calculations at ωB97X-D/6-31G(d,p)^[14] reveal that nucleophilic attack of the phosphorus atom of **3** at the ketene carbonyl carbon first gives adduct **9** ($\Delta G = 11.0 \text{ kcal mol}^{-1}$; $\Delta G_a = 13.1 \text{ kcal mol}^{-1}$; Scheme 4, top left), which affords **10** after ZnCl₂ transfer from P to O ($\Delta G = -18.3 \text{ kcal mol}^{-1}$; $\Delta G_a = 13.1 \text{ kcal mol}^{-1}$). Furthermore, we confirmed experimentally that intermediate **10** is able to react with either the C=C or C=O double bond of another equiv of diphenylketene to form **6** and **7** after extrusion of Zn complex **5**.

Acknowledgements

This work was supported by the European Union (Marie Curie ITN SusPhos, Grant Agreement No. 317404). Prof. Dr. David Scheschkewitz is gratefully acknowledged for fruitful discussions.

Conflict of interest

The authors declare no conflict of interest.

Keywords: adducts · Lewis acids · N-heterocyclic carbenes · phosphinidenes · phosphorus heterocycles

How to cite: *Angew. Chem. Int. Ed.* **2017**, *56*, 7948–7951
Angew. Chem. **2017**, *129*, 8056–8059

- [1] a) K. Lammertsma, *Top. Curr. Chem.* **2003**, *229*, 95–119; b) J. C. Slootweg, K. Lammertsma in *Science of Synthesis*, Vol. 42 (Eds.: B. M. Trost, F. Mathey), Thieme, Stuttgart, **2009**, pp. 15–36.
- [2] a) F. Mathey, N. H. Tran Huy, A. Marinetti, *Helv. Chim. Acta* **2001**, *84*, 2938–2957; b) R. Streubel, *Top. Curr. Chem.* **2003**, *223*, 91–109; c) K. Lammertsma, M. J. M. Vlaar, *Eur. J. Org. Chem.* **2002**, 1127–1138; d) H. Aktaş, J. C. Slootweg, K. Lammertsma,

- Angew. Chem. Int. Ed.* **2010**, *49*, 2102–2113; *Angew. Chem.* **2010**, *122*, 2148–2159.
- [3] Phosphinidene chalcogenides: C. M. E. Graham, T. E. Pritchard, P. D. Boyle, J. Valjus, H. M. Tuononen, P. J. Ragogna, *Angew. Chem. Int. Ed.* **2017**, *56*, 6236–6240; *Angew. Chem.* **2017**, *129*, 6332–6336.
- [4] A. Velian, C. C. Cummins, *J. Am. Chem. Soc.* **2012**, *134*, 13978–13981.
- [5] K. Hansen, T. Szilvási, B. Blom, S. Inoue, J. Epping, M. Driess, *J. Am. Chem. Soc.* **2013**, *135*, 11795–11798.
- [6] L. Weber, U. Lassahn, H.-G. Stammer, B. Neumann, *Eur. J. Inorg. Chem.* **2005**, 4590–4597.
- [7] A. J. Arduengo III, C. J. Carmalt, J. A. C. Clyburne, A. H. Cowley, R. Pyati, *Chem. Commun.* **1997**, 981–982.
- [8] CCDC 923521 (**1**), 1523933 (**2**), 1523907 (**3**), 1523934 (**4**), 1523939 (**5**), 1523936 (**6**), 1523935 (**8**) and 1523941 (**10**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre. For the experimental details of the X-ray crystal structure determinations, see the Supporting Information.
- [9] A. K. Adhikari, T. Grell, P. Lönnecke, E. Hey-Hawkins, *Eur. J. Inorg. Chem.* **2016**, 620–622.
- [10] O. Back, M. Henry-Ellinger, C. D. Martin, D. Martin, G. Bertrand, *Angew. Chem. Int. Ed.* **2013**, *52*, 2939–2943; *Angew. Chem.* **2013**, *125*, 3011–3015.
- [11] a) G. Frison, A. Sevin, *J. Organomet. Chem.* **2002**, 643–644, 105–111; b) R. R. Rodrigues, C. L. Dorsey, C. A. Arceneaux, T. W. Hudnall, *Chem. Commun.* **2014**, 50, 162–116.
- [12] A. J. Arduengo III, J. C. Calabrese, A. H. Cowley, H. V. R. Dias, J. R. Goerlich, W. J. Marshall, B. Riegel, *Inorg. Chem.* **1997**, *36*, 2151–2158.
- [13] Carbene–phenylphosphinidene adducts have been prepared by using free carbenes and (PPh)₃: a) Ref. [12]; b) A. J. Arduengo III, H. V. R. Dias, J. C. Calabrese, *Chem. Lett.* **1997**, 143–144; using dichlorophenylphosphine followed by reduction, see: c) ref. [10]; using alkali metal tetraphosphanediides, see: d) ref. [9]; using primary phosphines, see: e) K. Pal, O. B. Hemming, B. M. Day, T. Pugh, D. J. Evans, R. A. Layfield, *Angew. Chem. Int. Ed.* **2016**, *55*, 1690–1693; *Angew. Chem.* **2016**, *128*, 1722–1725; f) H. Schneider, D. Schmidt, U. Radius, *Chem. Commun.* **2015**, *51*, 10138–10141.
- [14] See the Supporting Information for further details.
- [15] So far only 1:1 complexes have been reported: a) T. G. Larocque, G. G. Lavoie, *New J. Chem.* **2014**, *38*, 499–502; b) D. Bockfeld, A. Doddi, P. G. Jones, M. Tamm, *Eur. J. Inorg. Chem.* **2016**, 3704–3713; For 1:2 complexes with coinage metals, see: c) V. A. K. Adiraju, M. Yousufuddin, H. V. R. Dias, *Dalton Trans.* **2015**, *44*, 4449–4454; d) A. Doddi, D. Bockfeld, A. Nasr, T. Bannenberg, P. G. Jones, M. Tamm, *Chem. Eur. J.* **2015**, *21*, 16178–16189.
- [16] Typical alkynes, alkenes, and dienes did not exhibit any reactivity toward **3**. We investigated the reactions of ^{Me}NHC=PPh (**1**) directly with organic electrophiles (see the Supporting Information for details).
- [17] P. J. Hammond, J. R. Lloyd, C. D. Hall, *Phosphorus Sulfur Relat. Elem.* **1981**, *10*, 47–51.
- [18] Zinc complexes of NHCs have been shown to be catalytically active in a number of processes: a) P. Jochmann, D. W. Stephan, *Angew. Chem. Int. Ed.* **2013**, *52*, 9831–9835; *Angew. Chem.* **2013**, *125*, 10014–10018; b) O. Jacquet, X. Frogneux, C. Das Neves Gomes, T. Cantat, *Chem. Sci.* **2013**, *4*, 2127–2131; c) A. Rit, T. P. Spaniol, L. Maron, J. Okuda, *Angew. Chem. Int. Ed.* **2013**, *52*, 4664–4667; *Angew. Chem.* **2013**, *125*, 4762–4765.
- [19] For a (NHC)AgBr 1D coordination polymer: A. M. Ortiz, P. Gómez-Sal, J. C. Flores, E. de Jesús, *Organometallics* **2014**, *33*, 600–603.
- [20] a) A. Zschunke, H. Meyer, K. Issleib, *Org. Magn. Reson.* **1975**, *7*, 470–474; b) H. Oehme, K. Issleib, E. Leissring, *J. Prakt. Chem.* **1972**, *314*, 66–70.
- [21] Derivatives of **6** and **7** have been shown to catalyze the 1,4-diboration of 1,3-dienes: C. H. Schuster, B. Li, J. P. Morken, *Angew. Chem. Int. Ed.* **2011**, *50*, 7906–7909; *Angew. Chem.* **2011**, *123*, 8052–8055.
- [22] a) K. Vaheesar, C. M. Kuntz, B. T. Sterenberg, *J. Organomet. Chem.* **2013**, 745–746, 347–355; For synthesis of the W(CO)₅ complexes, see: b) M. P. Duffy, Y. Lin, F. Ho, F. Mathey, *Organometallics* **2010**, *29*, 5757–5758; c) A. Marinetti, F. Mathey, *Organometallics* **1984**, *3*, 456–461; d) A. I. Arkhypchuk, A. Orthaber, V. Alina Mihali, A. Ehlers, K. Lammertsma, S. Ott, *Chem. Eur. J.* **2013**, *19*, 13692–13704.
- [23] Selected reviews: a) C. D. Martin, M. Soleilhavoup, G. Bertrand, *Chem. Sci.* **2013**, *4*, 3020–3030; b) D. J. D. Wilson, J. L. Dutton, *Chem. Eur. J.* **2013**, *19*, 13626–13637; c) M. N. Hopkinson, C. Richter, M. Schedler, F. Glorius, *Nature* **2014**, *510*, 485–496; d) Y. Wang, G. H. Robinson, *Inorg. Chem.* **2014**, *53*, 11815–11832.
- [24] A. M. Tondreau, Z. Benkő, J. R. Harmer, H. Grützmacher, *Chem. Sci.* **2014**, *5*, 1545–1554.

Manuscript received: April 9, 2017

Revised manuscript received: May 9, 2017

Accepted manuscript online: May 15, 2017

Version of record online: June 8, 2017