



UvA-DARE (Digital Academic Repository)

Gumbel statistics for entanglement spectra of many-body localized eigenstates

Buijsman, W.; Gritsev, V.; Cheianov, V.

DOI

[10.1103/PhysRevB.100.205110](https://doi.org/10.1103/PhysRevB.100.205110)

Publication date

2019

Document Version

Final published version

Published in

Physical Review B

[Link to publication](#)

Citation for published version (APA):

Buijsman, W., Gritsev, V., & Cheianov, V. (2019). Gumbel statistics for entanglement spectra of many-body localized eigenstates. *Physical Review B*, 100(20), Article 205110. <https://doi.org/10.1103/PhysRevB.100.205110>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Gumbel statistics for entanglement spectra of many-body localized eigenstates

Wouter Buijsman^{1,*}, Vladimir Gritsev^{1,2} and Vadim Cheianov³

¹*Institute for Theoretical Physics Amsterdam and Delta Institute for Theoretical Physics,
University of Amsterdam, P.O. Box 94485, 1090 GL Amsterdam, The Netherlands*

²*Russian Quantum Center, Skolkovo, Moscow 143025, Russia*

³*Instituut-Lorentz and Delta Institute for Theoretical Physics, Universiteit Leiden, P.O. Box 9506, 2300 RA Leiden, The Netherlands*



(Received 26 July 2019; revised manuscript received 7 October 2019; published 8 November 2019)

An entanglement spectrum encodes statistics beyond the entanglement entropy, of which several have been studied in the context of many-body localization. We numerically study the extreme value statistics of entanglement spectra of many-body localized eigenstates. The physical information encoded in these spectra is almost fully carried by the few smallest elements, suggesting the extreme value statistics to have physical significance. We report the surprising observation of Gumbel statistics. Our result provides an analytical, parameter-free characterization of many-body localized eigenstates.

DOI: [10.1103/PhysRevB.100.205110](https://doi.org/10.1103/PhysRevB.100.205110)

I. INTRODUCTION

Many-body localization (MBL) is understood as a distinct phase of matter that cannot be described by conventional statistical physics [1]. Driven by theoretical and experimental progress, there has been a surge of interest in the phenomenon over the past decade [2]. MBL appears in sufficiently strongly disordered interacting quantum many-body systems, where the appearance of local integrals of motion [3,4] leads to, e.g., emergent integrability [5,6], the absence of thermalization [7], and logarithmic growth of entanglement entropy in time after a quantum quench [8–10].

Thermal and many-body localized phases are separated by an MBL transition [11–13]. At the localized side of the transition, eigenstates obey area-law scaling of entanglement entropy, while volume-law scaling is observed at the thermal side [14]. Entanglement entropies can be extracted from entanglement spectra [15,16]. An entanglement spectrum encodes statistics beyond the entanglement entropy [17], of which several have been studied in the context of MBL [18–24].

The physical information encoded in the entanglement spectrum of a many-body localized eigenstate is almost fully carried by only a few elements, independent of system size [20]. This indicates the potential physical significance of the extreme value statistics [25] of entanglement spectra in the context of MBL. In this work, we study the extreme value statistics of entanglement spectra of many-body localized eigenstates.

Extreme value statistics display universal characteristics over a wide range of physically relevant conditions [26–29]. We report the surprising observation of Gumbel statistics [25,30]. These statistics, being observed in studies on various physical phenomena [31–35], apply to the extreme value of $n \rightarrow \infty$ independent samples drawn from a distribution with

a faster than power-law asymptotic decay. Our result provides an analytical, parameter-free characterization of many-body localized eigenstates.

II. GUMBEL STATISTICS

Following parts of Ref. [25], we here briefly discuss Gumbel statistics. Let X_i ($i = 1, 2, \dots, n$) be a sequence of independent and identically distributed random variables drawn from a distribution for which the distribution function (the probability that $X_i \leq x$) is given by $F(x)$,

$$P\{X_i \leq x\} = F(x). \quad (1)$$

Let M_n denote the largest element of the sequence. It follows from the independence of the X_i that the distribution function of M_n is given by

$$P\{M_n \leq x\} = F^n(x). \quad (2)$$

Two distribution functions F_1 and F_2 are said to be of the same type if, up to normalization,

$$F_2(x) = F_1(ax + b) \quad (3)$$

for some $a > 0$ and b . From the extremal types theorem, it follows that if

$$\lim_{n \rightarrow \infty} F^n(a_n x + b_n) = G(x) \quad (4)$$

for some a_n and b_n , then G is a distribution function of the same type as one of the three extreme value distributions. The distribution function of one of these extreme value distributions, relevant in the context of this work, is given by

$$G(x) = \exp(-e^{-x}). \quad (5)$$

This type emerges, e.g., for a density function $f = dF/dx$ asymptotically decaying faster than a power law, i.e., as

$$f(x) \sim \exp(-x^\alpha), \quad (6)$$

with $\alpha > 0$ a free parameter. Equation (6) covers, e.g., exponential ($\alpha = 1$) and Gaussian ($\alpha = 2$) decays.

*w.buijsman@uva.nl

The statistics of G given in Eq. (5) are referred to as Fisher-Tippett-Gumbel [27] or Gumbel [28] statistics. For these statistics, the rate of convergence depends nontrivially on the structure of F [36]. When comparing the statistics of a collection of extreme values with Gumbel statistics, it is convenient [26–29,31–35] to take a_n and b_n such that the distribution has mean zero and standard deviation 1. The distribution function of the same type as G given in Eq. (5) with these values for the mean and standard deviation is obtained through Eq. (3) with

$$a = \pi/\sqrt{6}, \quad b = \gamma, \quad (7)$$

where $\gamma \approx 0.577$ is Euler's constant. The corresponding density function is given explicitly in Eq. (17) below.

III. ENTANGLEMENT SPECTRA

Here, we review the concept of entanglement spectra. In the most general form, the setting is a quantum system divided into subsystems A and B with Hilbert space dimensions m and n . A pure state $|\psi\rangle$ of the composite system can be expanded in basis states $|a_i\rangle$ and $|b_j\rangle$ of the respective subsystems as

$$|\psi\rangle = \sum_{i,j} X_{ij} |a_i\rangle \otimes |b_j\rangle, \quad (8)$$

where X is an $m \times n$ matrix. Labeling the subsystems such that $m \geq n$, the Schmidt decomposition of X uniquely expands $|\psi\rangle$ as a linear combination of product states over the subsystems,

$$|\psi\rangle = \sum_{i=1}^n \sqrt{\lambda_i} |\alpha_i\rangle \otimes |\beta_i\rangle, \quad (9)$$

where $|\alpha_i\rangle$ and $|\beta_i\rangle$ are basis states for respectively subsystems A and B , and the λ_i ($\lambda_i \geq 0$) are the Schmidt coefficients. An element λ_i can be interpreted as the physical weight of the product state $|\alpha_i\rangle \otimes |\beta_i\rangle$, providing a contribution of $-\lambda_i \ln(\lambda_i)$ to the entanglement entropy. The elements e_i of the entanglement spectrum [15,16] are given by

$$e_i = -\ln(\lambda_i). \quad (10)$$

We remark that in some literature (e.g., Refs. [18,20,21]) the “quantum information definition” $e_i = \lambda_i$ of the entanglement spectrum is used. The smallest of the e_i carry the largest physical weight. In this work, the focus is on the statistics of the smallest of the e_i .

The Schmidt coefficients for ergodic (“random”) states [37] obey the eigenvalue statistics of the fixed-trace Wishart-Laguerre random matrix ensemble [38]. For this ensemble, the joint density function $P\{\lambda_{1,2,\dots,n} = x_{1,2,\dots,n}\}$ of the eigenvalues is proportional to

$$\prod_{i=1}^n x_i^{\alpha\beta/2} \prod_{j < k} |x_j - x_k|^\beta \delta\left(\sum_{i=1}^n x_i - 1\right), \quad (11)$$

where $\alpha = (1 + m - n) - 2/\beta$ and β is the Dyson index given by 1 or 2 if the eigenstate is for a system with or without time-reversal symmetry, respectively. The elements are strongly correlated, due to which Gumbel statistics do not apply. For the values of n relevant in the context of this work,

the extreme value statistics of the smallest $e_i = -\ln(\lambda_i)$ are close to Gaussian up to ~ 3 standard deviations around the mean value (verified numerically).

IV. PHYSICAL SETTING

We study the eigenstates of a spin chain with random onsite disorder. Let σ_i^α denote a Pauli matrix ($\alpha = x, y, z$) acting on site i , and let $S_i^\alpha = \sigma_i^\alpha/2$ denote the corresponding spin-1/2 operator. The Hamiltonian of the model under consideration is given by

$$H = \sum_{i=1}^L (\vec{S}_i \cdot \vec{S}_{i+1} + h_i S_i^z + \Gamma S_i^x). \quad (12)$$

We impose periodic boundary conditions $S_{L+1}^\alpha \equiv S_1^\alpha$, set $\Gamma = 0.1$, and sample h_i from the uniform distribution ranging over $[-W, W]$. We restrict the focus to eigenstates associated with eigenvalues close to the middle of the spectrum (quantified below) for system sizes $L = 10, 12, 14$. The model is studied in Ref. [18], where indications for an MBL transition at $W \approx 3.5$ are reported.

The numerical analysis for systems of size $L = 10$ or $L = 12$ involves the 10 eigenstates associated with energies closest to the middle $[\max(E_i) + \min(E_i)]/2$ of the spectrum E_i [$i = 1, 2, \dots, \dim(H)$], while for systems of size $L = 14$ this number is set to 50. Histograms are drawn from the data of at least 2.5×10^5 eigenstates, which corresponds to at least 25,000 disorder realizations for $L = 10$ and $L = 12$, or at least 5,000 disorder realizations for $L = 14$.

For the calculation of entanglement spectra, we split the chain into subsystems A and B covering respectively the first and last $L/2$ sites, such that $n = 2^{L/2}$. Note that for $\Gamma = 0$ the Hamiltonian reduces to the “standard model of MBL” [11,12,39]. Then, the total spin projection

$$S^z = \sum_{i=1}^L S_i^z \quad (13)$$

is a conserved quantity, due to which the entanglement spectrum is given by the union of independent subspectra labeled by either

$$S_A^z = \sum_{i \in A} S_i^z \quad \text{or} \quad S_B^z = \sum_{i \in B} S_i^z. \quad (14)$$

This phenomenon is reflected in, e.g., a block-diagonal structure of X in Eq. (8).

V. RESULTS

We here show the main result, namely the observation of Gumbel statistics for the entanglement spectra of many-body localized eigenstates. Let $e_{\min} = \min_i(e_i)$ denote the smallest element of an entanglement spectrum. Because Gumbel statistics are formulated for the largest element of a sequence, we study the statistics of $-e_{\min}$.

Let $\langle \cdot \rangle$ denote an expectation value, and let

$$\mu = \langle -e_{\min} \rangle, \quad \sigma^2 = \langle e_{\min}^2 \rangle - \langle e_{\min} \rangle^2 \quad (15)$$




FIG. 1. Density of \tilde{e}_{\min} for $W = 4$ (top) and $W = 5$ (bottom) at $L = 10, 12, 14$, combined with the densities $g(x)$ and $h(x)$. Note the logarithmic scales on the vertical axes.

denote respectively the mean and variance of the distribution of $-e_{\min}$. We define \tilde{e}_{\min} as

$$\tilde{e}_{\min} = \frac{-e_{\min} - \mu}{\sqrt{\sigma^2}}. \quad (16)$$

By construction, the distribution of \tilde{e}_{\min} has mean zero and standard deviation 1. We compare the density $P\{\tilde{e}_{\min} = x\}$ of \tilde{e}_{\min} with the density function

$$g(x) = \frac{\pi}{\sqrt{6}} \exp \left[- \left(\frac{\pi}{\sqrt{6}} x + \gamma \right) - e^{-\left(\frac{\pi}{\sqrt{6}} x + \gamma \right)} \right] \quad (17)$$

for Gumbel statistics having the same mean and standard deviation. For reference, we also compare it with the standard Gaussian, approximating the statistics of \tilde{e}_{\min} for ergodic states, for which the density function is given by

$$h(x) = \frac{1}{\sqrt{2\pi}} \exp \left(-\frac{1}{2} x^2 \right). \quad (18)$$




FIG. 2. Density of \tilde{e}_{\min} for $W = 2, 3, 4, 5$ at $L = 14$, combined with the densities $g(x)$ and $h(x)$. Note the logarithmic scales on the vertical axis.

Figure 1 compares the density of \tilde{e}_{\min} for $W = 4$ and $W = 5$ at $L = 10, 12, 14$ with $g(x)$ and $h(x)$. The density of \tilde{e}_{\min} is approximated by a histogram with bins of width 0.05, which is normalized to unit area to allow for a direct comparison with the (normalized) probability densities. We observe good agreement with $g(x)$ for both disorder strengths at $L = 12, 14$. Deviations from Gumbel statistics can presumably be attributed to finite-size effects. These effects play a role in both the physics of the eigenstates (becoming stronger localized with increasing system size), as well as in the approach towards the limit $n \rightarrow \infty$ (required to observe Gumbel statistics).

Figure 2 compares the density of \tilde{e}_{\min} with $g(x)$ and $h(x)$ for $W = 2, 3, 4, 5$ at $L = 14$. Qualitative similarities between the density of \tilde{e}_{\min} and $g(x)$ can be observed at all disorder strengths. The eigenstate entanglement spectra of Hamiltonian (12) are known to show statistics deviating from the expectation for ergodic states already at disorder strengths well below the MBL transition [18]. More generally, the thermal side of the MBL transition is not fully ergodic [40]. We were not able to draw conclusions on the convergence of the distribution of \tilde{e}_{\min} towards Gumbel statistics with increasing system size at the thermal side of the MBL transition, and remark that the statistics of \tilde{e}_{\min} cannot be used as a probe for the MBL transition for the system sizes under consideration. Note that at the thermal side of the MBL transition the physical significance of the statistics of \tilde{e}_{\min} is presumably limited due to the vanishing physical weight in the thermodynamic limit $L \rightarrow \infty$.

VI. DISCUSSION

The observation of Gumbel statistics suggests that the largest elements of an entanglement spectrum \tilde{e}_i are uncorrelated. To verify this, we study the presence of short-range correlations between the largest \tilde{e}_i . Short-range correlations can be probed through spacing statistics [41]. We order the \tilde{e}_i




FIG. 3. Densities of r_1 for $W = 2, 3, 4, 5$ at $L = 14$, combined with the densities for Poissonian and Wigner-Dyson statistics ($\beta = 1$).

in decreasing order (i.e., $\tilde{e}_i \geq \tilde{e}_{i+1}$), and focus on the ratio of consecutive spacings $r_1 \in [0, 1]$ given by

$$r_1 = \min\left(\frac{\tilde{e}_1 - \tilde{e}_2}{\tilde{e}_2 - \tilde{e}_3}, \frac{\tilde{e}_2 - \tilde{e}_3}{\tilde{e}_1 - \tilde{e}_2}\right). \quad (19)$$

In the absence of short-range correlations, the distribution of r_1 obeys Poissonian statistics, for which

$$P\{r_1 = x\} = \frac{2}{(1+x)^2}. \quad (20)$$

Ergodic states obey Wigner-Dyson spacing statistics [42]. For systems with time-reversal symmetry (Dyson index $\beta = 1$), the corresponding density of r_1 is well approximated [41] by

$$P\{r_1 = x\} \approx \frac{27}{8} \frac{x + x^2}{(1 + x + x^2)^{5/2}}. \quad (21)$$

Figure 3 compares the density of r_1 for $W = 2, 3, 4, 5$ at $L = 14$ with Poissonian and Wigner-Dyson spacing statistics. At all disorder strengths, the spacing statistics are close to Poissonian, indicating the (near) independence of the largest \tilde{e}_i .

VII. CONCLUSION AND OUTLOOK

In summary, we have provided numerical evidence that the entanglement spectra of many-body localized eigenstates obey Gumbel statistics. Because the physical weight of these spectra is almost fully carried by the few smallest elements, one might expect the extreme value statistics to have physical significance. Our result provides an analytical, parameter-free characterization of many-body localized eigenstates. We stress that no conclusions on the thermal side of the MBL transition can be drawn.

The main open question remaining is the physical mechanism responsible for the occurrence of Gumbel statistics, and the way it can be explained in terms of phenomenological models [5]. A possible starting point for further investigations might be provided by the notion that an entanglement spectrum can be interpreted as the eigenvalue spectrum of the entanglement Hamiltonian

$$H_{\text{ent}} = -\ln(\rho_B), \quad (22)$$

where $\rho_B = \text{Tr}_A(|\psi\rangle\langle\psi|)$ is the partial trace of the density matrix $|\psi\rangle\langle\psi|$ over basis states of subsystem A [15]. One might hypothesize that the statistics of the eigenvector associated with e_{\min} carry relevant information.

ACKNOWLEDGMENTS

We thank Maksym Serbyn for useful discussions. This work is part of the Delta-ITP consortium, a program of the Netherlands Organization for Scientific Research (NWO) that is funded by the Dutch Ministry of Education, Culture and Science (OCW).

-
- [1] R. Nandkishore and D. A. Huse, Many-body localization and thermalization in quantum statistical mechanics, *Annu. Rev. Condens. Matter Phys.* **6**, 15 (2015).
- [2] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, *Colloquium: Many-body localization, thermalization, and entanglement*, *Rev. Mod. Phys.* **91**, 021001 (2019).
- [3] M. Serbyn, Z. Papić, and D. A. Abanin, Local Conservation Laws and the Structure of the Many-Body Localized States, *Phys. Rev. Lett.* **111**, 127201 (2013).
- [4] V. Ros, M. Müller, and A. Scardicchio, Integrals of motion in the many-body localized phase, *Nucl. Phys. B* **891**, 420 (2015).
- [5] D. A. Huse, R. Nandkishore, and V. Oganesyan, Phenomenology of fully many-body-localized systems, *Phys. Rev. B* **90**, 174202 (2014).
- [6] J. Z. Imbrie, Diagonalization and Many-Body Localization for a Disordered Quantum Spin Chain, *Phys. Rev. Lett.* **117**, 027201 (2016).
- [7] M. Serbyn, Z. Papić, and D. A. Abanin, Quantum quenches in the many-body localized phase, *Phys. Rev. B* **90**, 174302 (2014).
- [8] M. Žnidarič, T. Prosen, and P. Prelovšek, Many-body localization in the Heisenberg XXZ magnet in a random field, *Phys. Rev. B* **77**, 064426 (2008).
- [9] J. H. Bardarson, F. Pollmann, and J. E. Moore, Unbounded Growth of Entanglement in Models of Many-Body Localization, *Phys. Rev. Lett.* **109**, 017202 (2012).
- [10] M. Serbyn, Z. Papić, and D. A. Abanin, Universal Slow Growth of Entanglement in Interacting Strongly Disordered Systems, *Phys. Rev. Lett.* **110**, 260601 (2013).
- [11] A. Pal and D. A. Huse, Many-body localization phase transition, *Phys. Rev. B* **82**, 174411 (2010).
- [12] D. J. Luitz, N. Laflorencie, and F. Alet, Many-body localization edge in the random-field Heisenberg chain, *Phys. Rev. B* **91**, 081103(R) (2015).

- [13] V. Khemani, S. P. Lim, D. N. Sheng, and D. A. Huse, Critical Properties of the Many-Body Localization Transition, *Phys. Rev. X* **7**, 021013 (2017).
- [14] P. T. Dumitrescu, R. Vasseur, and A. C. Potter, Scaling Theory of Entanglement at the Many-Body Localization Transition, *Phys. Rev. Lett.* **119**, 110604 (2017).
- [15] H. Li and F. D. M. Haldane, Entanglement Spectrum as a Generalization of Entanglement Entropy: Identification of Topological Order in Non-Abelian Fractional Quantum Hall Effect States, *Phys. Rev. Lett.* **101**, 010504 (2008).
- [16] N. Laflorencie, Quantum entanglement in condensed matter systems, *Phys. Rep.* **646**, 1 (2016).
- [17] P. Vivo, M. P. Pato, and G. Oshanin, Random pure states: Quantifying bipartite entanglement beyond the linear statistics, *Phys. Rev. E* **93**, 052106 (2016).
- [18] Z.-C. Yang, C. Chamon, A. Hama, and E. R. Mucciolo, Two-Component Structure in the Entanglement Spectrum of Highly Excited States, *Phys. Rev. Lett.* **115**, 267206 (2015).
- [19] S. D. Geraedts, R. Nandkishore, and N. Regnault, Many-body localization and thermalization: Insights from the entanglement spectrum, *Phys. Rev. B* **93**, 174202 (2016).
- [20] M. Serbyn, A. A. Michailidis, D. A. Abanin, and Z. Papić, Power-Law Entanglement Spectrum in Many-Body Localized Phases, *Phys. Rev. Lett.* **117**, 160601 (2016).
- [21] Z.-C. Yang, A. Hama, S. M. Giampaolo, E. R. Mucciolo, and C. Chamon, Entanglement complexity in quantum many-body dynamics, thermalization, and localization, *Phys. Rev. B* **96**, 020408(R) (2017).
- [22] F. Pietracaprina, G. Parisi, A. Mariano, S. Pascazio, and A. Scardicchio, Entanglement critical length at the many-body localization transition, *J. Stat. Mech.* (2017) 113102.
- [23] S. D. Geraedts, N. Regnault, and R. M. Nandkishore, Characterizing the many-body localization transition using the entanglement spectrum, *New J. Phys.* **19**, 113021 (2017).
- [24] J. Gray, S. Bose, and A. Bayat, Many-body localization transition: Schmidt gap, entanglement length, and scaling, *Phys. Rev. B* **97**, 201105(R) (2018).
- [25] M. R. Leadbetter, G. Lindgren, and H. Rootzén, *Extremes and Related Properties of Random Sequences and Processes*, Springer Series in Statistics Vol. 11 (Springer-Verlag, New York, 1983).
- [26] S. T. Bramwell, K. Christensen, J.-Y. Fortin, P. C. W. Holdsworth, H. J. Jensen, S. Lise, J. M. López, M. Nicodemi, J.-F. Pinton, and M. Sellitto, Universal Fluctuations in Correlated Systems, *Phys. Rev. Lett.* **84**, 3744 (2000).
- [27] T. Antal, M. Droz, G. Györgyi, and Z. Rácz, $1/f$ Noise and Extreme Value Statistics, *Phys. Rev. Lett.* **87**, 240601 (2001).
- [28] E. Bertin, Global Fluctuations and Gumbel Statistics, *Phys. Rev. Lett.* **95**, 170601 (2005).
- [29] A. Lakshminarayan, S. Tomsovic, O. Bohigas, and S. N. Majumdar, Extreme Statistics of Complex Random and Quantum Chaotic States, *Phys. Rev. Lett.* **100**, 044103 (2008).
- [30] E. J. Gumbel, *Statistics of Extremes* (Columbia University Press, New York, 1958).
- [31] S. T. Bramwell, J.-Y. Fortin, P. C. W. Holdsworth, S. Peysson, J.-F. Pinton, B. Portelli, and M. Sellitto, Magnetic fluctuations in the classical XY model: The origin of an exponential tail in a complex system, *Phys. Rev. E* **63**, 041106 (2001).
- [32] S. N. Majumdar and A. Comtet, Exact Maximal Height Distribution of Fluctuating Interfaces, *Phys. Rev. Lett.* **92**, 225501 (2004).
- [33] H. G. Katzgraber, M. Körner, F. Liers, M. Jünger, and A. K. Hartmann, Universality-class dependence of energy distributions in spin glasses, *Phys. Rev. B* **72**, 094421 (2005).
- [34] S. Hofferberth, I. Lesanovsky, T. Schumm, A. I. Imambekov, V. Gritsev, E. Demler, and J. Schmiedmayer, Probing quantum and thermal noise in an interacting many-body system, *Nat. Phys.* **4**, 489 (2008).
- [35] I. Lovas, B. Dóra, E. Demler, and G. Zaránd, Full counting statistics of time-of-flight images, *Phys. Rev. A* **95**, 053621 (2017).
- [36] G. Györgyi, N. R. Moloney, K. Ozogány, Z. Rácz, and M. Droz, Renormalization-group theory for finite-size scaling in extreme statistics, *Phys. Rev. E* **81**, 041135 (2010).
- [37] D. N. Page, Average Entropy of a Subsystem, *Phys. Rev. Lett.* **71**, 1291 (1993).
- [38] P. J. Forrester, *Log-Gases and Random Matrices*, London Mathematical Society Monographs Vol. 34 (Princeton University Press, Princeton and Oxford, 2010).
- [39] V. Oganesyan and D. A. Huse, Localization of interacting fermions at high temperature, *Phys. Rev. B* **75**, 155111 (2007).
- [40] D. J. Luitz and Y. Bar Lev, The ergodic side of the many-body localization transition, *Ann. Phys. (NY)* **529**, 1600350 (2017).
- [41] Y. Y. Atas, E. Bogomolny, O. Giraud, and G. Roux, Distribution of the Ratio of Consecutive Level Spacings in Random Matrix Ensembles, *Phys. Rev. Lett.* **110**, 084101 (2013).
- [42] M. L. Mehta, *Random Matrices*, 3rd ed., Pure and Applied Mathematics Vol. 142 (Elsevier, New York, 2004).