



UvA-DARE (Digital Academic Repository)

Random Matrix Ensemble for the Level Statistics of Many-Body Localization

Buijsman, W.; Cheianov, V.; Gritsev, V.

Published in:
Physical Review Letters

DOI:
[10.1103/PhysRevLett.122.180601](https://doi.org/10.1103/PhysRevLett.122.180601)

[Link to publication](#)

Citation for published version (APA):
Buijsman, W., Cheianov, V., & Gritsev, V. (2019). Random Matrix Ensemble for the Level Statistics of Many-Body Localization. *Physical Review Letters*, 122(18), [180601]. <https://doi.org/10.1103/PhysRevLett.122.180601>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Random Matrix Ensemble for the Level Statistics of Many-Body Localization

Wouter Buijsman,^{1,*} Vadim Cheianov,² and Vladimir Gritsev^{1,3}

¹*Institute for Theoretical Physics Amsterdam and Delta Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*

²*Instituut-Lorentz and Delta Institute for Theoretical Physics, Universiteit Leiden, P.O. Box 9506, 2300 RA Leiden, The Netherlands*

³*Russian Quantum Center, Skolkovo, Moscow 143025, Russia*



(Received 13 July 2018; published 10 May 2019)

We numerically study the level statistics of the Gaussian β ensemble. These statistics generalize Wigner-Dyson level statistics from the discrete set of Dyson indices $\beta = 1, 2, 4$ to the continuous range $0 < \beta < \infty$. The Gaussian β ensemble covers Poissonian level statistics for $\beta \rightarrow 0$, and provides a smooth interpolation between Poissonian and Wigner-Dyson level statistics. We establish the physical relevance of the level statistics of the Gaussian β ensemble by showing near-perfect agreement with the level statistics of a paradigmatic model in studies on many-body localization over the entire crossover range from the thermal to the many-body localized phase. In addition, we show similar agreement for a related Hamiltonian with broken time-reversal symmetry.

DOI: [10.1103/PhysRevLett.122.180601](https://doi.org/10.1103/PhysRevLett.122.180601)

Random matrix theory [1,2] provides an essential toolbox in nuclear [3–5], condensed matter [6–8] and mesoscopic [9,10] physics, and is used as well in, e.g., high energy physics [11–13]. In these fields, the physical interest for random matrix theory comes from the apparent universality of the local spectral statistics of quantum systems that are chaotic in the semiclassical limit [14]. Inspired by seminal works of Wigner [15] and Dyson [16], one typically compares local spectral statistics with the local eigenvalue statistics of random matrices taken from the Gaussian orthogonal (GOE), unitary (GUE), or symplectic (GSE) ensemble—depending on the type of transformation by which the Hamiltonian is diagonalized. These so-called Wigner-Dyson level statistics provide an excellent description of the local spectral statistics of a vast majority of the systems that are considered as quantum chaotic (ergodic) [17].

The GOE, GUE, and GSE are covered by to the Gaussian β ensemble [16,18]. Here, $\beta \in (0, \infty)$ is a continuous parameter which for the GOE, GUE, and GSE corresponds to $\beta = 1, 2, 4$, respectively. The Gaussian β ensemble also covers Poissonian level statistics ($\beta \rightarrow 0$), as typically observed for regular (nonergodic) systems [19,20]. The Gaussian β ensemble provides a smooth interpolation between Poissonian and Wigner-Dyson level statistics. Thanks to relatively recent progress made by Dumitriu and Edelman [21], the eigenvalue statistics of the Gaussian β ensemble can be sampled at low computational costs.

Physical systems displaying level statistics that can be tuned from Poissonian to Wigner-Dyson are of central interest in the field of many-body localization (MBL) [22,23]. Numerical studies provide evidence [24–27] for an intermediate phase characterized by, e.g., Griffiths

effects in between the thermal (corresponding to $\beta \approx 1$) and the MBL (corresponding to $\beta \approx 0$) phase at finite system sizes. In this Letter, we numerically study the level statistics of a standard model in studies on MBL. Remarkably, we find near-perfect agreement with the eigenvalue statistics of the Gaussian β ensemble over the entire crossover range, where β is a single fitting parameter. Additionally, we show that similar agreement holds for a related Hamiltonian with broken time-reversal symmetry. We interpret the eigenvalue statistics of the Gaussian β ensemble as generalized Wigner-Dyson level statistics. We show how the Gaussian β ensemble provides a smooth interpolation between Poissonian and Wigner-Dyson level statistics by a systematic investigation of the eigenvalue statistics for $\beta \in [0, 1]$.

Gaussian β ensemble.—An ensemble of random matrices T is described by a probability distribution $P(T)$ [1]. An example is the GOE. For this ensemble of real symmetric matrices, the probability distribution is given by

$$P(T) = C_n e^{-\text{Tr}(T^2)}, \quad (1)$$

where C_n is a normalization constant and $\text{Tr}(\cdot)$ denotes a trace. The GOE is invariant under transformations $T \rightarrow O^{-1}TO$ for real orthogonal matrices O . Similarly, the GUE and GSE are invariant under unitary and symplectic transformations, respectively. Because there are only three types of associative division algebras (real, complex, and quaternionic numbers), no invariant random matrix ensembles exist beyond the GOE, GUE, and GSE. The joint probability distribution for the eigenvalues $\{e_i\}$ of n -dimensional matrices from the Gaussian ensembles is given by

TABLE I. Numerically obtained values of $\langle r \rangle$ for the Gaussian β ensemble at various β (see main text for details). The value for $\beta = 0$ is obtained from the expression given in the main text.

β	0.00	0.01	0.05	0.10	0.15	0.20
$\langle r \rangle$	0.386(3)	0.389(0)	0.398(4)	0.408(9)	0.420(1)	0.429(5)
β	0.25	0.30	0.35	0.40	0.45	0.50
$\langle r \rangle$	0.438(1)	0.446(2)	0.453(6)	0.461(7)	0.469(3)	0.475(8)
β	0.55	0.60	0.65	0.70	0.75	0.80
$\langle r \rangle$	0.482(6)	0.489(0)	0.494(6)	0.500(8)	0.505(8)	0.511(2)
β	0.85	0.90	0.95	1.00	2.00	4.00
$\langle r \rangle$	0.516(4)	0.521(5)	0.526(2)	0.530(2)	0.599(7)	0.673(9)

Comparison with spectral statistics.—Here, we compare the level statistics of a standard model in studies on MBL with the eigenvalue statistics of the Gaussian β ensemble. We consider a disordered spin-1/2 XXZ chain, for which the Hamiltonian is given by

$$H = \sum_{i=1}^L (S_i^x S_{i+1}^x + S_i^y S_{i+1}^y + \Delta S_i^z S_{i+1}^z) + \sum_{i=1}^L h_i S_i^z, \quad (7)$$

with $S_i^\alpha = \frac{1}{2} \sigma_i^\alpha$, where σ_i^α are Pauli matrices ($\alpha = x, y, z$) acting on site i . During the last decade, the level statistics of this Hamiltonian have been studied extensively in, e.g., Refs. [7,8,34–38]. In particular, the intermediate level statistics between the thermal and the MBL phase have been studied by means of a two-stage flow picture in Ref. [8]. Following these references, we impose periodic boundary conditions $\sigma_{i+L}^\alpha \equiv \sigma_i^\alpha$, sample h_i from the uniform distribution ranging over $[-W, W]$, set $\Delta = 1$ (unless stated otherwise), and restrict the focus to the symmetry sector $\sum_i S_i^z = 0$. We set $L = 16$, for which $\dim(H) = 12870$. We consider at least 1000 disorder realizations for each value of W . For each value of W separately, we restrict the focus to the energy window containing the middle 10% of the union of all sampled spectra. The system exhibits a smooth crossover from Poissonian to Wigner-Dyson level statistics in the region $1.7 \lesssim W \lesssim 4.0$.

Figure 2 shows the distributions of r and s for the spectra of the Hamiltonian compared with the corresponding distributions for the Gaussian β ensemble, where β is estimated from $\langle r \rangle$. Note that, since r is independent of the average level spacing, no unfolding [33] is required for drawing the distribution of this quantity. Before drawing the histograms of s for the Hamiltonian, the spectra are unfolded by numerically estimating the smooth part of the density of states [39]. Remarkably, we observe near-perfect agreement between the spectral statistics of the Hamiltonian and the corresponding eigenvalue statistics of




FIG. 2. Numerically obtained distributions of r and s for the Hamiltonian at various W (solid lines) and the corresponding distributions for the Gaussian β ensemble (dashed lines, identical color scheme). The top and bottom left plots are for $\Delta = 1$, the bottom right one for $\Delta = 2$.

the Gaussian β ensemble at all disorder strengths. Similar agreement can be found for $\Delta = 2$, which is illustrated in the lower right panel.

In Fig. 3, we study the sensitivity to finite-size effects. The top panels show that the agreement between the level




FIG. 3. Numerically obtained distributions of r for the Hamiltonian at various W (solid lines) and the corresponding distributions for the Gaussian β ensemble (dashed lines, identical color scheme) for $L = 12, 14$ (top panels) and the estimated value of β for the spectra of the Hamiltonian as a function of L and W (bottom panels).




FIG. 4. Numerically obtained distribution of r for $H + H'$ at various W (solid lines) and the corresponding distributions for the Gaussian β ensemble (dashed lines, identical color scheme).

statistics of the Hamiltonian and the Gaussian β is near perfect also at $L = 12, 14$. The bottom panels show a flow towards Wigner-Dyson (Poissonian) level statistics for $W \lesssim 3$ ($W \gtrsim 3$) with increasing system size. For $L \rightarrow \infty$, the system is believed to display an MBL transition at $W \approx 3.6$ [34]. As there is a one-to-one relation between $\langle r \rangle$ and β , these results can in principle be appended with previous results from, e.g., Ref. [35]. Studying $\langle r \rangle$ as a function of the matrix dimension n for the Gaussian β ensemble indicates a difference of less than 1% between the value for $n = 500$ and $n = 10^5$ at all values $\beta \in [0, 1]$.

Breaking time-reversal symmetry.—Ergodic systems with broken time-reversal symmetry are characterized by Wigner-Dyson level statistics for $\beta = 2$ [1]. For the Hamiltonian given in Eq. (7), time-reversal symmetry can be broken in an experimentally relevant way by adding the three-body term

$$H' = \sum_{i=1}^L \vec{S}_i \cdot (\vec{S}_{i+1} \times \vec{S}_{i+2}), \quad (8)$$

where $\vec{S}_i = [S_i^x, S_i^y, S_i^z]^T$ [40]. For $\beta \gtrsim 1$, the distribution of the ratio of consecutive level spacings for the Gaussian β ensemble can be approximated with high precision [32] from Eq. (2) with $n = 3$, giving

$$P(r) \sim \frac{(r + r^2)^\beta}{(1 + r + r^2)^{1+3\beta/2}}. \quad (9)$$

In what follows, estimates of $\beta \geq 1$ from $\langle r \rangle$ are obtained by using Eq. (9). Figure 4 shows the distribution of r for the spectra of $H + H'$ at several values of W compared with the corresponding distributions for the Gaussian β ensemble, where β is estimated from $\langle r \rangle$. Again, we observe near-perfect agreement between corresponding curves at all disorder strengths.

Higher order spacing ratios.—Going beyond the study of the distribution of the ratios of consecutive level spacings




FIG. 5. Numerically obtained distributions of $r^{(n)}$ with $n = 2, 3, 4, 5$ for the Hamiltonian at various W (solid lines) and the corresponding distributions for the Gaussian β ensemble (dashed lines, identical color scheme).

and the level spacing distribution, we here study the higher order ratios $r^{(n)} \in [0, 1]$ of level spacings, for a spectrum $\{E_i\}$ sorted in ascending order defined as

$$r_i^{(n)} = \min \left(\frac{E_{i+2n} - E_{i+n}}{E_{i+n} - E_i}, \frac{E_{i+n} - E_i}{E_{i+2n} - E_{i+n}} \right). \quad (10)$$

Note that $r^{(1)} = r$. For the Gaussian β ensemble, it can be shown rigorously that the distribution of $r^{(n)}$ for $\beta = 2/(n+1)$ is equivalent to the distribution of $r^{(1)}$ for $\beta = 2(n+1)$ [41]. Evidence for a broader class of interrelations involving $\beta = 1, 2, 4$ has been provided recently in Ref. [42].

Figure 5 shows the distributions of $r^{(n)}$ for the Hamiltonian compared with the corresponding distributions for the Gaussian β ensemble, where the value of β is estimated from $\langle r \rangle$. No unfolding is applied to the spectra of the Hamiltonian. We observe qualitative agreement up to $n = 3$ (i.e., up to 6 level spacings) for all values of W . The algorithm used to unfold the spectra can be suboptimal for the system under consideration. Attempts to compare the spectra of the Hamiltonian and the Gaussian β ensemble on longer ranges by other measures such as the spectral rigidity [43], density-density correlation function [44], and the spectral form factor [45] did not provide conclusive results, presumably due to this effect.

Discussion and conclusions.—We have proposed a generalization of Wigner-Dyson level statistics from the discrete taxonomy $\beta = 1, 2, 4$ to the continuous one $\beta \in (0, \infty)$. Using the matrix model for the Gaussian β ensemble introduced in Ref. [21], we have shown how the Gaussian β ensemble provides a smooth interpolation between Poissonian and Wigner-Dyson level statistics.

We have studied the level statistics of a paradigmatic model in studies on MBL, and found near-perfect agreement with the corresponding statistics of the Gaussian β ensemble over the full crossover range between the thermal (corresponding to $\beta \approx 1$) and many-body localized (corresponding to $\beta \approx 0$) phase, where β is a single fitting parameter. We have shown that similar agreement holds for a related Hamiltonian with broken time-reversal symmetry.

We expect that this work paves a way for further investigations in various ways. Primarily, we believe it would be of significant interest to explore how universal the Gaussian β ensemble describes the spectral statistics of quantum systems that show intermediate level statistics between Poissonian and Wigner-Dyson. In view of this, we note that there are several known physical and mathematical models supporting intermediate level statistics, studied mostly in the context of either single-particle models of quantum chaos [46–49] or the Anderson localization transition for noninteracting systems [50–53]. A crossover between Poissonian and Wigner-Dyson level statistics for $\beta = 2$ has also been found recently in a generalized SYK model [54].

Next, we expect that our results are of relevance in the field of MBL. In this field, level statistics are a key ingredient in both numerical [7,35,37] and analytical [55] studies. The detailed quantitative characterization of the level statistics of the Hamiltonian provided in this work might be valuable in, e.g., the finite-size scaling analysis of the MBL transition [34,35] and studies on the intermediate phase separating the thermal from the MBL phase [26] at finite system sizes. Finally, we hope that this Letter can contribute to the ongoing studies [45,56,57] on the fundamental correspondence between classical and quantum chaos.

We thank Tomaž Prosen and Maksym Serbyn for very useful discussions. V.G. acknowledges support from the Erwin Schrödinger Institute in Vienna. This Letter is part of the Delta-ITP consortium, a program of the Netherlands Organization for Scientific Research (NWO) that is funded by the Dutch Ministry of Education, Culture, and Science (OCW).

* w.buijsman@uva.nl

- [1] M. L. Mehta, *Random Matrices*, 3rd ed., Pure and Applied Mathematics Vol. 142 (Elsevier, New York, 2004).
- [2] T. Guhr, A. Müller-Groeling, and H. A. Weidenmüller, Random-matrix theories in quantum physics: common concepts, *Phys. Rep.* **299**, 189 (1998).
- [3] G. E. Mitchell, E. G. Bilpuch, P. M. Endt, and J. F. Shrinier, Broken Symmetries and Chaotic Behavior in ^{26}Al , *Phys. Rev. Lett.* **61**, 1473 (1988).
- [4] H. A. Weidenmüller and G. E. Mitchell, Random matrices and chaos in nuclear physics: Nuclear structure, *Rev. Mod. Phys.* **81**, 539 (2009).
- [5] B. Dietz, A. Heusler, K. H. Maier, A. Richter, and B. A. Brown, Chaos and Regularity in the Doubly Magic Nucleus ^{208}Pb , *Phys. Rev. Lett.* **118**, 012501 (2017).
- [6] B. I. Shklovskii, B. Shapiro, B. R. Sears, P. Lambrianides, and H. B. Shore, Statistics of spectra of disordered systems near the metal-insulator transition, *Phys. Rev. B* **47**, 11487 (1993).
- [7] V. Oganesyan and D. A. Huse, Localization of interacting fermions at high temperature, *Phys. Rev. B* **75**, 155111 (2007).
- [8] M. Serbyn and J. E. Moore, Spectral statistics across the many-body localization transition, *Phys. Rev. B* **93**, 041424(R) (2016).
- [9] C. W. J. Beenakker, Universality in the Random-Matrix Theory of Quantum Transport, *Phys. Rev. Lett.* **70**, 1155 (1993).
- [10] C. W. J. Beenakker, J. M. Edge, J. P. Dahlhaus, D. I. Pikulin, S. Mi, and M. Wimmer, Wigner-Poisson Statistics of Topological Transitions in a Josephson Junction, *Phys. Rev. Lett.* **111**, 037001 (2013).
- [11] M. Giordano, T. G. Kovács, and F. Pittler, Universality and the QCD Anderson Transition, *Phys. Rev. Lett.* **112**, 102002 (2014).
- [12] T. Li, J. Liu, Y. Xin, and Y. Zhou, Supersymmetric SYK model and random matrix theory, *J. High Energy Phys.* **06** (2017) 111.
- [13] T. G. Kovács and R. Á. Vig, Localization transition in $\text{SU}(3)$ gauge theory, *Phys. Rev. D* **97**, 014502 (2018).
- [14] O. Bohigas, M. J. Giannoni, and C. Schmit, Characterization of Chaotic Quantum Spectra and Universality of Level Fluctuation Laws, *Phys. Rev. Lett.* **52**, 1 (1984).
- [15] E. P. Wigner, Characteristic vectors of bordered matrices with infinite dimensions, *Ann. Math.* **62**, 548 (1955).
- [16] F. J. Dyson, Statistical theory of the energy levels of complex systems. I, *J. Math. Phys. (N.Y.)* **3**, 140 (1962).
- [17] D. Poilblanc, T. Ziman, J. Bellissard, F. Mila, and G. Montambaux, Poisson vs GOE statistics in integrable and non-integrable quantum hamiltonians, *Europhys. Lett.* **22**, 537 (1993).
- [18] P. J. Forrester, *Log-Gases and Random Matrices*, London Mathematical Society Monographs Vol. 34 (Princeton University Press, Princeton and Oxford, 2010).
- [19] M. V. Berry and M. Tabor, Level clustering in the regular spectrum, *Proc. R. Soc. A* **356**, 375 (1977).
- [20] A. Relaño, J. Dukelsky, J. M. G. Gómez, and J. Retamosa, Stringent numerical test of the Poisson distribution for finite quantum integrable Hamiltonians, *Phys. Rev. E* **70**, 026208 (2004).
- [21] I. Dumitriu and A. Edelman, Matrix models for beta ensembles, *J. Math. Phys. (N.Y.)* **43**, 5830 (2002).
- [22] D. M. Basko, I. L. Aleiner, and B. L. Altshuler, Metal-insulator transition in a weakly interacting many-electron system with localized single-particle states, *Ann. Phys. (Amsterdam)* **321**, 1126 (2006).
- [23] D. A. Abanin and Z. Papić, Recent progress in many-body localization, *Ann. Phys. (Amsterdam)* **529**, 1700169 (2017).
- [24] Y. Bar Lev, G. Cohen, and D. R. Reichman, Absence of Diffusion in an Interacting System of Spinless Fermions on a One-Dimensional Disordered Lattice, *Phys. Rev. Lett.* **114**, 100601 (2015).

- [25] K. Agarwal, S. Gopalakrishnan, M. Knap, M. Müller, and E. Demler, Anomalous Diffusion and Griffiths Effects Near the Many-Body Localization Transition, *Phys. Rev. Lett.* **114**, 160401 (2015).
- [26] V. Khemani, S. P. Lim, D. N. Sheng, and D. A. Huse, Critical Properties of the Many-Body Localization Transition, *Phys. Rev. X* **7**, 021013 (2017).
- [27] D. J. Luitz and Y. Bar Lev, The ergodic side of the many-body localization transition, *Ann. Phys. (Amsterdam)* **529**, 1600350 (2017).
- [28] I. Dumitriu and A. Edelman, Global spectrum fluctuations for the β -Hermite and β -Laguerre ensembles via matrix models, *J. Math. Phys. (N.Y.)* **47**, 063302 (2006).
- [29] G. Le Caër, C. Male, and R. Delannay, Nearest-neighbour spacing distributions of the β -Hermite ensemble of random matrices, *Physica (Amsterdam)* **383A**, 190 (2007).
- [30] A. Relaño, L. Muñoz, J. Retamosa, E. Faleiro, and R. A. Molina, Power-spectrum characterization of the continuous Gaussian ensemble, *Phys. Rev. E* **77**, 031103 (2008).
- [31] P. Vivo and S. N. Majumdar, On invariant 2×2 β -ensembles of random matrices, *Physica (Amsterdam)* **387A**, 4839 (2008).
- [32] Y. Y. Atas, E. Bogomolny, O. Giraud, and G. Roux, Distribution of the Ratio of Consecutive Level Spacings in Random Matrix Ensembles, *Phys. Rev. Lett.* **110**, 084101 (2013).
- [33] F. Haake, *Quantum Signatures of Chaos*, 3rd ed., Springer Series in Synergetics, Vol. 54 (Springer-Verlag, Berlin Heidelberg, 2010).
- [34] A. Pal and D. A. Huse, Many-body localization phase transition, *Phys. Rev. B* **82**, 174411 (2010).
- [35] D. J. Luitz, N. Laflorencie, and F. Alet, Many-body localization edge in the random-field Heisenberg chain, *Phys. Rev. B* **91**, 081103(R) (2015).
- [36] C. L. Bertrand and A. M. García-García, Anomalous Thouless energy and critical statistics on the metallic side of the many-body localization transition, *Phys. Rev. B* **94**, 144201 (2016).
- [37] J. A. Kjäll, Many-body localization and level repulsion, *Phys. Rev. B* **97**, 035163 (2018).
- [38] K. Kudo and T. Deguchi, Finite-size scaling with respect to interaction and disorder strength at the many-body localization transition, *Phys. Rev. B* **97**, 220201(R) (2018).
- [39] We estimate the (scaled) smooth part of the density of states for each spectrum separately by using the Wolfram Research, Inc. *Mathematica* version 10.0 function `SmoothKernelDistribution` with bandwidth specifications `{"Adaptive", Automatic, 1}`, `"Biweight" (W = 2, 3)` or `{"Adaptive", Automatic, 1}` ($W = 4, 5$).
- [40] Y. Avishai, J. Richert, and R. Berkovits, Level statistics in a Heisenberg chain with random magnetic field, *Phys. Rev. B* **66**, 052416 (2002).
- [41] P. J. Forrester, A random matrix decimation procedure relating $\beta = 2/(r+1)$ to $\beta = 2(r+1)$, *Commun. Math. Phys.* **285**, 653 (2009).
- [42] S. H. Tekur, U. T. Bhosale, and M. S. Santhanam, Higher-order spacing ratios in random matrix theory and complex quantum systems, *Phys. Rev. B* **98**, 104305 (2018).
- [43] G. Montambaux, D. Poilblanc, J. Bellissard, and C. Sire, Quantum Chaos in Spin-Fermion Models, *Phys. Rev. Lett.* **70**, 497 (1993).
- [44] C. H. Joyner, S. Müller, and M. Sieber, Semiclassical approach to discrete symmetries in quantum chaos, *J. Phys. A* **45**, 205102 (2012).
- [45] P. Kos, M. Ljubotina, and T. Prosen, Many-Body Quantum Chaos: Analytic Connection to Random Matrix Theory, *Phys. Rev. X* **8**, 021062 (2018).
- [46] F. M. Izrailev, Intermediate statistics of the quasi-energy spectrum and quantum localisation of classical chaos, *J. Phys. A* **22**, 865 (1989).
- [47] T. Prosen and M. Robnik, Energy level statistics in the transition region between integrability and chaos, *J. Phys. A* **26**, 2371 (1993).
- [48] T. Prosen and M. Robnik, Semiclassical energy level statistics in the transition region between integrability and chaos: Transition from Brody-like to Berry-Robnik behaviour, *J. Phys. A* **27**, 8059 (1994).
- [49] E. B. Bogomolny, U. Gerland, and C. Schmit, Models of intermediate spectral statistics, *Phys. Rev. E* **59**, R1315 (1999).
- [50] M. Moshe, H. Neuberger, and B. Shapiro, Generalized Ensemble of Random Matrices, *Phys. Rev. Lett.* **73**, 1497 (1994).
- [51] V. E. Kravtsov and I. V. Lerner, Effective plasma model for the level correlations at the mobility edge, *J. Phys. A* **28**, 3623 (1995).
- [52] V. E. Kravtsov and K. A. Muttalib, New Class of Random Matrix Ensembles with Multifractal Eigenvectors, *Phys. Rev. Lett.* **79**, 1913 (1997).
- [53] S. Sorathia, F. M. Izrailev, V. G. Zelevinsky, and G. L. Celardo, From closed to open one-dimensional Anderson model: Transport versus spectral statistics, *Phys. Rev. E* **86**, 011142 (2012).
- [54] A. M. García-García, B. Loureiro, A. Romero-Bermúdez, and M. Tezuka, Chaotic-Integrable Transition in the Sachdev-Ye-Kitaev Model, *Phys. Rev. Lett.* **120**, 241603 (2018).
- [55] J. Z. Imbrie, On many-body localization for quantum spin chains, *J. Stat. Phys.* **163**, 998 (2016).
- [56] S. Müller, S. Heusler, P. Braun, F. Haake, and A. Altland, Semiclassical Foundation of Universality in Quantum Chaos, *Phys. Rev. Lett.* **93**, 014103 (2004).
- [57] W. Buijsman, V. Gritsev, and R. Sprik, Nonergodicity in the Anisotropic Dicke Model, *Phys. Rev. Lett.* **118**, 080601 (2017).