



UvA-DARE (Digital Academic Repository)

Radio non-detection during nearly-simultaneous Swift/XRT observations of MAXI J0911-635/Swift J0911.9-6452 in NGC 2808

Tudor, V.; Bahramian, A.; Sivakoff, G.; Chomiuk, L.; Heinke, C.; Li, R.; Maccarone, T.; Miller-Jones, J.; Plotkin, R.; Russell, T.; Strader, J.; Tetarenko, A.; Tremou, E.

Publication date

2016

Document Version

Final published version

Published in

The astronomer's telegram

License

Unspecified

[Link to publication](#)

Citation for published version (APA):

Tudor, V., Bahramian, A., Sivakoff, G., Chomiuk, L., Heinke, C., Li, R., Maccarone, T., Miller-Jones, J., Plotkin, R., Russell, T., Strader, J., Tetarenko, A., & Tremou, E. (2016). Radio non-detection during nearly-simultaneous Swift/XRT observations of MAXI J0911-635/Swift J0911.9-6452 in NGC 2808. *The astronomer's telegram*, 8914. <http://www.astronomerstelegram.org/?read=8914>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Outside

GCN
IAUCs

Other

ATel on [Twitter](#) and [Facebook](#)
ATELstream
ATel Community Site[[Previous](#) | [Next](#) | [ADS](#)]

Radio non-detection during nearly-simultaneous Swift/XRT observations of MAXI J0911-635/Swift J0911.9-6452 in NGC 2808

ATel #8914; *V. Tudor (ICRAR-Curtin), A. Bahramian, G. Sivakoff (Alberta), L. Chomiuk (Michigan State), C. Heinke (Alberta), R. Li (Michigan State), T. Maccarone (Texas Tech), J. Miller-Jones, R. Plotkin, T. Russell (ICRAR-Curtin), J. Strader (Michigan State), A. Tetarenko (Alberta), E. Tremou (Michigan State)*

on 6 Apr 2016; 09:09 UT

Credential Certification: James Miller-Jones (james.miller-jones@curtin.edu.au)

Subjects: Radio, X-ray, Binary, Globular Cluster, Neutron Star, Transient

Referred to by ATel #: [8971](#), [8986](#), [9738](#), [10425](#)

We report nearly-simultaneous radio and X-ray observations of the newly discovered X-ray transient MAXI J0911-635/Swift J0911.9-6452 in the globular cluster NGC 2808 (ATel #[8872](#), #[8884](#)). These observations indicate the accretor is likely a neutron star.

Following its detection, we triggered a NAPA program on the ATCA (C2902, PI:Sivakoff) and requested contemporaneous Swift observations. We carried out ATCA observations between 05:22 UT and 07:57 UT on 2016 April 4, for a total on-source integration time of 2.1h, using two 2-GHz basebands centred at 5.5 and 9 GHz. The array was in a compact configuration (H214), with a resolution of 4.2"x0.7" on the longest baselines. No source was detected within the 90% Swift/XRT error circle (ATel #[8884](#)), down to a 5 sigma upper limit of 70 μ Jy. Assuming a flat radio spectrum, this upper limit corresponds to a 5 GHz luminosity of $4.5e28$ erg/s ($L = 4 \pi d^2 v S_{\nu}$) at the distance of NGC 2808 (10.4kpc; Correnti et al. 2016, arXiv:1603.05254).

Our Swift/XRT observation was taken an hour after the ATCA observations, between 08:52:29 and 09:00:58 UT in WT mode, with the source having a 0.5-10 keV count rate of 3.72 \pm 0.10 cnt/s. The spectrum can be well fit by an absorbed power-law, typical of X-ray binaries in a hard accretion state. We find an absorption column density $N_{\text{H}} = (7.7\pm 1.5)e21$ cm⁻² (abundances from Wilms et al., 2000, ApJ 542, 914; cross sections from Verner et al., 1996, ApJ, 465, 487), photon index 1.85 \pm 0.16, and unabsorbed 1-10 keV X-ray flux of (2.40 \pm 0.15)e-10 erg cm⁻²s⁻¹

Related

- 13760 [NICER observes the tail of a long-duration X-ray burst in the direction of MAXI J0911-655](#)
- 13754 [MAXI detection of a long X-ray burst possibly from MAXI J0911-655/Swift J0911.9-6452](#)
- 12869 [NICER follow-up observations of the accreting millisecond X-ray pulsar MAXI J0911-655](#)
- 12846 [The transiently accreting millisecond X-ray pulsar Swift J0911.9-6452 returns to a high X-ray activity level](#)
- 12831 [The accreting millisecond X-ray pulsar Swift J0911.9-6452 decays in X-rays after ~3.2 years of activity](#)
- 10425 [INTEGRAL detection of continued hard X-ray emission from MAXI J0911-655](#)
- 9740 [Parkes upper limit on pulsed radio emission from MAXI J0911-655 at 20-cm](#)
- 9738 [INTEGRAL detection of MAXI J0911-655: still active](#)
- 8986 [INTEGRAL observation of MAXI J0911-655](#)
- 8971 [Chandra identification of the X-ray transient MAXI J0911-635/Swift J0911.9-6452 in NGC 2808](#)
- 8914 [Radio non-detection during nearly-simultaneous Swift/XRT observations of MAXI J0911-635/Swift J0911.9-6452 in NGC 2808](#)
- 8884 [MAXI J0911-655/Swift J0911.9-6452: Swift XRT localization of the new transient in NGC 2808](#)
- 8872 [MAXI/GSC and Swift/BAT detections of a new X-ray outburst from MAXI J0911-655/Swift J0911.9-6452](#)

(90% confidence; $\chi_{\text{red}}^2 = 0.9$, 46 dof). This corresponds to an X-ray luminosity of $\sim 3.1 \times 10^{36}$ erg/s in the 1-10 keV band at 10.4 kpc, or $\sim 1.8\%$ of the Eddington limit of a 1.4 solar mass neutron star.

Since the expected absorption column density towards NGC 2808 is $N_{\text{H}} = 1.9 \times 10^{21} \text{ cm}^{-2}$ (following Bahramian et al. 2015, MNRAS, 452, 3475), our X-ray spectrum indicates that either the source is intrinsically absorbed or that the spectrum is not an absorbed power-law. We therefore tried an absorbed disk blackbody model (more consistent with soft/thermal state accretion), finding an absorption column density $N_{\text{H}} = (3.8 \pm 1.0) \times 10^{21} \text{ cm}^{-2}$, disk temperature of 1.5 ± 0.2 keV, and unabsorbed 1-10 keV X-ray flux of $(1.9 \pm 0.1) \times 10^{-10} \text{ erg cm}^{-2} \text{ s}^{-1}$. The fit is statistically better ($\chi_{\text{red}}^2 = 0.8$ for 46 dof) than the powerlaw fit. The implied projected inner disk radius is small (1.4 ± 0.3 km), which would imply a high inclination angle. The flux corresponds to an X-ray luminosity of $\sim 2.5 \times 10^{36}$ erg/s in the 1-10 keV band at 10.4 kpc, or $\sim 1.4\%$ of the Eddington limit of a 1.4 solar mass neutron star.

Typical soft-to-hard transitions occur at bolometric luminosities of 2% of the Eddington luminosity (Maccarone 2003, A&A, 409, 697). Hard-to-soft transitions generally occur near the peaks of outbursts, except in the case of outbursts that do not have soft states. The luminosity here is slightly below 2% Eddington, albeit without bolometric corrections. Additionally, soft states are more susceptible to inclination angle effects than hard states, and this could be an indication that the source is a high inclination angle system. If the source continues to rise in X-ray flux, that would argue against this scenario, and instead favor a hard state with some absorption intrinsic to the source.

From the most recent plot of the radio/X-ray correlation for black hole and neutron star systems (Deller et al., 2015, ApJ, 809, 13) and the radio upper limit, we suggest that MAXI J0911-635 is either a typical neutron star low-mass X-ray binary or a radio-quiet accreting millisecond pulsar.

Further radio observations are planned.

We thank the ATNF and Swift staff for the rapid scheduling of the observations.

[[Telegram Index](#)]

R. E. Rutledge, Editor-in-Chief

Derek Fox, Editor

Mansi M. Kasliwal, Co-Editor

rrutledge@astronomerstelegam.org

dfox@astronomerstelegam.org

mansi@astronomerstelegam.org