



UvA-DARE (Digital Academic Repository)

Corrigendum: Comparison of 21-Plex PCR and API 20C AUX, MALDI-TOF MS, and rDNA Sequencing for a Wide Range of Clinically Isolated Yeast Species: Improved Identification by Combining 21-Plex PCR and API 20C AUX as an Alternative Strategy for Developing Countries

Arastehfar, A.; Daneshnia, F.; Kord, M.; Roudbary, M.; Zarrinfar, H.; Fang, W.; Hashemi, S.J.; Najafzadeh, M.J.; Khodavaisy, S.; Pan, W.; Liao, W.; Badali, H.; Rezaie, S.; Zomorodian, K.; Hagen, F.; Boekhout, T.

DOI

[10.3389/fcimb.2019.00176](https://doi.org/10.3389/fcimb.2019.00176)

Publication date

2019

Document Version

Final published version

Published in

Frontiers in Cellular and Infection Microbiology

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Arastehfar, A., Daneshnia, F., Kord, M., Roudbary, M., Zarrinfar, H., Fang, W., Hashemi, S. J., Najafzadeh, M. J., Khodavaisy, S., Pan, W., Liao, W., Badali, H., Rezaie, S., Zomorodian, K., Hagen, F., & Boekhout, T. (2019). Corrigendum: Comparison of 21-Plex PCR and API 20C AUX, MALDI-TOF MS, and rDNA Sequencing for a Wide Range of Clinically Isolated Yeast Species: Improved Identification by Combining 21-Plex PCR and API 20C AUX as an Alternative Strategy for Developing Countries. *Frontiers in Cellular and Infection Microbiology*, 9, Article 176. <https://doi.org/10.3389/fcimb.2019.00176>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).



Corrigendum: Comparison of 21-Plex PCR and API 20C AUX, MALDI-TOF MS, and rDNA Sequencing for a Wide Range of Clinically Isolated Yeast Species: Improved Identification by Combining 21-Plex PCR and API 20C AUX as an Alternative Strategy for Developing Countries

OPEN ACCESS

Approved by:

Frontiers Editorial Office,
Frontiers Media SA, Switzerland

*Correspondence:

Sadegh Khodavaisy
sadegh_7392008@yahoo.com
Weihua Pan
panweihua@smmu.edu.cn

†These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Clinical Microbiology,
a section of the journal
Frontiers in Cellular and Infection
Microbiology

Received: 21 April 2019

Accepted: 08 May 2019

Published: 28 May 2019

Citation:

Arastehfar A, Daneshnia F, Kord M,
Roudbary M, Zarrinfar H, Fang W,
Hashemi SJ, Najafzadeh MJ,
Khodavaisy S, Pan W, Liao W,
Badali H, Rezaie S, Zomorodian K,
Hagen F and Boekhout T (2019)
Corrigendum: Comparison of 21-Plex
PCR and API 20C AUX, MALDI-TOF
MS, and rDNA Sequencing for a Wide
Range of Clinically Isolated Yeast
Species: Improved Identification by
Combining 21-Plex PCR and API 20C
AUX as an Alternative Strategy for
Developing Countries.
Front. Cell. Infect. Microbiol. 9:176.
doi: 10.3389/fcimb.2019.00176

Amir Arastehfar^{1†}, Farnaz Daneshnia^{1†}, Mohammad Kord², Maryam Roudbary³,
Hossein Zarrinfar⁴, Wenjie Fang⁵, Sayed Jamal Hashemi^{2,6},
Mohammad Javad Najafzadeh⁷, Sadegh Khodavaisy^{2,8*}, Weihua Pan^{5*}, Wanqing Liao⁵,
Hamid Badali⁹, Sassan Rezaie², Kamiar Zomorodian¹⁰, Ferry Hagen¹ and
Teun Boekhout^{1,5,11}

¹ Yeast Biodiversity Department, Westerdijk Fungal Biodiversity Institute, Utrecht, Netherlands, ² Department of Medical Parasitology and Mycology, Tehran University of Medical Sciences, Tehran, Iran, ³ Department of Medical Mycology and Parasitology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran, ⁴ Allergy Research Center, Mashhad University of Medical Sciences, Mashhad, Iran, ⁵ Shanghai Key Laboratory of Molecular Medical Mycology, Department of Dermatology, Shanghai Institute of Medical Mycology, Shanghai Changzheng Hospital, Second Military Medical University, Shanghai, China, ⁶ Food Microbiology Research Center, Tehran University of Medical Sciences, Tehran, Iran, ⁷ Department of Parasitology and Mycology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran, ⁸ Zoonoses Research Center, Kurdistan University of Medical Sciences, Sanandaj, Iran, ⁹ Department of Medical Mycology, Invasive Fungi Research Center, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran, ¹⁰ Department of Medical Mycology and Parasitology, Basic Sciences in Infectious Diseases Research Center, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran, ¹¹ Yeast Biodiversity Department, Institute of Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, Netherlands

Keywords: API 20C AUX, 21-plex PCR, MALDI-TOF MS, LSU rDNA sequencing, developing countries

A Corrigendum on

Comparison of 21-Plex PCR and API 20C AUX, MALDI-TOF MS, and rDNA Sequencing for a Wide Range of Clinically Isolated Yeast Species: Improved Identification by Combining 21-Plex PCR and API 20C AUX as an Alternative Strategy for Developing Countries
by Arastehfar, A., Daneshnia, F., Kord, M., Roudbary, M., Zarrinfar, H., Fang, W., et al. (2019).
Front. Cell. Infect. Microbiol. 9:21. doi: 10.3389/fcimb.2019.00021

An author's name was incorrectly spelled as "Amir Aarstehfar." The correct spelling is "Amir Arastehfar".

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2019 Arastehfar, Daneshnia, Kord, Roudbary, Zarrinfar, Fang, Hashemi, Najafzadeh, Khodavaisy, Pan, Liao, Badali, Rezaie, Zomorodian, Hagen and Boekhout. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.