



UvA-DARE (Digital Academic Repository)

Static and Dynamic Predictors of General and Violent Criminal Offense Recidivism in the Forensic Outpatient Population: A Meta-Analysis

Eisenberg, M.J.; van Horn, J.E.; Dekker, J.M.; Assink, M.; van der Put, C.E.; Hendriks, J.; Stams, G.J.J.M.

DOI

[10.1177/0093854819826109](https://doi.org/10.1177/0093854819826109)

Publication date

2019

Document Version

Final published version

Published in

Criminal justice and behavior

[Link to publication](#)

Citation for published version (APA):

Eisenberg, M. J., van Horn, J. E., Dekker, J. M., Assink, M., van der Put, C. E., Hendriks, J., & Stams, G. J. J. M. (2019). Static and Dynamic Predictors of General and Violent Criminal Offense Recidivism in the Forensic Outpatient Population: A Meta-Analysis. *Criminal justice and behavior*, 46(5), 732-750. <https://doi.org/10.1177/0093854819826109>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, P.O. Box 19185, 1000 GD Amsterdam, The Netherlands. You will be contacted as soon as possible.
UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

STATIC AND DYNAMIC PREDICTORS OF GENERAL AND VIOLENT CRIMINAL OFFENSE RECIDIVISM IN THE FORENSIC OUTPATIENT POPULATION

A Meta-Analysis

MARA J. EISENBERG

JOAN E. VAN HORN

JUDITH M. DEKKER

de Waag

MARK ASSINK

CLAUDIA E. VAN DER PUT 

JAN HENDRIKS

GEERT JAN J. M. STAMS

University of Amsterdam

Several meta-analyses have examined the Central Eight risk factors for reoffending in the general forensic population. To our knowledge, this is the first meta-analysis to focus exclusively on the predictive validity of risk factors for recidivism in forensic outpatients. A multilevel meta-analysis was conducted of studies in which static and dynamic risk factors were investigated as predictors of violent and/or general recidivism in forensic outpatients. Twenty-seven studies were included, with 543 effect sizes in a unique population of 116,982 adult offenders. The Central Eight risk domains were found to be predictive of violent and general recidivism; however, these factors predicted recidivism with small-to-moderate effects. Overall, in the same domain, the dynamic risk factors were more strongly related to recidivism than the static risk factors. This knowledge may be used to guide future outpatient treatment.

Keywords: forensic outpatient; risk factor; recidivism; meta-analysis

INTRODUCTION

Since several studies have demonstrated the limited rehabilitative effect of prison sentences, forensic outpatient treatment has become increasingly important (Mears, Cochran,

AUTHORS' NOTE: *This work was partly funded by Dutch Kwaliteit Forensische Zorg (KFZ Call 2015-39) [Quality Forensic Care] of the Ministry of Security and Justice, The Netherlands. The Ministry of Justice had no involvement in decisions concerning the study design, data collection, analysis, and interpretation of the data, nor in the writing and publication of the report. J. M. Dekker is now at Pameijer, Rotterdam. Correspondence concerning this article should be addressed to Mara J. Eisenberg, Research Department, de Waag, P.O. Box 1362, 3500 BJ Utrecht, The Netherlands; e-mail: meisenberg@dfzs.nl.*

CRIMINAL JUSTICE AND BEHAVIOR, 2019, Vol. 46, No. 5, May 2019, 732–750.

DOI: 10.1177/0093854819826109

Article reuse guidelines: sagepub.com/journals-permissions

© 2019 International Association for Correctional and Forensic Psychology

& Bales, 2012; Nagin, Cullen, & Jonson, 2009). Our knowledge of effective intervention to reduce delinquent behavior is mainly based on research on mixed populations (inpatients, prisoners, and offenders under varying degrees of probation supervision). To optimize the effect of treatment of the growing group of outpatients (Bales, Burkes, Scaggs, & Clark, 2015), it is important to enhance our understanding of which risk and criminogenic factors should be targeted in treatment to reduce reoffending. The aim of the present study was, therefore, to meta-analytically examine the most important risk and needs factors for forensic outpatients. However, the term outpatient is defined in a number of different ways across various studies. For instance, offenders in some studies were labeled as forensic outpatients when they were solely under intensive community supervision, whereas in other studies forensic outpatients received varying interventions, including drug treatment, cognitive-behavioral therapy, or family treatment (Lawrence & Lyons, 2011; Patenaude, 2013; Ploeg, 2008; Subramanian & Shames, 2013). Moreover, the term “parolees” or “probationers” is also used to describe a population that can be interpreted as forensic outpatients in the sense that they have been offered the same interventions mentioned above. The common denominator seems to be that all of these patients live in the community. Hence, in this article, the term “forensic outpatients” will be used to describe the population of offenders who have received community supervision and/or any treatment intervention while living in the community.

In the last decade, the central focus in offender treatment has been the risk–need–responsivity (RNR) principles, as these principles prescribe the basics of effective outpatient forensic care. The RNR model is an offender rehabilitation model developed by Bonta and Andrews (2017). In this model, the Risk principle states that high-risk offenders should be treated more frequently and for longer than low-risk offenders. The Need principle describes the focus of the treatment, which should target dynamic (changeable) risk factors, while the third principle (Responsivity) describes the required qualities of the therapist (general responsivity) and the tailoring of the treatment to the specific (dis)abilities of the patient (specific responsivity). Bonta and Andrews (2017) studied the effect of adhering these principles to forensic treatment on recidivism, concluding that recidivism rates decreased as the number of principles applied in the treatment protocol increased. Adherence to all three RNR principles had a greater rehabilitative effect for outpatients compared with clinical patients ($r = .35$ for community-based treatment and $r = .17$ for custody; Bonta & Andrews, 2017).

To adequately implement the risk and needs principles, knowledge of the most predictive static and dynamic criminogenic risk factors for recidivism is necessary. Static risk factors are features of the offenders’ histories that predict recidivism but are not amenable to intervention, such as prior offenses. By contrast, dynamic risk factors are potentially changeable factors, such as Substance Abuse and Criminal Network. Given that dynamic risk factors are considered to be responsible for the increase in risk, they have also been referred to as criminogenic needs (Bonta & Andrews, 2017). Five (meta-analytic) studies of predictors of recidivism among general offenders (Olver, Stockdale, & Wormith, 2014), youthful offenders (Grieger & Hosser, 2014), mentally disordered offenders (Bonta, Blais, & Wilson, 2014), racial minorities (Gutierrez, Wilson, Rugge, & Bonta, 2013), and drug offenders (Wooditch, Tang, & Taxman, 2014) found support for the eight strongest static and dynamic predictors of recidivism, also referred to as the Central Eight risk/needs factors. In more detail, the Central Eight risk/needs factors are as follows: a History of Antisocial Behavior

(early involvement in criminal activity, a large number and variety of prior offenses, rule violations); Antisocial Personality Traits (impulsive or aggressive behavior, callous disregard for others); Antisocial Cognitions (attitudes, values, and rationalizations condoning antisocial behavior); Antisocial Friends (social support from friends with antisocial attitudes and/or who are involved in criminal activity); Family/Relationships (lack of social support of family members and/or a turbulent and violent relationship with a romantic partner); School/Work (quality of interpersonal relationships at school or work and low performance and involvement); Leisure (low levels of involvement and satisfaction in anticriminal leisure pursuits); and Substance abuse (problems with alcohol and/or drugs; Bonta & Andrews, 2017).

In other meta-analytic studies, moderator effects were found. Specifically, Substance Abuse and Personal/Emotional Problems were more predictive of recidivism in women than in men (Olver et al., 2014), while lack of Leisure was a better predictor of recidivism among ethnic/racial majority offenders than among offenders from ethnic/racial minority groups (Gutierrez et al., 2013; Olver et al., 2014; Wilson & Gutierrez, 2014). The offender's age was also found to significantly moderate the relationship between the predictive value of risk factors and recidivism, that is, as age increases, the predictive effect of risk factors decreases (van der Put, Deković, Stams, Hoeve, & van der Laan, 2012). Interestingly, Spruit, van der Put, Gubbels, and Bindels (2017) found a U-shaped relationship between risk factors and recidivism across age, with an increased effect among offenders below the age of 40 and a decreased effect among offenders aged 40 years and over. Given the relevance of gender, race/ethnicity, and age to the relationship between risk factors and recidivism, these variables were included in the current meta-analysis as moderators.

AIM OF THE CURRENT META-ANALYSIS

The central aim of this multilevel meta-analysis was to identify the risk and needs factors for the prediction of general and violent recidivism among forensic outpatients.

In several ways, this study adds to the existing body of forensic knowledge in general and to findings from previous meta-analyses of forensic populations in particular. First, it focuses on a more homogeneous forensic population (i.e., outpatients) instead of the more general forensic population of in- and outpatients (Olver et al., 2014), or a subgroup of severely mentally disordered offenders who are typically treated in inpatient forensic clinics (see, for example, Bonta et al., 2014). A more homogeneous forensic outpatient sample was achieved through excluding studies of former inpatients or ex-detainees. The difference between inpatients and outpatients emerges in their offense history and recidivism rates. Compared with forensic outpatients, forensic inpatients have committed more severe violent offenses, often leading to physical injuries, and suffer more from major mental disorders such as psychotic and personality disorders (Ministry of Security and Justice, 2017). Although outpatients also suffer from personality disorders, the prevalence of psychotic disorders is much lower, while the incidence of impulse control and substance abuse disorders is much higher (Henrichs, Bogaerts, Sijtsma, & Klerx-van Mierlo, 2015; Spruit et al., 2017). This could explain why the recidivism rates reported in previous meta-analyses (in which the proportion of inpatients was relatively large) vary widely from 39% to 55.5% for general recidivism and 13.7% to 31.6% for violent recidivism, with the lowest recidivism percentages being found among general offenders and the highest among ethnic/racial minorities (Gutierrez et al., 2013; Olver et al., 2014).

On average, studies of outpatient populations report recidivism rates closer to the lower end of the inpatient range noted in the abovementioned studies. For example, a study by Rhodes, Dyou, Kling, Hunt, and Luallen (2013) found an overall recidivism percentage of 24.9% among offenders on federal community supervision. Accordingly, the current meta-analysis may shed some light on the differences in recidivism rates when focusing solely on an outpatient population. Second, the outcomes of the present study will contribute to more meaningful practices in terms of the applicability of the RNR principles to outpatient forensic care. Moreover, by including gender, race/ethnicity, and age as moderators in this specific group of offenders, some light will be shed on their relevance in terms of responsivity, and thus on how the treatment response may be optimized. Third, a distinction is made between the static and dynamic nature of risk factors within specific risk domains. Previous studies have shown the importance of the changes in the dynamic risk factors as a means of monitoring treatment progress (Lewis, Olver, & Wong, 2012), and findings have suggested a relation between positive changes in the dynamic risk factors and lowered recidivism rates (De Vries Robbé, De Vogel, Douglas, & Nijman, 2014). However, the findings from these studies are of limited use for the formulation of specific treatment goals in clinical practice as they represent an aggregated score of dynamic risk factors (total dynamic risk score and total static risk score). More specifically, the outcomes tell us about the importance of change in dynamic risk factors and recidivism, but little about which specific dynamic risk factors bring about changes in recidivism and should thus be targeted in treatment. Hence, the comparison of unique static and dynamic risk factor scores will provide more useful information on their relative importance for the prediction of recidivism and formulation of specific treatment targets.

In line with previous (meta-analytic) studies on the predictors of recidivism (Bonta et al., 2014; Grieger & Hosser, 2014; Gutierrez et al., 2013; Olver et al., 2014; Wooditch et al., 2014), it was expected that the same risk factors for general recidivism and violent recidivism would emerge for outpatients as for the general forensic population, as outpatients are a part of this general population. However, the strength of the predictive value could differ due to the fact that effect sizes are influenced by recidivism base rates. When base rates deviate from 50%—for instance, in the case of low-base-rate events such as sexual recidivism, where base rates are often below 10%—effect sizes remain small, with $r < .20$ (Babchishin & Helmus, 2016). In general, outpatients recidivate at a lower rate than inpatients. Hence, smaller effect sizes were expected for predictors of general and violent recidivism among outpatients.

METHOD

STUDY INCLUSION CRITERIA

Several criteria were used to govern the inclusion of primary studies. First, these studies needed to examine offenders who were 18 years of age or older, male or female, not incarcerated, and either treated in a forensic outpatient treatment setting or supervised by a probation officer in the community. Studies were included when incarceration preceded outpatient treatment and/or community supervision. We further noted that there were some differences in the definitions of “outpatient” and “treatment,” as well as in related terminology, between the countries in which the primary studies were conducted; only studies in which community sentencing was operationalized as an imposed outpatient/community-based treatment (such

as a psychological or addiction treatment, probation, or supervision) were chosen for inclusion. In this operationalization, treatment length and/or movement restrictions could vary between primary studies, as community sentences are imposed by court judges working in different jurisdictions. In all samples of offenders, the offenders were allowed to have jobs, to live in a private home (i.e., not institutionalized), and to move freely without supervision. Offenders in the samples of primary studies could have been imprisoned at some point in their past or could have received treatment for mental illness. However, studies were excluded when the patients had been released from forensic inpatient treatment or when offenders had been released from prison without outpatient treatment or supervision.

Second, the included studies had to define recidivism as a new arrest, reconviction, or violation of judicial conditions. Third, the studies needed to examine at least one static or dynamic risk factor for recidivism of any offense type. As mentioned in the "Introduction" section, static risk factors are features of the offenders' histories that predict recidivism, but that are not amenable to intervention; by contrast, dynamic risk factors (such as having criminal friends) are changeable and can be targeted in interventions. Moreover, studies examining risk factors for child sexual abuse (and not other types of offenses) were excluded, as risk factors for sex offending differ from other types of offending behavior (Mann, Hanson, & Thornton, 2010). An additional reason to exclude risk factors for sexual offending is that these factors are not incorporated in the Central Eight risk factors proposed by Bonta and Andrews (2017), which form the focus of the current meta-analysis.

Risk factors could not be included when statistical information needed for calculating effect sizes was missing. Hazard ratios were not included, as these ratios refer to differences between groups in time until an event occurs, which cannot be transformed into a correlation. Fourth, only quantitative research was included, except for studies reporting on results of survival analyses, responsivity factors (e.g., treatment motivation), and predictors of sexual recidivism. Studies focusing on the validation of a risk assessment instrument were only included when associations between individual items and recidivism were reported. Finally, we only included studies written in English, Dutch, or German.

LITERATURE SEARCH

In our search for relevant primary studies, we applied three strategies (see Figure 1). First, all electronic databases ($n = 413$) accessible through the library of the University of Amsterdam were searched (the complete list of databases is available upon request). The most important databases we exhaustively searched were the following: ERIC (Education Resources Information Center), Pubmed, Web of Science, Medline, Cochrane Library, PsycINFO, PiCarta, JSTOR (Journal Storage), and Hein Online. In this electronic search, the syntax component "forensic outpatient" OR "forensic community based" OR "probation" OR "parole" OR "post release" AND "adult" OR "adult offender" OR "adult delinquent" was combined with the following syntax component: "risk factor" OR "predict*" OR "criminogenic need" OR "recidivis*" OR "re-offen*" OR "reoffen*" OR "repetition" OR "recurrence." This electronic search resulted in 15,482 hits consisting of primary studies covering a wide variety of topics and samples. To narrow down these hits to studies that were eligible for inclusion, the first and third author proceeded with reading titles, abstracts, and (if necessary) full article texts. Of the 15,482 primary studies that were initially found, 2,563 studies appeared to be relevant after following deduplication and screening of the




Figure 1: Predictors of Recidivism in Forensic Outpatients Meta-Analysis Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Flow Diagram

Source: Adapted from Moher, Liberati, Tetzlaff, Altman, and The PRISMA Group (2009).

titles. After reading the study abstracts, 2,097 studies were disregarded, leading to a sample of 466 primary studies. The full texts of these studies were then evaluated against all inclusion criteria. In the end, eight studies were included in the qualitative synthesis and 27 primary studies were included in the meta-analysis.

In our second search strategy, we manually screened the reference lists of the studies that were eligible for inclusion in the previous step. We also examined reference lists from other studies that are relevant in the context of the present review, such as Greiner, Law, and Brown (2015), Ostermann (2015), and Wikoff, Linhorst, and Morani (2012). In our final search strategy, we contacted several experts in risk factors for recidivism and requested published and unpublished studies that would be eligible for inclusion.

CODING OF STUDIES

To code the studies, we developed an electronic coding form. All static and dynamic risk factors for general and violent recidivism that could be retrieved from the included studies were clustered into various “risk domains,” which are groups of risk factors that are (more or less) similar in nature. A total of 16 risk domains were created for general recidivism, whereas 11 risk domains were created for violent recidivism (see Table 1). In the analyses, the risk domains were divided into dynamic and static risk factors and analyzed separately. The dynamic risk factors are as described in Table 1, whereas the static risk factors mainly refer to historical events in a particular domain. For descriptive purposes, information was collected on the following: country where the study was conducted, type of index offense, average age of the sample, recidivism outcome measure, gender (percentage of males in samples), percentage of mentally ill offenders, and race/ethnicity (percentage of subjects belonging to the racial/ethnic majority). The measure of recidivism outcome was also included in the present study, as Andersen and Skardhamar (2017) concluded that the difference in the occurrence of rearrests, reconviction, or reimprisonment has an effect on the comparability of studies. Rearrests occur more often than reconviction, which can result in a power difference that affects the strength of effect sizes. Therefore, recidivism outcome measure, gender, and race/ethnicity were tested as potential moderators of the effect of each risk domain. The percentage of mentally ill offenders in samples could not be tested as a potential moderator because only three studies provided this information. After both the first and third author coded seven of the 27 studies independently, an overall interrater reliability was calculated for each included study. This reliability ranged from *good* ($\kappa = .80$) to *very good* ($\kappa = 1$) according to the criteria put forward by Landis and Koch (1977).

STATISTICAL ANALYSES

Several formulas were used to transform the reported effect sizes (Cohen’s d , odds ratios, and area under the normal curve [AUC]) to Pearson’s correlation coefficients (r). The Rosenthal (1994) formula was used to transform Cohen’s d to r , whereas the Borenstein, Hedges, Higgins, and Rothstein (2009) formula was used to transform the odds ratio to Cohen’s d and the Ruscio (2008) formula was applied to transform AUC to r . Each effect size represented the relationship between a risk factor and recidivism. Effect sizes were transformed into z scores and checked for outliers to control for any disproportionate influence of outliers on the results. The standardized effect sizes all fell within the range of $-3.29 < z > 3.29$ (Tabachnik & Fidell, 2013), implying that no outliers were identified.

Most studies reported multiple effect sizes that could be classified under a single risk domain. Extracting more than one effect size from a single study violates an important assumption of meta-analysis, namely that all reported effect sizes should be independent. To deal with the dependency of effect sizes, a three-level random-effects meta-analytic model was used to analyze the combined effect sizes in each risk domain (Assink & Wibbelink, 2016). In this model, three levels of variance were accounted for: sampling variance of the observed effect sizes (Level 1), variance of effect sizes within studies (Level 2), and variance of effect sizes between studies (Level 3).

For the analyses, the “*rma.mv*” function of the “*metafor*” package (Viechtbauer, 2010) was used in the program R Studio (Version 3.2.2; R Core Team, 2015). The data set was built in SPSS version 20. For each risk domain, an overall effect was estimated in an

TABLE 1: Risk Domains and Examples of Static and/or Dynamic Risk Factors Classified in Each Domain

Risk domain	Static and/or dynamic	Examples of included risk factors
1. Criminal History	Static	Criminality in adulthood, early antisocial behavior, escape from prison/clinic, having spent time in prison, history of property offenses/sex offenses/violent behavior, previous violations of judicial conditions, adjustment problems in prison/clinic, previous convictions/offenses
2. Antisocial Pattern	Static	Antisocial pattern of behavior
3. Antisocial Attitude	Dynamic	Current general acceptance of criminal behavior
4. Criminal Friends	Dynamic	Procriminal attitude/orientation, lack of concern for others, antisocial scale score, abusive attitude
5. Substance Abuse	Static	Social influences toward criminal behavior, (general social) rejection/loneliness, antisocial companions, criminal friends, abusive friends
6. Education/Employment	Dynamic	History of alcohol/drug problems, (severe) drug abuse
7. Family/Partner	Static	Current alcohol/drug problems, (severe) drug abuse
8. Personal/Psychological Problems	Dynamic	History of dysfunctional behavior during education and/or work
9. Living Situation	Static	Current dysfunctional behavior during education and/or work
10. Treatment	Dynamic	History of: Marital or relational problems, lack of (family) support, marital dissatisfaction, family problems, relational instability, nonsupporting partner, lack of social support
11. Financial Problems	Static	History of: Personal problems, subjective well-being, ^a (un)fulfillment, stress, anxiety/depression symptoms, psychosis, social/mental health, emotional instability, emotional regulation, mental health problems, personality type: aggressive/dependent/neurotic
12. Leisure	Dynamic	Immigration and history of: Subjective well-being, ^a living environment, accommodation, residence, effects of neighborhood: disadvantaged neighborhood/stability of residence
13. Age at Risk	Static	Negative attitude toward treatment (in the past)
14. Gender	Dynamic	Noncooperation with treatment
15. Race/Ethnicity	Static	Subjective well-being ^a concerning finances, financial difficulties, income >US\$10,000
16. Self-esteem	Dynamic	Lack of prosocial leisure/recreation, subjective well-being, ^a leisure, and lack of social participation
	Static	Age at risk of recidivism
	Static	Being male
	Static	Belonging to the ethnic/racial majority
	Dynamic	Lack of self-esteem, self-confidence, self-efficacy or belief in ability to reach life goals

^aSubjective well-being refers to an instrument to measure subjective well-being reported by the offender over several domains used in the study to predict recidivism.

intercept-only model. Two one-sided log-likelihood ratio tests were performed to determine whether the variance of effect sizes within studies (Level 2) and between studies (Level 3) was significant. The Level 1 variance was treated as known and calculated using the

formula of Cheung (2014, p. 2015). In the event of significant variance within a risk domain, either at Level 2 or at Level 3, we proceeded with testing variables as potential moderators of the overall effect of a risk domain. For these moderator analyses, the intercept-only models were extended by including potential moderators (i.e., recidivism outcome measure, gender, and race/ethnicity) as covariates. In the moderator analysis, all categories of moderators were tested simultaneously using an omnibus test (F test). When the F test was significant, the regression coefficient (slope) of each category was tested via t test to determine the category for which the moderating effect was found. In the case of a continuous variable (moderator), the significance of the omnibus test is equal to the significance of the t test. In the case of a nonsignificant omnibus test (nonmoderating effect), it was not expected that the unexplained Level 2 and/or Level 3 variance would substantially decrease.

BIAS AND SENSITIVITY ANALYSES

To examine whether bias (such as publication bias; Rothstein, Sutton, & Borenstein, 2005) influenced our results, we applied the trim-and-fill method (Duval & Tweedie, 2000) using the “trimfill” function of the “metafor” package (Viechtbauer, 2010) in the program R Studio (Version 3.2.2; R Core Team, 2015). In this method, effect sizes were plotted against the standard error, and in the event that the “funnel” was asymmetric, outlying effect sizes were recoded to fit the normal distribution of effect sizes, while “missing” effect sizes were imputed to restore the symmetry of the funnel. Next, we added the “missing” effect sizes to the data set to reestimate the overall effect of each risk domain for which an asymmetric funnel plot was obtained.

RESULTS

Twenty-seven studies were included, reporting on a total of 543 effect sizes and in which a total of 116,982 (N) unique offenders participated. Six study samples comprised only female offenders ($n = 2,782$, 2.4%), eight study samples comprised only male offenders ($n = 5,583$, 4.8%), and 13 study samples comprised both male and female offenders, of which 86.4% ($n = 93,852$) were men. The average age of the participants was 33.5 years ($SD = 3.8$, range = 20.5–40.0). Only three studies ($n = 530$) provided information on the mental illnesses of the offenders; therefore, the mental illnesses of participants were not tested in the moderator analyses.

Sixteen studies were conducted in the United States, seven in Canada, two in Western European countries, and two in Australia. In 16 studies, forensic outpatient treatment consisted of supervision (combined with treatment) by probationary institutes, while offenders received forensic community-based/outpatient treatment in the remaining 11 studies. The offenders received forensic outpatient treatment for any form of delinquent behavior, except sexual offending behavior, as studies of sex offenders were excluded from the present analysis. The average general recidivism rate was 35.0% ($SD = 13.7$, range = 16.6–62.4), while the average violent recidivism rate was 17.1% ($SD = 3.1$, range = 12.6–20.3). The average follow-up time was 3.8 years ($SD = 2.7$, range = 0.7–10.2). Average effect sizes are reported in terms of r . The criteria for interpreting these effect sizes, given a 50% base rate, are $\leq .10$ for a small effect, $\leq .24$ for a medium effect, and $\leq .37$ for a large effect (Rice & Harris, 2005). For base rates other than 50%, these criteria for small, medium, and large effects can

be calculated using the conversion formulae (after Rosenthal, 1994) provided by Rice and Harris (2005). Using these formulae, the criteria for small, medium, and large effect sizes for a 35.0% base rate for general recidivism are .095, .232, and .357, respectively. For a 17.1% base rate for violent recidivism, these criteria are .075, .185, and .288, respectively.

RISK DOMAINS FOR GENERAL RECIDIVISM

The average effect sizes of the 13 risk domains for general recidivism were significant, implying that these risk domains were positively associated with general recidivism (see Table 2). An Antisocial Pattern of behavior correlated most strongly with general recidivism ($r = .324$) and showed a medium effect size. Ten out of the remaining 13 risk domains were significantly associated with general recidivism and yielded an effect that was small in size. The domain of race/ethnicity (i.e., belonging to an ethnic/racial majority) yielded a significant but very small effect size, with $r = .077$. When domains contained both static and dynamic risk factors and correlated significantly with general recidivism, only the dynamic risk factors contributed to a significant correlation with general recidivism. Only static risk factors in the Criminal History domain correlated significantly and moderately with general recidivism ($r = .202$). Table 2 provides an overview of the variance of the effect sizes within the same study (Level 2 variance) and the variance of the effect sizes between studies (Level 3 variance). When significant Level 2 and/or Level 3 variance was found within a risk domain, different variables may moderate the association between the risk domain and general recidivism.

RISK DOMAINS FOR VIOLENT RECIDIVISM

For violent recidivism, 11 out of the 16 risk domains were tested in the study (see Table 3). We found significant effects for the following risk domains: Criminal History, Antisocial Pattern, Antisocial Attitudes, Criminal Friends, Substance Abuse, Education/Employment, Family/Partner, Personal/Psychological Problems, Living Environment, and Leisure. We found that all risk domains except Financial Problems correlated significantly with violent recidivism, with the strongest effect being found for Antisocial Pattern ($r = .27$). Using the adjusted interpretation guidelines, we found a medium effect for the following risk domains on violent recidivism: Antisocial Pattern, Living Environment, Criminal History, and Education/Employment. We found a small effect for the following risk domains: Antisocial Attitudes, Criminal Friends, and Substance Abuse. For the Family/Partner domain, we found only a very small effect.

MODERATORS FOR GENERAL RECIDIVISM

As shown in the Supplemental Material (available in the online version of this article), gender, race/ethnicity, and type of recidivism outcome (rearrest or reconviction) were tested as moderators of the effect of risk domains on general recidivism. Gender seemed to moderate the effect of static risk factors in the Family/Partner and Substance Abuse domains, as well as the effect of dynamic risk factors in the Self-esteem domain. Race/ethnicity moderated only the effect of the Treatment domain. As the percentage of the ethnic/racial majority increased in samples, the effect of the Treatment domain increased. However, after a Bonferroni adjustment ($p < .002$), none of the abovementioned moderating effects were upheld.

TABLE 2: Overall Effect Sizes (Correlations) for the Risk Domains for General Recidivism

Domain	S/D	No. of studies	No. of ES	Mean <i>r</i> (SE)	95% CI	Sig. Mean <i>r</i>	% var.			% var. Level 3	
							Level 1	Level 2	Level 3		
1. Criminal History	S	18	57	.202 (.045)	[-.112, .292]	<.001	0.42	.142***	43.36	.162***	56.22
2. Antisocial Pattern	SD	8	12	.324 (.089)	[-.129, .520]	.004	0.27	.084***	11.18	.236***	88.54
Antisocial Pattern	S	3	4	.530 (.207)	[-.128, 1.19]	.083	1.77	.079	4.81	.346	93.42
Antisocial Pattern	D	5	8	.221 (.048)	[-.108, .333]	.002	0.91	.086***	58.95	.071***	40.14
3. Antisocial Attitudes	D	10	23	.204 (.049)	[-.102, .306]	<.001	0.72	.058***	14.49	.141***	84.80
4. Criminal Friends	D	11	24	.221 (.061)	[-.094, .347]	.001	0.49	.063***	9.65	.191***	89.86
5. Substance Abuse	SD	17	34	.193 (.029)	[-.134, .252]	<.001	1.40	.058***	25.18	.099***	73.42
Substance Abuse	S	3	6	.132 (.069)	[-.046, .310]	.115	3.83	.000	<0.000	.114	96.17
Substance Abuse	D	17	28	.198 (.029)	[-.139, .256]	<.001	1.37	.056***	24.81	.097***	73.82
6. Education/Employment	SD	20	45	.141 (.038)	[-.066, .217]	<.001	0.81	.114***	40.33	.138***	58.86
Education/Employment	S	11	18	.003 (.029)	[-.059, .065]	.915	7.61	.044	21.82	.078	70.58
Education/Employment	D	15	27	.211 (.051)	[-.106, .317]	<.001	0.50	.078***	15.93	.180***	83.56
7. Family/Partner	SD	20	60	.100 (.022)	[-.055, .145]	<.001	2.34	.098***	69.28	.063***	28.38
Family/Partner	S	4	13	.062 (.043)	[-.032, .156]	.177	4.39	.062**	41.09	.072**	54.53
Family/Partner	D	18	47	.105 (.024)	[-.056, .153]	<.001	2.14	.109***	77.72	.055***	20.14
8. Personal/Psychological Problems	SD	16	38	.119 (.027)	[-.064, .174]	<.001	1.66	.077***	46.72	.081***	51.63
Personal/Psychological Problems	D	16	34	.120 (.029)	[-.061, .179]	<.001	1.37	.076***	41.22	.090***	57.41
9. Living Environment	SD	8	20	.142 (.047)	[-.042, .241]	.008	1.41	.041***	9.71	.123***	88.89
Living Environment	D	6	16	.188 (.047)	[-.088, .288]	.001	1.92	.048***	17.84	.101***	80.24
10. Treatment	SD	8	15	.197 (.142)	[-.108, .501]	.188	1.70	.102***	6.38	.387***	91.92
Treatment	S	5	7	.301 (.224)	[-.247, .849]	.228	.002	.000	<0.000	.498	99.99
Treatment	D	4	8	.089 (.051)	[-.031, .209]	.122	21.66	.125***	78.34	.000***	<0.000
11. Financial Problems	D	8	16	.174 (.024)	[-.124, .225]	<.001	12.12	.013	7.32	.044	80.57
12. Leisure	D	10	22	.161 (.027)	[-.104, .217]	<.001	2.42	.045***	30.62	.067***	66.97
13. Age at Risk	S	14	23	-.008 (.024)	[-.058, .041]	.724	3.32	.077***	71.92	.045***	24.76
14. Gender	S	5	5	.089 (.036)	[-.010, .188]	.068	6.82	.051	46.59	.051	46.59
15. Race/Ethnicity	S	9	13	.077 (.017)	[-.039, .114]	<.001	12.43	.039***	58.31	.027***	29.25
16. Self-esteem	D	4	9	-.081 (.054)	[-.206, .044]	.174	40.11	.000	<0.000	.093	59.90

Note. S/D = static risk factors and/or dynamic risk factors; ES = effect sizes; Mean *r* = Mean Fisher's *z*; CI = confidence interval; Sig. = significance; % var. = percentage of variance explained; Level 2 var. = variance between effect sizes from the same study; Level 3 var. = variance between studies. **p* < .05. ***p* < .01. ****p* < .001.

TABLE 3: Overall Effect Sizes (Correlations) for the Risk Domains for Violent Recidivism

Domain	S/D	No. of studies	No. of ES	<i>M</i> (<i>SD</i>)	95% CI	Sig. Mean <i>r</i>	% var. Level 1	Level 2 variance	% var. Level 2	Level 3 variance	% var. Level 3
1. Criminal History	S	5	14	.221 (.033)	[.149, .293]	<.001	1.28	.088***	81.34	.041***	17.37
2. Antisocial Pattern	SD	4	5	.271 (.061)	[.102, .439]	.011	1.08	.048*	17.38	.104*	81.54
3. Antisocial Attitudes	D	5	9	.169 (.049)	[.056, .282]	.009	2.16	.074*	43.30	.083*	54.55
4. Criminal Friends	D	5	8	.147 (.043)	[.045, .249]	.011	1.98	.105***	90.68	.030***	7.34
5. Substance Abuse	D	5	7	.141 (.037)	[.051, .232]	.009	2.66	.087***	97.34	.000***	<0.000
6. Education/ Employment	D	5	6	.207 (.022)	[.150, .264]	<.001	8.94	.042**	91.06	.000**	0
7. Family/Partner	SD	7	20	.058 (.027)	[.002, .114]	.045	5.37	.098***	92.33	.015***	2.29
Family/Partner	D	7	14	.082 (.032)	[.013, .150]	.023	3.56	.106***	96.44	.000***	<0.000
8. Personal/ Psychological Problems	D	6	9	.135 (.016)	[.097, .172]	<.001	9.32	.038***	90.68	.000***	<0.000
9. Living Environment	D	3	3	.236 (.035)	[.086, .386]	.021	100	.000	<0.000	.000	<0.000
11. Financial Problems	D	3	3	.127 (.069)	[-.171, .426]	.208	29.26	.070	35.37	.070	35.37
12. Leisure	D	6	8	.170 (.004)	[.160, .180]	<.001	100	.000	<0.000	.000	0

Note. S/D = static risk factors and/or dynamic risk factors; ES = effect sizes; Mean *r* = Mean Fisher's *z*; CI = confidence interval; Sig. = significance; % var. = percentage of variance explained; Level 2 var. = variance between effect sizes from the same study; Level 3 var. = variance between studies.

p* < .05. *p* < .01. ****p* < .001.

MODERATORS FOR VIOLENT RECIDIVISM

The moderator analyses showed that the effects of Antisocial Pattern, Antisocial Attitudes, having Criminal Friends, and lack of Education/Employment on violent recidivism were moderated by gender and race/ethnicity (see Supplemental Material). As before, after a Bonferroni adjustment ($p < .004$), none of the abovementioned moderating effects were upheld. Finally, type of recidivism outcome (rearrest or reconviction) did not moderate the effect of any of the examined risk domains for violent recidivism. There was no difference between general recidivism and violent recidivism in terms of the strength of the risk domains.

PUBLICATION BIAS

The results of the trim-and-fill analyses are presented in the Supplemental Material. After the trim-and-fill analyses were completed, the effect of most of the risk domains became smaller, but did not change in terms of relative strength (small-to-moderate effects were found after trim and fill). However, the effects of most risk domains for violent recidivism decreased. Only the effects of Criminal Friends, Substance Abuse, and Education/Employment domains increased. Again, the relative strength of the effect sizes (small effects) did not change.

DISCUSSION

The central aim of this multilevel meta-analytic study was to shed light on the relative strength (expressed in effect size) of risk and needs factors for the prediction of general and violent recidivism in an outpatient offender population. In general, the present findings were supportive of Bonta and Andrews's (2017) study in which the Central Eight risk factors were related to general and violent recidivism. Breaking down the Central Eight risk domains into

static and dynamic factors, the present study showed that two static risk domains, namely, Criminal History and Antisocial Pattern, were the strongest predictors of both general and violent recidivism. This finding corroborates the results of previous meta-analyses (Bonta et al., 2014; Collins, 2010; Olver et al., 2014). However, when risk domains included both static and dynamic risk factors, the dynamic risk factors were more strongly predictive of general and violent recidivism compared with the static risk factors. This was found for Substance Abuse, Education/Employment, Family/Partner, and Personal/Psychological Problems. As far as we know, this is the first study of forensic outpatients that directly compares specific static and dynamic risk factors within the same risk domain (e.g., history of substance abuse compared with current substance abuse). Previous studies targeting this issue have presented an additional predictive effect of the total score on dynamic risk factors on top of the total score on static risk factors in a subgroup of severely mentally ill offenders who had been released from mandatory inpatient treatment (De Vries Robbé et al., 2014), and despite small effects similar results were found for sexual offenders (Hanson, Helmus, & Harris, 2015; van den Berg et al., 2018). Moreover, the small effect sizes found in the meta-analysis by van den Berg and colleagues (2018), as well as in the present study, could have been the result of the fluctuating number of studies, relatively long follow-up time, and low base rates, as pointed out by Babchishin and Helmus (2016). An adjustment of the interpretation guidelines for recidivism base rates (Rice & Harris, 2005) yielded more medium effect sizes for the prediction of violent recidivism in the present study.

In the RNR model, the relevant risk domains found in the current study were labeled as dynamic, indicating that they describe the current situation and are changeable through intervention. Nonetheless, these factors become static in nature when they describe the offenders' situation in the past. From our results, it can be concluded that it is not the dynamic risk factors per se that should be focused on in research, but rather the changes in these factors. A number of studies underline this conclusion; for instance, Pelissier, Jones, and Cadigan (2007) showed that substance abuse treatment had a positive effect on sobriety after successful completion of treatment, resulting in lower recidivism rates. Based on this new knowledge and previous studies of treatment effects among forensic (out)patients, more attention should be paid to (changes in) dynamic risk factors (Douglas & Skeem, 2005; Kraemer et al., 1997). With this in mind, substance abuse is of special interest, mainly because of its high prevalence among offenders. A study of a Dutch forensic outpatient sample showed that the prevalence of substance-related disorders was 61.5% among general violence offenders and 30.9% among intimate violence offenders (Kraanen, Scholing, & Emmelkamp, 2012). Moreover, 29.9% of these offenders had been intoxicated while committing an offense. Another Dutch study of sex offenders pointed out that substance abuse was a consistent predictor of general recidivism across age groups (Wilpert, van Horn, & Boonmann, 2018), while forensic outpatients with mental illness and substance abuse issues were found to have the highest dropout and recidivism rates compared with offenders with only substance abuse problems or other mental illnesses (van Horn, Eisenberg, Souverein, & Kraanen, 2018). Despite the high prevalence of substance abuse among the forensic outpatient population, we found only a small effect of substance abuse, in line with the findings of Bonta and Andrews (2017).

Apart from research on treatment that focuses on substance abuse, very few studies on the relationship between changes in other dynamic risk factors and recidivism have been published. In a study by Vedel, Emmelkamp, and Schippers (2008), a decrease in intimate

partner violence was found in men who were continuing to abstain from substance abuse following cognitive-behavioral therapy. Howard and Dixon (2013) studied the relationship between changes in risk factors and general and violent recidivism among 196,493 English and Welsh offenders. They found that, after treatment, the total risk factor scores decreased for all offenders, although this decrease was larger among nonrecidivists. Specifically, they found the strongest relationship between changes in substance abuse and recidivism. In addition, changes in accommodation, employability, and (criminal) attitude were all significantly associated with recidivism. By contrast, changes in recognizing the impact of offending and psychiatric treatment were not associated with recidivism. Although it could be concluded from the latter finding that treatment does not effectively reduce recidivism, the item in question only refers to being in treatment and not to how actively an offender participates in treatment (e.g., motivation for treatment). Wooditch and colleagues (2014) did study the effect of being in treatment and found a risk-reducing effect of treatment that focused on changes in the dynamic risk factors, such as substance abuse, stable accommodation, employment, leisure, social support of prosocial friends, and establishing meaningful family and partner relationships.

MODERATORS

According to the responsivity principle of the RNR model (Bonta & Andrews, 2017), race/ethnicity and gender should be indicators used to accommodate treatment to the client's cognitive abilities and learning style (the responsivity principle) and should also be regarded as noncriminogenic factors. However, the present results do not confirm this assumption: Gender and race/ethnicity had either no or a negligibly significant predictive value for general and violent recidivism, and no moderator effects of gender, race/ethnicity, or recidivism outcome (rearrests or reconviction) were found.

The studies in the current meta-analysis included intelligence and age as possible predictors of recidivism, but did not provide usable information on both factors to facilitate their inclusion as possible moderators. In theory, adapting treatment methods to patients' characteristics (e.g., gender, race/ethnicity, intelligence, and age in outpatient facilities; Responsivity) should increase the effectiveness of outpatient treatment (Bonta & Andrews, 2017). However, researchers do not often include these factors as moderators in their studies. The current limited number of studies that include these moderators hinders the formulation of specific advice on responsivity in outpatient offenders as far as gender, race/ethnicity, intelligence, and age are concerned.

LIMITATIONS AND FUTURE RESEARCH

Publication bias caused by human error and the article acceptance practices of journals is a widely recognized problem in meta-analyses (Thornton & Lee, 2000). The protocolled searches and coding schemes used in this study minimized registration errors to the greatest possible extent. However, even when combined with the efforts of the authors to contact the authors of unpublished research, it is unlikely that all available research was included in this meta-analysis. Therefore, a trim-and-fill analysis was used to detect any effect of possible publication bias. The results showed minor changes in the strength of the effect sizes of the risk domains, from which it can be concluded that the results of this meta-analysis are a reliable representation of the results found until now. Despite this, the predictive validity of static and dynamic


risk factors in forensic outpatients is influenced by low base rates and the small number of available studies (Babchishin & Helmus, 2016; Rice & Harris, 2005). Although the results point toward the conclusion that the effect sizes of this study underline the predictive validity of the static and dynamic risk factors, it should be noted that this is the first study to address this issue in an exclusively outpatient offender population. Hence, the present results need to be replicated in future and should be interpreted carefully for the time being.

The key results of this meta-analysis confirm the predictive value of the Central Eight risk domains for general and violent recidivism among forensic outpatients. Forensic outpatient treatment should thus focus on the dynamic risk factors of the following domains: Antisocial Pattern and Attitudes, Substance Abuse, Criminal Friends, Education/Employment, Family/Partner, Personal/Psychological Problems, Living Situation, Leisure, and Financial Problems. Future research should explore the relationship between changes in these dynamic risk factors and recidivism, along with the possible moderating effects of age, race/ethnicity, gender, and intelligence. Moreover, new meta-analysis on predictors of recidivism should be performed on the forensic residential and specified forensic outpatient subpopulations, preferably also in countries outside of Northern America.

SUPPLEMENTAL MATERIAL

Supplemental Material is available in the online version of this article at <https://journals.sagepub.com/home/cjb>.

ORCID ID

C. E. van der Put  <https://orcid.org/0000-0002-0448-962X>

REFERENCES

References marked with an asterisk indicate studies included in the meta-analysis.

- *Abrams, L. S., Terry, D., & Franke, T. M. (2011). Community-based juvenile reentry services: The effects of service dosage on juvenile and adult recidivism. *Journal of Offender Rehabilitation, 50*, 492-510. doi:10.1080/10509674.2011.596919
- Andersen, S. N., & Skardhamar, T. (2017). Pick a number: Mapping recidivism measures and their consequences. *Crime & Delinquency, 63*, 613-635. doi:10.1177/0011128715570629
- Assink, M., & Wibbelink, C. J. M. (2016). Fitting three-level meta-analytic models in R: A step-by-step tutorial. *The Quantitative Methods for Psychology, 12*, 154-174. doi:10.20982/tqmp.12.3.p154
- Babchishin, K. M., & Helmus, L.-M. (2016). The influence of base rates on correlations: An evaluation of proposed alternative effect sizes with real-world data. *Behavior Research Methods, 48*, 1021-1031. doi:10.3758/s13428-015-0627-7
- Bales, W. D., Burkes, K. J., Scaggs, S. J. A., & Clark, C. L. (2015). Recidivism. In J. D. Wright (Ed.), *International encyclopedia of the social & behavioral sciences* (2nd ed., Vol. 20, pp. 31-36). doi:10.1016/B978-0-08-097086-8.45079-8
- Bonta, J. L., & Andrews, D. A. (2017). *The psychology of criminal conduct* (6th ed.). New York, NY: Routledge.
- Bonta, J. L., Blais, J., & Wilson, H. A. (2014). A theoretically informed meta-analysis of the risk for general and violent recidivism for mentally disordered offenders. *Aggression and Violent Behavior, 19*, 278-287. doi:10.1016/j.avb.2014.04.014
- Borenstein, M., Hedges, L. V., Higgins, J. P. T., & Rothstein, H. R. (2009). *Introduction to meta-analysis*. Chichester, UK: Wiley.
- *Bouman, Y. H. A., Schene, A. H., & De Ruiter, C. (2009). Subjective well-being and recidivism in forensic psychiatric outpatients. *International Journal of Forensic Mental Health, 8*, 225-234. doi:10.1080/14999011003635647
- *Canales, D. D., Campbell, M. A., Wei, R., & Totten, A. E. (2014). Prediction of general and violent recidivism among mentally disordered adult offenders: Test of the level of service/risk-need-responsivity (LS/RNR) instrument. *Criminal Justice and Behavior, 41*, 971-991. doi:10.1177/0093854814523003
- Cheung, M. W. L. (2014). Modeling dependent effect sizes with three-level meta-analyses: A structural equation modeling approach. *Psychological Methods, 19*, 211-229. doi:10.1037/a0032968
- Collins, R. E. (2010). The effect of gender on violent and nonviolent recidivism: A meta-analysis. *Journal of Criminal Justice, 38*, 675-684. doi:10.1016/j.jcrimjus.2010.04.041
- *Degiorgio, L., & DiDonato, M. (2014). Predicting probationer rates of reincarceration using dynamic factors from the Substance Abuse Questionnaire-Adult Probation III (SAQ-Adult Probation III). *American Journal of Criminal Justice, 39*, 94-108. doi:10.1007/s12103-013-9198-2

- De Vries Robbé, M., De Vogel, V., Douglas, K. S., & Nijman, H. L. I. (2015). Changes in dynamic risk and protective factors for violence during inpatient forensic psychiatric treatment: Predicting reductions in postdischarge community recidivism. *Law and Human Behavior, 39*, 53-61. doi:10.1037/lhb0000089
- Douglas, K. S., & Skeem, J. L. (2005). Violence risk assessment: Getting specific about being dynamic. *Psychology, Public Policy and Law, 11*, 347-383.
- Duval, S., & Tweedie, R. (2000). Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics Journal of the International Biometric Society, 56*, 455-463. doi:10.1111/j.0006-341X.2000.00455.x
- *Fisher, W. H., Hartwell, S. W., Deng, X., Pinals, D. A., Fulwiler, C., & Roy-Bujnowski, K. (2014). Recidivism among released state prison inmates who received mental health treatment while incarcerated. *Crime & Delinquency, 60*, 811-832. doi:10.1177/0011128714541204
- *Gordon, H., Kelty, S. F., & Julian, R. (2015). An evaluation of the Level of Service/Case Management Inventory in an Australian community corrections environment. *Psychiatry, Psychology and Law, 22*, 247-258. doi:10.1080/13218719.2014.941090
- Greiner, L. E., Law, M. A., & Brown, S. L. (2015). Using dynamic factors to predict recidivism among women: A four-wave prospective study. *Criminal Justice and Behavior, 42*, 457-480. doi:10.1177/0093854814553222
- Grieger, L., & Hosser, D. (2014). Which risk factors are really predictive? An analysis of Andrews and Bonta's "Central Eight" risk factors for recidivism in German youth correctional facility inmates. *Criminal Justice and Behavior, 41*, 613-634.
- Gutierrez, L., Wilson, H. A., Rugge, T., & Bonta, J. (2013). The prediction of recidivism with aboriginal offenders: A theoretically informed meta-analysis. *Canadian Journal of Criminology and Criminal Justice, 55*, 55-99. doi:10.3138/cjccj.2011.E.51
- Hanson, R. K., Helmus, L.-M., & Harris, A. J. R. (2015). Assessing the risk and needs of supervised sexual offenders. *Criminal Justice and Behavior, 42*, 1205-1224. doi:10.1177/0093854815602094
- *Hanson, R. K., & Wallace-Capretta, S. (2004). Predictors of criminal recidivism among male batterers. *Psychology, Crime & Law, 10*, 413-427. doi:10.1080/10683160310001629283
- *Henderson, H., & Miller, H. A. (2013). The (twice) failure of the Wisconsin Risk Need Assessment in a sample of probationers. *Criminal Justice Policy Review, 24*, 199-221. doi:10.1177/0887403411422410
- Henrichs, J., Bogaerts, S., Sijtsema, J., & Klerx-van Mierlo, F. (2015). Intimate partner violence perpetrators in a forensic psychiatric outpatient setting: Criminal history, psychopathology, and victimization. *Journal of Interpersonal Violence, 30*, 2109-2128. doi:10.1177/0886260514552272
- Howard, P. D., & Dixon, L. (2013). Identifying change in the likelihood of violent recidivism: Causal dynamic risk, factors in the OASys violence predictor. *Law and Human Behavior, 37*, 163-174. doi:10.1037/lhb0000012
- *Hsu, C.-I., Caputi, P., & Byrne, K. (2009). The Level of Service Inventory-revised (LSI-r): A useful risk assessment measure for Australian offenders? *Criminal Justice and Behavior, 36*, 728-740. doi:10.1177/0093854809335409
- *Huebner, B. M., & Berg, M. T. (2011). Examining the sources of variation in risk for recidivism. *Justice Quarterly, 28*, 146-173. doi:10.1080/07418820903365213
- *Huebner, B. M., DeJong, C., & Cobbina, J. (2010). Women coming home: Long-term patterns of recidivism. *Justice Quarterly, 27*, 225-254. doi:10.1080/07418820902870486
- Kraanen, F. L., Scholing, A., & Emmelkamp, P. M. G. (2012). Substance use disorders in forensic psychiatry: Differences among different types of offenders. *International Journal of Offender Therapy and Comparative Criminology, 56*, 1201-1219. doi:10.1177/0306624X11420252
- Kraemer, H. C., Kazdin, A. E., Offord, D. R., Kessler, R. C., Jensen, P. S., & Kupfer, D. J. (1997). Coming to terms with the terms of risk. *Archives of General Psychiatry, 54*, 337-343.
- *Kropp, P. R., & Hart, S. D. (2000). The Spousal Assault Risk Assessment (SARA) guide: Reliability and validity in adult male offenders. *Law and Human Behavior, 24*, 101-118. doi:10.1023/A:1005430904495
- Landis, J. R., & Koch, G. G. (1977). The measurement of observer agreement for categorical data. *Biometrics, 33*, 159-174. doi:10.2307/25293102529310843571
- Lawrence, A., & Lyons, D. (2011). *Principles of effective state sentencing and corrections policy. A report of the NCSL sentencing and corrections work group*. Washington, DC: National Conference of State Legislatures. Available from <http://www.ncsl.org>
- Lewis, K., Olver, M. E., & Wong, S. C. P. (2012). The Violence Risk Scale: Predictive validity and linking changes in risk with violent recidivism in a sample of high-risk offenders with psychopathic traits. *Assessment, 22*, 150-164. doi:10.1177/1073191112441242
- Mann, R. E., Hanson, R. K., & Thornton, D. (2010). Assessing risk for sexual recidivism: Some proposals on the nature of psychologically meaningful risk factors. *Sexual Abuse: A Journal of Research and Treatment, 22*, 191-217. doi:10.1177/1079063210366039
- Mears, D. P., Cochran, J. C., & Bales, W. D. (2012). Gender differences in the effects of prison on recidivism. *Journal of Criminal Justice, 40*, 370-378. doi:10.1016/j.jcrimjus.2012.06.009

- *Miller, H., & Miller, J. M. (2010). Community in-reach through jail reentry: Findings from a quasi-experimental design. *Justice Quarterly*, 27, 893-910. doi:10.1080/07418825.2010.482537
- Ministry of Security and Justice. (2017). *Forensische zorg in getal 2012-2016 [Forensic Care in number 2012-2016]*. Retrieved from <https://www.tbsnederland.nl>
- Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G. & The PRISMA Group. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Medicine*, 6(7). doi:10.1371/journal.pmed.1000097
- Nagin, D. S., Cullen, F. T., & Jonson, C. L. (2009). Imprisonment and reoffending. *Crime and Justice*, 38, 115-200. doi:10.1086/599202
- Olver, M. E., Stockdale, K. C., & Wormith, J. S. (2014). Thirty years of research on the Level of Service Scales: A meta-analytic examination of predictive accuracy and sources of variability. *Psychological Assessment*, 26, 156-176. doi:10.1037/a0035080
- Ostermann, M. (2015). How do former inmates perform in the community? A survival analysis of rearrests, reconvictions, and technical parole violations. *Crime & Delinquency*, 61, 163-187. doi:10.1177/0011128710396425
- Patenaude, A. L. (2013). Intermediate sanctions and community-based corrections. In J. Winterdyk & M. Weinrath (Eds.), *Adult corrections in Canada: A comprehensive overview* (pp. 225-248). Whitby, Ontario, Canada: de Sitter Publications.
- Pelissier, B., Jones, N., & Cadigan, T. (2007). Drug treatment aftercare in the criminal justice system: A systematic review. *Journal of Substance Abuse Treatment*, 32, 311-320. doi:10.1016/j.jsat.2006.09.007
- Ploeg, G. (2008). Norway. In A. Kalmthout & I. Durnescu (Eds.), *Probation in Europe* (pp. 503-512). Oisterwijk, The Netherlands: Wolf Legal Publishers. Available from <http://www.cep-probation.org>
- *Prendergast, M. L., Messina, N. P., Hall, E. A., & Warda, U. S. (2011). The relative effectiveness of women-only and mixed-gender treatment for substance-abusing women. *Journal of Substance Abuse Treatment*, 40, 336-348. doi:10.1016/j.jsat.2010.12.001
- R Core Team. (2015). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- *Reingle, J. M., Striley, C. W., Small, E., Crecelius, R., O'Leary, C. C., & Cottler, L. B. (2013). Can courtroom behavior predict recidivism? An assessment of the courtroom behavior check list for women presenting in drug court. *American Journal of Criminal Justice*, 38, 520-534. doi:10.1007/s12103-012-9190-2
- *Rettinger, L. J., & Andrews, D. A. (2010). General risk and need, gender specificity, and the recidivism of female offenders. *Criminal Justice and Behavior*, 37, 29-46. doi:10.1177/0093854809349438
- Rhodes, W., Dyou, C., Kling, R., Hunt, D., & Luallen, J. (2013). *Recidivism of offenders on federal community supervision*. Washington, DC: U.S. Department of Justice. Available from <https://www.ncjrs.gov>
- Rice, M. E., & Harris, G. T. (2005). Comparing effect sizes in follow-up studies: ROC area, Cohen's *d*, and *r*. *Law and Human Behavior*, 29, 615-620. doi:10.1007/s10979-005-6832-7
- Rosenthal, R. (1994). Parametric measures of effect size. In H. Cooper & L. V. Hedges (Eds.), *The handbook of research synthesis* (pp. 231-244). New York, NY: SAGE.
- Rothstein, H., Sutton, A. J., & Borenstein, M. (2005). *Publication bias in meta-analysis: Prevention, assessment and adjustments*. Chichester, UK: Wiley.
- Ruscio, J. (2008). A probability-based measure of effect size: Robustness to base rates and other factors. *Psychological Methods*, 13, 19-30. doi:10.1037/1082-989X.13.1.19
- *Salisbury, E. J., & Van Voorhis, P. (2009). Gendered pathways: A quantitative investigation of women probationers' paths to incarceration. *Criminal Justice and Behavior*, 36, 541-566. doi:10.1177/0093854809334076
- *Spjeldnes, S., Jung, H., Maguire, L., & Yamatani, H. (2012). Positive family social support: Counteracting negative effects of mental illness and substance abuse to reduce jail ex-inmate recidivism rates. *Journal of Human Behavior in the Social Environment*, 22, 130-147. doi:10.1080/10911359.2012.646846
- Spruit, A., Van der Put, C. E., Gubbels, J., & Bindels, A. (2017). Age differences in the severity, impact and relative importance of dynamic risk factors for recidivism. *Journal of Criminal Justice*, 50, 69-77.
- *Stalans, L. J., & Lurigio, A. J. (2015). Parenting and intimate relationship effects on women offenders' recidivism and non-compliance with probation. *Women & Criminal Justice*, 25, 152-168. doi:10.1080/08974454.2014.909764
- *Steiner, B., Makarios, M. D., Travis, L. F., III, & Meade, B. (2012). Examining the effects of community-based sanctions on offender recidivism. *Justice Quarterly*, 29, 229-257. doi:10.1080/07418825.2011.555413
- Subramanian, R., & Shames, A. (2013). *Sentencing and prison practices in Germany and the Netherlands: Implications for the United States*. New York, NY: Vera Institute of Justice.
- Tabachnik, B. G., & Fidell, L. S. (2013). *Using multivariate statistics* (6th ed.). Boston, MA: Allyn & Bacon.
- Thornton, A., & Lee, P. (2000). Publication bias in meta-analysis: Its causes and consequences. *Journal of Clinical Epidemiology*, 53, 207-216.
- *Tillyer, M. S., & Vose, B. (2011). Social ecology, individual risk, and recidivism: A multilevel examination of main and moderating influences. *Journal of Criminal Justice*, 39, 452-459. doi:10.1016/j.jcrimjus.2011.08.003

- van den Berg, J. W., Smid, W., Schepers, K., Wever, E., van Beek, D., Janssen, E., & Gijs, L. (2018). The predictive properties of dynamic sex offender risk assessment instruments: A meta-analysis. *Psychological Assessment, 30*, 179-191. doi:10.1037/pas0000454
- van der Put, C. E., Deković, M., Stams, G. J. J. M., Hoeve, M., & van der Laan, P. H. (2012). Het belang van vroegtijdig ingrijpen bij jeugdcriminaliteit [The importance of early intervening in youth crime]. *Kind en Adolescent, 33*(1), 2-20.
- van Horn, J. E., Eisenberg, M., Souverein, F. A., & Kraanen, F. (2018). The predictive value of the Central Eight criminogenic risk factors: A multi-group comparison of dually diagnosed violent offenders with other subgroups of violent offenders. *HSOA Journal of Addiction & Addictive Disorders, 5*, Article 014. doi:10.24966/AAD-7276/100014
- *VanVoorhis, P., Spiropoulos, G., Ritchie, P. N., Seabrook, R., & Spruance, L. (2013). Identifying areas of specific responsiveness in cognitive-behavioral treatment outcomes. *Criminal Justice and Behavior, 40*, 1250-1279. doi:10.1177/0093854813494182
- *Van Voorhis, P., Wright, E. M., Salisbury, E., & Bauman, A. (2010). Women's risk factors and their contributions to existing risk/needs assessment: The current status of a gender-responsive supplement. *Criminal Justice and Behavior, 37*, 261-288. doi:10.1177/0093854809357442
- Vedel, E., Emmelkamp, P. M. G., & Schippers, G. M. (2008). Individual cognitive-behavioral therapy and behavioral couples therapy in alcohol use disorder: A comparative evaluation in community-based addiction treatment centers. *Psychotherapy and Psychosomatics, 77*, 280-288. doi:10.1159/000140087
- Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software, 36*(3), 1-48.
- Wikoff, N., Linhorst, D. M., & Morani, N. (2012). Recidivism among participants of a reentry program for prisoners released without supervision. *Social Work Research, 36*, 289-299. doi:10.1093/swr/svs021
- Wilpert, J., van Horn, J. E., & Boonmann, C. (2018). Comparing the Central Eight risk factors: Do they differ across age groups of sex offenders? *International Journal of Offender Therapy and Comparative Criminology, 62*, 4278-4294. doi:10.1177/0306624X18758899
- Wilson, H. A., & Gutierrez, L. (2014). Does one size fit all? A meta-analysis examining the predictive ability of the Level of Service Inventory (LSI) with Aboriginal offenders. *Criminal Justice and Behavior, 41*, 196-219. doi:10.1177/0093854813500958
- Wooditch, A., Tang, L. L., & Taxman, F. S. (2014). Which criminogenic need changes are most important in promoting desistance from crime and substance use? *Criminal Justice and Behavior, 41*, 276-299. doi:10.1177/0093854813503543
- *Wormith, J. S., Hogg, S., & Guzzo, L. (2012). The predictive validity of a general risk/needs assessment inventory on sexual offender recidivism and an exploration of the professional override. *Criminal Justice and Behavior, 39*, 1511-1538. doi:10.1177/0093854812455741
- *Wormith, J. S., Hogg, S. M., & Guzzo, L. (2015). The predictive validity of the LS/CMI with Aboriginal offenders in Canada. *Criminal Justice and Behavior, 42*, 481-508. doi:10.1177/0093854814552843

Mara J. Eisenberg, MSc, is a researcher at de Waag, a Dutch forensic psychiatric outpatient treatment center and PhD student at the Research Institute of Child Development and Education of the University of Amsterdam. Her research is focused on risk factors and risk assessment in the forensic outpatient population.

Joan E. van Horn is head of the Research Department of de Waag. She initiates, conducts, and supervises (granted) scientific research projects aimed at enhancing the effectiveness and efficiency of diagnostics and treatment through which community safety and relapse prevention are targeted. Major fields of interest are eHealth/mHealth, risk assessment, recidivism, and treatment effectiveness.

Judith M. Dekker, MSc, is pedagogue at Pameijer where residential and outpatient care for adults with a mental and/or psychological disability is provided. She coaches patients in different areas of their lives.

Mark Assink, PhD, is a postdoctoral researcher at the University of Amsterdam, The Netherlands. His research is focused on developing and validating instruments for risk and needs assessment, examining risk factors for juvenile delinquency and different forms of child abuse, and examining the effectiveness of (preventive) psychosocial interventions. He has substantial expertise in meta-analytic research.

Claudia E. van der Put is an assistant professor at the Research Institute of Child Development and Education of the University of Amsterdam. Her research interest is risk assessment and prevention of adverse developmental outcomes such as juvenile delinquency and problematic child-rearing situations.

Jan Hendriks is a forensic and clinical psychologist. He is professor of forensic psychiatry and psychology at the VU University of Amsterdam, and professor of forensic orthopedagogics at the University of Amsterdam. His research interests are juvenile and adult sex offenders, female offenders, and interventions for delinquent youth.

Geert Jan J. M. Stams is a full professor of forensic child and youth care sciences at the University of Amsterdam (The Netherlands). His research interests are in a broad range of aspects relevant to forensic child and youth care, such as moral reasoning of juvenile delinquents, attachment between parents and juvenile delinquents, instrument development and validation, and effectiveness of (preventive) interventions for juvenile antisocial behavior and child maltreatment.