UvA-DARE (Digital Academic Repository)

Work ability related to job demands and functional limitations in various occupational populations

van Schaaik, A.

Publication date

2020

Document Version

Final published version

License

Other

[Link to publication](#)

Citation for published version (APA):

van Schaaik, A. (2020). *Work ability related to job demands and functional limitations in various occupational populations*. [Thesis, fully internal, Universiteit van Amsterdam].

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

*Work ability related to job demands
and functional limitations in various
occupational populations*

Art van Schaijk

**Work ability related to job demands and functional limitations in
various occupational populations**

Art van Schaijk

Colofon:

The research project Work ability related to job demands and functional limitations in various occupational populations was funded by the Amsterdam UMC Coronel Institute of Occupational health and Stichting Fonds Scholing en Ordening voor het besloten busvervoer (FSO) (2016EUSF20164).

The studies in this dissertation were carried out at Amsterdam UMC, University of Amsterdam, Department of Public and Occupational Health, Coronel Institute of Occupational Health, Amsterdam Public Health research institute, the Netherlands

Cover design: Art van Schaaijk

Layout: Art van Schaaijk

Printed by: Ipskamp Printing, proefschriften.net

ISBN: 978-94-6421-060-6

© Art van Schaaijk, 2020

All rights reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording or any information storage and retrieval system, without written permission of the author.

**Work ability related to job demands and functional limitations in various
occupational populations**

ACADEMISCH PROEFSCHRIFT

ter verkrijging van de graad van doctor

aan de Universiteit van Amsterdam

op gezag van de Rector Magnificus

prof. dr. ir. K.I.J. Maex

ten overstaan van een door het College voor Promoties ingestelde commissie,

in het openbaar te verdedigen in de Aula der Universiteit

op woensdag 4 november 2020, te 17.00 uur

door Art van Schaaijk

geboren te 's-Hertogenbosch.

PROMOTIECOMMISSIE

Promotor	Prof. dr. M.H.W. Frings-Dresen	AMC-UvA
Copromotor	Dr. K. Nieuwenhuijsen	AMC-UvA
Overige leden	Prof dr. U. Bültmann	Rijksuniversiteit Groningen
	Prof dr. N. van Dijk	AMC-UvA
	Prof dr. R.H.H. Engelbert	AMC-UvA
	Prof dr. A.E. Kunst	AMC-UvA
	Prof dr. W. van Rhenen	Nyenrode Business Universiteit

Faculteit der Geneeskunde

TABLE OF CONTENTS

Chapter 1	General introduction	7
Chapter 2	Reproducibility of work ability and work functioning instruments <i>Occupational Medicine, 2018, 68(2), 116-119.</i>	27
Chapter 3	Work ability and work functioning: measuring change in individuals recently returned to work. <i>International archives of occupational and environmental health, 2018, 92(3), 423-433.</i>	37
Chapter 4	Work ability and percentage of hours worked related to limitations in patients with upper extremity musculoskeletal disorders: a cross-sectional cohort study. <i>BMC Musculoskeletal Disorders, 2020;21(1), 389-401.</i>	59
Chapter 5	Mediating factors for the relationship between stress and work ability over time in young adults. <i>International Journal of Environmental Research and Public Health, 2020, 17(7), 2530.</i>	85
Chapter 6	Appraisal of work ability in relation to job-specific health requirements in ambulance workers. <i>International archives of occupational and environmental health, 2017, 90(1), 123-131.</i>	127
Chapter 7	Work Ability and Vitality in Coach Drivers: An RCT to Study the Effectiveness of a Self-Management Intervention during the Peak Season. <i>International Journal of Environmental Research and Public Health, 2019, 16(12), 2214.</i>	145
Chapter 8	General discussion	175
Chapter 9	Summary	199
Chapter 10	Samenvatting	207
Chapter 11	Appendices	215
I.	List of abbreviations	216
II.	Curriculum Vitae	218
III.	List of publications	219
IV.	Co-authors and their affiliations	221
V.	PhD Portfolio	222
VI.	Acknowledgments/Dankwoord	224

1

General introduction

IMPORTANCE OF WORK AND ITS RELATION TO HEALTH

A large part of adult life is spent working. The expected duration of working life in the Netherlands is more than 40 years, around half of life expectancy, with an increasing trend in the lengthening of the expected duration of working life [1, 2]. Therefore work is an important part of the quality of people's lives [3, 4]. Being able to work is a meaningful factor in people's wellbeing and self-esteem [5]. This knife cuts both ways. On the one side, working and earning an income gives people the feeling of being useful and contributing to society, which increases the quality of life [6]. On the other side, work can negatively influence health, decreasing the quality of life and resulting in work-related health complaints. Health complaints caused either at work or outside can in turn influence the ability to fulfil job demands (often also called work demands), which also depends on the individual's personal resources to work and recover from work strain [7]. When an imbalance occurs between job demands and a worker's resources, it can impact their overall ability to work [8]. Thus, the ability to work is a constant balance between an individual's resources and specific job demands, as each worker and each job position vary [9].

WORK ABILITY

The ability to perform work and meet job demands is called 'work ability,' a term introduced in 1981 by the Finnish Institute of Occupational Health. Work ability is the balance between job demands and the worker's resources, and this construct has been defined over the years by Ilmarinen et al. (1997) as 'the quality of workers at present and in the near future to be able to do their job satisfactorily with respect to job demands, health and mental resources' [10].

DETERMINANTS OF WORK ABILITY

Firstly, work ability depends on one's job. Different jobs have different demands, health requirements and resources, all of which constitute the work environment. In jobs with more physical demands, physical work ability is generally of greater importance than mental work ability, while in jobs with more mental demands, such as office workers, mental work ability is of greater importance [7]. Physical functional limitations will have a larger impact on the work ability of employees with greater physical job demands, than on the work ability of employees with greater mental job demands and vice versa. Job demands, in combination with health of the workers, have an impact on the work ability and work functioning of employees [11, 12].

In addition to work and health factors, many other aspects can influence the ability to work. To study the determinants of work ability, Ilmarinen conceptualized work ability in a model that describes known determinants of work and life linked directly to work

ability. As defined in this model, work ability is constantly changing because the value of each aspect is subject to change per individual. Ilmarinen visualized the constituents of work ability as being a house. During later amendments, the house included a balcony, garden, social and operational environment, and even digital (social) media influences (Figure 1) [13].

Figure 1: Dimensions of work ability, depicted in the Work Ability House model (latest version) [13].

According to Ilmarinen's work ability house, the base starts with sufficient physical and mental functional capacity, meaning that a worker's physical and mental health need to be satisfactory in order to perform work [7, 14]. This base has consequences on all other levels of the house. On the next level, skills and knowledge are important in how an individual is able to do his or her job. This floor of skills, knowledge, and the competencies needed to fulfil a job satisfactorily are based on the health and physical and mental functional capacities of the first floor. This floor is under constant renovation as individuals constantly update their knowledge and skills.

On the next floor, norms and values are represented. This level not only represents the norms and values of the working individual, but also the values of the company the individual works for. When these competencies are adequate, the attitude and motivation to work come into play.

The first three floors represent the worker's resources, and the top floor consists of job demands and other work aspects, which include the work environment, work content, job demands, and the organization. All these contribute to good work ability when managed in the right way. All other floors need to be in balance with this fourth floor to maintain sufficient work ability.

However, work ability can still be influenced from factors outside the house, including family, the immediate social environment, the external operational environment, and society. All these aspects influence the work ability of a person. In this dissertation, we focus on the relationship of health and job demands with work ability.

All aspects of the work ability house model are subject to change and depend highly on the worker's personal situation. Because each change has a different or distinct effect on work ability, it is important to map these changes in different populations. This change can happen on an individual level, e.g. functional limitations, but they also occur on a company or environmental and societal level, e.g. changed work environment or job demands.

Although much research has been conducted on work ability, relatively little is known [15-22]. In this dissertation, some aspects will be studied in greater detail. First, the clinimetric properties of a work ability instrument are tested. To determine if and how work ability is affected, knowing measurement properties is a priority. To use an instrument that measures workability in research and practice, the ability of that instrument to detect changes needs to be established. Although the work ability house is an extensive model of work ability, not all properties of the measurement instruments of work ability have been determined.

Next, this conceptual house model of work ability is a global overview of aspects that might influence work ability in an individual. But when it comes to knowledge of specific working populations and the strength of effects, many aspects are still unknown. Studying work ability in specific working populations can contribute to greater knowledge of which job demands and health requirements are important for work ability. In this way, we aim to study work ability in different circumstances in order to tackle work ability issues in different working populations.

Lastly, work ability in individuals is hard to determine since each individual has a different personal and work environment. However, in specific populations, work ability can be affected similarly. In what way and to what extent work ability is affected

in populations with specific job demands or functional limitations remains unknown in many populations. Studying these aspects can aid in developing interventions to help workers remain being able to meet job demands. Therefore, we aim to determine work ability and related aspects in specific working populations.

MEASUREMENT OF WORK ABILITY

Multiple ways of measuring work ability have been developed as work ability can be interpreted and measured in different ways. The two most frequently used measurement instruments are the Work Ability Index (WAI), and its single item derivate, the Work Ability Score (WAS) [23, 24]. These instruments were developed to quantify the ability to meet job demands.

Work ability research started with the Work Ability Index (WAI) [7]. This tool consists of seven domains of self-assessed work ability. The WAI itself is long and comprehensive. Therefore, a single item measure was developed to assess work ability. This measure was called the Work Ability Score (WAS) and consisted solely of the first item of the WAI [24]. This item assessed self-appraised work ability on a scale of zero to ten with a single question, where workers themselves assessed their own work ability compared to their lifetime best: What is your current work ability to fulfil your job demands, compared with your lifetime best (where 0 represents no work ability at all, and 10 represents the best work ability ever experienced). This single-item measurement showed sufficient convergent validity with the complete WAI and was found to be suitable for systematic screening of work ability in employee surveys and research in the field of occupational health [25, 26]. Because the WAS can be measured quickly and is less complex to answer, the WAS is used in the studies included in this dissertation. Next to the general work ability, the physical and mental/emotional work ability can be measured with a single-item measurement in the same way.

Monitoring work ability can give insight into imminent problems with working in good health until retirement age. To be able to monitor work ability, it is important that the instrument, besides showing sufficient convergent validity, is capable of measuring changes (improvements and deteriorations) in the work ability of workers. To be able to use the WAS, it is essential that this instrument is validated and that its clinimetric properties have been mapped. With a valid instrument, the course of work ability can be monitored, and alternatively improvements or deteriorations of workers' work ability can be detected. In order to determine change in work ability, the reproducibility and responsiveness of the WAS needs to be determined. However, these properties of the WAS are still unknown.

WORK ABILITY IN OCCUPATIONAL HEALTH PRACTICE

Work ability is of growing importance in life. Because of increasing life expectancy, retirement age is rising and the participation rate of older workers has risen in the past decades [27, 28]. In many countries (among which The Netherlands) the pensionable age will be linked to the life expectancy [28]. Consequently, the expected age at retirement is also rising [29]. This means that the workforce needs to work and remain in good health until a later age. This means the negative influences of work will last longer. In an ageing population, being able to work until a later age, and thus having sufficient work ability one's entire working life, becomes increasingly important as we know both health and work ability decline with increasing age, and older workers need more time to recover from labor [4, 7, 30-33].

Occupational health professionals aim to sustain or improve the work ability of workers during the entire course of their working life, as well as maintain their sustainable employability. Sustainable employability can be described as being able to contribute in a valuable way at work until a later age, while protecting the workers physical and mental health [34]. An important aspect of sustainable employability is that workers have the capabilities and resources that allow them to carry out their work while safeguarding their health and welfare. Monitoring work ability in workers can help occupational health professionals detect when sustainable employment is at risk.

PREDICTIVE VALUE OF WORK ABILITY

Work ability is a known predictor of long-term sick leave and disability pension [12, 16, 35-47]. A decrease in work ability can therefore be a sign that the possibility to work in good health until retirement age is at risk. Measuring and monitoring work ability over the entire working life span can help detect problems in a timely manner. Preventing a decrease in work ability can be achieved by timely intervention, therefore decreasing the risk for future disability pension.

Even after sick leave occurred, the level of work ability can still predict the duration of time to return to work and future work participation [45, 48]. On the other hand, good work ability predicts the ability to stay in good health and meet job demands until retirement age [49]. Next to future sick leave and disability pension, work ability also has a large effect on work motivation [50]. Workers who feel they can meet their job demands, have better motivation to work. Maintaining work ability can improve a worker's motivation to work. Additionally, work ability predicts work status: people with higher work ability are more likely to be offered a job as they are more attractive to employers [51]. This also underlines the need for maintaining work ability in workers and intervening when their work ability has decreased.

WORK ABILITY IN SPECIFIC POPULATIONS

Although the focus in work ability research now mostly lies on improving the work ability of older workers, monitoring work ability in young workers may be beneficial for work ability in later life [52]. Monitoring work ability over the course of working life can help detect early declines in work ability. This monitoring can already start in the beginning of one's career. Relatively little is known about work ability in young workers [53]. Monitoring work ability in young adults can be a valuable strategy to predict future sick leave and identify problems in subsequent years [38, 53]. Therefore, studying work ability in young working populations can contribute to maintaining work ability over the course of working life.

Next to monitoring young workers, identifying risks for work ability loss in specific working populations can contribute towards maintaining work ability throughout one's working life [24]. Despite individual differences, work ability can be affected in a common way in working populations with similarities, as in populations with specific functional limitations, or populations with comparable job demands [24, 54]. Two of the largest contributors to work ability are job demands and health requirements [24]. Therefore, we zoom in on populations with functional limitations, and specific job demands.

Work ability related to functional limitations

As the base of the work ability house consists of an individual's health, functional limitations can influence work ability [55, 56]. In what way they decrease work ability, and which limitations have the largest influence, is still unknown. Functional limitations can cause work ability to change accordingly, depending on the strength of the association between the specific limitation and the change in work ability. Therefore, it can be useful studying which limitations affect work ability most, to target interventions or adjust job demands to bypass these limitations [57]. In that way, the worker is still able to work and meet job demands, maintaining sustainable employability. In this dissertation, we study two examples of limitations. One physical limitation, namely Upper Extremity MusculoSkeletal Disorder (UEMSD) complaints, and the second is stress, as an example of mental functional limitations. It is known that both UEMSD and stress have a negative impact on work ability, but it remains unclear and what other factors influence this relationship [58-61].

Work ability related to job demands

Changes in work ability can indicate difficulties in meeting certain job demands. Certain job demands can have a stronger relationship with work ability than others [62, 63]. Meeting these job demands is therefore critical to maintaining good work ability [11, 64]. In specific occupations, certain job demands are more important than

others to function satisfactorily in that job [64, 65]. In a high-demand profession such as ambulance worker, an individual's physical fitness must be sufficient to carry out rescue tasks [66]. Other examples of occupations where job demands have a strong relationship with work ability are occupations with seasonal or periodic changes in workload, e.g. in farming or the leisure industry. These peak loads can take their toll on work ability. In such occupations, these periods of peak load can bring along adverse health effects, and possibly a permanent decrease in work ability [15, 67]. In occupations with a changing workload over time, the worker is required to cope with these demands. One example of such an occupation is coach driver. Coach drivers have to cope with a substantial seasonal change in workload throughout the year. This can affect the ability to recover from work, in turn affecting their work ability [68].

Consequently, not being able to meet these job demands has a large impact on the work ability of workers in these occupations. It is therefore important to map which job demands are most strongly related to work ability. Once these are known, the ability to meet these demands can be closely monitored. In addition, prevention can be focused on being able to meet these job demands. By implementing preventive measures, groups can be helped to work longer in good health [69]. Some jobs necessitate specific health requirements [11, 64, 70]. Although meeting health requirements is essential executing job tasks, it remains unclear how the ability to meet demands is related to work ability.

How interventions can help maintain work ability in different working populations needs to be tested [24]. To improve sustainable employability and prevent decreases in work ability, the factors affecting work ability still need to be studied since it is still unknown how work ability is affected by peak loads. Additionally, the effect of interventions needs to be evaluated to improve interventions and occupational healthcare, especially for those occupations at risk of declining work ability. In this thesis, an example of a self-management intervention is evaluated during the peak season of the coach sector.

INTERVENTIONS TO PREVENT DECLINES IN WORK ABILITY

By improving and maintaining work ability during increased workload, workers are able to stay in good health while being able to meet job demands. The ability to contribute in a valuable way to work until a later age while protecting workers' physical and mental health is an important goal in occupational health [71]. In order to achieve this, workers need to be enabled to reach this goal, which can be done by adjusting work or the workplace, and keeping track of workers' needs [72]. Interventions can be aimed at multiple aspects of the work ability house. A worker can be given resources in the sense of autonomy or improved supervisory support, to

decrease the risk of sick leave or disability pension [73-75]. Improving general health increases the resources of a worker to meet increased demands.

Alternatively, monitoring work ability, especially in vulnerable populations can give timely indications that intervention is desired [76]. By intervening in a timely manner, work ability might be preserved [77-79]. Relating job demands and functional limitations to work ability can shed light on important parameters and reveal problem areas in these populations. When work ability is studied in various populations, occupations and workers at risk of decreasing sustainable employment can be more quickly identified and given special attention in the form of preventive measures or work modification [24].

In coach drivers, it is essential to determine which resources or changes in work and behavior are effective in maintaining work ability over time. Knowing that work ability relates to multiple aspects of work and private life, it makes sense that interventions are aimed at these aspects that characterize an occupation or population. Determining which aspects of work ability could or should be improved is the first step in developing interventions. Especially in coach drivers, interventions can be aimed at multiple aspects of health and work as it remains a challenge to maintain a healthy lifestyle during peak season, when drivers are away from home most of the time.

Multiple interventions have been developed with different strategies, of which the majority focuses on physical work ability, with varying results [80-89]. It still remains a challenge to develop interventions that are able to improve work ability in workers as single-targeted work ability interventions have at most, if any, only slight benefits for work ability. More extensive interventions that focus on multiple aspects of work ability are needed to improve work ability. From previous research, we know that to improve the work ability of an individual, interventions should be aimed at multiple factors, not only work [7, 21]. These aspects need to relate to the work ability house of the group the intervention is being applied to.

These interventions need to be developed and tested with vulnerable working populations, such as coach drivers. Interventions will preferably be specifically designed to tackle problems experienced by workers in this occupation. After testing and evaluating these interventions, they can then be used to improve and contribute to sustainable employment. It still remains unclear which interventions work in which population, and how they should be administered [80, 90]. A population of coach drivers is particularly suitable for performing an intervention with the goal of maintaining work ability during peak season because there are multiple aspects related to work ability that can be addressed.

SCIENTIFIC GAPS

In summary, in this dissertation we aim to add scientific knowledge to fill the following gaps. Before using the WAS instrument to measure change, its reproducibility and responsiveness needs to be determined. This is essential information in determining if and when an individual has experienced a change in work ability. In this way, the course of work ability can be followed and improvements and deteriorations can be mapped. The quantification of significant and relevant change is currently lacking. Therefore a stable population, as well as a population with changing work ability, needs to be studied.

In working populations, work ability is influenced by certain functional limitations. These limitations can take its toll on work ability. Which limitations are related to work ability, and in what way they affect work ability is still unknown in many populations, among which UEMSD patients. In addition to physical functional limitations, mental functional limitations can influence work ability. The pathway of how work ability is influenced and what factors affect this relationship is still unknown in young workers.

High demand occupations, such as ambulance workers are characterized by special job-specific requirements, which are often a prerequisite in meeting the job's demands. Studying which of these demands affect work ability most can help develop preventive strategies. Other occupations with seasonal peak demands, such as coach drivers, need to be guided to prevent declines in work ability during peak season. Whether it is possible to maintain work ability over this period is unknown. In both occupations, more information about demands is needed and possible preventive strategies still need to be studied.

AIMS

The aim of this dissertation was to increase the knowledge about the construct of work ability. This aim was divided into three objectives to fulfil the gaps in scientific and societal knowledge:

1. To gather knowledge about characteristics of work ability measurement in groups of healthy employees and in groups of employees with health complaints and related functional limitations;
2. To gather knowledge about the association between functional limitations and work ability in working populations;
3. To gather knowledge about the course of work ability in work situations characterized by aspects of high workload or in work situations with seasonal changes in workload.

Based on these aims, the following study questions were composed:

- I. What are the reproducibility and responsiveness of the work ability score of workers who return to work after sick leave?
- II. How are functional limitations related to work ability in working populations with specific health complaints?
- III. What is the course of work ability in occupations characterized by specific health requirements and peak loads, and to what extent can this be influenced?

CONTENT OF THIS DISSERTATION

In this thesis, the above-mentioned aims and research questions are answered in multiple studies using diverse populations. The topics that were studied are shown schematically in Figure 2.

Figure 2: Content of this dissertation.

First, the measurement properties of the WAS are determined to facilitate studying the course of work ability in populations (orange). In Chapter 2, the reproducibility of the WAS and a recently developed Work Functioning instrument are determined for an average working population. The course of work ability of individuals returning to work after being on sick leave for at least two weeks are determined in the third

Chapter, where the responsiveness of the WAS and the Work Functioning instrument were tested. The course of work ability in individuals returning to work is studied to determine the distinctive capacity of the WAS and the Work Functioning instrument. This is essential information when monitoring work ability in individuals.

Building on that information, the functional limitations related to work ability are studied (green). In Chapter 4, functional limitations in UEMSD patients are related to work ability. Next, appraised work ability is related to the percentage of worked hours (at the current time) to study to which extent they correspond. To gain information on how the relationship between stress and work ability is mediated, several factors that can influence this relationship are studied in young adults at the beginning of their working life, which is represented in the fifth Chapter of this thesis.

The work ability of specific populations can be analyzed, to study the contributions and influences of job demands and specific limitations on work ability (red). Chapter 6 studies these influences of job-specific health requirements on work ability in ambulance workers. In Chapter 7, an interactive toolbox was developed and evaluated aimed at improving work ability and vitality in coach drivers during peak season where high workload arises.

These studies are linked to prevention and intervention as way of improving occupational healthcare and mapping pathways on how work ability is influenced via several pathways (blue).

In the general discussion (Chapter 8), the main findings of these studies are summarized, and discussed in light of their societal relevance and relevance for (future) research.

REFERENCES

1. Eurostat. Duration of working life - statistics. 2020.
2. OECD. Health status - Life expectancy at birth (indicator). 2020.
3. Milosevic M, Golubic R, Knezevic B, Golubic K, Bubas M, Mustajbegovic J. Work ability as a major determinant of clinical nurses' quality of life. *J Clin Nurs*. 2011;20(19-20):2931-8.
4. Tavakoli-Fard N, Mortazavi SA, Kuhpayehzadeh J, Nojomi M. Quality of life, work ability and other important indicators of women's occupational health. *Int J Occup Med Environ Health*. 2016;29(1):77-84.
5. Waddell G, Burton AK. Is work good for your health and well-being?: The Stationery Office; 2006.
6. Ward SJ, King LA. Work and the good life: How work contributes to meaning in life. *Research in Organizational Behavior*. 2017;37:59-82.
7. Ilmarinen J, Tuomi K, Seitsamo J, editors. New dimensions of work ability. International congress series; 2005: Elsevier.
8. Airila A, Hakanen JJ, Schaufeli WB, Luukkonen R, Punakallio A, Lusa S. Are job and personal resources associated with work ability 10 years later? The mediating role of work engagement. *Work & Stress*. 2014;28(1):87-105.
9. Van den Broeck A, Elst TV, Baillien E, Sercu M, Schouteden M, De Witte H, et al. Job Demands, Job Resources, Burnout, Work Engagement, and Their Relationships: An Analysis Across Sectors. *J Occup Environ Med*. 2017;59(4):369-76.
10. Ilmarinen J, Tuomi K, Klockars M. Changes in the work ability of active employees over an 11-year period. *Scand J Work Environ Health*. 1997;23 Suppl 1:49-57.
11. Sluiter JK. High-demand jobs: age-related diversity in work ability? *Appl Ergon*. 2006;37(4):429-40.
12. McGonagle AK, Fisher GG, Barnes-Farrell JL, Grosch JW. Individual and work factors related to perceived work ability and labor force outcomes. *J Appl Psychol*. 2015;100(2):376-98.
13. Ilmarinen J. From Work Ability Research to Implementation. *Int J Environ Res Public Health*. 2019;16(16).
14. Ilmarinen J. Towards a longer worklife! Ageing and the quality of worklife in the European Union: Finnish Institute of Occupational Health, Ministry of Social Affairs and Health, Helsinki; 2006. 476 p.
15. Tuomi K, Eskelinen L, Toikkanen J, Jarvinen E, Ilmarinen J, Klockars M. Work load and individual factors affecting work ability among aging municipal employees. *Scand J Work Environ Health*. 1991;17 Suppl 1:128-34.
16. Tuomi K, Ilmarinen J, Seitsamo J, Huuhtanen P, Martikainen R, Nygard CH, et al. Summary of the Finnish research project (1981-1992) to promote the health and work ability of aging workers. *Scand J Work Environ Health*. 1997;23 Suppl 1:66-71.
17. Karttunen JP, Rautiainen RH. Work Ability Index among Finnish dairy farmers. *J Agric Saf Health*. 2009;15(4):353-64.

18. Feldt T, Hyvonen K, Makikangas A, Kinnunen U, Kokko K. Development trajectories of Finnish managers' work ability over a 10-year follow-up period. *Scand J Work Environ Health*. 2009;35(1):37-47.
19. Lindbohm ML, Taskila T, Kuosma E, Hietanen P, Carlsen K, Gudbergsson S, et al. Work ability of survivors of breast, prostate, and testicular cancer in Nordic countries: a NOCWO study. *J Cancer Surviv*. 2012;6(1):72-81.
20. Amorim JS, Salla S, Trelha CS. Factors associated with work ability in the elderly: systematic review. *Rev Bras Epidemiol*. 2014;17(4):830-41.
21. van den Berg TI, Elders LA, de Zwart BC, Burdorf A. The effects of work-related and individual factors on the Work Ability Index: a systematic review. *Occup Environ Med*. 2009;66(4):211-20.
22. Kuoppala J, Lamminpää A. Rehabilitation and work ability: a systematic literature review. *J Rehabil Med*. 2008;40(10):796-804.
23. Tuomi K, Ilmarinen J, Jahkola A, Katajarinne L, Tulkki A. Work ability index: Institute of Occupational Health Helsinki; 1994.
24. Gould R, Ilmarinen J, Järvisalo J, Koskinen S. Dimensions of work ability. Helsinki: Finnish Centre for Pensions (ETK) The Social Insurance Institution (Kela) National Public Health Institute (KTL) Finnish Institute of Occupational Health (FIOH); 2008.
25. El Fassi M, Bocquet V, Majery N, Lair ML, Couffignal S, Mairiaux P. Work ability assessment in a worker population: comparison and determinants of Work Ability Index and Work Ability score. *BMC Public Health*. 2013;13:305.
26. Ebener M, Hasselhorn HM. Validation of Short Measures of Work Ability for Research and Employee Surveys. *Int J Environ Res Public Health*. 2019;16(18).
27. OECD. Pensions at a Glance 2011- Retirement-income Systems in OECD and G20 Countries. Paris: OECD Publishing; 2011.
28. Finnish Centre for Pensions. Retirement ages in different countries 2019 16-3-2020]. Available from: <https://www.etk.fi/en/the-pension-system/international-comparison/retirement-ages/> (Accessed 16th March 2020).
29. Finnish Centre for Pensions. Age at retirement continues to rise 2019 16-3-2020]. Available from: <https://www.etk.fi/en/tiedote/age-at-retirement-continues-to-rise/> (Accessed 16th March 2020).
30. Ilmarinen JE. Aging workers. *Occup Environ Med*. 2001;58(8):546-52.
31. Boissonneault M, de Beer J. Work Ability Trajectories and Retirement Pathways: A Longitudinal Analysis of Older American Workers. *J Occup Environ Med*. 2018;60(7):e343-e8.
32. Costa G, Sartori S. Ageing, working hours and work ability. *Ergonomics*. 2007;50(11):1914-30.
33. Tuomi K, Jarvinen E, Eskelinen L, Ilmarinen J, Klockars M. Effect of retirement on health and work ability among municipal employees. *Scand J Work Environ Health*. 1991;17 Suppl 1:75-81.

34. van der Klink JJ, Bultmann U, Burdorf A, Schaufeli WB, Zijlstra FR, Abma FI, et al. Sustainable employability--definition, conceptualization, and implications: A perspective based on the capability approach. *Scand J Work Environ Health*. 2016;42(1):71-9.
35. Alavinia SM, de Boer AG, van Duivenbooden JC, Frings-Dresen MH, Burdorf A. Determinants of work ability and its predictive value for disability. *Occup Med (Lond)*. 2009;59(1):32-7.
36. Jaaskelainen A, Kausto J, Seitsamo J, Ojajarvi A, Nygard CH, Arjas E, et al. Work ability index and perceived work ability as predictors of disability pension: a prospective study among Finnish municipal employees. *Scand J Work Environ Health*. 2016;42(6):490-9.
37. Roelen CA, van Rhenen W, Groothoff JW, van der Klink JJ, Twisk JW, Heymans MW. Work ability as prognostic risk marker of disability pension: single-item work ability score versus multi-item work ability index. *Scand J Work Environ Health*. 2014;40(4):428-31.
38. Kujala V, Tammelin T, Remes J, Vammavaara E, Ek E, Laitinen J. Work ability index of young employees and their sickness absence during the following year. *Scand J Work Environ Health*. 2006;32(1):75-84.
39. Lundin A, Kjellberg K, Leijon O, Punnett L, Hemmingsson T. The Association Between Self-Assessed Future Work Ability and Long-Term Sickness Absence, Disability Pension and Unemployment in a General Working Population: A 7-Year Follow-Up Study. *J Occup Rehabil*. 2016;26(2):195-203.
40. Schouten LS, Joling CI, van der Gulden JW, Heymans MW, Bultmann U, Roelen CA. Screening manual and office workers for risk of long-term sickness absence: cut-off points for the Work Ability Index. *Scand J Work Environ Health*. 2015;41(1):36-42.
41. Schouten LS, Bultmann U, Heymans MW, Joling CI, Twisk JW, Roelen CA. Shortened version of the work ability index to identify workers at risk of long-term sickness absence. *Eur J Public Health*. 2016;26(2):301-5.
42. Alavinia SM, van den Berg TI, van Duivenbooden C, Elders LA, Burdorf A. Impact of work-related factors, lifestyle, and work ability on sickness absence among Dutch construction workers. *Scand J Work Environ Health*. 2009;35(5):325-33.
43. Ahlstrom L, Grimby-Ekman A, Hagberg M, Dellve L. The work ability index and single-item question: associations with sick leave, symptoms, and health--a prospective study of women on long-term sick leave. *Scand J Work Environ Health*. 2010;36(5):404-12.
44. Bethge M, Gutenbrunner C, Neuderth S. Work Ability Index predicts application for disability pension after work-related medical rehabilitation for chronic back pain. *Arch Phys Med Rehabil*. 2013;94(11):2262-8.
45. Bethge M, Spanier K, Peters E, Michel E, Radoschewski M. Self-Reported Work Ability Predicts Rehabilitation Measures, Disability Pensions, Other Welfare Benefits, and Work Participation: Longitudinal Findings from a Sample of German Employees. *J Occup Rehabil*. 2018;28(3):495-503.
46. Roelen CA, Heymans MW, Twisk JW, van der Klink JJ, Groothoff JW, van Rhenen W. Work Ability Index as tool to identify workers at risk of premature work exit. *J Occup Rehabil*. 2014;24(4):747-54.

47. Lundin A, Leijon O, Vaez M, Hallgren M, Torgen M. Predictive validity of the Work Ability Index and its individual items in the general population. *Scand J Public Health*. 2017;45(4):350-6.
48. de Boer AG, Verbeek JH, Spelten ER, Uitterhoeve AL, Ansink AC, de Reijke TM, et al. Work ability and return-to-work in cancer patients. *Br J Cancer*. 2008;98(8):1342-7.
49. Karttunen JP, Rautiainen H. Risk factors and prevalence of declined work ability among dairy farmers. *J Agric Saf Health*. 2011;17(3):243-57.
50. Feissel A, Swart E, March S. The Impact of Work Ability on Work Motivation and Health: A Longitudinal Study Based on Older Employees. *J Occup Environ Med*. 2018;60(5):e238-e44.
51. Rueda S, Chambers L, Wilson M, Mustard C, Rourke SB, Bayoumi A, et al. Association of returning to work with better health in working-aged adults: a systematic review. *Am J Public Health*. 2012;102(3):541-56.
52. Ilmarinen J. Work ability--a comprehensive concept for occupational health research and prevention. *Scand J Work Environ Health*. 2009;35(1):1-5.
53. Boström M, Holmgren K, Sluiter JK, Hagberg M, Grimby-Ekman A. Experiences of work ability in young workers: an exploratory interview study. *Int Arch Occup Environ Health*. 2016;89(4):629-40.
54. Bugajska J, Sagan A. Chronic musculoskeletal disorders as risk factors for reduced work ability in younger and ageing workers. *Int J Occup Saf Ergon*. 2014;20(4):607-15.
55. van Muijen P, Schellart AJM, Duijts SFA, van der Beek AJ. The mediating role of coping between self-reported health complaints and functional limitations, self-assessed work ability and work status of long-term sick-listed cancer survivors. *Eur J Cancer Care (Engl)*. 2019;28(1):e12928.
56. Faber E, Kuiper JI, Burdorf A, Miedema HS, Verhaar JA. Treatment of impingement syndrome: a systematic review of the effects on functional limitations and return to work. *J Occup Rehabil*. 2006;16(1):7-25.
57. Skovlund SV, Blafoss R, Sundstrup E, Andersen LL. Association between physical work demands and work ability in workers with musculoskeletal pain: cross-sectional study. *BMC Musculoskelet Disord*. 2020;21(1):166.
58. Goedhard RG, Goedhard WJA, editors. *Work ability and perceived work stress*. International Congress Series; 2005: Elsevier.
59. Gharibi V, Mokarami H, Taban A, Yazdani Aval M, Samimi K, Salesi M. Effects of Work-Related Stress on Work Ability Index among Iranian Workers. *Saf Health Work*. 2016;7(1):43-8.
60. Monteiro MS, Alexandre NM, Ilmarinen J, Rodrigues CM. Work ability and musculoskeletal disorders among workers from a public health institution. *Int J Occup Saf Ergon*. 2009;15(3):319-24.
61. Boschman JS, Noor A, Lundstrom R, Nilsson T, Sluiter JK, Hagberg M. Relationships between work-related factors and musculoskeletal health with current and future work ability among male workers. *Int Arch Occup Environ Health*. 2017;90(6):517-26.

62. Oliv S, Noor A, Gustafsson E, Hagberg M. A Lower Level of Physically Demanding Work Is Associated with Excellent Work Ability in Men and Women with Neck Pain in Different Age Groups. *Saf Health Work*. 2017;8(4):356-63.
63. van der Doef MP, Schelvis RMC. Relations Between Psychosocial Job Characteristics and Work Ability in Employees with Chronic Headaches. *J Occup Rehabil*. 2019;29(1):119-27.
64. Boschman JS, Hulshof CTJ, Frings-Dresen MHW, Sluiter JK. Job-specific mandatory medical examinations for the police force. *Occup Med (Lond)*. 2017;67(6):469-76.
65. Punakallio A. Balance abilities of workers in physically demanding jobs: With special reference to firefighters of different ages. *Journal of Sports Science and Medicine*. 2005;4(8):1-47.
66. Rue CA, Rayson MP, Walker EF, Doherty J, Thompson J, Myers SD, et al. A job task analysis to describe the physical demands of specialist paramedic roles in the National Ambulance Resilience Unit (NARU). *Work*. 2019;63(4):547-57.
67. Hakkarainen P, Moilanen L, Hanninen V, Heikkinen J, Rasanen K. Work ability among Finnish workers with type 1 diabetes. *Occup Med (Lond)*. 2016;66(6):446-53.
68. Sluiter JK, van der Beek AJ, Frings-Dresen MHW. The influence of work characteristics on the need for recovery and experienced health: a study on coach drivers. *Ergonomics*. 1999;42(4):573-83.
69. Oakman J, Neupane S, Proper KI, Kinsman N, Nygard CH. Workplace interventions to improve work ability: A systematic review and meta-analysis of their effectiveness. *Scand J Work Environ Health*. 2018;44(2):134-46.
70. Johnson RW, Mermin GB, Resseger M. Job demands and work ability at older ages. *J Aging Soc Policy*. 2011;23(2):101-18.
71. McGonagle AK, Barnes-Farrell JL, Di Milia L, Fischer FM, Hobbs BB, Iskra-Golec I, et al. Demands, resources, and work ability: A cross-national examination of health care workers. *European Journal of Work and Organizational Psychology*. 2014;23(6):830-46.
72. Johansson G, Hultin H, Moller J, Hallqvist J, Kjellberg K. The impact of adjustment latitude on self-assessed work ability in regard to gender and occupational type. *Scand J Occup Ther*. 2012;19(4):350-9.
73. de Croon EM, Sluiter JK, Nijssen TF, Kammeijer M, Dijkmans BA, Lankhorst GJ, et al. Work ability of Dutch employees with rheumatoid arthritis. *Scand J Rheumatol*. 2005;34(4):277-83.
74. Bernburg M, Vitzthum K, Groneberg DA, Mache S. Physicians' occupational stress, depressive symptoms and work ability in relation to their working environment: a cross-sectional study of differences among medical residents with various specialties working in German hospitals. *BMJ Open*. 2016;6(6):e011369.
75. Smither JW, London M, Reilly RR. Does performance improve following multisource feedback? A theoretical model, meta-analysis, and review of empirical findings. *Personnel psychology*. 2005;58(1):33-66.

76. Sell L, Bultmann U, Rugulies R, Villadsen E, Faber A, Sogaard K. Predicting long-term sickness absence and early retirement pension from self-reported work ability. *Int Arch Occup Environ Health*. 2009;82(9):1133-8.
77. Schwarze M, Egen C, Gutenbrunner C, Schriek S. Early Workplace Intervention to Improve the Work Ability of Employees with Musculoskeletal Disorders in a German University Hospital-Results of a Pilot Study. *Healthcare (Basel)*. 2016;4(3).
78. Parkka J, Merilahti J, Mattila EM, Malm E, Antila K, Tuomisto MT, et al. Relationship of psychological and physiological variables in long-term self-monitored data during work ability rehabilitation program. *IEEE Trans Inf Technol Biomed*. 2009;13(2):141-51.
79. de Boer AG, Burdorf A, van Duivenbooden C, Frings-Dresen MH. The effect of individual counselling and education on work ability and disability pension: a prospective intervention study in the construction industry. *Occup Environ Med*. 2007;64(12):792-7.
80. Cloostermans L, Bekkers MB, Uiters E, Proper KI. The effectiveness of interventions for ageing workers on (early) retirement, work ability and productivity: a systematic review. *Int Arch Occup Environ Health*. 2015;88(5):521-32.
81. Pohjonen T, Ranta R. Effects of worksite physical exercise intervention on physical fitness, perceived health status, and work ability among home care workers: five-year follow-up. *Prev Med*. 2001;32(6):465-75.
82. Hoving JL, Bartelds GM, Sluiter JK, Sadiraj K, Groot I, Lems WF, et al. Perceived work ability, quality of life, and fatigue in patients with rheumatoid arthritis after a 6-month course of TNF inhibitors: prospective intervention study and partial economic evaluation. *Scand J Rheumatol*. 2009;38(4):246-50.
83. Kettunen O, Vuorimaa T, Vasankari T. 12-mo intervention of physical exercise improved work ability, especially in subjects with low baseline work ability. *Int J Environ Res Public Health*. 2014;11(4):3859-69.
84. Gram B, Holtermann A, Bultmann U, Sjogaard G, Sogaard K. Does an exercise intervention improving aerobic capacity among construction workers also improve musculoskeletal pain, work ability, productivity, perceived physical exertion, and sick leave?: a randomized controlled trial. *J Occup Environ Med*. 2012;54(12):1520-6.
85. Gao Y, Nevala N, Cronin NJ, Finni T. Effects of environmental intervention on sedentary time, musculoskeletal comfort and work ability in office workers. *Eur J Sport Sci*. 2016;16(6):747-54.
86. Sjögren T. Effectiveness of a workplace physical exercise intervention on the functioning, work ability, and subjective well-being of office workers: a cluster randomised controlled cross-over trial with a one-year follow-up: University of Jyväskylä; 2006.
87. Nurminen E, Malmivaara A, Ilmarinen J, Ylostalo P, Mutanen P, Ahonen G, et al. Effectiveness of a worksite exercise program with respect to perceived work ability and sick leaves among women with physical work. *Scand J Work Environ Health*. 2002;28(2):85-93.

88. Oude Hengel KM, Bosmans JE, Van Dongen JM, Bongers PM, Van der Beek AJ, Blatter BM. Prevention program at construction worksites aimed at improving health and work ability is cost-saving to the employer: results from an RCT. *Am J Ind Med*. 2014;57(1):56-68.
89. Ting JZR, Chen X, Johnston V. Workplace-Based Exercise Intervention Improves Work Ability in Office Workers: A Cluster Randomised Controlled Trial. *Int J Environ Res Public Health*. 2019;16(15).
90. Hoefsmit N, Houkes I, Nijhuis FJ. Intervention characteristics that facilitate return to work after sickness absence: a systematic literature review. *J Occup Rehabil*. 2012;22(4):462-77.

2

Reproducibility of work ability and work functioning instruments

A. van Schaijk, K. Nieuwenhuijsen, M. H. W. Frings-Dresen and J. K. Sluiter

Occupational Medicine, 2018, 68(2), 116-119.

ABSTRACT

Background Work ability (WA) and work functioning (WF) instruments can be useful in occupational health practice. The reproducibility of both instruments is important to their relevance for daily practice. Clinimetrics concerns the methodological and statistical quality of instruments and their performance in practice.

Aims: To assess the reproducibility of WA and WF instruments.

Methods Dutch workers completed a questionnaire containing WA questions and the WF questionnaire twice with a 7-day interval between. The questionnaire included an appraisal of current general, physical and mental/emotional WA (0–10) and the composite WF questionnaire of 49 items. We measured reproducibility, reliability and agreement by calculating the intraclass correlation coefficient (ICC), the standard error of measurement (SEM) and the smallest detectable change (SDC).

Results The answers of 104 respondents were available for analysis. General, physical and mental/emotional WA had ICC values of 0.52, 0.69 and 0.56, respectively. For WF, the ICC value was 0.85. For general WA, the SEM was 0.71. For physical and mental/emotional WA, the SEMs were 0.75 and 0.74, respectively. For general, physical and mental/emotional WA, the SDC was 1.98, 2.09 and 2.05 respectively. The SEM of the WF score was 4.78, and the SDC was 13.25.

Conclusions The WA questions showed moderate reliability, while the WF instrument showed good reliability. Occupational health professionals can use the SDCs of the instruments to monitor changes in WA and WF in workers over time.

Key words: Anchor question; Health Surveys; Reproducibility of results; Surveys and questionnaires; work ability appraisal; Work Ability Score (WAS); Work performance

INTRODUCTION

Work ability (WA) and work-related health have become increasingly important in recent years. WA is the extent to which people are capable of doing their job satisfactorily with respect to their job demands and their health [1]. A closely related concept is work functioning (WF) which is described as the relationship between health-related capacities and the ability to fulfil obligations to meet expectations in the workplace [2]. The working population is ageing. Workers experience health complaints more often, which increases their risk of lower work ability [3, 4].

Various instruments are available to occupational health professionals in research and practice to assess WA, WF and other work-related health issues. Two of these are the WA questions of the Work Ability Index (WAI) [5] and a recently developed WF instrument [6]. Both instruments are designed to identify workers who function insufficiently in their job, taking into account the interaction between their health and the specific demands of their job. The first three questions of the WAI can be used for a quick general assessment of physical and mental abilities to work. The WF questionnaire is more time-consuming and can be used for in-depth analysis. It considers four domains 'Quality of work performance', 'Recovery from work', 'Quantity of work' and 'Capacity to work' in an integrated way [7].

Studies using the first question of the WAI, the appraisal of general WA, (also called the Work Ability Score [WAS] [4, 8]), showed sufficient convergent validity with the seven-item WAI questionnaire ($r = 0.63$) [9]. Little variation in WAS was seen in healthy workers with a mean score between 8 and 9 [4, 9, 10]. The smallest detectable change (SDC) of the first question of the WAI in a test-retest situation was 2.0 in a study of elderly construction workers and 1.9 in a group of 29 young adults [10-12]. The reproducibility characteristics of these WA and WF instruments will indicate whether they are useful and applicable for monitoring over time. If so, they may be useful for workers' health surveillance or monitoring return-to-work after sick leave. Reproducibility can be measured using reliability and agreement measures in a stable working population in a stable working environment to eliminate changes from outside sources [13]. Reliability shows how well workers can be distinguished from each other regardless of measurement error, taking variability between subjects into account. Agreement assesses how well repeated measurements agree, taking systematic changes into account. Agreement can be measured and displayed with the standard error of measurement (SEM), which represents measurement error, and the SDC [13, 14]. The aim of this study was to evaluate the reproducibility of WA and WF instruments. Our research question was: *What is the reproducibility of work ability and work functioning?*

METHODS

We recruited subjects from the Dutch working population aged between 15 and 75, living in the Netherlands, with a paid job and working at least 12 h a week. We only included those who had been working in the same job role for the last 4 weeks. We used convenience sampling through social media and flyers with a website link to the participant information and informed consent page in June and July 2016. We emailed the second questionnaire 7 days after completion of the first questionnaire, to minimize differences in weekday-related working fluctuation. We planned to stop recruitment after 101 participants had completed the questionnaires twice, 1 week apart. We based this on the recommendations for adequate sample size of a checklist for quality: the COnsensus-based Standards for the selection of health Measurement INstruments (COSMIN) [15]. We conducted the research in accordance with the declaration of Helsinki [16]. The research proposal was approved by the Medical Ethical Committee of the Academic Medical Center, who judged that a comprehensive evaluation was not required since this study was not subject to the Medical Research Involving Human Subjects Act (W16_153#16.177). We asked an anchor question before the second questionnaire to establish stability during interval between the questionnaires [17]: *Did anything significantly change in the way you perform your work compared to the previous time you filled out this questionnaire?* Response options were: ‘improved’, ‘not changed’ and ‘deteriorated’. We only analysed data from those who reported ‘not changed’.

Three WA questions appraised respondents’ current general, physical and mental/emotional WA compared with lifetime best, on a scale of 0–10, with 0 meaning ‘no work ability at all’, and 10 meaning ‘the best work ability ever experienced’ [5, 10]. We adapted the second and third questions of the WAI for physical and mental/emotional WA to resemble the first question by changing the scale from 1–5 to 0–10. We used the WAS as they were recorded by respondents on the 0–10 scale.

The composite WF questionnaire developed and validated by Boezeman et al. contains 49 questions in four domains: ‘Quality of work performance’, ‘Recovery from work’, ‘Quantity of work’ and ‘Capacity to work’ [6, 7]. For WF, we transformed domain scores for the four domains to a 0–100 scale, provided that <50% was missing [7]. We used these 0–100 scores to calculate the WF overall score as: $((\text{domain score quality of work performance} \times 1.66) + (\text{domain score recovery from work} \times 1.15) + (\text{domain score quantity of work} \times 1.11) + (\text{domain score capacity to work} \times 0.77)) / 4.69$ [7].

We determined the reliability of both instruments using the ICC_{agreement}, which is the best parameter for continuous variables [14]. We used the two-way random setting,

looking at single measures. An Intraclass Correlation Coefficient (ICC) value ≥ 0.90 is considered to be excellent reliability. A value in the range of ≥ 0.70 to < 0.90 is considered to be good, while values ≥ 0.50 to < 0.70 and ≥ 0.30 to < 0.50 are considered moderate and poor, respectively. A value < 0.30 indicates very poor reliability [18]. We determined the SEM to assess the accuracy of both instruments using the formula: ' $SEM_{agreement} = SD \times \sqrt{1 - ICC_{agreement}}$ ' [13]. The SDC is the smallest measurable within-person change above the measurement error and calculated as ' $1.96 \times \sqrt{2} \times SEM$ ' [13]. Agreement parameters are directly interpretable since the outcome is presented on the same scale as the original measure. We performed all statistical analyses using IBM SPSS Statistics 23 except for the calculation of the SEM and SDC, which we calculated using Microsoft Excel 2010.

RESULTS

One hundred and forty-three respondents consented and completed the first questionnaire (65% of those that visited the information and informed consent page). Of these, 121(85%) completed the second questionnaire 1 week later. Of these, the answers of 104 respondents could be used for analysis. We excluded data from 16 respondents because they reported a change in their work circumstances, i.e. vacation, reporting sick leave or a job change. We excluded data from one respondent due to $\geq 50\%$ missing values in one domain of the WF instrument. The remaining 104 respondents were diverse in occupation and age. Occupations ranged from domestic help to psychologist in the health and social work sector, from saleswoman to managing director in the retail business, from host to chef cook in the hospitality sector, and from secretary to public prosecutor in the legal sector. Forty-three percent of respondents were male; 37% were 18–30 years old, 22%, 31–40, 17%, 41–50 and 24%, ≥ 51 . Relatively high WAS, good WF scores and minimal changes were reported. Absolute mean scores and change scores for the main outcome measures are shown in Table 1.

Table 1: Mean and standard deviation (SD) for Work Ability (WA) and Work Functioning (WF) scores.

	Mean T1 (SD)		Mean T2 (SD)		Mean change (SD)		Mean absolute change (SD)	
General WA (0–10)	8.1	(0.98)	8.0	(1.08)	0.06	(1.01)	0.58	(0.83)
Physical WA (0–10)	8.3	(1.39)	8.4	(1.30)	−0.10	(1.07)	0.73	(0.78)
Mental/emotional WA (0–10)	8.0	(1.19)	8.0	(1.06)	−0.06	(1.05)	0.67	(0.81)
WF (0–100)	17.2	(12.42)	16.4	(12.22)	0.76	(6.75)	4.93	(4.65)

General WA = WAS

Table 2 shows that reliability for general, physical and mental/emotional WA was moderate, and good for WF. The SDCs for overall WA and for physical and mental/emotional WA show that a change of two points represents measurable change rather than measurement error. The SDC for WF shows that a change of 13.25 points represents 95% certainty of measurable change.

Table 2: Reliability (ICC) and agreement (SEM and SDC) for WAS and WF scores.

	Reliability	Agreement	
	ICC	SEM	SDC
General WA (0–10)	0.52	0.71	1.98
Physical WA (0–10)	0.69	0.75	2.09
Mental/emotional WA (0–10)	0.56	0.74	2.05
WF (0–100)	0.85	4.78	13.25

DISCUSSION

Reliability was moderate for WA questions and good for the WF questionnaire in a population of Dutch workers of various occupations and ages. The SDC for WA questions was 2 on a scale of 0–10, and 13 points for the WF questionnaire on a scale of 0–100.

A strength of this study is that all the quality protocols of the COSMIN checklist were followed. In contrast to other studies of reproducibility, we confirmed stability of work situation of respondents by using an anchor question. This method is often used in studies assessing the minimal important change [19, 20]. Another strength is that we compared two instruments in one population, where other studies have usually assessed the reproducibility of one instrument. Bias is unlikely because all educational levels are represented in the population. A potential limitation is that we only included those currently working and in the same role for at least 4 weeks. Including work-impaired workers of those on (part-time) sick leave, might have produced more

variation in WAS and higher ICC scores. However, WF is probably best assessed in those currently working.

Our results for ICC, SEM and SDC for general WA are similar to those found in a previous study with a 4-week interval between the questionnaires [10, 11]. In addition to reliability, we explored the agreement characteristics of both instruments. Agreement is more stable over different populations and is more of an instrument characteristic than the reliability, which is more population-dependent [13].

In practice, assessing change in WA and WF scores can be useful for sick leave prevention or monitoring return-to-work, giving opportunities for intervention if return-to-work does not progress as desired. Occupational health professionals can use the SDC of the instruments to assess whether measurable change has occurred in repeated assessments. We suggest that further research focuses on determining the clinical relevance of the SDC.

KEY POINTS

- This study found that reliability of work ability was moderate, and reliability of the work functioning instrument was good.
- The smallest detectable change for work ability questions was 2 on a scale of 0–10, and for the work functioning questionnaire 13 points on a scale of 0–100.
- These instruments can be used by occupational professionals and researchers to assess changes in work ability and work functioning scores over time in sickness prevention and in monitoring return-to-work.

Acknowledgments The authors thank all respondents that took part in this study.

Competing interests None declared

REFERENCES

1. Ilmarinen J, Tuomi K, Klockars M. Changes in the work ability of active employees over an 11-year period. *Scand J Work Environ Health*. 1997;23 Suppl 1:49-57.
2. Lagerveld SE, Bultmann U, Franche RL, van Dijk FJ, Vlasveld MC, van der Feltz-Cornelis CM, et al. Factors associated with work participation and work functioning in depressed workers: a systematic review. *J Occup Rehabil*. 2010;20(3):275-92.
3. Zoer I, Frings-Dresen MH, Sluiter JK. Are musculoskeletal complaints, related work impairment and desirable adjustments in work age-specific? *Int Arch Occup Environ Health*. 2014;87(6):647-54.
4. Gould R, Ilmarinen J, Järvisalo J, Koskinen S. Dimensions of work ability. Helsinki: Finnish Centre for Pensions (ETK) The Social Insurance Institution (Kela) National Public Health Institute (KTL) Finnish Institute of Occupational Health (FIOH); 2008.
5. Tuomi K, Ilmarinen J, Jahkola A, Katajarinne L, Tulkki A. Work ability index, 2nd revised edition. Helsinki: Finnish Institute of Occupational Health; 1998.
6. Boezeman EJ, Sluiter JK, Nieuwenhuijsen K. Measuring Work Functioning: Validity of a Weighted Composite Work Functioning Approach. *J Occup Rehabil*. 2015;25(3):537-42.
7. Boezeman EJ, Nieuwenhuijsen K, de Bekker-Grob EW, van den Akker-van Marle ME, Sluiter JK. The relative importance of the domains of work functioning: evaluations of health-impaired employees, healthy employees, and employers. *J Occup Environ Med*. 2015;57(4):361-6.
8. Schouten LS, Bultmann U, Heymans MW, Joling CI, Twisk JW, Roelen CA. Shortened version of the work ability index to identify workers at risk of long-term sickness absence. *Eur J Public Health*. 2015:301-5.
9. El Fassi M, Bocquet V, Majery N, Lair ML, Couffignal S, Mairiaux P. Work ability assessment in a worker population: comparison and determinants of Work Ability Index and Work Ability score. *BMC Public Health*. 2013;13:305.
10. de Zwart BC, Frings-Dresen MH, van Duivenbooden JC. Test-retest reliability of the Work Ability Index questionnaire. *Occup Med (Lond)*. 2002;52(4):177-81.
11. Boschman JS, van der Molen HF, Frings-Dresen MH, Sluiter JK. The impact of common mental disorders on work ability in mentally and physically demanding construction work. *Int Arch Occup Environ Health*. 2014;87(1):51-9.
12. Boström M, Sluiter JK, Hagberg M. Changes in work situation and work ability in young female and male workers. A prospective cohort study. *BMC Public Health*. 2012;12:694.
13. de Vet HC, Terwee CB, Knol DL, Bouter LM. When to use agreement versus reliability measures. *J Clin Epidemiol*. 2006;59(10):1033-9.
14. Terwee CB, Bot SD, de Boer MR, van der Windt DA, Knol DL, Dekker J, et al. Quality criteria were proposed for measurement properties of health status questionnaires. *J Clin Epidemiol*. 2007;60(1):34-42.

15. Terwee CB, Mokkink LB, Knol DL, Ostelo RW, Bouter LM, de Vet HC. Rating the methodological quality in systematic reviews of studies on measurement properties: a scoring system for the COSMIN checklist. *Qual Life Res.* 2012;21(4):651-7.
16. WMA. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA.* 2013;310(20):2191-4.
17. de Vet HC, Terwee CB, Ostelo RW, Beckerman H, Knol DL, Bouter LM. Minimal changes in health status questionnaires: distinction between minimally detectable change and minimally important change. *Health Qual Life Outcomes.* 2006;4:54.
18. van Lummel RC, Walgaard S, Hobert MA, Maetzler W, van Dieen JH, Galindo-Garre F, et al. Intra-Rater, Inter-Rater and Test-Retest Reliability of an Instrumented Timed Up and Go (iTUG) Test in Patients with Parkinson's Disease. *PLoS One.* 2016;11(3):e0151881.
19. de Vet HC, Ostelo RW, Terwee CB, van der Roer N, Knol DL, Beckerman H, et al. Minimally important change determined by a visual method integrating an anchor-based and a distribution-based approach. *Qual Life Res.* 2007;16(1):131-42.
20. Gartner FR, Nieuwenhuijsen K, van Dijk FJ, Sluiter JK. Interpretability of change in the Nurses Work Functioning Questionnaire: minimal important change and smallest detectable change. *J Clin Epidemiol.* 2012;65(12):1337-47.

3

Work ability and work functioning: measuring change in individuals recently returned to work

A. van Schaijk, K. Nieuwenhuijsen, M.H.W. Frings-Dresen and J.K. Sluiter

International archives of occupational and environmental health, 2018, 92(3), 423-433.

ABSTRACT

Purpose: To assess: (1) whether work ability and work-functioning instruments can detect relevant changes in their respective parameters following a return to work (RTW) and (2) what proportion of those returning to work show changes in their work ability and work functioning.

Methods: A total of 1073 workers who returned to work after at least 2 weeks of sick leave were invited to fill out three questionnaires in the first 8 weeks after RTW. These consisted of an appraisal of general, physical, and mental/emotional work ability (scores 0–10) and a work-functioning questionnaire (scores 0–100). Minimal Important Change (MIC) was defined to determine the proportion of people, whose scores had changed at weeks 5 and 8 following RTW. The Smallest Detectable Change (SDC) was determined to put the MIC in perspective of measurement error.

Results: Of all participants, 235 were eligible for the analysis. All MIC values were below the SDC and thus not suitable for use. The SDC for work ability was 2.2 and 19.9 for work functioning. In the first 5 weeks after RTW, 10–15% showed a relevant, measurable improvement in work ability, and work functioning based on the SDC margins.

Conclusions: Both instruments were unable to identify change after RTW adequately. We can conclude that 10–15% of individuals showed improvement in work ability and work functioning in the first 5 weeks after RTW when SDC is used.

Keywords: Minimally clinical important difference, Occupational physician, Return to work, Smallest detectable change, Work Ability Score

INTRODUCTION

Work ability and work functioning are important parameters in a working population. Both concepts relate to a workers' assessment of being capable of carrying out the work given his or her health condition. Work ability is a generic term that includes all aspects of the ability to work at once, and work functioning focusses on several domains that have influence on the functioning of a worker. When workers are signed off sick, the work ability and work functioning are affected. The goal for occupational physicians and other occupational health professionals is to allow them to return to work (RTW) quickly while optimizing their health.

Several strategies have been developed to improve the RTW of workers returning from sick leave [1-3]. In many of these strategies, an employee returning from an extended sick leave will be reintroduced back into the workplace in steps, e.g., for periods of fewer hours, in a graded RTW programme or with modified requirements of workplace activities [3-5]. Central to these strategies is that the functioning of the employees on RTW—their performance, productivity, quality, quantity, and capacity—must be sufficient in those hours and activities. When working fewer hours, the employee has more time to recover from work before the next workday starts. With modified activity, the worker executes only the tasks that they can fulfil given their health situation. It is likely that, in the weeks following the RTW, an individual experiences a gradual increase in work ability and work functioning [6]. If this increase can be identified, the occupational health professional or manager can intervene if improvement does not occur. This can prevent recurrent sick leave by adjusting the work according to the workload capacity growth or loss.

For a successful RTW, a worker needs sufficient work ability. It is important to monitor individual work ability in the RTW process [7]. Different instruments to assess work ability and work functioning can be found in the literature [8-18]. Two of these instruments are the *appraised work ability* and *composite work-functioning* instrument [17, 18]. The appraised work ability tool is a single-item selfappraisal of workers' current work ability compared to their lifetime best, derived from the Work Ability Index [19]. The composite work-functioning tool is a 49-item domain-based work-functioning questionnaire. For this study, these two instruments were chosen, because the Work Ability Score is a single-item question and may be useful for quick assessment, while the work-functioning score can add valuable information on the different domains that need attention for the RTW process.

It is unknown whether the RTW process is linked to a relevant improvement in work ability and work functioning. We expect that after RTW, work participation gradually returns to the original level or a new maximum, and should thus be associated with a positive change in work ability and work functioning over time. Currently,

occupational physicians in The Netherlands rarely use these existing instruments to assess the quality of RTW after absenteeism. One of the reasons may be that the use of these instruments in this context is not yet tested, and the magnitude for relevant change in these instruments is not yet established. Besides this, the time frame in which any relevant change should have occurred after RTW still needs to be defined. Quality of RTW is assessed as the extent to which an employee can fully function again based on his health and the requested work demands. For these instruments to be able determine any relevant changes following an RTW, and establish the optimal time frame in which to deploy the instruments, the trajectory of work ability and work functioning after a period of absenteeism needs to be studied.

It is unknown if the existing instruments for measuring work ability and work functioning are able to detect a change in work ability and work functioning in the early weeks after absenteeism. Therefore, the primary goal of this study is to assess whether these two instruments to monitor work ability and work functioning can detect a change over the first 2–8 weeks of RTW following absenteeism of at least 2 weeks. This leads to the first research question:

Q1 To what extent are the work ability and work-functioning instrument able to detect relevant change after RTW irrespective of the origin of absence?

Subsequently, we aim to use both instruments (the appraised work ability tool and composite work-functioning questionnaire) to determine the trajectories of both work ability and work-functioning in the early weeks after RTW. If we can determine what a relevant change is for both instruments, we can determine the optimal time to administer the questionnaires. Towards this end, we assess at what time after returning to work the majority of workers have made a relative improvement in work ability or work functioning. We envision that this will allow occupational professionals to monitor RTW and have the ability to identify—and subsequently intervene—if improvement in work ability or work functioning does not occur. This leads to the second research question:

Q2 What proportion of employees show a relevant change (as determined in the previous research question) in work ability and work functioning, in the first 8 weeks after RTW?

STUDY POPULATION AND METHODS

Ethics statement

The research was conducted in accordance with the declaration of Helsinki [20]. The research proposal was approved by the Medical Ethical Committee of the Academic Medical Center, who judged that a comprehensive evaluation was not required, since

this study was not subject to the Medical Research Involving Human Subjects Act (W16_154#16.178).

Informed consent

Informed consent was obtained from all individual participants included in the study.

Participants

We planned to include workers for a time frame of 11 months, from August 2016 until June 2017. In this period, a total of 1214 e-mail invitations were sent out by the occupational health service of an applied university, university, and academic hospital in The Netherlands to employees returning from sick leave or maternity leave of at least 2 weeks. The email invitations contained participant information and an informed consent form. After giving informed consent, the participants entered their contact information and date of RTW. Any personal or medical information of people that were invited for participation was not disclosed to the researchers, and the occupational health service received no information of participation, to preserve the privacy of the patients.

Measurements and study design

The participants received three identical questionnaires at weeks 2, 5, and 8 after RTW, including an anchor question in the second questionnaire. The first questionnaire was sent out by email 2 weeks after the date of RTW. The second and third questionnaires were sent out 5 and 8 weeks after RTW, respectively. This questionnaire consisted of two parts. In the first part, the participant appraised their work ability on a scale of 0–10: 0 signifying no work ability at all, and 10 meaning the best work ability ever experienced by the participant. This is the first question of the Work Ability Index, also known as the Work Ability Score (WAS) [18, 21]. Work ability was sub-divided into general, physical, and mental/emotional work ability. The second part of the questionnaire consisted of the weighted composite work-functioning questionnaire [17]. This work-functioning questionnaire consists of 49 items spread over four domains: “Quality of work performance,” “Recovery from work,” “Quantity of work”, and “Capacity to work.” Scores for each of the domains were converted to a 0–100 score. Using these scores and a weighting factor, an end score representative of work functioning was calculated—also on a scale of 0–100—with a score of 0 meaning no problems with work functioning and a score of 100 meaning maximum limitation in work functioning [22]. At 5 weeks after RTW, the following anchor question was asked: *did anything significantly change in the way you are able to carry out all your work satisfactorily since you returned to work?* The three possible answers were: *Improved, not changed or deteriorated.*

Statistical analyses

Before the MIC calculation, the group averages on different time intervals after return to work were calculated and compared by means of a T test to describe the general improvement or deterioration of the participants over time.

To determine the relevant change in work ability and work functioning, the Minimal Important Change (MIC) was calculated for the study population. To put this in perspective of measurement error, we also calculated the Smallest Detectable Change (SDC). To answer the second research question, we determined the proportion of people who changed in their work ability and work functioning according to these statistical parameters.

First, the scores of the work ability questions and the work-functioning instrument were separated per answer on the anchor question. This separation was performed to distinguish between the different groups of people who can experience a different trajectory of work functioning and work ability, which may influence an overall average. This anchor question was used to determine the relevant change of the measurements. The MIC is defined by the CONsensus-based STANDards for the selection of health Measurement INSTRuments (COSMIN) as the smallest change in score that is relevant to the patient [23]. The COSMIN checklist was used to improve the methodological quality of this study on measurement properties. The MIC was determined in two ways using the anchor-based method. These anchor-based methods use an external criterion, which is the relevant change as perceived by the patient/worker.

The first method we used to determine the MIC was the global ratings of change (GRC) or mean change method [24]. The mean change method was used to determine the difference between workers who experienced an improved work ability or work functioning and workers who reported that no change has occurred. In this method, the mean change score is calculated by first determining the individual change scores. The mean of the change scores of the group that has no improvement is subtracted from the mean of the change scores of the group that improved to find the difference. This difference represents the MIC based on the global ratings of change method. The same is done for the *deteriorated* and *not changed* groups to find the MIC of deterioration.

The second anchor-based method to determine the MIC is called the anchor-based MIC distribution [25]. This method assesses the MIC by a visual method integrating the Receiver Operating Characteristic (ROC) as recommended by de Vet et al. [25]. The MIC was defined as the optimal cut-off point on the ROC curve, closest to the upper left corner. The optimal cut-off point is the point with the minimal misclassification of patients (where the sum of the percentages of type 1 and type 2

error is the lowest) [25]. This point was calculated as the Youden's J-statistic; $Sensitivity + Specificity - 1$ [26]. In the formula, the sensitivity and specificity are valued equally. This analysis was used to determine the optimal cut-off point of the area under the ROC curve (AUC) with sensitivity and specificity valued equally. The ROC plots the true-positive rate (sensitivity) against the false positive rate (1-specificity). Sensitivity is the chance of correctly classifying a change as being *improved*. The specificity is the chance of correctly classifying an individual as *not changed* (true negative rate). A larger AUC indicates a greater degree of correct classification. The AUC and the optimal cut-off point of the ROC were determined for both instruments in this study.

For both instruments, the SDC was calculated to check if the MIC was greater than the SDC [27]. This SDC is based on measurement error is calculated with the formula: $1.96 \times \sqrt{2} \times SEM$ [28]. The SDC is the smallest measurable change within a person that is not attributable to measurement error. As standards for acceptable AUC of ROC statistics, we used: values <0.70 as inadequate, ≥ 0.7 to <0.80 as acceptable, between ≥ 0.80 to <0.90 as excellent, and ≥ 0.90 as outstanding [29]. An AUC of at least 0.70 will be considered as adequate.

To answer the second research question, the change scores of the participants have been related to both MIC values and the anchor itself. These standards can be used to calculate the percentage of persons that have a change greater than the calculated cut-off points [30]. In this way, it is possible to track the course of RTW. The percentage of participants who reported a positive change in RTW (according to the different MIC values) were calculated for each instrument and timepoint in this study. This analysis was done to evaluate the use of these two instruments in determining the percentages of workers who experience an increase in their work functioning or work ability in weeks 2–8 after RTW.

All analyses have been performed using IBM SPSS statistics 24. Global ratings of change and SDC were calculated using Microsoft Excel.

RESULTS

Of the 1073 e-mail invitations sent out by the occupational health service, a total of 308 people gave their informed consent. In total, 235 participants were eligible for participation in this research, after 73 people were excluded, because their date of RTW was over 2 weeks ago, or they did not return to their original workplace. A total of 170 persons completed the first questionnaire 2 weeks after RTW. Of the 170 participants, 85% were female, since we also invited people returning from pregnancy leave. A significant proportion (43%) of the population completing the questionnaire

was over 51 years of age. The age categories 18–30, 31–40 and 41–50, constituted 10, 31, and 16%, respectively.

At weeks 5 and 8 following RTW, 161 and 155 people filled out the work ability questions, respectively. For 120 of these, it was possible to calculate a work-functioning problems score. The average baseline scores for work ability were 7.4, 7.6, and 7.3 for general, physical, and mental/emotional work ability, respectively, and 27.4 for work-functioning problems. The average baseline and follow-up scores are shown in Tables 1 and 2. Overall, although significant, the group scores barely differed in the early weeks after RTW.

On a group level, there are significant improvements on T2 compared to T1 for general and physical work ability, and for the work-functioning problems score. On T3, there was no significant difference in work ability compared to T2. The work-functioning score increases from T1 to T2 but decreases again on T3. This decrease was only significant for work-functioning.

Table 1: Mean group scores and standard deviations (SD) of work ability and work functioning in the early weeks after RTW.

	T1 (2 weeks after RTW)		T2 (5 weeks after RTW)		T3 (8 weeks after RTW)	
	Mean	SD	Mean	SD	Mean	SD
General work ability (0–10) (higher is better)	7.4	(1.28)	7.6 *	(1.26)	7.6	(1.46)
Physical work ability (0–10) (higher is better)	7.6	(1.44)	7.8**	(1.36)	7.7	(1.40)
Mental/emotional work ability (0–10) (higher is better)	7.3	(1.66)	7.4	(1.52)	7.4	(1.52)
Work functioning problems (0–100) (lower is better)	27.4	(16.39)	23.1 *	(16.27)	26.1**	(18.71)

The number of workers is absent ($n = 120\text{--}170$)

* Sign. different from prior measurement ($p < 0.01$)

** Sign. different from prior measurement ($p < 0.05$)

In Table 2, the scores of the respondents at 2, 5, and 8 weeks after return to work are shown. The respondents are grouped by their answer on the research question (*deteriorated*, *not changed* or *improved* on T2 when compared to T1) to give more insight to the trajectory of return to work. The three groups all experience a difference in their trajectory. The numbers at T2 are higher than the numbers at both other timepoints, because the file was split by answer on the anchor question. This anchor question was asked at T2. Therefore, it was possible to have missed the first ($n = 28$)

or last ($n = 21\text{--}29$) measurement. If T2 was missed, it was not possible to classify the participant to one of the *improved*, *not changed* or *deteriorated* groups.

Table 2: Mean and standard deviations (SD) for the three time intervals after RTW, grouped answer to the anchor question ($n=99\text{--}142$).

	T1 (2 weeks after RTW)			T2 (5 weeks after RTW)			T3 (8 weeks after RTW)		
	<i>n</i>	Mean	SD	<i>n</i>	Mean	SD	<i>n</i>	Mean	SD
Deteriorated									
General work ability	13	6.5	1.05	14	5.9	1.49	11	5.5	1.92
Physical work ability	13	7.6	1.50	14	7.1	1.69	11	6.2	2.14
Mental/emotional work ability	13	6.5	1.76	14	5.9	2.06	11	5.4	1.57
Work functioning problems	13	36.8	13.31	10	45.7	13.03	7	52.9	15.86
Not changed									
General work ability	70	7.5	1.33	82	7.7	1.19	59	7.6	1.29
Physical work ability	69	7.8	1.34	82	7.8	1.33	59	7.6	1.35
Mental/emotional work ability	70	7.4	1.67	82	7.3	1.52	59	7.2	1.61
Work functioning problems	66	26.2	16.88	59	21.2	16.13	46	28.1	20.70
Improved									
General work ability	59	7.4	1.25	65	7.8	1.04	56	8.0	1.32
Physical work ability	59	7.5	1.47	65	7.9	1.29	56	7.9	1.32
Mental/emotional work ability	59	7.2	1.75	65	7.8	1.18	56	7.8	1.23
Work functioning problems	59	28.1	16.88	51	20.8	13.65	46	20.6	14.52

The first method used to calculate the MIC was the GRC method, using the group averages of persons, where a change score could be calculated. These values are shown in Table 3. The second method is the anchor-based MIC distribution method. As a first step, the visual representations of change scores are shown in figures 1a–d. These are the scores separated into the three groups *deteriorated*, *not changed* and *improved*, according to the anchor question.

Fig. 1: Distribution of change scores of work ability and work functioning (a–d), separated by answer on the anchor question (*deteriorated*, *not changed* and *improved*).

As part of the second step, the ROC curves for improvement are calculated and are shown in figures 2a–d. All AUCs of improvement were classified as inadequate. Although the AUC of deterioration (not shown) was higher than the AUC for improvement, only the AUC of work functioning was excellent with an AUC of 0.87. The AUC's for deterioration of general, physical, and mental/emotional work ability were 0.66, 0.61, and 0.66, respectively, which can be classified as inadequate [29].

Fig. 2: ROC curve with AUC values of improvement for work ability and work functioning.

The results of both MIC calculation methods for both improvement and deterioration, and the calculated SDC, are shown in Table 3. The scores for work functioning are inverted when compared to the work ability questions due to the scale of measurement. A higher score is positive for work ability but negative for work functioning. The point of minimal misclassification is the point closest to the upper left corner of the graph and is calculated using the Youden J-statistic [26]. The corresponding value to the highest J-statistic is shown in Table 3 as the MIC based on the AUC method.

Table 3: Minimal Important Change (MIC) calculated in two ways and Smallest Detectable Change (SDC) of general, physical and mental/emotional work ability and work functioning.

	MIC based on GRC method		MIC based on ROC (AUC) method		SDC
	<i>Improv.</i>	<i>Deter.</i>	<i>Improv.</i>	<i>Deter.</i>	
General work ability	0.08	– 0.66	0.5	– 0.5	2.20
Physical work ability	0.19	– 0.33	0.5	– 0.5	1.93
Mental/emotional work ability	0.45	– 0.74	0.5	– 0.5	2.11
Work-functioning problems	– 4.77	15.07	– 5.96	3.47	19.90

As can be concluded from Table 3, there is effectively no difference between the GRC method and the ROC method for the working ability questions, because they are all smaller than 1 (or – 1). This is the minimum difference that a person can fill in on the questionnaire. According to the MIC calculated with both methods, 1 point difference signifies a relevant change with respect to a previous measurement. However, the SDC is over 2, meaning that the MIC falls within the range of measurement error. A change of one point on the work ability questions can thus still be attributed to measurement error.

The MIC for work-functioning problems calculated with the GRC method is – 4.77 for improvement and 15.07 for deterioration (due to the inverted scale of this instrument). When calculated with the ROC method, these values are – 5.96 and 3.47, respectively. However, both these values fall within the measurement error of the SDC, which is 19.90 point on a 100 point scale.

To assess if the SDC classifies the same amount of workers as changed (as according to the respondents’ answer on the anchor question), the calculated SDCs were used to classify participants in practice [31]. The percentage of workers that are *improved* or *deteriorated* according to the different SDC cut-off values is shown in Table 4. This table describes the percentages of participants that show improvement when the different SDC cut-off values for change are used. As shown, these cut-off values are conservative compared to the anchor question.

Table 4: Percentage (%) and number (n) of people deteriorated, not changed, and improved as classified by the different cut-off values as calculated in the first research question.

	Changed between week 2 and 5		Changed between week 5 and 8		Changed between week 2 and 8	
	%	(n)	%	(n)	%	(n)
Anchor (only between T1 and T2)						
–	10	(16)			–	–
=	50	(84)			–	–
+	40	(67)			–	–
Total	100	(167)			–	–
SDC of General Work Ability WA (Based on a SDC of 2 or larger)						
–	4	(5)	6	(7)	6	(7)
=	86	(120)	88	(111)	82	(102)
+	10	(14)	6	(8)	12	(15)
Total	100	(139)	100	(126)	100	(124)
SDC of Physical Work Ability WA (Based on a SDC of 2 or larger)						
–	5	(7)	6	(8)	8	(10)
=	83	(115)	86	(108)	80	(98)
+	12	(16)	8	(10)	12	(15)
Total	100	(138)	100	(126)	100	(123)
SDC of Mental Work Ability (Based on a SDC of 2 or larger)						
–	5	(7)	8	(10)	8	(10)
=	80	(111)	83	(105)	77	(95)
+	15	(21)	9	(11)	15	(19)
Total	100	(139)	100	(126)	100	(124)
SDC of Work Functioning (Based on a SDC of 19,9 or larger)						
–	1	(1)	6	(5)	5	(5)
=	89	(89)	94	(85)	88	(84)
+	10	(10)	0	(0)	6	(6)
Total	100	(100)	100	(90)	100	(95)

(SDC= Smallest Detectable Change, – indicates *deteriorated*, = indicates *not changed* and + indicates *improved* according to the different SDC cut-off values)

Eight weeks after RTW, 12–15% of the study population improved when the SDC of work ability is used as the cut-off point, compared to 2 weeks after RTW. When looked at the same statistic for work functioning, only 6% showed improvement. The percentage of people that improve after week 5 barely increases when looked at the SDC. A deterioration is more likely to occur between weeks 5 and 8 than between weeks 2 and 5.

DISCUSSION

Key results

The responsiveness of three work ability questions and the work-functioning questionnaire in a sample of workers returning to work was inadequate. The workers that have improved cannot reliably be distinguished from persons that have not improved with these two instruments. Because the MIC values fall within the measurement error, the SDC values must be used to determine change [30, 32]. A change value above the SDC means that a relevant change has occurred that has a 95% certainty of not being attributable to measurement error. In this population, 10–15% showed a significant and measurable increase in work ability and work functioning over the first 5 weeks of RTW. Therefore, on a group level, work ability, and work-functioning increase minimally after RTW.

The responsiveness for deterioration is better than for improvement, but still inadequate, with the exception of the work-functioning instrument, which showed excellent responsiveness for the *deteriorated* group. However, it must be noted that this statistic was calculated using only 12 subjects; therefore, it must be interpreted with care. This is in line with the previous findings, where the AUC of the work ability question was found to be inadequate for distinguishing between persons at high or low risk for disability pension [33]. Another study by Abma et al. (2013)—studying the Work Role Functioning Questionnaire 2.0—showed similar results for their instrument [34]. We found no instruments measuring similar constructs with excellent responsiveness.

Our assumption that people returning to work will have a lower Work Ability Score and a higher work-functioning problems score than the regular working population, which will then gradually return to normal values in the early weeks after RTW, seems only partly correct. This assumption was based on various models of RTW [3, 6, 35]. In these models, the employee is assumed to recommence work while at suboptimal working capacity. The goal in this phase is to let the worker gradually learn and function towards their full (new) capacity. These first weeks after RTW can be seen as the “action phase” [6] or “re-entry phase” [36], in which the worker recommences work and attempts to meet the workplace demands. This phase is seen as a critical phase in the RTW process because of the high risk for relapse [6]. Arends et al. (2014) state that analysis of work-functioning trajectories after RTW may help identify workers at risk for relapse and support these individuals to remain in work [37]. In this study, we did not find a large increase in work ability or a substantial decrease in work-functioning problems over the first few weeks of RTW on an individual level. This may be due to the high baseline values (2 weeks after RTW). The average population in this study returned to work with values close to that of a normal working population.

Compared to a normal working population [38], the baseline Work Ability Score was 0.7 lower, and the baseline work-functioning problems score was 10.2 higher, but the population returning to work does not reach the values of a normal working population in the first 8 weeks after RTW.

Although these work ability and work-functioning instruments are not suitable for determining change on an individual level, the scores at a group level showed that there is compliance with the described models. The minor increase in work ability and decrease in work-functioning problems at a group level at 5 weeks after RTW shows that the average population is coping with their work demands upon RTW (Table 1). The next step in a successful RTW is maintaining or even increasing work ability and work functioning. This phase is called “maintenance” or “stay-at-work” [36, 39]. For a sustainable RTW, it is essential to detect workers whose re-entry does not go as planned or workers who show a decrease in work ability and work functioning or who do not reach normal values. A decrease in these parameters, or not achieving the normal values of healthy working people, could be a reason for an occupational health professional to contact the worker with the aim of preventing a relapse. We found that 5–8% of the population in this study showed a decrease in workability and increased work-functioning problems in the first 8 weeks after RTW. This population may be at risk of relapsing to sick leave again.

Strengths and limitations

A strength of this study is that the anchor incorporates a meaningful change as a subjective outcome reported by the patient [40]. Another strength is that the population is followed over a time frame over which a change in work ability and work functioning could be expected. To our knowledge, this is the first study examining work ability and work-functioning trajectories in the first weeks after RTW. A drawback of this study is that a distinction between people returning from different causes of sick leave cannot be made. We chose to include all diagnoses, because we wanted the population to be heterogeneous and representative of the population of workers returning to work—thus various scenarios resulting in a RTW are included. In addition, in The Netherlands, pregnancy leave falls within the scope of sick leave, and these individuals are, therefore, included in our analyses. Therefore, this study cannot make statements regarding specific populations or diagnoses. Another drawback is that a large number of participants who gave informed consent did not fill out the questionnaires at the different timepoints. It could be that the invitation to fill out the questionnaire has ended up in the spam filter. Considering this, the dropout over the three timepoints is minimal.

Interpretation

The results of this study show that the responsiveness of the work ability questions and the work-functioning questionnaire is insufficient to determine a change in the early phase of RTW in this population of workers based on the MIC. The authors suggest to refrain from using the single item of the Work Ability Index, or the WAS score— for individual use, in this context, since the measurement error is high compared to the MIC.

Because the MIC is below the SDC, whenever a minimal important change occurs, this falls within the measurement error, meaning that it cannot be said with certainty that a relevant and measurable change has occurred. Therefore, although workers considered a small change as *improved* or *deteriorated*, the measurement error on this instrument makes it unable to detect this change. The SDC should thus be used as the cut-off value to conclude that a relevant and measurable change has occurred. For this reason, the SDC is used as the cut-off point to determine change in the second research question.

Large changes need to occur to be able to determine with certainty that a relevant, measurable change has occurred. With better instruments lacking, occupational professionals can use the SDC as a reference for change in populations returning to work—to determine a relevant and measurable change—bearing in mind that people not meeting this score can also experience an improvement or deterioration that falls within the measurement error. Only extreme cases will show up as *improved* or *deteriorated*. The best way for now to monitor the RTW process in a heterogeneous sample remains assessment by an occupational physician during a regular consultation, or a preventative medical examination. This underlines the complexity of monitoring work ability in (recently returned to work) workers.

Prior research has shown working people to generally appraise their work ability between seven and nine out of ten [18, 21, 33, 41, 42]. A change of at least 2 for work ability and 20 for work functioning is hard to achieve and will rarely occur. Workers, therefore, can experience a change in these parameters without this being picked up by the instruments used. This factor can go some way to explain the large discrepancy between the number of people changed on anchor question and the appraised work ability or work functioning as presented in Table 4. Due to there being little variation in this population, the SDC is high compared to the MIC, making it hard to distinguish between people (as evidenced by the overlap in Fig. 1). Contributing to the low variation is the high baseline score at 2 weeks after RTW, signifying that workers returning to work at this point view themselves as being well capable of doing their job well, which is clearly good news.

Separating the scores over time by their answer on the anchor question sheds light on the reasons for the small increase in work ability and work functioning. The average scores change when the population is separated by the answer to the anchor question (Table 2). When workers grouped as *decreased* are excluded, and the analysis only includes people who score *not changed* and *improved* on the anchor question, it can be seen that scores do reach those of healthy workers around 8 weeks after RTW. This may indicate that the period after RTW is critical for occupational professionals to detect these people who are deteriorating and prevent these workers from falling back into sick leave again. This is an important group, and future research needs to focus on detecting people who are at risk of relapsing.

Generalizability

Because the population studied are those returning to work after sickness, we expected it to have a larger variance in work ability and work-functioning scores than regular working populations. This variance makes it easier to distinguish persons from each other. However, even in this population, the majority of people score between a 7 and 9 on work ability (75% at T1). As expected, the increase in work ability and work functioning is significant on a group level, though only minimally. This increase directly after RTW plateaus towards week 8 after RTW as the total population scores are moving back towards the baseline score of 2 weeks after RTW. As hypothesized above, this may be attributable to the *deteriorated* group, who may have been signed off sick in a normal population, but remain in this population, because they have just returned to work. The calculated SDC and MIC may be generalizable to other heterogeneous populations in different workplaces, because the mean and standard deviation of the measurements in this study are known to be similar for different populations [18, 21, 33, 38, 41, 42]. However, it must be noted that in specific homogeneous populations, work ability and work functioning scores and their variation among the study population can be different, resulting in different MIC and SDC scores, and therefore, different thresholds for increases or decreases for that specific population.

Conclusion

Since the AUC for the work ability questions and the work-functioning instrument are inadequate, we advise refraining from using the single-item Work Ability Score to distinguish between individual workers who improve or deteriorate over the RTW process. The capacity of the work-functioning instrument to detect change in people returning to work is also inadequate, but better than the single-item work ability questions, especially for the *deteriorated* group. Because the MIC values of both instruments fall within that of measurement error, the SDC should be used to assess measurable, relevant change. This leads to the conclusion that 10–15% of the individuals in this study returning to work show an improvement greater than the SDC.

Acknowledgments: The authors thank the occupational health service for their cooperation, and all the respondents for participation in this research. The authors did not receive any funding or financial compensation for writing this article and do not have any conflicting interests.

Author contributions: AS, KN, MF and JS were all involved in the development of the study protocol and designing the informed consent and patient information form as well as the writing process. AS did the data collection and performed the statistical analyses and KN performed data checks.

Conflict of interest: The authors declare that they have no conflict of interest.

Ethical approval: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

OpenAccess: This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

REFERENCES

1. Schaafsma FG, Whelan K, van der Beek AJ, van der Es-Lambeek LC, Ojajarvi A, Verbeek JH. Physical conditioning as part of a return to work strategy to reduce sickness absence for workers with back pain. *Cochrane Database Syst Rev*. 2013(8):CD001822.
2. Schneider U, Linder R, Verheyen F. Long-term sick leave and the impact of a graded return-to-work program: evidence from Germany. *Eur J Health Econ*. 2016;17(5):629-43.
3. Hogelund J, Holm A, McIntosh J. Does graded return-to-work improve sick-listed workers' chance of returning to regular working hours? *J Health Econ*. 2010;29(1):158-69.
4. Viikari-Juntura E, Kausto J, Shiri R, Kaila-Kangas L, Takala EP, Karppinen J, et al. Return to work after early part-time sick leave due to musculoskeletal disorders: a randomized controlled trial. *Scand J Work Environ Health*. 2012;38(2):134-43.
5. Viikari-Juntura E, MacEachen E. Work modification as a treatment for low-back pain. *Scand J Work Environ Health*. 2015;41(3):219-21.
6. Franche RL, Krause N. Readiness for return to work following injury or illness: conceptualizing the interpersonal impact of health care, workplace, and insurance factors. *J Occup Rehabil*. 2002;12(4):233-56.
7. Ilmarinen J, Tuomi K, Seitsamo J, editors. New dimensions of work ability. International congress series; 2005: Elsevier.
8. Lerner D, Amick BC, 3rd, Rogers WH, Malspeis S, Bungay K, Cynn D. The Work Limitations Questionnaire. *Med Care*. 2001;39(1):72-85.
9. Kessler RC, Barber C, Beck A, Berglund P, Cleary PD, McKenas D, et al. The World Health Organization Health and Work Performance Questionnaire (HPQ). *J Occup Environ Med*. 2003;45(2):156-74.
10. Shikar R, Halpern MT, Rentz AM, Khan ZM. Development of the Health and Work Questionnaire (HWQ): an instrument for assessing workplace productivity in relation to worker health. *Work*. 2004;22(3):219-29.
11. van Roijen L, Essink-Bot ML, Koopmanschap MA, Bonsel G, Rutten FF. Labor and health status in economic evaluation of health care. The Health and Labor Questionnaire. *Int J Technol Assess Health Care*. 1996;12(3):405-15.
12. Endicott J, Nee J. Endicott Work Productivity Scale (EWPS): a new measure to assess treatment effects. *Psychopharmacol Bull*. 1997;33(1):13-6.
13. Reilly MC, Zbrozek AS, Dukes EM. The validity and reproducibility of a work productivity and activity impairment instrument. *Pharmacoeconomics*. 1993;4(5):353-65.
14. Burton WN, Conti DJ, Chen CY, Schultz AB, Edington DW. The role of health risk factors and disease on worker productivity. *J Occup Environ Med*. 1999;41(10):863-77.
15. Koopman C, Pelletier KR, Murray JF, Sharda CE, Berger ML, Turpin RS, et al. Stanford presenteeism scale: health status and employee productivity. *J Occup Environ Med*. 2002;44(1):14-20.
16. Lerner D, Amick BC, 3rd, Lee JC, Rooney T, Rogers WH, Chang H, et al. Relationship of employee-reported work limitations to work productivity. *Med Care*. 2003;41(5):649-59.

17. Boezeman EJ, Sluiter JK, Nieuwenhuijsen K. Measuring Work Functioning: Validity of a Weighted Composite Work Functioning Approach. *J Occup Rehabil.* 2015;25(3):537-42.
18. El Fassi M, Bocquet V, Majery N, Lair ML, Couffignal S, Mairiaux P. Work ability assessment in a worker population: comparison and determinants of Work Ability Index and Work Ability score. *BMC Public Health.* 2013;13:305.
19. Tuomi K, Ilmarinen J, Jahkola A, Katajarinne L, Tulkki A. Work ability index, 2nd revised edition. Helsinki: Finnish Institute of Occupational Health; 1998.
20. WMA. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA.* 2013;310(20):2191-4.
21. Gould R, Ilmarinen J, Järvisalo J, Koskinen S. Dimensions of work ability. Helsinki: Finnish Centre for Pensions (ETK) The Social Insurance Institution (Kela) National Public Health Institute (KTL) Finnish Institute of Occupational Health (FIOH); 2008.
22. Boezeman EJ, Nieuwenhuijsen K, de Bekker-Grob EW, van den Akker-van Marle ME, Sluiter JK. The relative importance of the domains of work functioning: evaluations of health-impaired employees, healthy employees, and employers. *J Occup Environ Med.* 2015;57(4):361-6.
23. de Vet H, Terwee, C., Mokkink, L., Knol, D. *Measurement in Medicine: A Practical Guide (Practical Guides to Biostatistics and Epidemiology)*: Cambridge University Press; 2011.
24. Crosby RD, Kolotkin RL, Williams GR. Defining clinically meaningful change in health-related quality of life. *J Clin Epidemiol.* 2003;56(5):395-407.
25. de Vet HC, Ostelo RW, Terwee CB, van der Roer N, Knol DL, Beckerman H, et al. Minimally important change determined by a visual method integrating an anchor-based and a distribution-based approach. *Qual Life Res.* 2007;16(1):131-42.
26. Fluss R, Faraggi D, Reiser B. Estimation of the Youden Index and its associated cutoff point. *Biom J.* 2005;47(4):458-72.
27. de Vet HC, Terwee CB, Ostelo RW, Beckerman H, Knol DL, Bouter LM. Minimal changes in health status questionnaires: distinction between minimally detectable change and minimally important change. *Health Qual Life Outcomes.* 2006;4:54.
28. de Vet HC, Terwee CB, Knol DL, Bouter LM. When to use agreement versus reliability measures. *J Clin Epidemiol.* 2006;59(10):1033-9.
29. Hosmer DW, Lemeshow S. *Applied Logistic Regression*, 2nd Ed. John Wiley and Sons, New York, NY 2000;Chapter 5.2.4:160-4.
30. van Kampen DA, Willems WJ, van Beers LW, Castelein RM, Scholtes VA, Terwee CB. Determination and comparison of the smallest detectable change (SDC) and the minimal important change (MIC) of four-shoulder patient-reported outcome measures (PROMs). *J Orthop Surg Res.* 2013;8:40.
31. Fritz JM, Hebert J, Koppenhaver S, Parent E. Beyond minimally important change: defining a successful outcome of physical therapy for patients with low back pain. *Spine (Phila Pa 1976).* 2009;34(25):2803-9.
32. Rysstad T, Roe Y, Haldorsen B, Svege I, Strand LI. Responsiveness and minimal important change of the Norwegian version of the Disabilities of the Arm, Shoulder and Hand

- questionnaire (DASH) in patients with subacromial pain syndrome. *BMC Musculoskeletal Disord.* 2017;18(1):248.
33. Roelen CA, van Rhenen W, Groothoff JW, van der Klink JJ, Twisk JW, Heymans MW. Work ability as prognostic risk marker of disability pension: single-item work ability score versus multi-item work ability index. *Scand J Work Environ Health.* 2014;40(4):428-31.
 34. Abma FI, van der Klink JJ, Bultmann U. The work role functioning questionnaire 2.0 (Dutch version): examination of its reliability, validity and responsiveness in the general working population. *J Occup Rehabil.* 2013;23(1):135-47.
 35. Corrigan PW, McCracken SG. Place first, then train: an alternative to the medical model of psychiatric rehabilitation. *Soc Work.* 2005;50(1):31-9.
 36. Young AE, Roessler RT, Wasiak R, McPherson KM, van Poppel MN, Anema JR. A developmental conceptualization of return to work. *J Occup Rehabil.* 2005;15(4):557-68.
 37. Arends I, van der Klink JJ, van Rhenen W, de Boer MR, Bultmann U. Predictors of recurrent sickness absence among workers having returned to work after sickness absence due to common mental disorders. *Scand J Work Environ Health.* 2014;40(2):195-202.
 38. van Schaaik A, Nieuwenhuijsen K, Frings-Dresen MHW, Sluiter JK. Reproducibility of work ability and work functioning instruments. *Occup Med (Lond).* 2018;68(2):116-9.
 39. Seing I, MacEachen E, Stahl C, Ekberg K. Early-return-to-work in the context of an intensification of working life and changing employment relationships. *J Occup Rehabil.* 2015;25(1):74-85.
 40. McGlothlin AE, Lewis RJ. Minimal clinically important difference: defining what really matters to patients. *JAMA.* 2014;312(13):1342-3.
 41. Schouten LS, Bultmann U, Heymans MW, Joling CI, Twisk JW, Roelen CA. Shortened version of the work ability index to identify workers at risk of long-term sickness absence. *Eur J Public Health.* 2016;26(2):301-5.
 42. de Zwart BC, Frings-Dresen MH, van Duivenbooden JC. Test-retest reliability of the Work Ability Index questionnaire. *Occup Med (Lond).* 2002;52(4):177-81.

4

Work ability and percentage of
hours worked related to
limitations in patients with
upper extremity
musculoskeletal disorders: a
cross-sectional cohort study

A. van Schaijk, K. Nieuwenhuijsen and M.H.W. Frings-Dresen

BMC Musculoskeletal Disorders, 2020, 21(1), 389-401.

ABSTRACT

Background: The aim of this study was to assess the relationship between self-reported work ability and hours worked at the current time in Upper Extremity Musculoskeletal Disorders (UEMSD) patients. To further investigate this relationship, the association of work ability and working hours with several limitations in daily and working life were explored.

Methods: In this cross-sectional cohort study, a questionnaire was sent out to members of the UEMSD patient organisation, containing self-reported work ability, questions on working hours and limitations in work due to UEMSD. Limitations were measured with the Disabilities of Arm Shoulder and Hand questionnaire, ShortForm-36 subscales, and common hand grasps or grips. Work ability was measured with the work ability score, while worked hours were operationalised as the percentage of hours worked compared to fulltime. The correlation between worked hours and work ability was tested with the Pearson correlation coefficient. Variance in work ability and the hours worked were explained by limitations and assessed with two linear regression analyses.

Results: Based on data of 794 respondents a moderate correlation was found between work ability and worked hours $r = 0.46$; 95% CI [0.40, 0.53]. Models including limitations explained 52 and 21% of total variance in work ability and worked hours, respectively. Variance in both can be explained by the degree of difficulties performing daily activities at work, limitations in daily activities as a consequence of health issues and the ability to perform a precision grip. Additionally, work ability can be explained by limitations at work and other daily activities due to physical health issues, while the percentage of hours can additionally be explained by the ability to grasp a large object with one hand, the ability to use a keyboard, and the subject's gender.

Conclusions: The number of worked hours does not fully match the work ability. Although they share three predictors, work ability and worked hours seem to be based on different aspects. Compared to work hours, work ability is more strongly related to limitations in daily activities and work. Taking self-reported work ability into account can improve the fit between work limitations and work hours.

Key words: Functioning, Limitations, Repetitive strain injury (RSI), UEMSD, Work ability appraisal, Working hours, WRUED

INTRODUCTION

Work ability reflects the extent to which people can do their job satisfactorily with respect to their job demands and their (physical and mental) health [1]. Work ability is an important factor in working life and is used as an indicator for the ability to perform work at a specific time, and can predict the future need for a work disability pension [2-7]. Although work ability is an abstract concept, it is seen as a standard for determining whether someone is able to work and is often used as a marker for the current ability of that person to perform in a job [8-10]. The predictive value of work ability has been extensively studied, however, the relationship with the current situation is less well-known. The relationship between the self-reported work ability, and the actual ability to work several hours at that same timepoint remains unknown.

Work ability can be determined in several ways. One of the ways to assess work ability in practice is with jobspecific testing in preventive medical examinations or observing work activities by occupational physicians [11]. Another more frequently used way to assess work ability is by letting workers appraise their own work ability [12]. In selfreported work ability, workers are asked to appraise their own work ability relating to their job demands at the current time. The first question of the Work Ability Index (WAI), also called the Work Ability Score (WAS), is often used for this purpose. The validity of the WAS has been established in relation to future sickness absences, but it is unclear whether the self-reported instrument WAS is a valuable and representative measure of what an employee can do at the current time in practice [5-7, 13]. Conceptually, one would expect a relationship between self-reported work ability and the actual hours that an employee has been able to do his/her job. If self-reported work ability is indeed representative of actual working hours, the WAS can aid occupational professionals in estimating the number of hours a worker can be required to work. When WAS and the hours worked do not concur, it is important to gain more information on why these measures differ and what causes the differences between the two estimates.

Workers with a chronic disease constitute a suitable population for studying the relationship between self-reported and actual work hours. Having a chronic disease with corresponding problems or limitations can lead to decreased mental and physical capacities, therefore threatening work ability and working hours [12, 14-21]. One such chronic condition is Upper Extremity Musculoskeletal Disorder (UEMSD) or Repetitive Strain Injury (RSI). In Europe, the one-year prevalence of neck and upper limb pain in the working population is almost 45% [22]. Although there is substantial variation in prevalence between countries, that number shows that at any given time, a major part of the workforce experiences musculoskeletal pain and accompanying limitations in the upper body. The pain and accompanying limitations performing daily tasks

caused by (chronic) UEMSD during daily and working life can decrease the current work ability in UEMSD patients and cause absences due to illness [23].

To determine to what extent the self-reported work ability corresponds to the ability to actually work at the current time in a population of UEMSD patients, it is vital to assess to which extent self-reported work ability is related to the number of hours worked. Therefore, we aimed to assess the coherence between self-reported work ability and the hours worked at the current time.

To further study the potential differences between work ability and worked hours, experienced limitations during daily activities and work might be important. Limitations during daily activities requiring the use of the upper extremities can also indicate problems during work hours because most work is being performed with the use of arms and hands [24]. Gold et al. found that the highest prevalence of chronic UEMSD complaints are mild complaints. In the group of workers with mild complaints, 76% continued working [25]. Even in the most severe cases of chronic UEMSD, 50% of patients continued working. This means that a large group of chronic UEMSD patients is still engaged in work activities while experiencing complaints of the upper extremities, leading to limitations during daily activities. It is unknown to what extent these limitations influence both self-reported work ability and hours worked in a different way. Limitations occur when a worker is unable to perform a certain daily or work activity. These limitations determine the ability to meet certain job demands.

Insight into how limitations during work are related to work ability and worked hours is important because it can aid occupational professionals in adapting the work or workplace to better fit the ability to perform work of a worker with UEMSD complaints. However, it is still unknown what type of limitations during daily and work activities lead to decreased work ability and a decreased number of working hours in UEMSD patients. This indicates a need for studying work ability and the hours worked and their relation to UEMSD-related limitations. In summary, our aim is to assess the relationship between self-reported work ability and the hours worked at the current time and additionally assess how limitations in daily and work life can explain variance in work ability and the amount of hours worked in UEMSD patients. These aims lead to the following two research questions:

What is the relationship between self-reported work ability and the hours worked at the current time in UEMSD patients? And

How are self-reported work ability and the hours worked at the current time related to limitations and activities in daily life?

METHODS

Design, participants, setting and procedure

In this cross-sectional study, a questionnaire was sent out in April 2005 to all 3250 members of the Dutch National UEMSD patient association, irrespective of the severity of complaints. Participants that were unemployed or on disability pension were excluded. Each questionnaire ('RSI: Complaints, health and possibilities') was accompanied by a participant information form. Filled out forms were returned anonymously and directly to the research institute by an enclosed return envelope. After 1 week, the members received a reminder to fill out and return the questionnaire. All patients participated on a voluntary basis and data was stored anonymously. The study was conducted in accordance with the declaration of Helsinki [26]. The research proposal for this secondary analyses of the data was submitted to and approved by the Medical Ethical Committee of the Academic Medical Center, which decreed that a comprehensive evaluation was not required as this study was not subject to the Medical Research Involving Human Subjects Act (W19_211#19.256).

Instruments

Respondent characteristics and basic information were gathered by a questionnaire. Besides the basic information about work, several instruments were used to measure the work hours, complaints and the work ability of UEMSD-patients [27]. The questionnaire further contained questions on (health) complaints, health and health perceptions, work and limitations [28].

The primary outcome, self-reported work ability, was measured with the Work Ability Score (WAS), the first question of the Work Ability Index (WAI). This single item WAI was included to assess the appraised work ability at the current time (according to the patient) on a scale of 0 to 10. This single item question showed sufficient convergent validity to the complete WAI [29]. A score of 0 meant *not being able to work* and a score closer to 10 meant increasing work ability compared to *their lifetime best* (score 10). Hours worked, was operationalized as the percentage of hours worked compared to fulltime employment (36 h). To calculate this variable, participants filled out the number of hours they actually worked in the past week. This value was chosen rather than the percentage of worked hours compared to the contract hours because multiple respondents were on a 0-h or flexible contract basis while working multiple hours. Many other respondents worked a numerous amount more than their contract hours, leading to abnormal values, which made the contract hours unsuitable for these analyses.

To answer our second research question, the limitations were operationalised in three ways: The degree of difficulties in performing daily activities in the last week was assessed by the complete Disability of the Arm, Shoulder and Hand (DASH) questionnaire, including the work and sport module [30, 31]. The DASH scores are scaled with 0 indicating no disability and 100 indicating most disability.

A second way to assess limitations was to ask whether patients have problems performing certain frequent grips of both hand and wrist. Illustrations of these common grasps and movements were included in the questionnaire, accompanied by a question on the limitations of performing these in daily work life. These illustrations are shown in the Appendix. These grasps and activities were, among others, the cylindrical grip, the lateral key grip, the precision grip or spherical grip, the hook grip or medium wrap, and common activities at work (i.e. using a computer mouse or keyboard, pushing with the hand or finger, and bending the wrist) [27, 32]. An example question next to an illustration was: When you have to perform activities during work using your hand as displayed in the illustration above, do you experience problems? The answer options to these questions were: 'Yes, almost always', 'Yes, sometimes', 'No, no problems', or 'No, I don't have to do this'. The outcome was divided into either 'yes, almost always' scored as 1, and all the other answer options grouped together scored as 0. This means that a score of 1 indicates problems with performing this movement compared to 'no' or 'only sometimes' problems, which was coded as 0.

The third way was to assess limitations in the health-related quality of life by using the Short-Form-36 (SF-36) [33-35]. Of the nine subscales, only 'physical functioning', 'pain', and 'physical role functioning' were used in the analysis. The subscale of physical functioning measured 'limitations in daily activities as a consequence of health issues' during the last 4 weeks. The pain and physical role functioning subscales represent 'pain and limitations due to pain' and 'limitations in work and other daily activities due to physical health issues', respectively [35]. These scores are measured on a scale between 0 and 100, where a score closer to 100 indicates a better general health status. Averages for a Dutch population aged 35–44 were 90 (± 14.4), 84 (± 21.7) and 83 (± 32.0) for physical functioning, pain and physical role functioning, respectively [35].

Additionally, demographic variables, among which age and gender were included, were used in the analyses to describe the study population and differentiate possible influencing characteristics. Males were coded as 1 where females were coded as 0.

Statistical analyses

Before the analyses were performed, cases with over 50% missing variables were deleted. Cases with one or both primary variables missing were also deleted from the

analyses. Before the analyses, the MCAR test was executed to establish the randomness of missing values. If the MCAR test was significant, no imputation has taken place. First, the self-reported work ability was related to the percentage of hours worked using a Pearson correlation analysis. In this analysis, the coherence between both outcomes was analysed.

Second, the variables, SF-36 subscales, DASH scores, the limitations in frequent grasps of hand and wrist, and age and gender were related to both self-reported work ability, and worked hours. In these analyses, the self-reported work ability and percentage of hours worked were entered as dependent variables, and the SF-36 subscales, the DASH scores and the limitations in frequent grasps of hand and wrist were included as independent variables. Two separate models were created, one for self-reported work ability and the other for the percentage of worked hours compared to fulltime employment, with only significant contributors to the explained variance. This makes assessing the limitations that explain the most variance in work ability and the percentage of hours worked possible. In this way, the most probable limitations that cause people to not be able to fully function in their job can be identified. Variance in work ability and the hours worked were explained by these limitations and assessed with two linear regression analyses. A model was created using backward selection. Non-significant variables were sequentially deleted from the model, creating a new model, until all remaining variables in the model had a p-value smaller than 0.05, and the model was complete.

As a final check, we tested both models for heteroscedasticity (through a visual inspection of the residual plots) and collinearity (through collinearity statistics: tolerance and Variance Inflation Factor (VIF) score). A tolerance ≤ 0.4 and a corresponding VIF score over 2.5 was set as cause for concern of collinearity, with tolerance calculated as $1-R^2$ and VIF as $1/\text{tolerance}$. The statistical analyses were done performing multiple regression analyses in IBM SPSS Statistics version 25.

RESULTS

Population characteristics

In total, 1129 UEMSD patients responded (35%), of which 794 were eligible for secondary analyses as they filled out both questions regarding work ability and hours worked and were currently employed [28]. Individuals who received disability pension, or did not have a job were excluded from the sample. The inclusion flowchart is shown in Fig. 1. The MCAR test showed that some missing values are not randomly missing. As a result, no imputation took place, and no data is listwise deleted.

Figure 1: Flowchart of inclusion process and number of included participants.

The mean age at the time of participation was 40.0 (± 8.5) years and one-third of all participants were male. Over 90% of included participants reported they were still working. The mean duration of complaints was 5.6 (± 3.1) years, and the mean extensiveness of pain regions was 5.8 (± 3.6) on a scale of 0 to 16 regions. Participant characteristics (of all respondents) are shown in Table 1. Population characteristics divided per job category according to the International Standard Classification of Occupations [36] are shown in the Appendix.

Table 1: Percentage (%) or mean $\pm$ standard deviation (SD) of participant characteristics of work ability and percentage of hours worked.

	%	Mean	SD	N
Age at time of participation		40.0	8.5	794
Gender male	37%			794
Education level				794
Primary school	1%			
Lower vocational education	1%			
Secondary vocational education	26%			
Higher vocational education	34%			
University	32%			
Other	6%			
% participants with pain	80%			794
% participants with stiffness	49%			794
% participants with tingling feelings or numbness	42%			794
Main cause of the start of complaints				742
A single cause cannot be identified	17%			
Overload during leisure	2%			
Short-term overload during work	5%			
Long-term overload during work	70%			
Accident, strain	1%			
Other(e.g. stress, posture, furniture)	53%			
Complaint duration in years		5.6	3.1	755
Work ability score (WAS) (0–10)		6.4	2.0	794
% worked of Fulltime employment (0–100)		77.0	35.0	794
DASH score (0–100)		28.3	19.0	668
DASH work module (0–100)		41.0	25.8	784
SF-36 subscale physical functioning (0–100)		80.0	15.3	780
SF-36 subscale pain (0–100)		59.3	20.1	793
SF-36 subscale physical role functioning (0–100)		41.9	39.3	788

The percentage of hours worked correlates significantly ($p < 0.001$) with the self-reported work ability ($n = 794$). The correlation coefficient is $r = 0.46$; 95% CI [0.40, 0.53] $p < 0.001$, which is considered a modest or moderate correlation [37].

Results of the univariate analyses are shown in the Appendix. In the multivariate analyses, the prediction model for the WAS contained four variables. These variables were able to explain 52% of total variance in WAS. The DASH work module score, the SF-36 physical functioning and physical role functioning domains and the precision grip were predictors in this model. Multicollinearity was not of concern in the WAS model as there were no VIF-scores over 2.5.

The model that explained 21% of total variance in percentage of hours worked contained five variables, namely: SF-36 physical functioning domain, the DASH score, the DASH work module, limitations using a keyboard and gender. Multicollinearity was a concern in the worked hours model where the VIF-score of the DASH was 3.4. When this variable was removed from the model, there was no more concern for collinearity. However, the variables ‘ability to hold a precision grip’ and ‘ability to grasp a large object with one hand’ became significant contributors and were added to the model again in accordance with the methods. The explained variance of the new model, without the DASH score, but including the re-entered variables as predictors, is also 21%. A higher score of the B-coefficient means a more positive contribution to the work ability score. Both final models are displayed in Tables 2 and 3. A plot of the residuals was analysed for a check on heteroscedasticity, but there were no reasons to believe this was of concern in either model.

Table 2: Model with statistically significant predictors of explained variance in WAS.

Model	Unstandardized Coefficients			95% Confidence Interval for B	
	B	Std. Error	p-value	Lower Bound	Upper Bound
Constant	5.487	0.592	<0.001	4.322	6.652
DASH work module	-0.034	0.004	<0.001	-0.042	-0.025
SF36 physical functioning	0.025	0.006	<0.001	0.014	0.037
SF36 physical role functioning	0.008	0.003	0.002	0.003	0.014
Precision grasp	-0.489	0.215	0.024	-0.913	-0.065

Explained variance of this model: $R^2=0.519$

Table 3: Model with statistically significant predictors of variance in worked hours.

Model	Unstandardized Coefficients			95% Confidence Interval for B	
	B	Std. Error	p-value	Lower Bound	Upper Bound
Constant	52.439	11.561	<0.001	29.725	75.153
DASH work module	-0.418	0.068	<0.001	-0.550	-0.285
SF36 physical functioning	0.318	0.115	0.006	0.092	0.544
Precision grip	13.007	4.210	0.002	4.737	21.278
Grasping a large object with one hand	-9.092	4.622	0.050	-18.174	-0.011
Using a keyboard	13.539	5.020	0.007	3.677	23.402
Gender	14.366	3.030	<0.001	8.412	20.320

Explained variance of this model: $R^2=0.207$

In this model, a higher SF-36 score, having problems with using a keyboard, having problems performing a precision grip and being male resulted in a higher percentage of worked hours, while a higher score on the DASH work module and limitations in grasping a large object with one hand predict a lower percentage of worked hours in individuals. In the univariate analyses, limitations using a keyboard and the ability to hold a precision grip are not significant. However, in the multiple regression, these variables become significant indicating suppression effects (or negative confounding) between variables. Limitation in using a keyboard is unsuppressed by the presence of the DASH work module.

DISCUSSION

Self-reported work ability and the percentage of hours worked correlate moderately, with a higher work ability score corresponding to a higher percentage of hours worked. Despite this moderate correlation, work ability and worked hours are explained to a different extent by limitations. Explained variance by limitations is higher for work ability than for worked hours. The degree of difficulties performing daily activities in work (DASH work module), the limitations in daily activities as a consequence of health issues (SF-36 physical functioning) and limitation in performing a precision grasp or grip explain the variance for both selfreported work ability and worked hours. The limitations in work and other daily activities due to physical health issues (SF-36 physical role functioning) explain variance in the self-reported work ability model, but not in the model of percentage of worked hours. Variance in the

percentage of worked hours is additionally explained by the ability to grasp a large object with one hand, limitations in using a keyboard and gender.

The WAS may be informative when establishing a reintegration plan since its relation to limitations is stronger than for the number of worked hours at the current time. Variance in the current work hours are only modestly explained by variables. The work hours do not seem related to many aspects of work and health included in our study. However, the WAS is an individual's estimation of their ability to meet their job demands that takes the limitations that an employee experiences into account. Consulting WAS can help to better match work hours to the limitations in work and daily life in order to prevent sick leave. By monitoring WAS, it is possible to timely adjust working hours as a preventive measure.

Gender did not explain variance in work ability in this population, while it did play a role in the estimation of working hours. Male participants have a 14% point higher percentage of working hours when experiencing the same limitations. The difference between men and women in the percentage of working hours might be explained by the fact that the overall number of women working fulltime is lower—28% vs. 58% of men. In this study, fulltime was set at 36 h a week. The difference between men and women might be explained by the fact that more women work part-time than men [38]. This, in turn, can explain the difference in explained variance by gender in the number of hours worked at the current time and the lower mean percentage of worked hours in women.

Comparison with other studies

Work ability has not been related to current work hours in populations with limitations in studies before. We could not identify other studies that studied this specific relationship. Leijten et al. (2014) did find that WAS and productivity at work (0–10), a concept closely related to working hours, were positively correlated in older employees with chronic diseases ($r = 0.23$, $p < 0.01$) [39]. Whereas Leijten studied the influence of chronic limitations on work ability 1 year later, our study provides knowledge on limitations at the current time in relation to work ability and worked hours at the same moment. Another study by Oakman et al. (2019) studied the relationship between the number of pain sites and work ability trajectories [40]. They found that the number of pain sites was associated with a decreasing work ability trajectory over time. A study by Kamaleri et al. (2009) showed the strong predictive power of the number of pain sites on work disability 14 years later [41]. In our study, pain was included as a potential predictor of workability, but was not retained in the final model. This means that not only pain predicts work ability, but that limitations in work might help better predict current work ability. However, these limitations are only moderately related to the hours worked. Although future work disability and

current work ability represent different concepts, we would argue—based on our study—that not only the number of pain sites or number of limitations is important, but also the type of limitations in cohesion with the type of work.

Strengths and limitations

A strength of this study is that we studied a population with a large variation in work ability due to the existence of limitations/health problems during work and daily life. This study population showed a larger variance in WAS in comparison to a general working population [42]. When studying the coherence between work ability and hours worked, a population with a large variation in work ability would be beneficial in order to study correlations between the two variables.

A limitation of this study is that the type of work that participants do was not taken into account. However, given the fact that participants have UEMSD complaints, it is plausible that the work of these participants involves working with arms and hands. All questions regarding movements and hours concerned the current job of the respondent, as is the assessment of self-reported work ability. If participants did not have to do certain activities during work, these were regarded as not being a limitation, and therefore not affecting work, and this should therefore not be a large source of bias. However, we expect some selection bias. Although the entire population of the UEMSD patient association received an invitation to participate in this study, it can be expected that the more severe cases were represented more strongly in the UEMSD patients association.

An aspect that might explain the difference in explained variance by limitations between work ability and the percentage of hours worked is the common method principle due to the cross-sectional nature of this study [43]. This principle suggests that two measurements are more strongly related to each other than a third variable due to the type of measurement. In this case, the limitations and WAS are both assessed by a questionnaire filled in by the workers themselves. The number of hours, however, is more factual and not only determined by the workers themselves, as the occupational physician and their employer play a role in determining the optimal number of work hours.

The outcome of hours worked was used to represent the impact of UEMSD-related limitations on the ability to perform work. However, since the variance in worked hours is only explained to a moderate extent by the factors in our model, we can conclude that hours worked is influenced by other factors than just these limitations. The family situation, social security system, psychological state, need for income generation or informal care can all influence the reason to work [44]. This can explain

the moderate correlation between work ability and the hours worked at the current time.

Implications

It seems that the number of hours worked does not fully match with self-reported work ability. When determining the work hours of a UEMSD patient, it can be profitable to better align these work hours with work ability, as workers take their limitations at work into account when assessing their own work ability. In this way, the working hours of a UEMSD patient correspond better with their self-reported ability to meet the job demands. Limitations are individual and job-specific. Workers can estimate the influence of limitations on their work best because they know every details about their job demands in relation to the limitations they encounter. Therefore, the appraisal of their work ability can be used to better match work hours to their limitations in order to prevent workers from going on sick leave. In the Netherlands for instance, an occupational physician can be consulted when a worker is unable to meet the work demands. This occupational physician advises in consultation with the employee and employer how and if work can be continued. One potential outcome of the consultation is that the employer is advised to change the work demands or work hours. When this proves impossible, the occupational physician can advise other jobs within or outside the company, that can be met by the employee without compromising their health.

Occupational physicians can use the appraisal of UEMSD patients' work ability to better match the ability to meet job demands with the working hours, keeping the limitations in mind. Occupational physicians can advise their patients to have their work hours and demands match their abilities. The results of this study are relevant for occupational professionals such as occupational physicians as they give an indication of which functional limitations affect the work ability most, for instance the ability to hold a cylindrical grip, precision grip and hook grip. The importance of the capability to perform certain actions can aid occupational professionals in determining the next steps in the treatment, to decrease the impact of limitations on work life. To confirm the additional value of taking work ability into account when determining work hours, a longitudinal study could be performed. Based on the univariate analyses, limitations in bending the wrist are quite important in some individuals, but not significant on a group level. For which individuals this limitation is of importance needs to be studied in future research. Future research can focus on differences between types of job demands. A division can be made between more physical demands using shoulders, or more desk work using fingers and hands. With this information, future work disability or sick leave might be prevented by using the right strategies to ensure work hours match work ability. Furthermore, the ability to use a

keyboard, the ability to hold a precision grip and the ability to grasp a large object with one hand are only related to the amount of hours worked when the DASH work module was present in the final model, and the B-coefficient changed direction. Since this indicates paradoxical confounding, the relationship between these variables needs to be investigated further.

Conclusions

Self-reported work ability is moderately correlated with the percentage of hours worked at the current time. Variance of both aspects is explained by limitations in a different way. Work ability is more strongly related to limitations in work than work hours themselves. Taking self-reported work ability into account can improve the fit between work limitations and work hours.

List of abbreviations

WAI: Work Ability Index

WAS: Work ability score

RSI: Repetitive Strain Injury

UEMSD: Upper Extremity Musculoskeletal Disorders

DASH: Disability of the Arm Shoulder and Hand

SF-36: Short Form 36

Declarations:

Ethics approval and consent to participate: The research proposal was submitted to and approved by the Medical Ethical Committee of the Academic Medical Center, which decreed that a comprehensive evaluation was not required as this study was not subject to the Medical Research Involving Human Subjects Act (W19_211#19.256). In the patient information letter was included that consent for analysing the data was given by returning the questionnaire. Therefore, by returning the questionnaire participants gave implicit consent for analysis of the entered data.

Consent for publication: Not applicable

Availability of data and material: Data are available at the first author upon reasonable request.

Competing interests: All authors declare that they have no conflict of interest.

Funding: No funding was obtained for this study.

Authors' contributions: AS was involved in the design, conceptualisation, analysis, interpretation and writing of the manuscript. KN was involved in the conceptualisation, data checks and review of the drafted manuscript. MF was involved

in the design, conceptualisation, coordination and review of the drafted manuscript. All authors read and approved the final manuscript.

Acknowledgments: The authors thank all participants in this study and the Dutch RSI patient association for their cooperation.

Appendix 1

a. Cylindrical grip or power grip

b. Lateral key grip

c. Precision grip or spherical grip

d. Hook grip or medium wrap

e. Using a computer mouse

f. Using a keyboard

g. Pushing with a flat hand

h. Pushing with one finger

i. Bending the wrist

j. Grasping a large object with one hand

Fig. 2: Illustrations of common grasps of hand and wrist in the questionnaire; When you have to perform activities during work using your hand as displayed in the illustration above, do you experience problems? The answer options to these questions were: 'Yes, almost always', 'Yes, sometimes', 'No, no problems', or 'No, I don't have to do this'. (Illustrations derived from the questionnaire 'RSI: Complaints, health and possibilities' with permission to publish.)

Table 4: Results of the univariate analyses of explained variance in Work Ability and hours worked (n=794).

Variable	Work Ability Score (0–10)			Percentage worked of fulltime (0–100)		
	p-value	Adj R ²	B	p-value	Adj R ²	B
Cylindrical grip	<0.001	0.12	–1.69	<0.001	0.04	–16.53
Lateral key grip	<0.001	0.09	–1.58	0.003	0.01	–11.41
Precision grip	<0.001	0.10	–1.61	0.062	0.00	–6.95
Hook grip	<0.001	0.10	–1.36	<0.001	0.03	–12.45
Using a computer mouse	0.001	0.02	–1.04	0.437	0.00	4.362
Using a keyboard	0.002	0.02	–0.92	0.683	0.00	2.16
Pushing with a flat hand	<0.001	0.07	–1.51	0.008	0.01	–10.84
Pushing with one finger	<0.001	0.02	–0.91	<0.001	0.02	–7.68
Bending the wrists	0.082	0.14	–2.44	0.279	0.02	–20.15
Grasping a large object with one hand	<0.001	0.07	–1.43	<0.001	0.03	–17.88
DASH	<0.001	0.36	–0.06	<0.001	0.09	–0.56
DASH work module	<0.001	0.42	–0.05	<0.001	0.08	–0.39
DASH hobby/sport module	<0.001	0.20	–0.03	<0.001	0.05	–0.26
SF-36 % Physical functioning	<0.001	0.25	0.07	<0.001	0.10	0.75
SF-36 % Physical role functioning	<0.001	0.29	0.03	<0.001	0.06	0.22
SF-36 % Pain	<0.001	0.34	0.06	<0.001	0.07	0.48
Age in 2005	0.008	0.01	–0.02	0.035	0.00	–0.31
Gender	0.002	0.01	0.45	<0.001	0.08	20.02

Table 5: International Standard Classification of Occupations (ISCO) of participants for males and females (n=794).

	Female		Male		Total	
	<i>N</i>	Percent	<i>N</i>	Percent	<i>N</i>	Percent
Cat 1. Managers	59	11.8	44	15.0	103	13.0
Cat 2. Professional	78	15.6	51	17.4	129	16.2
Cat 3. Technicians and associate professionals	18	3.6	67	22.9	85	10.7
Cat 4. Clerical support workers	205	40.9	76	25.9	281	35.4
Cat 5. Service and sales workers	44	8.8	19	6.5	63	7.9
Cat 7 or 8. Craft and related trades workers & plant and machine operators, and assemblers	64	12.8	20	6.8	84	10.6
Student	8	1.6	8	2.7	16	2.0
Not classifiable	25	5.0	8	2.7	33	4.2
Total	501	100	293	100	794	100

REFERENCES

1. Ilmarinen J, Tuomi K, Klockars M. Changes in the work ability of active employees over an 11-year period. *Scand J Work Environ Health*. 1997;23 Suppl 1:49-57.
2. Jaaskelainen A, Kausto J, Seitsamo J, Ojajarvi A, Nygard CH, Arjas E, et al. Work ability index and perceived work ability as predictors of disability pension: a prospective study among Finnish municipal employees. *Scand J Work Environ Health*. 2016;42(6):490-9.
3. Lindberg P, Josephson M, Alfredsson L, Vingard E. Comparisons between five self-administered instruments predicting sick leaves in a 4-year follow-up. *Int Arch Occup Environ Health*. 2009;82(2):227-34.
4. Bethge M, Gutenbrunner C, Neuderth S. Work Ability Index predicts application for disability pension after work-related medical rehabilitation for chronic back pain. *Arch Phys Med Rehabil*. 2013;94(11):2262-8.
5. Lundin A, Kjellberg K, Leijon O, Punnett L, Hemmingsson T. The Association Between Self-Assessed Future Work Ability and Long-Term Sickness Absence, Disability Pension and Unemployment in a General Working Population: A 7-Year Follow-Up Study. *J Occup Rehabil*. 2016;26(2):195-203.
6. Sell L, Bultmann U, Rugulies R, Villadsen E, Faber A, Sogaard K. Predicting long-term sickness absence and early retirement pension from self-reported work ability. *Int Arch Occup Environ Health*. 2009;82(9):1133-8.
7. Ahlstrom L, Grimby-Ekman A, Hagberg M, Dellve L. The work ability index and single-item question: associations with sick leave, symptoms, and health—a prospective study of women on long-term sick leave. *Scand J Work Environ Health*. 2010;36(5):404-12.
8. Gould R, Ilmarinen J, Järvisalo J, Koskinen S. Dimensions of work ability. 2008.
9. Tuomi K, Huuhtanen P, Nykyri E, Ilmarinen J. Promotion of work ability, the quality of work and retirement. *Occup Med (Lond)*. 2001;51(5):318-24.
10. Notenbomer A, Groothoff JW, van Rhenen W, Roelen CA. Associations of work ability with frequent and long-term sickness absence. *Occup Med (Lond)*. 2015;65(5):373-9.
11. van Schaaik A, Boschman JS, Frings-Dresen MH, Sluiter JK. Appraisal of work ability in relation to job-specific health requirements in ambulance workers. *Int Arch Occup Environ Health*. 2017;90(1):123-31.
12. Roelen CA, van Rhenen W, Groothoff JW, van der Klink JJ, Twisk JW, Heymans MW. Work ability as prognostic risk marker of disability pension: single-item work ability score versus multi-item work ability index. *Scand J Work Environ Health*. 2014;40(4):428-31.
13. Kinnunen U, Natti J. Work ability score and future work ability as predictors of register-based disability pension and long-term sickness absence: A three-year follow-up study. *Scand J Public Health*. 2018;46(3):321-30.
14. van den Berg S, Burdorf A, Robroek SJW. Associations between common diseases and work ability and sick leave among health care workers. *Int Arch Occup Environ Health*. 2017.

15. Bugajska J, Sagan A. Chronic musculoskeletal disorders as risk factors for reduced work ability in younger and ageing workers. *Int J Occup Saf Ergon*. 2014;20(4):607-15.
16. Koolhaas W, van der Klink JJ, de Boer MR, Groothoff JW, Brouwer S. Chronic health conditions and work ability in the ageing workforce: the impact of work conditions, psychosocial factors and perceived health. *Int Arch Occup Environ Health*. 2014;87(4):433-43.
17. de Vries HJ, Reneman MF, Groothoff JW, Geertzen JH, Brouwer S. Self-reported work ability and work performance in workers with chronic nonspecific musculoskeletal pain. *J Occup Rehabil*. 2013;23(1):1-10.
18. Bubas M, Milosevic M, Delic-Brkljacic D. Occupational diseases, working ability and employment status in the working population of Croatia. *Coll Antropol*. 2008;32(3):677-80.
19. Oberlinner C, Yong M, Nasterlack M, Pluto RP, Lang S. Combined effect of back pain and stress on work ability. *Occup Med (Lond)*. 2015;65(2):147-53.
20. Alavinia SM, van den Berg TI, van Duivenbooden C, Elders LA, Burdorf A. Impact of work-related factors, lifestyle, and work ability on sickness absence among Dutch construction workers. *Scand J Work Environ Health*. 2009;35(5):325-33.
21. Lahelma E, Laaksonen M, Lallukka T, Martikainen P, Pietilainen O, Saastamoinen P, et al. Working conditions as risk factors for disability retirement: a longitudinal register linkage study. *BMC Public Health*. 2012;12:309.
22. Farioli A, Mattioli S, Quagliari A, Curti S, Violante FS, Coggon D. Musculoskeletal pain in Europe: the role of personal, occupational, and social risk factors. *Scand J Work Environ Health*. 2014;40(1):36-46.
23. Feleus A, Miedema HS, Bierma-Zeinstra SM, Hoekstra T, Koes BW, Burdorf A. Sick leave in workers with arm, neck and/or shoulder complaints; defining occurrence and discriminative trajectories over a 2-year time period. *Occup Environ Med*. 2017;74(2):114-22.
24. van den Heuvel SG, Ijmker S, Blatter BM, de Korte EM. Loss of productivity due to neck/shoulder symptoms and hand/arm symptoms: results from the PROMO-study. *J Occup Rehabil*. 2007;17(3):370-82.
25. Gold JE, Piligian G, Glutting JJ, Hanlon A, Frings-Dresen MH, Sluiter JK. Cluster analysis of symptoms among patients with upper extremity musculoskeletal disorders. *J Occup Rehabil*. 2010;20(4):526-36.
26. WMA. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*. 2013;310(20):2191-4.
27. Sluiter JK, Rest KM, Frings-Dresen MH. Criteria document for evaluating the work-relatedness of upper-extremity musculoskeletal disorders. *Scand J Work Environ Health*. 2001;27 Suppl 1:1-102.
28. Sluiter JK, Frings-Dresen MH. Quality of life and illness perception in working and sick-listed chronic RSI patients. *Int Arch Occup Environ Health*. 2008;81(4):495-501.

29. El Fassi M, Bocquet V, Majery N, Lair ML, Couffignal S, Mairiaux P. Work ability assessment in a worker population: comparison and determinants of Work Ability Index and Work Ability score. *BMC Public Health*. 2013;13:305.
30. Veehof MM, Slegers EJ, van Veldhoven NH, Schuurman AH, van Meeteren NL. Psychometric qualities of the Dutch language version of the Disabilities of the Arm, Shoulder, and Hand questionnaire (DASH-DLV). *J Hand Ther*. 2002;15(4):347-54.
31. van Eck ME, Lameijer CM, El Moumni M. Structural validity of the Dutch version of the disability of arm, shoulder and hand questionnaire (DASH-DLV) in adult patients with hand and wrist injuries. *BMC Musculoskelet Disord*. 2018;19(1):207.
32. Liu X, Zhan Q. Description of the human hand grasp using graph theory. *Med Eng Phys*. 2013;35(7):1020-7.
33. Brazier JE, Harper R, Jones NM, O'Cathain A, Thomas KJ, Usherwood T, et al. Validating the SF-36 health survey questionnaire: new outcome measure for primary care. *BMJ*. 1992;305(6846):160-4.
34. Aaronson NK, Muller M, Cohen PD, Essink-Bot ML, Fekkes M, Sanderman R, et al. Translation, validation, and norming of the Dutch language version of the SF-36 Health Survey in community and chronic disease populations. *J Clin Epidemiol*. 1998;51(11):1055-68.
35. Van der Zee K, Sanderman R. Het meten van de algemene gezondheidstoestand met de RAND-36, een handleiding [Measuring the general health with the RAND-36, a manual]. In: SHARE RI, editor.: Groningen: Northern Centre for Health Care Research, University of Groningen, the Netherlands; 1993. p. 28.
36. International Labour Office. International standard classification of occupations ISCO-08 - Structure, group definition and correspondence tables. Geneva 2012. 2010 p.
37. Taylor R. Interpretation of the correlation coefficient: a basic review. *Journal of diagnostic medical sonography*. 1990;6(1):35-9.
38. Damaske S, Frech A. Women's Work Pathways Across the Life Course. *Demography*. 2016;53(2):365-91.
39. Leijten FR, van den Heuvel SG, Ybema JF, van der Beek AJ, Robroek SJ, Burdorf A. The influence of chronic health problems on work ability and productivity at work: a longitudinal study among older employees. *Scand J Work Environ Health*. 2014;40(5):473-82.
40. Oakman J, Neupane S, Prakash KC, Nygard CH. What Are the Key Workplace Influences on Pathways of Work Ability? A Six-Year Follow Up. *Int J Environ Res Public Health*. 2019;16(13).
41. Kamaleri Y, Natvig B, Ihlebaek CM, Bruusgaard D. Does the number of musculoskeletal pain sites predict work disability? A 14-year prospective study. *Eur J Pain*. 2009;13(4):426-30.
42. van Schaaik A, Nieuwenhuijsen K, Frings-Dresen MHW, Sluiter JK. Reproducibility of work ability and work functioning instruments. *Occup Med (Lond)*. 2018;68(2):116-9.

43. Podsakoff PM, MacKenzie SB, Lee JY, Podsakoff NP. Common method biases in behavioral research: a critical review of the literature and recommended remedies. *J Appl Psychol.* 2003;88(5):879-903.
44. de Vries HJ, Brouwer S, Groothoff JW, Geertzen JH, Reneman MF. Staying at work with chronic nonspecific musculoskeletal pain: a qualitative study of workers' experiences. *BMC Musculoskelet Disord.* 2011;12:126.

5

Mediating Factors for the Relationship Between Stress and Work Ability over Time in Young Adults

A. van Schaaik, A. Noor Baloch, S. Thomée, M.H.W. Frings-Dresen, M. Hagberg and
K. Nieuwenhuijsen

International Journal of Environmental Research and Public Health, 2020, 17(7),
2530.

ABSTRACT

Aim: Stress can affect work ability. The aim of this study was to identify how this pathway is mediated over time in young adults.

Methods: Participants of the Work Ability in Young Adults cohort were selected. A theoretical framework was built, which led to a statistical model. Selected dimensions of mediators were recovery, work demands affecting private life, feelings of control over private life, and physical activity in leisure time. A quadruple serial mediation model was built with four mediators.

Results: The total effect of stress on work ability was -0.3955 , 95% CI $[-0.4764, -0.3146]$. The total indirect effect amounted to 81% with an effect of -0.3182 , 95% CI $[-0.3750, -0.2642]$. The relationship between stress and Work Ability Score five years later in young adults was mediated by stress five years later, work demands affecting private life, feelings of control over private life and feeling well-rested upon waking.

Conclusions: These results indicate that work demands affecting private life and feelings of control over private life are important mediators of the relationship between stress and work ability in young adults. A well-balanced relationship between work and private life can counteract the influence of stress on work ability in this age group.

Keywords: Mediation; sustainable employability; work ability score; work-private life balance

INTRODUCTION

Work ability is the extent to which people can do their job satisfactorily without doing harm to their mental and physical health and is an important indicator of sustainable employability of workers [1]. Low work ability has been known to be a predictor of absenteeism and early retirement [2-5]. Stress is one of the most common occupational health problems that influences work ability. In recent years, workloads have increased, which has been accompanied by a growing number of workers reporting severe stress complaints. In Europe, the prevalence of men and women reporting to have work-related stress “always” or “most of the time” is 26% and 27%, respectively [6].

Ilmarinen (2009) stated that young adults need special attention when it comes to work ability. He indicated that it is important to ensure good work ability because at the start of working life workers face high expectations [1]. However, the work ability of young workers is at risk, as it has been reported that this age group has the highest prevalence of stress complaints [6, 7]. Young workers are people below the age of 35 [8]. An important feature of young workers is that their working career has just begun. The transition from studies to work in young people may be laden with factors that can affect their health [9]. In this life phase they are often also starting families, for which they need to realize a balance between work and private life. Stress complaints at an early stage of working life can lead to burnout, depression and unfavorable employment outcomes at a later age, according to the life course perspective [10, 11]. This life course perspective, an emerging perspective in occupational health, underlines the importance of events during previous life experiences [12].

A poor start to working life has indeed been found to have direct and long-term economic, personal, and social consequences [13]. Timely intervention for young workers at risk due to stress may be beneficial for enhancing and maintaining work ability during the rest of their working life [14].

Studies in workers at the beginning of their working career with new exposures can contribute to new knowledge in occupational epidemiology [15]. There are indications that—with regard to work stress—early detection and intervention is important when it comes to young people, as work stress precipitates depression and anxiety disorders in young workers [10, 16]. Young workers can profit their entire working life from the benefits of early intervention and improve their resilience to workplace stressors. Early intervention may thus promote work participation and prevent permanent exit from the labor market [14]. It has been suggested that interventions may be improved if there is a more in-depth understanding of stress and its pathways [17].

In previous research this relationship between stress and work ability has been well-established in different populations [18-22]. Several models have been developed to study the causes of stress in an occupational setting [23, 24]. To determine what happens after stress occurs, and how this affects work ability, it is important that factors affecting this relationship are mapped. It is known that appraised health mediates the effect of stress on work ability in older workers [18]. The mediating effect of health becomes stronger in older workers. However, from the literature it remains unclear what factors mediate the relationship between stress and work ability in young workers.

The need for a better understanding of the relationship between work ability and mental health underlines the urgency to study the role of factors that influence the relationship between stress and work ability in young workers. As it is hard to eliminate all stress, these factors are important in preventing stress from having a large impact on work ability.

In Sweden, the prevalence of stress in 2018 was highest among those aged 16–29, with 35% of women and 18% of men in the population indicating feeling stressed [25]. When stress complaints manifest at an early age, the work ability over the course of the working life can be affected as work stress can lead to later in the working life.

The Swedish Work Ability in Young Adults (WAYA) cohort allows us to study several factors that might influence the relationship between stress and work ability through mediation. In mediation, part of the relationship between stress and work ability can be explained by the presence of another variable/dimension (the mediator). A requirement for mediation is that the mediating factor is related to both stress and work ability. Based on previous research, possible mediating factors present in the WAYA cohort data that are associated with stress and work ability were selected for the current study. These mediating factors can be subdivided into four dimensions: recovery from work, work-home interference, control over private life, and physical activity in leisure time. These four dimensions have all been linked to both stress and work ability, and therefore might mediate this relationship.

The first dimension, recovery from work, can be influenced by stress and the social relationships at work [26]. Stress and the social situation at work are strongly related to disturbed sleep and impaired awakening [27]. In a representative sample of the Swedish working population, 53% of workers reported sleep disorders due to conditions at work [28]. Both stress and work ability are associated with aspects of recovery from work/work-related fatigue and events in private life [29, 30].

The second dimension, work to private life interference, is mostly caused by job strain and time-interference [31]. This dimension entails the spillover of negative aspects from work to private life. Stress is among others associated with work demands that

can interfere with private life [28, 32]. This increased negative influence of job demands on private life is associated with reduced work ability in young adults [32].

The third dimension, control over private life, can be affected by (occupational) stress. This entails feeling in control over private life and feeling that one has an influence on desired outcomes i.e., health, interpersonal relations, or financial situations. When stress is high, it can have a negative effect on private life [33]. A lack of control at home is associated with negative health effects, and private life can influence work ability [34, 35].

The fourth dimension entails physical activity in leisure time. Physical activity in leisure time differs from physical activity during work [36]. Lower physical activity in leisure time is known to be related to higher stress levels in individuals [37]. A lack of vigorous activity during leisure time is associated with a lower work ability, and the level of physical activity during leisure time is related to work ability [38].

Based on these aforementioned existing relationships in the literature, we expect a mediating effect on the relationship between stress and work ability by the selected dimensions. Intervening on known mediators may help improve sustainable work ability by reducing the negative effect of stress on future work ability. Therefore, results from this study can possibly help occupational professionals monitor work ability in the context of stress and enable work-related interventions. Subsequently, the aim of this study was to map out whether these factors mediate the relationship between stress and work ability over time in younger workers. To enable employers and young workers to eventually maintain sustainable employment during their entire working life, we aim to determine which dimension mediates the relationship between stress and work ability in young adults, and to what extent. Therefore, we drafted the following research question:

How is the relationship between stress and work ability mediated in young adults over time by recovery, the influence of job demands on private life, feelings of control over private life and physical activity during leisure time?

MATERIALS AND METHODS

Participants and Design

We used questionnaire data from 2012 and 2017 that had been collected in the WAYA cohort of Swedish young adults. The cohort started in 2007 using 20,000 young adults (equally divided between men and women) of 20–24 years old randomly selected from the total population registry held by the Swedish Tax Agency. The first of four questionnaires was sent out in 2007 with follow-up questionnaires in 2008, 2012, and 2017. In these questionnaires, questions on work demands and general health related

to these work demands were asked in order to assess occupational health in young adults [39]. The questionnaires were sent out by mail and could be answered by mail or online. Two reminders were sent out per follow-up time-interval. Each questionnaire was accompanied by a lottery ticket worth €1, regardless of whether the questionnaire was filled out and returned or not. During first inspection of respondents, the percentage of young adults working full-time rose from 38% in 2007 to 44%, 67%, and 80% in 2008, 2012 and 2017, respectively. As few young adults had full-time jobs in 2007 and 2008, the time points used for mediation analyses were 2012 and 2017. Persons who answered that they were neither working nor studying ($n = 250$ in 2012 and $n = 235$ in 2017) did not answer the questions on work ability and were excluded from this sample (see Appendix C).

Ethical Declarations

This research was conducted in accordance with the Declaration of Helsinki [40]. The research proposal was submitted to and approved by the Medical Ethical Committee of the Academic Medical Center and Swedish regional ethical committee Gothenburg Sweden, who decreed that a comprehensive evaluation was not required since this study was not subject to the Medical Research Involving Human Subjects Act (W19_432#19.498; T862-17). The original research proposal was reviewed and approved by the Central Ethical Review Board of the University of Gothenburg, under number T876-11.

Theoretical Framework

When mediation occurs, the relationship between two variables (stress and work ability) can be (partly) explained by another variable (the mediator). A theoretical framework was developed to confirm possible mediating factors. This framework included four possible dimensions of mediators: Recovery, work–home interference, control over private life, and physical activity. In the second part of the analysis, the theoretical framework was expanded with relationships between mediating variables. The mediation pathways were explored based on this theoretical framework.

The mediating variables of interest for these dimensions were as follows: Feeling well-rested upon waking (recovery), demands at work affecting private life (work–home interference), feelings of control over private life (control over private life), and physical activity in leisure time (physical activity). From the theoretical framework, gender was expected to play a role in the pathways of mediation with potentially different relationships between X (independent variable, stress), Y (dependent variable, work ability) and M_i (possible mediators).

Variables

Work ability was assessed with the Work Ability Score (WAS) i.e., the first question of the Work Ability Index (WAI), where respondents appraised their own work ability on a scale of 0 (no work ability at all) to 10 (best work ability ever experienced). Convergent validity compared to the complete WAI was more than acceptable ($r = 0.63$; $p < 0.001$) [41]. Stress was assessed by a single-item question that was validated against the Maslach burnout inventory ($r = 0.51$; $p < 0.001$) and the mental health subscale of the Short-Form-36 (SF-36) ($r = -0.63$; $p < 0.001$) and was found satisfactory for measuring stress in various work-life situations [42]. An explanation was included with the question, “Stress means a situation in which a person feels tense, restless, nervous, or anxious, or is unable to sleep at night because his/her mind is troubled all the time. Do you feel this kind of stress these days?”. Response options on this question were recorded on a 5-point scale where a score of 1 corresponded to “not at all” and a score of 5 with “very much”.

The dimension of recovery was assessed by a question on recovery by sleep: “How often in the last 30 days have you woken up feeling well-rested?”. The answer options were “never”, “once/a few times per month”, “several times per week”, and “almost every day”, which were coded as 1 to 4, respectively.

A question on demands at work that negatively affect private life were used for the work-home interference dimension: “Do the requirements of work/studies affect your personal life (leisure, home, and family life) in a negative way?”. There were five answer options were “very rarely”, “quite rarely”, “sometimes”, “quite often”, and “very often”, which were coded as 1 to 5, respectively.

Control over private life was assessed with one question statement: “I feel I have control over and can handle things that happen in my private life”. Respondents could indicate how well this statement matched one of four answer options: “poorly”, “rather poorly”, “well”, or “very well”, which were coded as 1 to 4, respectively.

The physical activity dimension was measured as physical activity during leisure time, and was measured with a single question assessing the average activity level during leisure time in the last twelve months i.e., “How physically active are you and how much do you move in your spare time?”. Four answer options were available, ranging from sedentary leisure time at one extreme (reading, watching TV, computer use or cinema visits) to hard training or competitive sports in competition at the other (running, skiing, biathlon) [43-45].

Statistical Analyses

The theoretical model was developed into a mediation model by adding mediators to the statistical model. After the causal steps of Baron and Kenny were confirmed,

several models were formed and tested [46]. The first model consisted of one mediating dimension, and the final model consisted of four mediators [47]. The exact way how the statistical model and mediators were formed is based on a new and reliable method, found in Appendix A [48].

This final model leads to a quadruple serial mediation model. The total effect in the quadruple mediation in series is the sum of all pathways in the model and can be written as follows:

Total effect (c) = Indirect effects + Direct effect (c')

$$c = a_1b_1 + a_2b_2 + a_3b_3 + a_4b_4 + a_1d_1b_2 + a_1d_4b_3 + a_1d_6b_4 + a_2d_2b_3 + a_2d_5b_4 + a_3d_3b_4 + a_1d_1d_2b_3 + a_1d_1d_5b_4 + a_1d_4d_3b_4 + a_2d_2d_3b_4 + a_1d_1d_2d_3b_4 + c'$$

This total effect was the sum of all possible pathways leading from X to Y , direct (c') or via mediating pathways. By adding mediators as regressors, we controlled for the presence of this mediator. These regression analyses were performed for each path, regressing the dependent variable (Y) on both the predictor (X) and the mediator (M). In multiple mediation, the effect of a_2 was calculated by regressing M_2 on both X and M_1 . In the final model, the effects were calculated by regressing the outcome (Y) on all the connected mediators (M_1 , M_2 , M_3 , and M_4) and the independent variable (X). The outcome variable was regressed on all previous predictors to calculate the contribution of each path.

Statistical analyses were performed using IBM SPSS Statistics for Windows, Version 25.0 (IBM Corp, Armonk, NY, USA), in combination with the PROCESS version 3.4 macro by Andrew F. Hayes [49]. Percentages of indirect effects were calculated using Microsoft Office 365 Excel 2016 (Microsoft, Redmont, WA, USA).

RESULTS

The total number of participants that filled out the questionnaires at 2012 and 2017 was 1733. Of these, 1432 filled out the stress and Work Ability Score on both time-points and were selected for analyses. A flowchart can be found in Appendix C.

The percentages of males in the population was 40% ($n = 575$) and 60% were female ($n = 857$). The means and standard deviations of the WAS at baseline (2012) were 8.2 (± 1.7), and five years later these were 7.9 (± 1.8). The stress scores of the population at both time points and other baseline population characteristics are shown in Table 1. Descriptive statistics of mediating variables separate for gender in 2012 and 2017 can be found in Appendix D.

Table 1: Demographic variables (mean and standard deviation (SD) of the study population that answered the questions on stress and work ability in 2012 and 2017 ($n = 1432$).

		2012		2017	
		Mean	SD	Mean	SD
Age		27.09	1.39	32.09	1.39
Weight (in kg)		72.96	14.53		
Height (in cm)		173.25	9.42		
WAS		8.22	1.67	7.93	1.77
Stress	Not at all	15.2%	($n = 217$)	13.0%	($n = 186$)
	Just a little	33.0%	($n = 473$)	30.7%	($n = 440$)
	To some extent	28.3%	($n = 405$)	29.7%	($n = 426$)
	Pretty much	18.3%	($n = 262$)	19.8%	($n = 284$)
	Very much	5.2%	($n = 75$)	6.7%	($n = 96$)
Work, study	Work or internship	73.3%	($n = 1050$)	87.1%	($n = 1247$)
	Study	12.0%	($n = 172$)	3.9%	($n = 56$)
	Both	14.7%	($n = 210$)	9.0%	($n = 129$)
Type of employment	Indefinite contract	68.2%	($n = 854$)	87.9%	($n = 1208$)
	Probationary period	4.7%	($n = 59$)	2.5%	($n = 35$)
	Fixed-term contract	9.2%	($n = 115$)	2.7%	($n = 37$)
	Other fixed-term contract (seasonal or project basis)	17.9%	($n = 224$)	6.8%	($n = 94$)
Highest completed education	Primary school (9 years)	1.1%	($n = 16$)	1.0%	($n = 14$)
	High School (12 years)	37.1%	($n = 531$)	24.9%	($n = 356$)
	University or tertiary, less than 3 years	11.8%	($n = 169$)	11.8%	($n = 168$)
	University or tertiary, more than 3 years	50.0%	($n = 715$)	62.3%	($n = 890$)
Family situation	Cohabiting, married, partnership	60.9%	($n = 869$)	75.1%	($n = 1073$)
	Girlfriend/boyfriend (not living together)	12.0%	($n = 172$)	6.3%	($n = 90$)
	Single	27.1%	($n = 387$)	18.6%	($n = 265$)

WAS: Work Ability Score

All steps of assumptions of mediation according to Baron and Kenny [46] were met for all four dimensions allowing the analysis to proceed to the single mediation step. The relationship between stress and work ability is shown in Table 2.

Table 2: Spearman correlations between stress and work ability variables in the mediation analysis for the years 2012 and 2017 for the total group and separate for males and females ($n = 1432$, $n = 575$, $n = 857$).

		Total	Males	Females
Stress 2012	Stress 2017	0.400	0.388	0.384
	WAS 2012	-0.329	-0.320	-0.322
	WAS 2017	-0.246	-0.231	-0.225
Stress 2017	WAS 2017	-0.429	-0.445	-0.397
WAS 2012	WAS 2017	0.364	0.393	0.337

All correlations were significant ($p < 0.01$).

All single mediation relationships were statistically significant with exception of the indirect effect (a_1b_1) of physical activity in 2017 (with stress in 2012 as predictor (X) and WAS in 2017 as outcome variable (Y); For all pathways, see Appendix A). The percentages of the indirect effect (a_1b_1) of the single mediation model with X_{2012} on Y_{2017} mediated by M_{2017} were 14.0%, 31.9%, 29.1%, and 1.9% for feeling well-rested upon waking, work demands affecting private life, feelings of control over private life, and physical activity in leisure time, respectively. In the model, when stress in 2017 was selected as a mediating variable, the relationship between stress in 2012 and the Work Ability Score in 2017 that went via the indirect pathway (a_1b_1) was 66% ($p < 0.001$), meaning that there was still a direct effect (c') of stress in 2012 towards work ability in 2017. Stress in 2017 was a strong mediator of the relationship between stress in 2012 and Work Ability Score in 2017.

Starting from the single mediation model, mediators were subsequently added to come to a final model with four mediators (see Appendix A for intermediate steps). Adding pathways through the fourth mediator (M_4) raised the number of indirect effects (X to Y) from 7 to 15 (see Figure A2d,e). All indirect effects except for indirect effect 14 were statistically significant. The final mediation model consisted of stress in 2017, demands at work affecting private life, feelings of control over private life, and feeling well-rested upon waking. These mediators were connected to each other by paths creating a quadruple serial mediation model. The effects of each pathway are shown in Table 3.

Table 3: Direct and indirect effects with 95% Confidence Interval (95% CI) for mediating pathways of the relationship between stress in 2012 and work ability in 2017 ($n = 1432$).

	Pathway	Effect	95% CI	% of Total Effect Passed Through Pathway
Total effect	c	-0.3955	[-0.4764, -0.3146]	100
Direct effect	c'	-0.0772 ^{ns}	[-0.1568, 0.0023] ^{ns}	19.5 ^{ns}
Total indirect effect	sum of all below	-0.3182	[-0.3750, -0.2642]	80.5
Indirect effect 1	a_1b_1	-0.1626	[-0.2075, -0.1205]	41.1
Indirect effect 2	a_2b_2	-0.0165	[-0.0308, -0.0048]	4.2
Indirect effect 3	a_3b_3	-0.0268	[-0.0469, -0.0089]	6.8
Indirect effect 4	a_4b_4	-0.009	[-0.0195, -0.0006]	2.3
Indirect effect 5	$a_1d_1b_2$	-0.0402	[-0.0592, -0.0235]	10.2
Indirect effect 6	$a_1d_4b_3$	-0.0385	[-0.0544, -0.0251]	9.7
Indirect effect 7	$a_1d_6b_4$	-0.0062	[-0.0123, -0.0017]	1.6
Indirect effect 8	$a_2d_2b_3$	-0.0024	[-0.0052, -0.0005]	0.6
Indirect effect 9	$a_2d_5b_4$	-0.0018	[-0.0039, -0.0004]	0.5
Indirect effect 10	$a_3d_3b_4$	-0.0014	[-0.0031, -0.0003]	0.4
Indirect effect 11	$a_1d_1d_2b_3$	-0.0059	[-0.0104, -0.0021]	1.5
Indirect effect 12	$a_1d_1d_5b_4$	-0.0045	[-0.0076, -0.0019]	1.1
Indirect effect 13	$a_1d_4d_3b_4$	-0.002	[-0.0039, -0.0007]	0.5
Indirect effect 14	$a_2d_2d_3b_4$	-0.0001 ^{ns}	[-0.0003, 0.0000] ^{ns}	0.0 ^{ns}
Indirect effect 15	$a_1d_1d_2d_3b_4$	-0.0003	[-0.0007, -0.0001]	0.1

All pathways were significant ($p < 0.05$) with exception of pathways marked with ^{ns} (not significant).

In the final model, the direct effect (c') was negative, but not statistically significant, meaning that the effect of stress in 2012 on work ability in 2017 was almost fully mediated by M_1 , M_2 , M_3 , and M_4 ($p = 0.0570$, 95% Confidence Interval (CI) [-0.1568, 0.0023], Degrees of Freedom (DF) (1426)). The selected mediators in the model were feelings of stress, feeling well-rested upon waking, work demands affecting private life, and feelings of control over private life in 2017. All effects for each individual path are shown in Figure 1.

Figure 1: Quadruple mediation with individual effect per path. All paths are significant $p < 0.05$, except c' .

The contribution for each mediator was tested in a parallel format where the d_i paths were eliminated, and the mediating effect only went via a_i and b_i paths through the corresponding mediators. This made it possible to compare the effects of each mediator in the model. The effect which went via pathway a_1b_1 was -0.1626 (95% CI $[-0.2068, -0.120]$), accumulating to 41%. The effects of a_2b_2 , a_3b_3 and a_4b_4 were -0.0567 (95% CI $[-0.0835, -0.0331]$), -0.0737 (95% CI $[-0.1020, -0.0498]$), and -0.0253 (95% CI $[-0.0411, -0.0117]$), respectively, corresponding to 14%, 19% and 6% of the total effect, indicating that the dimension of feelings of control over private life had the strongest mediating effect. Although M_3 had a larger mediating effect than M_2 , the theoretical framework suggested that work demands affecting private life preceded feelings of control over private life, resulting in this sequence of mediators in the final serial mediation model.

All four mediators explained 87% of the longitudinal relationship between stress in 2012 and work ability in 2017 for males and 79% for females. In both genders, adding these mediators led to the direct effect (c') not being significant anymore. The mediating effect of M_1 , M_2 , M_3 , and M_4 in the quadruple parallel mediation model are displayed in Table 4 for males and Table 5 for females. A slight difference in mediating pathways between both genders becomes clear from Table 4 and Table 5, where the mediating effect of all mediators combined is larger in males. In Appendix E, the serial quadruple mediation model stratified for gender is shown with the effect of each path and pathway.

Table 4: Direct and indirect effects for mediating pathways for males of the relationship between stress in 2012 and work ability in 2017 in a parallel design ($n = 575$).

Males			
Pathway	Effect	95% CI	% of Total Effect Passed Through Pathway
Total effect (c)	-0.3825	$[-0.5006, -0.2643]$	100
Direct effect (c')	-0.0498^{ns}	$[-0.1604, 0.0608]^{ns}$	13.0 ^{ns}
Total indirect effect	-0.3326	$[-0.4256, -0.2462]$	87.0
a_1b_1	-0.1809	$[-0.2490, -0.1178]$	47.3
a_2b_2	-0.0505	$[-0.0874, -0.0203]$	13.2
a_3b_3	-0.0855	$[-0.1322, -0.0469]$	22.4
a_4b_4	-0.0158	$[-0.0344, -0.0005]$	4.1

All pathways were significant ($p < 0.05$) with exception of pathways marked with: ^{ns} (not significant).

Table 5: Direct and indirect effects for mediating pathways for females of the relationship between stress in 2012 and work ability in 2017 in a parallel design ($n = 857$).

Females			
Pathway	Effect	95% CI	% of Total Effect Passed Through Pathway
Total effect (c)	-0.3577	[-0.4677, -0.2477]	100
Direct effect (c')	-0.0760 ^{ns}	[-0.1861, 0.0341] ^{ns}	21.2 ^{ns}
Total indirect effect	-0.2817	[-0.3547, -0.2154]	78.8
a ₁ b ₁	-0.1404	[-0.1991, -0.0839]	39.3
a ₂ b ₂	-0.0515	[-0.0874, -0.0213]	14.4
a ₃ b ₃	-0.0624	[-.0989, -0.0320]	17.4
a ₄ b ₄	-0.0273	[-.0500, -0.0089]	7.6

All pathways were significant ($p < 0.05$) with exception of pathways marked with: ^{ns} (not significant).

DISCUSSION

Findings

The relationship between stress in 2012 and Work Ability Score five years later, i.e., 2017, in young adults was mediated by stress in 2017, work demands affecting private life in 2017, feelings of control over private life in 2017, and feeling well-rested upon waking in 2017, which accounted for 81% of this relationship. This indicates that events at work that affect private life and feelings of control over private life are important for maintaining work ability in young adults with stress complaints. A well-balanced work–private life relationship can counteract the influence of stress on work ability.

Comparison to Other Studies

Not many studies have been carried out that consider mediators related to work ability. The stress and work ability relationship has been found before, but these studies did not go in-depth regarding other possibly related variables that mediated or moderated this [50]. In our study, the correlation between stress and work ability five years later was found to be moderate and significant, which is in accordance with a previous study that found a trend of long-term stress influencing work ability [51]. Our results show that part of this relationship can be explained by the selected mediating dimensions.

A study by Boström et al. (2012), with the same cohort but at earlier time points, found that job demands that affect private life in a negative way were associated with a reduced work ability [32]. Our study found that this relationship explains part of the stress–work ability relation. This means that part of the relation between stress and work ability can be explained by a spillover of work demands into private life. A study by Geurts et al. (1999) found that negative work–home interference was associated with a decrease in work-related health and general health indicators, which in turn might explain part of the relationship with work ability [52].

Although studies argue that work–home interference can be seen as a cause of stress, our study investigates the influence of demands at work affecting private life and feelings of control over private life on the relationship between stress and work ability [30]. Stress as measured in our study can be caused by other aspects than work–family interference. Geurts et al. (2005) found that higher levels of work–home interference were associated with job pressure, job support, and job control [31]. Peeters et al. (2004) found that job stressors lead to work–home interference and not the other way around [53]. However, two other studies by Dikkers et al. (2006) and Demerouti et al. (2004) found that workload (indicating stress) can be a precursor and consequence of work–home interference [54, 55]. Therefore, it still remains unclear how the interaction between stress and work–home interference is constituted.

Yang et al. found that appraised health was a mediator in the stress to work ability relationship for older workers [18]. In a report on psychosocial risks in Europe, it was found that older workers report better work–life balance than young workers. The authors suggested that workers with more experience in the labor market have better working conditions, allowing them to better balance work with other activities [6]. Our study focused on young workers who still need to balance work and private life, showing that the spillover of work into private life and decreased feelings of control over private life should be avoided. Although Geurts et al. (2005) found that home control was not significantly associated with home–work interference, we did see that there was an influence of feelings of control over private life on work ability [31].

Strengths and Limitations

A strength of this study is that we used longitudinal data to study mediation over time in order to make statements about the relationship between stress and work ability over time. Another strength is the depth in which the statistical model was studied. Whereas other studies in this field commonly use two mediators, this study incorporated four mediators while checking the relevance of the serial model. This quadruple mediation model was thoroughly considered by first developing a theoretical framework.

To the best of our knowledge, a study on mediators for the relationship between stress and work ability over time has not been performed before. Therefore, this study adds valuable information regarding prolonged stress and the relationship with work ability.

A limitation of this study is that some of the variables were not validated, as these single item questions represented an entire concept. A consequence can be that the measured concepts do not completely resemble the intended purpose. However, these questions approach the represented dimension as closely as possible. Another possible weakness is that some of the mediator variables were ordinal, but the PROCESS macro uses ordinary least squares regression to handle these variables and predict linearity [56]. All participants were randomly selected from the population to minimize selection bias. Selection bias because of employment status could not have occurred since although persons without a job were invited to participate in the cohort, they did not answer the question on work ability, and were consequently excluded from these analyses. Men were less likely to respond which may put limitations on the generalizability of the findings since there may be participation bias. However, since we conducted an analysis for men and women separately, we have insight in the effect of gender on our findings. A further analysis of people who dropped out between baseline and follow-up showed no differences between the drop-out and continued group, and therefore the risk of attrition bias is low.

Practical Implications

Stress is a common phenomenon in young workers. This study adds information on pathways of how stress can influence work ability in young workers. These pathways can help inform the implementation of interventions in this early phase of a working life. Such interventions may prevent a decreasing trajectory of work ability from regressing [57]. This study adds knowledge about the previously unknown pathways between stress and later work ability in young workers. We now know which mediators for this relationship can be targeted to prevent stress from influencing work ability over time. The results of this study indicate that the relation between stress and work ability is mostly mediated by the spillover of work demands into private life and the feelings of control over private life, indicating that these aspects pose a threat for the sustainable employability of young workers. Therefore, there is an indication that the personal life is affected by stress in the occupational setting, and that the personal life is important for work ability. Policy makers can improve their policies by acknowledging the impact of stress on work ability and their mediating factors. With our results confirming that the private situation is important in the relationship between stress and work ability, it is advisable that both the employer and employee

take care of the prevention of and coping with occupational stress, and avoiding work demands to negatively influence personal life.

An important role is therefore placed in boundary management to improve the work–private life balance. Both employer and employee should aim to prevent the work demands affecting private life in a negative way, to prevent declines in work ability over the working life course. Since stress is a common phenomenon in young workers, employees should discuss this issue and possible interventions with their employer at an early stage to prevent decreases in work ability. One such method can be to set boundaries and limits for accessibility outside work hours to aim for a higher feeling of control over their personal life, inhibiting the effect of stress on work ability [39, 58, 59]. Interventions can be aimed at increasing workplace autonomy and social support at work, which are known to be related to lower work stress levels and less spillover of the negative effect of work demands into private life [60, 61].

Early interventions may prevent declines in work ability at later age. When the mediating aspects are managed at the start of and during working life, it can possibly lead to an improved sustainable employment at a later age. An early intervention targeting known mediators can lead to an inhibition of the negative effect of stress on work ability.

Although there were no large differences between men and women, it may be desirable to pay attention to the differences in pathways between men and women when forming strategies to inhibit the effect of stress on work ability.

Implications for further research are that, in the next step, interventions aimed at the mediating factors are tested over time to confirm the preventive efficiency of the interventions to prevent declines in work ability over time. Future research studying work ability needs to take the mediators found in this study into account when mapping out work ability, or when studying work ability in different working populations of all ages.

CONCLUSIONS

The relationship between stress in 2012 and Work Ability Score five years later, i.e., 2017, in young adults was mediated by stress in 2017, work demands affecting private life in 2017, feelings of control over private life in 2017, and feeling well-rested upon waking in 2017, which accounted for 81% of this relationship. Since stress in 2017 mediated a large effect of the longitudinal relationship between stress and work ability, we can conclude that repeated stress is a risk factor for a lower work ability in young adults if not correctly managed.

Author Contributions Conceptualization, A.v.S., M.F-D., M.H., and K.N.; data curation, A.v.S. and A.N.B.; formal analysis, A.v.S. and A.N.B.; funding acquisition, M.H. and M.F-D.; investigation, A.v.S., A.N.B., S.T., M.F-D., M.H., and K.N.; methodology, A.v.S. and A.N.B.; project administration, K.N.; resources, M.H. and M.F-D.; software, A.v.S. and A.N.B.; supervision, M.F-D., M.H. and K.N.; visualization, A.v.S.; writing—original draft preparation, A.v.S.; writing—review and editing, A.N.B., S.T., M.F-D., M.H., and K.N.; All authors have read and agreed to the published version of the manuscript.

Funding This research was funded by AFA Insurance. The funder did not have any involvement in the content of this study.

Acknowledgments We are very thankful to the participants of the WAYA (Work Ability Young Adults) cohort, and we acknowledge AFA Insurance (AFA Försäkring) for research funding (Occupational and Environmental Medicine, Sahlgrenska Academy at University of Gothenburg).

Conflicts of Interest All authors declare having no conflict of interest. Financial support was provided by AFA insurance. Data availability: Data cannot be shared publicly as information about health and personal factors are regarded as sensitive information, and when sharing such data there has to be an approval from a Swedish Ethical committee (according to Swedish law) even if data is anonymized. Data is available with an approval from an ethical review board. For data requests, contact: School of Public Health and Community Medicine, Gothenburg University, Box 414, 405 30, Gothenburg, Sweden. The name of the data set is “WAYA” (Work Ability Young Adults).

Appendix A. Methods of Shaping the Statistical Model

Based on the statistical model as shown in Figure A1, a statistical model was built according to the following steps following the criteria for mediation of Baron and Kenny (1986) [46].

Figure A1: Primary theoretical framework consisting of four dimensions of possible mediating factors of the relationship between stress and work ability.

First, the requirements for mediation are tested in regression analyses. The assumptions of mediation have to be met to move to the next step. In the first step of our statistical analysis, the requirements for mediation by Baron and Kenny (1986) were tested. The requirements are also known as the causal steps approach [46]. The relationship between X and Y (c), X and M (a_1), and M and Y (b_1) that are required criteria to determine possible mediation, were studied. The first assumption is that there is a causal relation between the X and Y variable, where the X variable precedes the Y variable. In mediation, the X variable also needs to precede the mediator variable and a causal relationship needs to be present as well. The second assumption of mediation is that the mediator precedes the Y variable and that there is a causal relationship between these two. In that case, the relationship between X and Y might be mediated via mediator M . The models used in different steps can be seen in Figure A2a.

In the second step, statistical analyses regarding single mediation were performed using the PROCESS version 3.4 macro for IBM SPSS Statistics by Andrew F. Hayes [49]. Every analysis performed using this macro used 10,000 bootstraps for calculation of the indirect effects and the 95% confidence interval. All the significance levels tested were with an α of 0.05. The model used in this macro was model 4 (parallel mediation) [49]. In this step, different models were developed: Cross-sectional models for 2012 and 2017, longitudinal models where the mediator was from 2012 and the outcome

was the Work Ability Score in 2017, and another longitudinal model where both the mediator and outcome were from 2017. In this last model, the macro tests for the significance of the mediating effect by calculating the relationship between X_{2012} on Y_{2017} that is mediated by M_{2017} . The effect of c' was calculated by dividing the direct effect (c') by the total effect (c) to calculate the percentage of direct effect. The percentage of the total effect (c) that is explained by the pathway via mediator M_{2017} was calculated by dividing the indirect effect (a_1b_1) by the total effect to estimate the percentage of mediation. Where the relationship between X and Y was smaller when the mediator was present in the model, the direct effect was smaller than the total effect, and where the indirect effect was significant, mediation of the selected mediator was established. This analysis was performed separately for each mediator (feeling well-rested upon waking, work demands affecting private life, feelings of control over private life, and physical activity in leisure time).

After the contribution of each mediator was analyzed, the mediators that were significantly relevant were grouped in a double mediator analysis to form a model with two mediating variables. In this third step, only the mediators that showed a significant indirect effect were selected. Only mediator and outcome variables of 2017 were selected. In this step, all possible combinations of mediators were explored to discover the strongest mediation combination. This multiple mediator model was tested with PROCESS macro with Model 6 (serial mediation).

In the fourth step, the strongest combination of two mediators derived from the third step was added after adding stress in 2017 as the first mediator. The sequence of mediators was based on a theoretical model with interference and causal relationships. The sequence is important because in a serial mediation model, the pathway via M_1 can continue via M_2 , but not the other way around. Therefore, the dimension that comes first and can affect others, is entered first.

In the fifth and final step of the model building process, the fourth mediator of the theoretical model was added. According to the theoretical model, this mediator was a mediator between the relationship of another mediator and the outcome variable. To test this model, the fourth mediator variable was added in the series. All possible pathways were analyzed and described in the results section.

Following these mediation analyses, the final model was tested with all mediators in parallel to the X and Y variable to see if pathways explained different parts of mediation. This makes it possible to check if pathways follow multiple mediators in series, or that pathways only go through one mediator (see Figure A2). For the final model, the effect of each pathway (including 95% confidence intervals) were reported, as were the effects of each individual path and percentages for each pathway.

(a): Step 1: Causal relationships, causal steps of Baron and Kenny (1986).

(b): Step 2: Single mediation.

(c): Step 3: Double mediation in series.

(d): Step 4: Triple mediation in series.

(e): Step 5: Quadruple mediation in series.

(f): Final quadruple mediation model in parallel.

Figure A2: Different mediation steps testing the relationships of the variables.

The total effect in the quadruple mediation is series is the sum of all pathways in the model and can be written as follows:

Total effect (c) = Indirect effects + Direct effect (c')

$$c = a_1b_1 + a_2b_2 + a_3b_3 + a_4b_4 + a_1d_1b_2 + a_1d_4b_3 + a_1d_6b_4 + a_2d_2b_3 + a_2d_5b_4 + a_3d_3b_4 + a_1d_1d_2b_3 + a_1d_1d_5b_4 + a_1d_4d_3b_4 + a_2d_2d_3b_4 + a_1d_1d_2d_3b_4 + c'$$

This total effect is the sum of all possible pathways leading from X to Y , direct (c'), or via mediating pathways. The effects of each pathway are calculated by regressing the dependent variable on the independent variable. By adding mediators as regressors, we control for the presence of this mediator. These regression analyses are performed for each pathway, regressing the dependent variable (Y) on both the predictor (X) and the mediator (M). In the multiple mediation model, the effect of a_1 is obtained by regressing M_1 on X , and the effect of d_1 is obtained by regressing M_2 on M_1 . In multiple mediation, the effect of a_2 is calculated by regressing both X and M_1 on M_2 . In the final model, the effects are calculated by regressing all the connected mediators (M_1 , M_2 , M_3 , and M_4) and independent variable (X) on the outcome (Y).

Appendix B. Results of Steps towards the Statistical Model

Step 1: Assumptions of Mediation (Relationships between Variables)

All steps of assumptions of mediation according to Baron and Kenny [46] were met for all four dimensions, except for physical activity in leisure time. We found no evidence for a relationship between stress condition and physical activity in leisure time in 2012, but did find a significant relation in 2017. We can therefore say that the assumptions are met to move to the single mediation step. The relationship between stress and work ability is shown in Table A1.

Table A1: Spearman correlations between X and Y variables in the mediation analysis for the years 2012 and 2017 for the total group and separate for males and females ($n = 1432$, $n = 575$, $n = 857$).

		Total	Males	Females
Stress 2012	Stress 2017	0.400	0.388	0.384
	WAS 2012	-0.329	-0.320	-0.322
	WAS 2017	-0.246	-0.231	-0.225
Stress 2017	WAS 2017	-0.429	-0.445	-0.397
WAS 2012	WAS 2017	0.364	0.393	0.337

WAS: Work Ability Score. All correlations were significant ($p < 0.01$).

Step 2: Single Mediation

In the second step, 17 single mediation models were tested, (four for each theoretical model and five for the last, as stress in 2017 was included as a mediator): Cross-sectional of 2012, cross-sectional of 2017, longitudinal where only the outcome was from 2017, and longitudinal where both the mediator and the outcome were from 2017. All relationships were statistically significant with exception of the indirect effect (a_1b_1) of physical activity in 2017 (with stress in 2012 WAS in 2017 as predictor (X) and outcome variable (Y)). The percentages of the indirect effect (a_1b_1) of the last theoretical model were 14.0%, 31.9%, 29.1% and 1.9% for feeling well-rested upon waking, work affecting private life, feelings of control over private life, and physical activity in leisure time, respectively.

In the model when stress in 2017 was selected as a mediating variable, the relationship between stress in 2012 and the Work Ability Score in 2017 that went via the indirect pathway (a_1b_1) was 65.8% ($p < 0.001$), meaning that there was still a direct effect (c') of stress in 2012 towards work ability in 2017. Stress in 2017 was a strong mediator of the relationship between stress in 2012 and Work Ability Score in 2017.

We concluded that, based on the theoretical framework, the stress in 2012 would be maintained as a predictor variable to see whether the relationship between stress in 2012 and future work ability was mediated by the selected mediators of 2017.

Step 3: Double Mediation

The third step included mediation of two mediators in series as shown in the Appendix in Figure A2c. Sixteen models were tested: Stress in 2017 as the first mediator, with all four mediators as second mediator, and the other models consisted of all combinations of the four mediators as first and second mediator (4*3 models).

In these models the pathways between stress in 2012 and the physical activity in leisure time in 2017 were no longer statistically significant. In all models, work demands affecting private life and feelings of control over private life were the largest mediating pathways, with the exception of stress. A model with feelings of control over private life and work demands affecting private life as M_1 and M_2 , went for 50% via indirect pathways, meaning that the relationship between stress in 2012 and work ability in 2017 is explained by pathways via these two mediators. For the percentage, it does not matter if we use parallel or series, as the effects of X on M_1 and M_2 , and the effects of M_1 and M_2 on Y , are still the same. Only the pathway of b_1 goes partly via d_1 and b_2 . In all models, the d_1 pathway was significant, indicating a relationship between both mediators, advocating the parallel design.

Step 4: Triple Mediation

In the triple mediation, stress was selected as the first mediating variable (M_1) having the largest mediating effect. In the theoretical framework, stress is seen as a precursor for the other mediator variables. Therefore, stress in 2017 is a precursor for the mediating variables in the remainder of the model. Stress in 2017 is fixed in the models as M_1 . In the triple mediation, work demands affecting private life was selected as M_2 , and feelings of control over private life was selected as M_3 . These two mediators were the largest mediators in the single mediation model, and the combination of these two was the largest in the double mediation model (when stress was not included). In the theoretical model, M_2 is preceding M_3 , as the feelings of control over private life can be affected by work demands affecting private life. In this model, all direct (c') and indirect (a_1b_1 , a_2b_2 , a_3b_3 , $a_1d_1b_2$, $a_1d_3b_3$, $a_2d_2b_3$, and $a_1d_1d_2b_3$) pathways as shown in figure A2d were significant. The total indirect effect was -0.3092 (95% CI $[-0.3648, -0.2569]$), meaning that with a total effect of -0.3995 (95% CI $[-0.4764, -0.3146]$), 78% of the effect of stress in 2012 on work ability in 2017 is mediated by M_1 , M_2 , and M_3 .

Step 5: Quadruple Mediation

Since sleep quality was related to work ability and stress, this could function as a mediator for the relationship between stress in 2012 and work ability in 2017. However, in the theoretical model sleep quality can be a mediator for the other mediators as well. This means that the relationship between work demands affecting private life and work ability can be mediated by whether or not people felt well-rested upon waking. Therefore, the variable of feeling well-rested upon waking in 2017 is

added as last mediator (M_4) to the model. Adding this pathway raised the number of indirect effects from 7 to 15 (see Figure A2e). All indirect effects except for indirect effect 14 were statistically significant. The effects are shown in Table A2.

Table A2: Direct and indirect effects for mediating pathways of the relationship between stress in 2012 and work ability in 2017 ($n = 1432$).

	Pathway	Effect	95% CI	% of Total Effect Passed Through Pathway
Total effect	c	-0.3955	[-0.4764, -0.3146]	100
Direct effect	c'	-0.0772 ^{ns}	[-0.1568, 0.0023] ^{ns}	19.5 ^{ns}
Total indirect effect	sum of all below	-0.3182	[-0.3750, -0.2642]	80.5
Indirect effect 1	a_1b_1	-0.1626	[-0.2075, -0.1205]	41.1
Indirect effect 2	a_2b_2	-0.0165	[-0.0308, -0.0048]	4.2
Indirect effect 3	a_3b_3	-0.0268	[-0.0469, -0.0089]	6.8
Indirect effect 4	a_4b_4	-0.009	[-0.0195, -0.0006]	2.3
Indirect effect 5	$a_1d_1b_2$	-0.0402	[-0.0592, -0.0235]	10.2
Indirect effect 6	$a_1d_4b_3$	-0.0385	[-0.0544, -0.0251]	9.7
Indirect effect 7	$a_1d_6b_4$	-0.0062	[-0.0123, -0.0017]	1.6
Indirect effect 8	$a_2d_2b_3$	-0.0024	[-0.0052, -0.0005]	0.6
Indirect effect 9	$a_2d_5b_4$	-0.0018	[-0.0039, -0.0004]	0.5
Indirect effect 10	$a_3d_3b_4$	-0.0014	[-0.0031, -0.0003]	0.4
Indirect effect 11	$a_1d_1d_2b_3$	-0.0059	[-0.0104, -0.0021]	1.5
Indirect effect 12	$a_1d_1d_5b_4$	-0.0045	[-0.0076, -0.0019]	1.1
Indirect effect 13	$a_1d_4d_3b_4$	-0.002	[-0.0039, -0.0007]	0.5
Indirect effect 14	$a_2d_2d_3b_4$	-0.0001 ^{ns}	[-0.0003, 0.0000] ^{ns}	0.0 ^{ns}
Indirect effect 15	$a_1d_1d_2d_3b_4$	-0.0003	[-0.0007, -0.0001]	0.1

All pathways were significant ($p < 0.05$) with exception of pathways marked with ^{ns}: (not significant).

In the final model, the direct effect (c') was negative, but not statistically significant ($p = 0.0570$, 95% Confidence Interval (CI) [-0.1568, 0.0023], Degrees of Freedom (DF) (1426)), irrespective of the effect of feelings of stress, feeling well-rested upon waking, work demands affecting private life, feelings of control over private life in 2017. All effects for each individual path are shown in Figure A31.

Figure A3: Quadruple mediation with individual effect per path. All paths are significant $p < 0.05$, except c' .

Step 6: Check Series and Parallel

The contribution for each mediator was tested in a parallel format where the d-paths were eliminated. This made it possible to compare the effects of each mediator in the model. The effect that went via pathway a_1b_1 was -0.1626 (95% CI $[-0.2068, -0.120]$), meaning 41.1%. This is the same as in the model with four mediators in series, because all of this effect goes through M_1 in both models. For M_2 , M_3 , and M_4 , this percentage changes between both models. In the parallel model the d_i paths are not present and the mediating effect of stress in 2012 goes only via a_i and b_i paths through the corresponding mediators. In the parallel model, only the direct mediation of the relationship between stress in 2012 and WAS in 2017 is studied. The effects of a_2b_2 , a_3b_3 and a_4b_4 were -0.0567 (95% CI $[-0.0835, -0.0331]$), -0.0737 (95% CI $[-0.1020, -0.0498]$), and -0.0253 (95% CI $[-0.0411, -0.0117]$), respectively, corresponding to 14.3%, 18.6%, and 6.4% of the total effect. Although M_3 has a larger mediating effect than M_2 , the theoretical framework suggested that work demands affecting private life preceded feelings of control over private life resulting in this sequence of mediators in the final serial mediation model. In Appendix D, the serial quadruple mediation model stratified for gender is shown.

Step 7: Mediation Separated for Gender Results

Analyses separated for gender showed that the effect of the four mediators in parallel design was larger in males than in females. All four mediators explain 87.0% of the longitudinal relationship between stress in 2012 and work ability in 2017 for males, and 78.8% for females. In both genders, adding these mediators lead to the direct effect (c') no longer being significant. The mediating effect of M_1 , M_2 , M_3 , and M_4 are displayed in Table A3 for males and A4 for females. A slight difference in mediating pathways between both genders is shown in Tables A3 and A4.

Table A3: Direct and indirect effects for mediating pathways for males of the relationship between stress in 2012 and work ability in 2017 in a parallel design ($n = 575$).

Males			
Pathway	Effect	95% CI	% of Total Effect Passed Through Pathway
Total effect (c)	-0.3825	[-0.5006, -0.2643]	100
Direct effect (c')	-0.0498 ^{ns}	[-0.1604, 0.0608] ^{ns}	13.0 ^{ns}
Total indirect effect	-0.3326	[-0.4256, -0.2462]	87.0
a ₁ b ₁	-0.1809	[-0.2490, -0.1178]	47.3
a ₂ b ₂	-0.0505	[-0.0874, -0.0203]	13.2
a ₃ b ₃	-0.0855	[-0.1322, -0.0469]	22.4
a ₄ b ₄	-0.0158	[-0.0344, -0.0005]	4.1

All pathways were significant ($p < 0.05$) with exception of pathways marked with: ^{ns}

Table A4: Direct and indirect effects for mediating pathways for females of the relationship between stress in 2012 and work ability in 2017 in a parallel design ($n = 857$).

Females			
Pathway	Effect	95% CI	% of Total Effect Passed Through Pathway
Total effect (c)	-0.3577	[-0.4677, -0.2477]	100
Direct effect (c')	-0.0760 ^{ns}	[-0.1861, 0.0341] ^{ns}	21.2 ^{ns}
Total indirect effect	-0.2817	[-0.3547, -0.2154]	78.8
a ₁ b ₁	-0.1404	[-0.1991, -0.0839]	39.3
a ₂ b ₂	-0.0515	[-0.0874, -0.0213]	14.4
a ₃ b ₃	-0.0624	[-0.0989, -0.0320]	17.4
a ₄ b ₄	-0.0273	[-0.0500, -0.0089]	7.6

All pathways were significant ($p < 0.05$) with exception of pathways marked with: ^{ns}

Appendix C. Flow-Chart of Participation

Figure A4: Flowchart of inclusion of the Work Ability in Young Adults (WAYA) cohort.

Appendix D. Descriptive Statistics of Mediator Variables

Table A5: Descriptive statistics of mediator variables for males and females in 2012 and 2017.

	Males (<i>n</i> = 575)				Females (<i>n</i> = 857)			
	2012		2017		2012		2017	
	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>
Stress in 2012								
<i>Not at all</i>	19.8	114	16.5	95	12.0	103	10.6	91
<i>Just a little</i>	36.2	208	35.1	202	30.9	265	27.8	238
<i>To some extent</i>	16.1	150	27.0	155	29.8	255	31.6	271
<i>Pretty much</i>	14.3	82	15.8	91	21.0	180	22.5	193
<i>Very much</i>	3.7	21	5.6	32	6.3	54	6.3	64
Work demands affecting private life in a negative way								
<i>Very rarely</i>	30.3	174	20.9	120	17.1	146	13.1	112
<i>Quite rarely</i>	20.7	119	26.3	151	19.3	165	21.0	180
<i>Sometimes</i>	30.7	176	32.7	188	34.2	292	38.7	332
<i>Quite often</i>	14.6	84	16.7	96	22.5	192	19.7	169
<i>Very often</i>	3.7	21	3.5	20	7.0	60	7.5	64
Feelings of control over private life								
<i>Matches very bad</i>	2.6	15	2.8	16	3.5	30	2.7	23
<i>Matches pretty bad</i>	14.1	81	14.6	84	15.9	136	19.1	164
<i>Matches pretty good</i>	54.5	312	55.5	319	53.5	547	56.7	486
<i>Matches very good</i>	28.8	165	27.1	156	27.1	232	21.5	184
Well-rested upon waking in the last 30 days								
<i>Never</i>	8.2	47	10.1	58	7.8	67	14.9	128
<i>Any/a few times per month</i>	49.7	285	50.1	288	52.8	452	53.2	456
<i>Multiple times per week</i>	32.4	186	29.4	169	29.2	250	25.1	215
<i>Almost every day</i>	9.8	56	10.4	60	10.2	87	6.8	58
WAS (0-10)								
<i>Mean, SD</i>	8.38	1.54	8.26	1.60	8.11	1.74	7.71	1.84

Appendix E. Mediation Pathways for Males

Table A6: Direct and indirect effects for mediating pathways of the relationship between stress in 2012 and work ability in 2017 in males ($n = 575$).

	Pathway	Effect	95% CI	% of Total Effect Passed Through Pathway
Total effect	c	-0.3825	[-0.5006, -0.2643]	100
Direct effect	c'	-0.0498 ^{ns}	[-0.1604, 0.0608] ^{ns}	13.0 ^{ns}
Total indirect effect	sum of all below	-0.3326	[-0.4277, -0.2466]	87.0
Indirect effect 1	a ₁ b ₁	-0.1809	[-0.2492, -0.1180]	43.7
Indirect effect 2	a ₂ b ₂	-0.0174	[-0.0399, -0.0014]	4.5
Indirect effect 3	a ₃ b ₃	-0.0306 ^{ns}	[-0.0705, 0.0043]	8.0 ^{ns}
Indirect effect 4	a ₄ b ₄	-0.0073 ^{ns}	[-0.0208, 0.0029]	1.9 ^{ns}
Indirect effect 5	a ₁ d ₁ b ₂	-0.0331	[-0.0580, -0.0135]	8.7
Indirect effect 6	a ₁ d ₄ b ₃	-0.0417	[-0.0697, -0.0205]	10.9
Indirect effect 7	a ₁ d ₆ b ₄	-0.0002 ^{ns}	[-0.0055, 0.0048]	0.1 ^{ns}
Indirect effect 8	a ₂ d ₂ b ₃	-0.0045	[-0.0114, -0.0001]	1.2
Indirect effect 9	a ₂ d ₅ b ₄	-0.0015 ^{ns}	[-0.0042, 0.0001]	0.4 ^{ns}
Indirect effect 10	a ₃ d ₃ b ₄	-0.0014 ^{ns}	[-0.0045, 0.0003]	0.4 ^{ns}
Indirect effect 11	a ₁ d ₁ d ₂ b ₃	-0.0086	[-0.0167, -0.0021]	2.2
Indirect effect 12	a ₁ d ₁ d ₅ b ₄	-0.0029	[-0.0067, -0.0001]	0.8
Indirect effect 13	a ₁ d ₄ d ₃ b ₄	-0.0018 ^{ns}	[-0.0047, 0.0000]	0.5 ^{ns}
Indirect effect 14	a ₂ d ₂ d ₃ b ₄	-0.0002 ^{ns}	[-0.0007, 0.0000]	0.1 ^{ns}
Indirect effect 15	a ₁ d ₁ d ₂ d ₃ b ₄	-0.0004 ^{ns}	[-0.0011, 0.0000]	0.1 ^{ns}

All pathways were significant ($p < 0.05$) with exception of pathways marked with ^{ns}

Figure A5: Quadruple mediation with individual effect per path for males. (All paths are significant $p < 0.05$, except a_4 , b_4 , d_6 , and c')

Appendix F. Mediation Pathways for Females

Table A7: Direct and indirect effects for mediating pathways of the relationship between stress in 2012 and work ability in 2017 in females ($n = 857$).

	Pathway	Effect	95% CI	% of Total Effect Passed Through Pathway
Total effect	c	-0.3577	[-0.4677, -0.2477]	100
Direct effect	c'	-0.076 ^{ns}	[-0.1861, 0.0341]]	21.2 ^{ns}
Total indirect effect	sum of all below	-0.2817	[-0.3549, -0.2147]	78.8
Indirect effect 1	a ₁ b ₁	-0.1404	[-0.1999, -0.0853]	39.3
Indirect effect 2	a ₂ b ₂	-0.0119 ^{ns}	[-0.0294, 0.0024]	3.3 ^{ns}
Indirect effect 3	a ₃ b ₃	-0.0234	[-0.0468, -0.0044]	6.5
Indirect effect 4	a ₄ b ₄	-0.0077 ^{ns}	[-0.0224, 0.0042]	2.2 ^{ns}
Indirect effect 5	a ₁ d ₁ b ₂	-0.0396	[-0.0672, -0.0159]	11.1
Indirect effect 6	a ₁ d ₄ b ₃	-0.0340	[-0.0545, -0.0172]	9.5
Indirect effect 7	a ₁ d ₆ b ₄	-0.0113	[-0.0224, -0.0031]	3.2
Indirect effect 8	a ₂ d ₂ b ₃	-0.0012 ^{ns}	[-0.0036, 0.0004]	0.3 ^{ns}
Indirect effect 9	a ₂ d ₅ b ₄	-0.0013 ^{ns}	[-0.0039, 0.0002]	0.4 ^{ns}
Indirect effect 10	a ₃ d ₃ b ₄	-0.0009 ^{ns}	[-0.0029, 0.0001]	0.3 ^{ns}
Indirect effect 11	a ₁ d ₁ d ₂ b ₃	-0.0039 ^{ns}	[-0.0089, 0.0001]	1.1 ^{ns}
Indirect effect 12	a ₁ d ₁ d ₅ b ₄	-0.0045	[-0.0090, -0.0012]	1.3
Indirect effect 13	a ₁ d ₄ d ₃ b ₄	-0.0014 ^{ns}	[-0.0036, 0.0002]	0.4 ^{ns}
Indirect effect 14	a ₂ d ₂ d ₃ b ₄	0.0000 ^{ns}	[-0.0002, 0.0000]	0.0 ^{ns}
Indirect effect 15	a ₁ d ₁ d ₂ d ₃ b ₄	-0.0002 ^{ns}	[-0.0005, 0.0000]	0.1 ^{ns}

All pathways were significant ($p < 0.05$) with exception of pathways marked with ^{ns}

Figure A6: Quadruple mediation with individual effect per path for females. All paths are significant $p < 0.05$, except a_2 , a_4 , d_3 , and c' .

REFERENCES

1. Ilmarinen J. Work ability--a comprehensive concept for occupational health research and prevention. *Scand J Work Environ Health*. 2009;35(1):1-5.
2. Kinnunen U, Natti J. Work ability score and future work ability as predictors of register-based disability pension and long-term sickness absence: A three-year follow-up study. *Scand J Public Health*. 2018;46(3):321-30.
3. Notenbomer A, Groothoff JW, van Rhenen W, Roelen CA. Associations of work ability with frequent and long-term sickness absence. *Occup Med (Lond)*. 2015;65(5):373-9.
4. Reeuwijk KG, Robroek SJ, Niessen MA, Kraaijenhagen RA, Vergouwe Y, Burdorf A. The Prognostic Value of the Work Ability Index for Sickness Absence among Office Workers. *PLoS One*. 2015;10(5):e0126969.
5. Roelen CA, van Rhenen W, Groothoff JW, van der Klink JJ, Twisk JW, Heymans MW. Work ability as prognostic risk marker of disability pension: single-item work ability score versus multi-item work ability index. *Scand J Work Environ Health*. 2014;40(4):428-31.
6. Vargas O, Flintrap J, Hassard J, Irastorza X, Milczarek M, Miller J, et al. Psychosocial risks in Europe: prevalence and strategies for prevention. Eurofound and EU-OSHA. 2014;Luxembourg: Publications Office of the European Union.
7. Hsu HC. Age Differences in Work Stress, Exhaustion, Well-Being, and Related Factors From an Ecological Perspective. *Int J Environ Res Public Health*. 2018;16(1).
8. Hill EJ, Erickson JJ, Fellows KJ, Martinengo G, Allen SM. Work and family over the life course: Do older workers differ? *Journal of Family and Economic Issues*. 2014;35(1):1-13.
9. Law PCF, Too LS, Butterworth P, Witt K, Reavley N, Milner AJ. A systematic review on the effect of work-related stressors on mental health of young workers. *Int Arch Occup Environ Health*. 2020:611-22.
10. Melchior M, Caspi A, Milne BJ, Danese A, Poulton R, Moffitt TE. Work stress precipitates depression and anxiety in young, working women and men. *Psychol Med*. 2007;37(8):1119-29.
11. Veldman K, Reijneveld SA, Verhulst FC, Ortiz JA, Bultmann U. A life course perspective on mental health problems, employment, and work outcomes. *Scand J Work Environ Health*. 2017;43(4):316-25.
12. Amick BC, McLeod CB, Bultmann U. Labor markets and health: an integrated life course perspective. *Scand J Work Environ Health*. 2016;42(4):346-53.
13. Bradley S, Nguyen AN. 13 The school-to-work transition. *International handbook on the economics of education*. 2004:484.
14. Lallukka T, Kronholm E, Pekkala J, Jappinen S, Blomgren J, Pietilainen O, et al. Work participation trajectories among 1,098,748 Finns: reasons for premature labour market exit and the incidence of sickness absence due to mental disorders and musculoskeletal diseases. *BMC Public Health*. 2019;19(1):1418.

15. Checkoway H, Pearce N, Kriebel D. Selecting appropriate study designs to address specific research questions in occupational epidemiology. *Occupational and environmental medicine*. 2007;64(9):633-8.
16. Boersma K, Lindblom K. Stability and change in burnout profiles over time: A prospective study in the working population. *Work & Stress*. 2009;23(3):264-83.
17. Grebner S, Elfering A, Semmer NK, Kaiser-Probst C, Schlapbach M-L. Stressful situations at work and in private life among young workers: An event sampling approach. *Social Indicators Research*. 2004;67(1-2):11-49.
18. Yang T, Liu T, Lei R, Deng J, Xu G. Effect of Stress on the Work Ability of Aging American Workers: Mediating Effects of Health. *Int J Environ Res Public Health*. 2019;16(13).
19. Bernburg M, Vitzthum K, Groneberg DA, Mache S. Physicians' occupational stress, depressive symptoms and work ability in relation to their working environment: a cross-sectional study of differences among medical residents with various specialties working in German hospitals. *BMJ Open*. 2016;6(6):e011369.
20. Habibi E, Dehghan H, Safari S, Mahaki B, Hassanzadeh A. Effects of work-related stress on work ability index among refinery workers. *J Educ Health Promot*. 2014;3:18.
21. Marklund S, Mienna CS, Wahlstrom J, Englund E, Wiesinger B. Work ability and productivity among dentists: associations with musculoskeletal pain, stress, and sleep. *Int Arch Occup Environ Health*. 2020;93(2):271-8.
22. Yong M, Nasterlack M, Pluto RP, Lang S, Oberlinner C. Occupational stress perception and its potential impact on work ability. *Work*. 2013;46(3):347-54.
23. Bakker AB, Demerouti E. The job demands-resources model: State of the art. *Journal of managerial psychology*. 2007:309-28.
24. Karasek Jr RA. Job demands, job decision latitude, and mental strain: Implications for job redesign. *Administrative science quarterly*. 1979:285-308.
25. Public Health Agency of Sweden. Statistik över vuxnas psykiska hälsa [Statistics of adults' mental health]. Solna: <http://www.folkhalsomyndigheten.se/>; 2019.
26. Demerouti E, Bakker AB, Geurts SAE, Taris TW. Daily recovery from work-related effort during non-work time. *Research in occupational stress and well-being: Vol 7 Current perspectives on job-stress recovery*: JAI Press/Emerald Group Publishing; 2009. p. 85–123.
27. Akerstedt T, Knutsson A, Westerholm P, Theorell T, Alfredsson L, Kecklund G. Sleep disturbances, work stress and work hours: a cross-sectional study. *J Psychosom Res*. 2002;53(3):741-8.
28. Arbetsmiljöverket, Sveriges Officiella statistik. Arbetsorsakade besvär 2018 - Arbetsmiljöstatistik Rapport 2018:3 - Work-Related Disorders 2018. 2018. Report No.: ISSN 1652-1110.
29. Sluiter JK, de Croon EM, Meijman TF, Frings-Dresen MHW. Need for recovery from work related fatigue and its role in the development and prediction of subjective health complaints. *Occup Environ Med*. 2003;60 Suppl 1:i62-70.

30. Peeters MCW, Montgomery AJ, Bakker AB, Schaufeli WB. Balancing work and home: how job and home demands are related to burnout. *International Journal of Stress Management*. 2005;12(1):43-61.
31. Geurts SAE, Taris TW, Kompier MAJ, Dikkers JSE, Van Hooff MLM, Kinnunen UM. Work-home interaction from a work psychological perspective: Development and validation of a new questionnaire, the SWING. *Work & Stress*. 2005;19(4):319-39.
32. Boström M, Sluiter JK, Hagberg M. Changes in work situation and work ability in young female and male workers. A prospective cohort study. *BMC Public Health*. 2012;12:694.
33. Rout UR, Rout JK. Occupational stress. *Stress Management for Primary Health Care Professionals*. 2002:25-39.
34. Boström M, Holmgren K, Sluiter JK, Hagberg M, Grimby-Ekman A. Experiences of work ability in young workers: an exploratory interview study. *Int Arch Occup Environ Health*. 2016;89(4):629-40.
35. Eriksson J, Dellve L, Eklof M, Hagberg M. Early inequalities in excellent health and performance among young adult women and men in Sweden. *Gend Med*. 2007;4(2):170-82.
36. Coenen P, Huysmans MA, Holtermann A, Krause N, van Mechelen W, Straker LM, et al. Do highly physically active workers die early? A systematic review with meta-analysis of data from 193 696 participants. *Br J Sports Med*. 2018;52(20):1320-6.
37. Wijndaele K, Matton L, Duvigneaud N, Lefevre J, De Bourdeaudhuij I, Duquet W, et al. Association between leisure time physical activity and stress, social support and coping: A cluster-analytical approach. *Psychology of Sport and Exercise*. 2007;8(4):425-40.
38. van den Berg TI, Elders LA, de Zwart BC, Burdorf A. The effects of work-related and individual factors on the Work Ability Index: a systematic review. *Occup Environ Med*. 2009;66(4):211-20.
39. Thomee S, Harenstam A, Hagberg M. Mobile phone use and stress, sleep disturbances, and symptoms of depression among young adults--a prospective cohort study. *BMC Public Health*. 2011;11:66.
40. World Medical Association. Declaration of Helsinki - Ethical principles for medical research involving human subjects. 2013;Adopted by the 18th WMA General Assembly, Helsinki, Finland, June 1964(amended by the 64th WMA General Assembly, Fortaleza, Brazil October).
41. El Fassi M, Bocquet V, Majery N, Lair ML, Couffignal S, Mairiaux P. Work ability assessment in a worker population: comparison and determinants of Work Ability Index and Work Ability score. *BMC Public Health*. 2013;13:305.
42. Elo AL, Leppanen A, Jahkola A. Validity of a single-item measure of stress symptoms. *Scand J Work Environ Health*. 2003;29(6):444-51.
43. Saltin B, Grimby G. Physiological analysis of middle-aged and old former athletes: comparison with still active athletes of the same ages. *Circulation*. 1968;38(6):1104-15.
44. Aires N, Selmer R, Thelle D. The validity of self-reported leisure time physical activity, and its relationship to serum cholesterol, blood pressure and body mass index. A population

- based study of 332,182 men and women aged 40-42 years. *Eur J Epidemiol*. 2003;18(6):479-85.
45. Rodjer L, Jonsdottir IH, Rosengren A, Bjorck L, Grimby G, Thelle DS, et al. Self-reported leisure time physical activity: a useful assessment tool in everyday health care. *BMC Public Health*. 2012;12:693.
 46. Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc Psychol*. 1986;51(6):1173-82.
 47. Hayes AF, Rockwood NJ. Regression-based statistical mediation and moderation analysis in clinical research: Observations, recommendations, and implementation. *Behav Res Ther*. 2017;98:39-57.
 48. Hayes AF, Montoya AK, Rockwood NJ. The analysis of mechanisms and their contingencies: PROCESS versus structural equation modeling. *Australasian Marketing Journal (AMJ)*. 2017;25(1):76-81.
 49. Hayes AF. Introduction to mediation, moderation, and conditional process analysis: A regression-based approach. New York, USA: Guilford Publications; 2017.
 50. Goedhard RG, Goedhard WJA, editors. Work ability and perceived work stress. International Congress Series; 2005: Elsevier.
 51. Lindegård A, Larsman P, Hadzibajramovic E, Ahlborg G, Jr. The influence of perceived stress and musculoskeletal pain on work performance and work ability in Swedish health care workers. *Int Arch Occup Environ Health*. 2014;87(4):373-9.
 52. Geurts S, Rutte C, Peeters M. Antecedents and consequences of work-home interference among medical residents. *Soc Sci Med*. 1999;48(9):1135-48.
 53. Peeters MCW, de Jonge J, Janssen PPM, van der Linden S. Work-home interference, job stressors, and employee health in a longitudinal perspective. *International Journal of Stress Management*. 2004;11(4):305-22.
 54. Dikkers JSE, Geurts SAE, Kompier MAJ, Taris TW, Houtman ILD, van den Heuvel F. Does workload cause work-home interference or is it the other way around? *Stress and Health: Journal of the International Society for the Investigation of Stress*. 2007;23(5):303-14.
 55. Demerouti E, Bakker AB, Bulters AJ. The loss spiral of work pressure, work-home interference and exhaustion: Reciprocal relations in a three-wave study. *Journal of Vocational behavior*. 2004;64(1):131-49.
 56. Hayes AF, Rockwood NJ. Conditional Process Analysis: Concepts, Computation, and Advances in the Modeling of the Contingencies of Mechanisms. *Am Behav Sci*. 2020;64(1):19-54.
 57. Oakman J, Neupane S, Prakash KC, Nygard CH. What Are the Key Workplace Influences on Pathways of Work Ability? A Six-Year Follow Up. *Int J Environ Res Public Health*. 2019;16(13).
 58. Derks D, van Duin D, Tims M, Bakker AB. Smartphone use and work-home interference: The moderating role of social norms and employee work engagement. *Journal of Occupational and Organizational Psychology*. 2015;88(1):155-77.

59. Mellner C. After-hours availability expectations, work-related smartphone use during leisure, and psychological detachment: The moderating role of boundary control. *International Journal of Workplace Health Management*. 2016;9(2):146-64.
60. Grawitch MJ, Ballard DW, Erb KR. To be or not to be (stressed): The critical role of a psychologically healthy workplace in effective stress management. *Stress and Health*. 2015;31(4):264-73.
61. French KA, Dumani S, Allen TD, Shockley KM. A meta-analysis of work-family conflict and social support. *Psychol Bull*. 2018;144(3):284-314.

6

Appraisal of work ability in relation to job-specific health requirements in ambulance workers

A. van Schaaik, J.S. Boschman, M.H.W. Frings-Dresen and J.K. Sluiter

International archives of occupational and environmental health, 2017, 90(1), 123-131.

ABSTRACT

Purpose: To gain insight into which job-specific health requirements relate to work ability, the following two research questions were formulated: Which job-specific health requirements are associated with the appraisal of work ability in ambulance drivers and paramedics? How are appraisals of physical and mental work ability associated with the appraisal of overall work ability in ambulance drivers and paramedics?

Method: Workers Health Surveillance cross-sectional data of 506 ambulance workers (236 drivers and 270 paramedics) were used. The tests for specific job requirements were divided into six categories. Work ability was appraised as overall, physical and mental/emotional. Multiple linear stepwise regression analyses were used to model the associations.

Results: Outcomes in 'raised alertness and judgment ability' ($R^2=0.09$), 'job-specific physical abilities' ($R^2=0.10$) and 'emotional peak load' ($R^2=0.07$) significantly explained appraised overall, physical and mental/emotional work ability. Physical and mental/emotional work ability together explained 48.3% of the variance of overall work ability. The explained variance by physical and mental/emotional work ability was almost 4% higher in drivers than in paramedics.

Conclusions: Overall work ability was significantly explained by outcomes in 'raised alertness and judgment ability' and 'emotional peak load'. Physical work ability was significantly explained by 'job-specific physical abilities' and 'raised alertness and judgment ability' outcomes, while 'emotional peak load' and 'raised alertness and judgment ability' outcomes significantly explained mental/emotional work ability. Physical and mental/emotional work ability explain the same proportion of variance in overall work ability.

Keywords: Work ability; Ambulance; Job-specific; Workers Health Surveillance (WHS); Job-specific requirements

INTRODUCTION

Work ability in ambulance workers is endangered due to the high physical and mental/emotional demands placed on them. The attribution to appraised work ability by job-specific health requirements has not been determined in ambulance workers. Evidence shows that ambulance workers have an injury rate of 3-5 times higher than the general working population [1, 2], and a higher health risk [3-5]. The risk for musculoskeletal disorders in the lower back is 13 times higher than for health professionals, while the risk for mental disorders is almost 14 times higher [2]. Ambulance work endangers the physical and mental health of ambulance workers and demands adequate health.

Consequently, it is clear that monitoring the health and work ability of ambulance workers is important to prevent deterioration of health and prevent hazardous situations. A strategy that can be used to monitor the health of ambulance workers is a job-specific Workers Health Surveillance (WHS) [6, 7].

The construct of work ability has been defined over the years by Ilmarinen et al. [8] as 'the quality of workers at present and in the near future to be able to do their job satisfactorily with respect to work demands, health and mental resources.' Work ability is especially important in jobs with specific requirements because of the high job demands. Ambulance work is such an occupation, with both high mental and physical demands [2, 5, 9] and a higher risk for lower work ability compared to other health professionals [2, 3, 10, 11].

Specific psychological demands are being able to drive with flashing lights and sirens, working under time pressure with critical and sometimes aggressive bystanders when human life is at stake, and making complicated (medical) decisions involving a person's health in perilous situations [6]. Besides psychological health demands, the psychosocial health demands are an important part of work ability. The emotional demands that accompany with an emergency call and coping with traumatic experiences lead to stress reactions [12] and may result in post-traumatic complaints and depression [3, 13]. Under demanding circumstances, the ambulance worker needs to remain a professional and must be able to make rational decisions. The work of ambulance workers is also physically strenuous. Ambulance workers need to be able to maintain a certain position for a period of time to be able to treat a patient. Their physique is also challenged when they need to take challenging or unconventional routes to reach a victim or patient.

To monitor all these specific requirements of ambulance workers, a job-specific WHS has been developed in the Netherlands in collaboration with the ambulance sector organization [14]. The WHS screens on physical capacity, work-related fatigue, sleepiness, post-traumatic stress complaints, and complaints of depression and

anxiety. These are categorized as ‘the ability to perceive and communicate’, ‘be able to have raised alertness and judgment ability’, ‘be capable of handling emotional peak load’, ‘job-specific physical abilities’, and ‘having respiratory complaints’ or ‘having skin complaints’ that affect working. In addition to these categories, the appraised overall, physical and mental/emotional work ability was scored. It is still unknown whether these health requirements are associated with the appraisal of work ability.

To satisfactorily function in work and guarantee a high quality of healthcare, it is important that ambulance workers comply to minimum standards. Although the job requirements for drivers and paramedics are similar, it is not known how health complaints or problems in work functioning relate to work ability in these different occupations. In the Netherlands in 2012, 2218 ambulance paramedics (1445 male, 773 female) and 1949 ambulance drivers (1712 male, 237 female) were employed who carried out 1,100,419 rides, of which 500,835 were emergency calls [15].

To date, no study has gathered information on job-specific aspects of health and work ability so specifically. Work ability can be divided into two aspects: mental and physical. Although it is known that mental and physical health has a different impact on self-reported work ability [16], it is still unclear how the self-reported mental and physical work ability relates to the appraisal of overall work ability. According to a study of Klasan et al. [10], the aspect that predicts low work ability in ambulance workers best is the amount of strenuous physical activity during work. But which job-specific health complaints greatly reduce work ability? It is still unknown where interventions should be aimed at, to expect the greatest positive change in work ability.

To explore the composition of appraised work ability and to study whether work ability appraisal has an additional value in the WHS, this study assesses the explained variance of job-specific mental and physical health requirements and ambulance workers’ appraisal of their own current work ability. This leads to the following research question: *Which job-specific health requirements are associated with the appraisal of work ability in ambulance drivers and paramedics?*

We expect physical job requirements to have a bigger impact on self-reported work ability than the mental requirements and will show up as more important in the model. We expect the highest explained variance of appraised overall work ability to be physical work ability for both occupations, based on previous studies in ambulance workers and other occupations [10, 16]. However, we expect that the mental demands are more important for ambulance drivers than for paramedics, since raised alertness is more frequently demanded for drivers. The second research question is: *How are appraisals of physical and mental work ability associated with the appraisal of overall work ability in ambulance drivers and paramedics?*

METHODS

We used Dutch WHS data collected in the ambulance sector over the period October 20, 2011 until January 22, 2013. Cross-sectional data of the WHS among 506 ambulance paramedics (n=270) and drivers (n=236) were available for analyses. All employees participated on a voluntary basis. The WHS was conducted according to a national protocol, ensuring the safety and anonymity of the participants. An occupational physician and/or assistant conducted all tests. The collective labor agreement of the ambulance sector 2011-2013 included an agreement which stated that the data of the WHS were stored in a national database and that data were to be analyzed anonymously by the Academic Medical Center (AMC) for further research [17]. Ambulance workers were reminded of this agreement in writing before participating in the WHS. The Medical Ethical Commission of the AMC authorized this study and decided that a full application was not required.

Specific job requirements are considered important requirements to be met by ambulance workers to satisfactorily fulfill their work and cope with their demanding working conditions without impairing their own health or the safety of others. Important requirements for ambulance work that can affect work ability were distinguished in six categories: 'to perceive and communicate'; 'having raised alertness and judgment ability'; 'dealing with emotional peak load'; 'job-specific physical abilities'; 'respiratory complaints'; and 'skin complaints.'

To assess 'the ability to perceive and communicate,' hearing and vision tests were performed. Necessary eyesight for perception consisted of color perception and acuity tests. Physicians tested color perception with the Ishihara test for color blindness [18]. The Landolt C-ring test was used to test visual acuity. Three yes/no questions on eyesight were about whether the employee had problems with vision during work, at night and with reading. The whisper-speech test tested the ability to hear human speech in conversation [19]. Two signal questions were asked about problems with the ability to hear inside and outside the ambulance.

'Raised alertness and judgment ability' were assessed by measuring the magnitude of work-related fatigue, mental complaints, alertness and sleepiness. Work-related fatigue was measured by the need for recovery after working time scale (NFR) [20]. Mental complaints were measured using the Brief Symptoms Inventory (BSI) subscale of depression. Alertness and sleepiness were measured using the Epworth Sleepiness Scale (ESS) [21]. Five questions on having problems concerning thinking, fatigue, depression, alertness and sleep complaints were included in yes/no items. Examples of signal questions are: 'Have you had difficulty in determining what successive sub-activities you had to perform on a patient recently?' and 'Would you consider yourself as someone with sleeping complaints?'

The third group of requirements addressed health complaints related to 'the ability to handle emotional peak load' and were addressed using the Impact of Event Scale to assess the level of post-traumatic complaints [22]. Anxiety level was addressed through the BSI subscale of anxiety [23, 24]. Five signal questions on traumatic experiences were included in this category. Limitations due to traumatic experiences and experienced emotional load and aggression were asked about in yes/no items, e.g., 'Do you encounter any limitations in your work functioning due to severe traumatic events that you experienced recently?' and 'Did you experience any aggression during your work towards you or a colleague recently?'.

'Job-specific physical abilities' were assessed through physical performance tests and four signal questions to assess musculoskeletal complaints. The ability to lift, maintain balance, and grip strength was measured using an ambulance balance test [25]. The ability to work in a kneeled position was assessed by means of a manual cardiopulmonary resuscitation (CPR) test on a dummy on floor level (Coronel Institute of Occupational health 2013). The ability to clamber and climb, lift, maintain balance, grip strength and energetic peak load was measured in the ambulance stair climb test [26, 27]. Signal questions for limitations in physical functioning were asked concerning physical complaints, e.g., 'Have you encountered any complaints in joint or muscle which resulted in limitations in your work during the last month?'

Both 'skin complaints' and 'respiratory complaints' were assessed with one signal question each. The questions were: 'Did you experience any needle stick or bite accidents in the past 5 years?' and 'Did you experience any breathing or airway complaints after incidental or repeated exposure to a high concentration of inhaled dust, fumes, gasses, or vapor in the last six months?'

The appraisal of overall work ability, physical work ability and mental/emotional work ability were assessed by means of one question each [28]. Workers had to assign a number between zero (not able to do ambulance work) and ten (the best work ability ever experienced for ambulance work) to their current ability to perform ambulance work, compared to their best ever [11]. These appraisals of work ability were used as dependent variables in the statistical analyses.

In addition to the above-mentioned categories that possibly relate to appraised work ability, personal data were included in the WHS. Occupation and gender were taken into account, being tested for significant differences. If significant differences were found, we corrected the models for this demographic and added it to each model.

Individual scores on all questions and tests were converted to a score for that specific part of the WHS as described above. Higher scores on the independent variables were indicative of worse performance or more health complaints or problems, and thus a theoretical negative effect on work ability. All missing values were on the

dichotomous variables (3%) and have been replaced by zero (non-case). In all six categories, signal questions were asked with a yes or no outcome. If one of the signal questions per category or sub-category was answered with yes, the variable for this subject was categorized as a sign of reduced work ability and scored with a one. The pass or fail score on the Landolt C-ring test was determined, as all scores on three distances (with both eyes) had to be sufficient (≥ 0.8) to pass the test [14]. A dichotomous variable was created for the Ishihara color test [29]. To pass the whisper-speech test, no more than three mistakes were allowed per ear [19]. Need for recovery, BSI subscale scores, the ESS and the Impact of Event Scale were registered as sum score [20-22, 24]. In the job-specific physical abilities category, to pass the balance test, (where an ambulance worker needed to walk over a wooden plank with an ambulance case in each hand,) the ambulance workers' three best times out of five needed to be within 30 s. Those three best results still needed to be within 30 s when the penalty seconds were added. No more than six mistakes were allowed in total in these three fastest tests combined. The CPR test had to be completed without having to stop or get up. In the stair climb test, the end heart rate was measured using a Polar heart rate monitor and registered at the last step. The test criteria were not met when the test was not completed within 90 s, or the heart rate stayed below 85% of heart rate maximum (calculated as $220 - \text{age}$), unless the test was completed within 60 s. If the criteria of any test was not met, the test was scored as one and regarded as determinant of reduced work ability.

The normality of the scores on the three indicators for work ability appraisal questions (dependent variables) was checked by means of a visual inspection of the histogram graphs. First, differences between drivers and paramedics, and men and women on appraised work ability were tested to verify which demographic variables the analysis should be controlled for. Second, to assess which job-specific health requirement categories are associated with appraisals of work ability, six (multiple) blockwise linear regression analyses were performed per dependent variable to assess the explained variance per category. All variables of one of the six requirement categories were entered simultaneously in the model in order to be able to make a distinction between categories and their impact on appraised work ability. A model for overall work ability was created where the category with the highest explained variance was entered first, followed by the next highest, and so on, until adding a category no longer led to significant change in explained variance. The final model was based on the previous model and included a correction for gender.

To answer the *second research question* and address the contributions of physical and mental work ability to the overall appraised work ability, a linear regression was performed with overall work ability as dependent variable and the appraisal of physical and mental work ability as independent variables. As a final check, each

model in the above analyses was tested for heteroscedasticity (through a visual inspection of the residual plots) and collinearity (through collinearity statistics and VIF score). A VIF score over 2.5 was set as cause for concern of collinearity. All statistical analyses were performed using the IBM SPSS Statistics 22 software.

RESULTS

Of all 509 ambulance workers that participated in the WHS, the data of 506 could be used for statistical analyses. Of the drivers, 205 were male and 31 were female, and of the paramedics, 196 were male and 74 were female. Three percent of 11,638 values were missing. Sample demographics are shown in Table 1. The average appraised of overall, mental and physical appraised work ability was respectively 8.5 (SD= 0.88), 8.5 (SD= 1.04) and 8.7 (SD= 0.91). There was no reason to assume that heteroscedasticity or collinearity occurred in any of the tests.

Table 1: Number (n) and percentage (%) of occupation, gender and mean and standard deviation of age of ambulance workers (n=506).

	Mean	Std. Deviation	N	%
Occupation				
<i>Driver</i>			236	47
<i>Paramedic</i>			270	53
Gender				
<i>Men</i>			401	79
<i>Women</i>			105	21
Age (in years)	43.1	8.6		

The number of cases and the sum scores on the dependent and independent variables are shown in Table 2 for descriptive purposes.

No statistically significant differences in the three work ability appraisals were found between ambulance drivers and paramedics (p-values of 0.18, 0.72 and 0.81). Gender explained a significant part of variance of appraised work ability (p≤0.001) and was therefore controlled for in the later models. The category that significantly predicts overall work ability most was ‘raised alertness and judgment ability’ in both occupations. When corrected for gender, 10.4% of the variance in overall work ability was explained. When comparing occupations, ‘raised alertness and judgment ability’ explained significant variability in both occupations, with 16% for ambulance drivers and an 8% for paramedics. The categories in order of explained variance are shown in Table 3.

Table 2: Outcomes (in mean (+SD) or cases (n%)) of 506 ambulance workers in appraised work ability and health requirements in six categories.

	Mean	Std. Deviation	Cases n*	Cases %*
Dependent variables				
Overall work ability	8.5	0.88	0	0.0
Physical work ability	8.5	1.04	5	1.0
Mental/emotional work ability	8.7	0.91	1	0.2
Categories of independent variables				
1. Vision problems			81	16.0
Hearing problems			36	7.1
Acuity problems			47	9.3
Color vision problems			38	7.5
Hearing human speech problems			16	3.2
2. Raised alertness & judgment complaints			62	12.3
Work-related fatigue (0-11)	0.6	1.26	7	1.4
Symptoms of depression (0-24)	0.5	1.15	36	7.1
Alertness and sleepiness problems (0-24)	2.9	2.31	4	0.8
3. Limitations due traumatic events			102	20.2
Symptoms of PTSD (0-75)	3.9	6.21	15	3.0
Symptoms of anxiety disorder (0-24)	0.7	1.13	38	7.5
4. Musculoskeletal complaints			107	21.1
Reduced balance			10	2.0
Reduced manual CPR abilities			20	4.0
Reduced stair climb abilities			100	19.8
5. Respiratory complaints			6	1.2
6. Skin complaints			11	2.2

* Based on the cut-off values in the WHS [30].

The categories of 'gender', 'raised alertness and judgment ability' and 'emotional peak load', were added to build the final model that explained 12.2% of the variance of the appraised overall work ability.

'Job-specific physical abilities' were the highest contributing category for physical work ability after correction for gender ($R^2 = 0.13$, $p \leq 0.001$). The model that included 'gender,' 'job-specific physical abilities' and 'raised alertness and judgment ability'

explained 15.3% of the variance of the appraised physical work ability. There were no differences between ambulance drivers and paramedics.

‘Emotional peak load’ was the highest contributing category of health requirements for mental work ability when corrected for gender ($R^2 = 0.09$, $p \leq 0.001$). The final model contained the categories of ‘gender,’ ‘emotional peak load’ and ‘raised alertness and judgment ability,’ and explained 11.0% of the variance of the appraised mental/emotional work ability. In ambulance drivers, to explain variability mental/emotional work ability, the most important category was ‘raised alertness and judgment ability’ followed by ‘job-specific physical abilities’ and ‘emotional peak load.’ In ambulance paramedics, a different sequence of importance was observed. The most important category for paramedics was ‘emotional peak load’ followed by ‘raised alertness and judgment ability’ and ‘skin complaints.’

Table 3: Order of explained variance per job requirement category per dependent variable from highest to lowest.

	Overall work ability	Physical work ability	Mental work ability
1	<i>Raised alertness and judgment ability*</i>	<i>Job-specific physical abilities*</i>	<i>Emotional peak load*</i>
2	<i>Emotional peak load*</i>	<i>Raised alertness and judgment ability*</i>	<i>Raised alertness and judgment ability**</i>
3 ^{ns}	<i>Job-specific physical abilities</i>	<i>Emotional peak load</i>	<i>Job-specific physical abilities</i>
4 ^{ns}	<i>Perceive and communicate</i>	<i>Perceive and communicate</i>	<i>Respiratory complaints</i>
5 ^{ns}	<i>Skin complaints</i>	<i>Respiratory complaints</i>	<i>Perceive and communicate</i>
6 ^{ns}	<i>Respiratory complaints</i>	<i>Skin complaints</i>	<i>Skin complaints</i>

*: Significant improvement in model indicated by F-change at $p < 0.01$

** : Significant improvement in model indicated by F-change at $p < 0.05$

^{ns}: No significant improvement

For the second question, appraised physical work ability explained a statistically significant proportion of 37.4% of the variance on the total score of overall work ability, while appraised mental work ability explained a statistical significant proportion of 39.1% (Table 4). Both variables together explained 48.3% of the total variance on overall work ability appraisal. The regression model for overall work ability for drivers had an explained variance of 0.51: $Y = 2.198 + (-0.141 \text{ 'Gender'}) + (0.310 \text{ 'Physical work ability'}) + (0.412 \text{ 'Mental/emotional work ability'})$. For paramedics, the model for overall work ability had an explained variance of 0.47: $Y = 2.584 + (-0.004 \text{ 'Gender'}) + (0.318 \text{ 'Physical work ability'}) + (0.373 \text{ 'Mental/emotional work ability'})$. Gender was coded as zero for men and one for women.

Table 4: Explained variance in appraised overall work ability by physical or/and mental/emotional work ability.

	Overall work ability all ambulance workers		Overall work ability drivers		Overall work ability paramedics	
	R- square	Sig.	R- square	Sig.	R- square	Sig.
<i>Physical work ability</i>	0.374	0.000	0.397	0.000	0.360	0.000
<i>Mental/emotional work ability</i>	0.391	0.000	0.416	0.000	0.378	0.000
<i>Together</i>	0.483	0.000	0.507	0.000	0.470	0.000

DISCUSSION

Outcomes in ‘raised alertness and judgment ability’ and ‘emotional peak load’ explained statistically significant variability in appraised overall work ability. Outcomes in ‘job-specific physical abilities’ and ‘raised alertness and judgment ability’ explained statistically significant variability in physical work ability. Outcomes in ‘emotional peak load’ and ‘raised alertness and judgment ability’ explained significant variability in mental/emotional work ability. The only difference between occupations occurs for mental/emotional work ability. For drivers, ‘raised awareness and judgment ability’ explains most variance, while ‘emotional peak load’ outcomes explain most variance in paramedics’ mental/emotional work ability.

Physical and mental/emotional work ability explained almost half of the variance of appraised overall work ability. Physical and appraised mental/emotional work ability explained the same proportion of variance in overall work ability. The explained variance of overall work ability by either physical and mental/emotional work ability is almost four percent higher in drivers than in paramedics.

Although hypothesized, no significant differences were found between ambulance drivers and ambulance paramedics. Physical and mental/emotional work ability together explained only 3% more variance of overall work ability for ambulance drivers than for paramedics. It was expected that physical work ability would explain more variance than mental/emotional work ability. In contrast to studies by Klasan et al. [10] and van de Vijfeijke et al. [16], this study found that appraised mental/emotional work ability explained more variance of overall work ability than appraisal of physical work ability did. These differences can partly be explained by the fact that we included emotional work ability in the question on mental work ability. This might cause the shift to increased importance for appraised mental/emotional work ability. Another plausible explanation for the difference in findings could lie in the different type of occupations that were studied. Van de Vijfeijke et al. studied an

internet panel, with unknown occupations. Klasan et al. studied all occupations at an Institute of Emergency Medicine and only found a significant relation between the physical domain and a lower work ability on follow-up. No significant relationship between the psychological domain and lower work ability on follow-up was found. The reason for this may lie in the fact that also nurses/medical technicians were included in the study population, with possible other work demands. Finally, the differences in results can be explained by the fact that the physical and mental/emotional aspects in both studies were not measured in a job-specific way as was in ours. The WHS is a preventive strategy to monitor the health of ambulance workers and map signals that could lead to health problems and a loss of work ability. Participation in the WHS is restricted to ambulance workers actively working as ambulance driver or paramedic at the time. This selection bias causes little variation and high grades for the three work ability appraisal questions. Another possible source for selection bias is that participation in the WHS is voluntary, which can also lead to ambulance workers who have more complaints or think they are at risk, being more likely to participate.

One limitation of this study can be found in the fact that emotional work ability and mental work ability cannot be split to determine their separate contribution. Furthermore, our cross-sectional design could be regarded as a limitation. However, in this study we did not aim to find predictors for future low work ability and causal relationships, but merely how work ability relates to the job-specific health requirements in ambulance workers. Therefore, a cross-sectional design was appropriate.

A strength of the current study is that information about job-specific health requirements is gathered using job-specific instruments. By doing so, we assessed job requirements that are focused on ambulance work, while most studies on work ability focused on general aspects of health and work ability usually, in the general working population. We used the single item questions to assess work ability. The single question used to quickly assess the appraised work ability has sufficient convergent validity compared to the seven-item WAI-questionnaire [28] [31]. This method was suggested to be used during medical examinations in occupational healthcare because of its user-friendliness. The appraisal of work ability was chosen as our dependent variable to be able to gain insight in how job-specific requirement categories contribute to appraised general, physical and mental/emotional work ability.

To our knowledge, this is the first study reporting the association with appraised work ability and job-specific requirement categories in contrast to specific instruments. This approach was chosen to determine the association of different job-specific health requirement categories that are important for ambulance work and overcome the problem that separate instruments only assess a small part of a job requirement. By

categorizing variables that affect work ability, insight was gained into what aspects are assessed as important job-specific requirements to be able to perform ambulance work in a safe way. By clustering variables, information was obtained on the important requirements that have to be met by ambulance workers to satisfactorily fulfill their work and cope with their demanding working conditions without impairing their health.

Other studies reported the relation between physical and mental health and work ability, whereas this study researched the association between job-specific requirements and work ability [16, 32]. This creates an opportunity for occupational health professionals to put the different scores in job-specific health tests into perspective. Occupational physicians can start interventions where insufficient scores on important categories are observed, to improve work ability. However, insufficient scores on other categories should still be treated.

The results of this study suggest that in addition to the job-specific requirement categories, there are gender differences in the experience of work ability in ambulance workers. But although significant differences were found between men and women, this difference can be confounded by the division of gender between the occupations in ambulance work. Therefore, models have been developed for the entire group and separately for drivers and paramedics, and these were corrected for gender.

Appraised work ability is often used as a quick selfreport measure to assess the work ability in working populations. In this study, while only 1.2% of the ambulance workers graded one aspect of their own work ability as insufficient, based on the tests the number of people scoring insufficient on one or more job-specific functional tests was 68%. This is indicative that the self-appraisal of work ability might be overrated compared to measurable job-specific health requirements. On the other hand, all outcomes together explain only 17% of the variance of appraised work ability which indicates that the appraisal of work ability includes factors other than these job-specific tests. Evidently, both appraised and measured work ability outcomes contribute in different ways to the assessment of work ability of ambulance workers.

In this study, the association of different categories of requirements with the appraisal of work ability was studied. Although not more than 17% of the variance can be explained, this study shows the relative importance of the contribution of each category per work ability outcome. 'Raised alertness and judgment ability' was significant and thus present in each model without being the most important aspect for physical and mental/emotional work ability. However, it was the most important category for overall work ability. This means that measuring the magnitude of work-

related fatigue, mental complaints, alertness and sleepiness seems important to prevent a loss of overall work ability.

Compliance with Ethical Standards

Conflict of Interest: The authors declare that they have no conflict of interest.

Ethical Approval: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed Consent: Informed consent was obtained from all individual participants included in the study.

REFERENCES

1. Maguire BJ, Smith S. Injuries and fatalities among emergency medical technicians and paramedics in the United States. *Prehosp Disaster Med.* 2013;28(4):376-82.
2. Roberts MH, Sim MR, Black O, Smith P. Occupational injury risk among ambulance officers and paramedics compared with other healthcare workers in Victoria, Australia: analysis of workers' compensation claims from 2003 to 2012. *Occup Environ Med.* 2015;72(7):489-95.
3. Sterud T, Ekeberg O, Hem E. Health status in the ambulance services: a systematic review. *BMC Health Serv Res.* 2006;6:82.
4. Hansen CD, Rasmussen K, Kyed M, Nielsen KJ, Andersen JH. Physical and psychosocial work environment factors and their association with health outcomes in Danish ambulance personnel - a cross-sectional study. *BMC Public Health.* 2012;12:534.
5. Boreham CA, Gamble RP, Wallace WF, Cran GW, Stevens AB. The health status of an ambulance service. *Occup Med (Lond).* 1994;44(3):137-40.
6. Sluiter JK. High-demand jobs: age-related diversity in work ability? *Appl Ergon.* 2006;37(4):429-40.
7. Sluiter JK, Weel ANH, Hulshof C. Leidraad Preventief medisch onderzoek van werkenden [Guidance preventive medical examination of workers]. KMKA, NVAB, Coronel Institute of Occupational Health, Academic Medical Center, Amsterdam; 2013. Report No.: 13-07.
8. Ilmarinen J, Tuomi K, Klockars M. Changes in the work ability of active employees over an 11-year period. *Scand J Work Environ Health.* 1997;23 Suppl 1:49-57.
9. Pek E, Fuge K, Marton J, Banfai B, Gombos GC, Betlehem J. Cross-sectional survey on self-reported health of ambulance personnel. *Scand J Trauma Resusc Emerg Med.* 2015;23:14.
10. Klasan A, Madzarac G, Milosevic M, Mustajbegovic J, Keleuva S. Predictors of lower work ability among emergency medicine employees: the Croatian experience. *Emerg Med J.* 2013;30(4):275-9.
11. van den Berg TI, Elders LA, de Zwart BC, Burdorf A. The effects of work-related and individual factors on the Work Ability Index: a systematic review. *Occup Environ Med.* 2009;66(4):211-20.
12. Sluiter JK, van der Beek AJ, Frings-Dresen MH. Medical staff in emergency situations: severity of patient status predicts stress hormone reactivity and recovery. *Occup Environ Med.* 2003;60(5):373-4; discussion 5.
13. van der Ploeg E, Kleber RJ. Acute and chronic job stressors among ambulance personnel: predictors of health symptoms. *Occup Environ Med.* 2003;60 Suppl 1:i40-6.
14. Sluiter JK, Frings-Dresen MHW. Aanstellingskeuring, en inhoud en organisatie van een periodiek arbeidsgezondheidkundige monitor voor de ambulance sector [Pre-employment medical examination and on-employment periodical monitor for the ambulance sector]. Coronel Institute of Occupational and Environmental Health, Academic Medical Center, Amsterdam; 2005. Report No.: 05-06.
15. Boers I, Duijf P, Leerkes G, van Rhijn J, van der Werff H. Ambulance in-zicht 2012 [Ambulance in-sight 2012]. Ambulance care the Netherlands; 2013. Report No.: 12.

16. van de Vijfeijke H, Leijten FR, Ybema JF, van den Heuvel SG, Robroek SJ, van der Beek AJ, et al. Differential effects of mental and physical health and coping style on work ability: a 1-year follow-up study among aging workers. *J Occup Environ Med*. 2013;55(10):1238-43.
17. AZN. Collectieve arbeidsovereenkomst sector ambulancezorg [Labor agreement ambulance care]. 2011. p. 1-117.
18. Thiadens AA, Hoyng CB, Polling JR, Bernaerts-Biskop R, van den Born LI, Klaver CC. Accuracy of four commonly used color vision tests in the identification of cone disorders. *Ophthalmic Epidemiol*. 2013;20(2):114-21.
19. Eekhof JAH, Dijk M, van der Linden AJH, Bos JC, Goossens EM, Springer MP. De kunst van het fluisteren: de fluisterspraaktest [The art of whispering: the whisper speech test]. *Huisarts en Wetenschap*. 1997;40(2):52-3.
20. van Veldhoven MJPM, Meijman TF. Het meten van psychosociale arbeidsbelasting met een vragenlijst: de vragenlijst beleving en beoordeling van de arbeid (VBBA) [Measuring psychosocial workload with a questionnaire: the questionnaire on the experience and assessment of work (QEEW)]. Conditions DfW, editor. Amsterdam: Nederlands Instituut voor Arbeidsomstandigheden NIA; 1994.
21. Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep*. 1991;14(6):540-5.
22. van der Ploeg E, Mooren TT, Kleber RJ, van der Velden PG, Brom D. Construct validation of the Dutch version of the impact of event scale. *Psychol Assess*. 2004;16(1):16-26.
23. Derogatis LR, Spencer PM. Brief symptom inventory: BSI: Pearson Upper Saddle River, NJ; 1993.
24. Meijer RR, de Vries RM, van Bruggen V. An evaluation of the Brief Symptom Inventory-18 using item response theory: which items are most strongly related to psychological distress? *Psychol Assess*. 2011;23(1):193-202.
25. Punakallio A. Balance abilities of different-aged workers in physically demanding jobs. *J Occup Rehabil*. 2003;13(1):33-43.
26. Plat MJ, Frings-Dresen MH, Sluiter JK. Reproducibility and validity of the stair-climb test for fire fighters. *Int Arch Occup Environ Health*. 2010;83(7):725-31.
27. Teh KC, Aziz AR. A stair-climb test of cardiorespiratory fitness for Singapore. *Singapore Med J*. 2000;41(12):588-94.
28. El Fassi M, Bocquet V, Majery N, Lair ML, Couffignal S, Mairiaux P. Work ability assessment in a worker population: comparison and determinants of Work Ability Index and Work Ability score. *BMC Public Health*. 2013;13:305.
29. Cole BL. Assessment of inherited colour vision defects in clinical practice. *Clin Exp Optom*. 2007;90(3):157-75.
30. Coronel Institute of Occupational Health. Periodiek Arbeidsgezondheidskundige Monitor (PAM) voor de Ambulancesector [Workers Health Surveillance (WHS) for the Ambulance Sector]. Coronel Institute of Occupational and Environmental Health, Academic Medical Center, Amsterdam commissioned by AZN; 2013. Report No.: 10-13.

31. Olsen E, Bjaalid G, Mikkelsen A. Work climate and the mediating role of workplace bullying related to job performance, job satisfaction, and work ability: A study among hospital nurses. *J Adv Nurs*. 2017;73(11):2709-19.
32. Koolhaas W, van der Klink JJ, de Boer MR, Groothoff JW, Brouwer S. Chronic health conditions and work ability in the ageing workforce: the impact of work conditions, psychosocial factors and perceived health. *Int Arch Occup Environ Health*. 2014;87(4):433-43.

7

Work Ability and Vitality in Coach Drivers: An RCT to Study the Effectiveness of a Self- Management Intervention during the Peak Season

A. van Schaijk, K. Nieuwenhuijsen and M.H.W. Frings-Dresen

International Journal of Environmental Research and Public Health, 2019, 16(12),

2214.

ABSTRACT

Background: This randomized controlled trial (RCT) evaluates the effectiveness of a self-management toolbox designed to maintain work ability and vitality in coach drivers over their peak season.

Methods: The intervention group received a self-management intervention providing advice aimed at increasing work ability and vitality. These suggestions targeted three specific domains: work–recovery–rest balance, food and drink intake, and physical activity. At the beginning (March), middle (July), and end (October) of the coach sector peak season, work ability, vitality, work-related fatigue, psychosomatic health, sleep complaints, and perceived mental exertion of coach drivers were assessed through questionnaires.

Results: A total of 96 drivers participated in the study. Access to the toolbox did not result in significant differences between groups. Work ability and vitality decreased significantly in both groups, falling from 7.8 ± 1.3 to 7.3 ± 1.6 and from 63 ± 16.7 to 55 ± 18.7 , respectively. Work-related fatigue increased from 35 ± 31.9 to 52 ± 35.3 . Psychosomatic health complaints, sleep complaints, and perceived mental exertion also increased significantly.

Conclusions: The uptake of the intervention was too low to determine if this toolbox can maintain work ability and vitality in coach drivers when compared with a control group. Overall work ability and vitality decrease significantly as the peak season progresses, while work-related fatigue accumulates. Other interventions should be explored to ensure sustainable employability in this population.

Keywords: e-health; health promotion; prevention; sustainable employment

INTRODUCTION

Work ability is of growing importance for individual workers, employers, and sector organizations. The concept of “work ability” is one that is central to research into sustainable employment strategies. The occupational health demands created by a rising pensionable age across many European countries led to growing interest and urgency in this field of study [1]. Work ability describes the capability for satisfactory employee functioning at work while maintaining adequate physical and mental well-being. Sustainable work ability can be threatened if workers struggle or are unable to meet the work demands placed on them because of impaired health, often due to advancing age. Early intervention is, therefore, of critical importance if seeking to optimize work ability; strategies targeting a declining work ability must exert an effect before workers become too incapacitated to function. To prevent the reduction of work ability, preventive efforts should be aimed at an active workforce [2].

Coach drivers are a typical example of a working population at risk of work ability losses. Sustainable good health can be at risk during the annual coach sector peak season, characterized by increased work demands due to long working days starting and ending at irregular hours. During the peak season, the number of rides increases while the number of drivers is limited. According to a study conducted by Schuring et al., the number of working hours per week in the peak season increases by 70% compared to the off-season [3]. During peak seasons, drivers may begin their working day early in the morning one day and late at night the next. The irregular work and sleep patterns that this kind of shift work entails place high demands on the work–recovery–rest balance of drivers [4, 5]. This adds to the stress levels already inherent in operating in a road network that is becoming increasingly busy and congested [6]. Additionally, coach drivers must also work within a strict and inflexible schedule when transporting passengers to their destination.

It is, therefore, difficult for drivers to maintain adequate physical and mental health due to the irregular living and eating habits inherent in this working schedule [7, 8]. The scope for a regular or healthy eating pattern is small, and this can lead to health complaints [9]. The availability and convenience of unhealthy food on the road makes it challenging for drivers to find healthy food or maintain a healthy diet. Also, due to tight schedules and extended working hours, the window for physical activity is limited. On the road, there may be little time—or opportunity—for leisure activities, and the energy level of a coach driver after a long day is typically too low to be conducive to much physical activity.

The work demands placed on drivers in the coach sector are the same for both younger and older workers; however, the majority of the drivers are over 50 years of age. It is well established that work ability decreases with age; however, in the coach

sector, the same work demands have to be met by an aging population of coach drivers [10-12].

In addition to reduced load-bearing capabilities, the need for recovery from work is known to be higher in older workers [13, 14]. Need for recovery from work was shown to be an indicator of work-related fatigue [15]. Also, importantly, the risk of accidents increases as work-related fatigue increases. It is, therefore, vital to monitor these parameters in persons responsible for the safety of large numbers of people on public roads [16, 17]. During the peak season, it can be expected that work ability will decrease when an increase in working hours is ineluctable. However, since work ability is linked to vitality and work-related fatigue [18, 19], it can further be predicted that, next to work ability, vitality will drop and work-related fatigue will increase as the peak season progresses and the workload increases [20].

In order to achieve sustainable employability, it is necessary to develop preventive strategies aimed at maintaining work ability and vitality over the peak season. Implementation of such strategies in the coach sector is geographically complicated through drivers being on the road and, as such, it becomes difficult to organize group meetings. Prior research suggested the utility of preventative, tailored interventions that could be applied to each subgroup of drivers [17]. One such tailored strategy, which aims to maintain work ability in coach drivers, is employing the use of a self-management toolbox. Self-management instruments are being more commonly utilized because of technical developments and the increased use of smart devices, allowing them to be used by those without a fixed workplace [21].

An increasingly important field in preventative healthcare strategies is that of “e-health”. Application of this broad discipline to preventative healthcare involves using internet technologies to change behaviors associated with ill health, to deliver preventative healthcare strategies to target populations, and to share healthcare-related information [22]. An example of e-health is development of a specific application as unguided self-management interventions that create awareness of public health messages and distribute information. The goal of unguided self-management interventions is to change behavior and support people without the need for organized meetings or personal contact. These are low-cost interventional methods with the scope to reach large groups of employees. Such methods aim to provide workers with a tool that allows self-optimization of work–life balance and can be conducted at any time and place suitable to the user [23-25]. This type of intervention is particularly suited to the working environment of coach drivers and may be able to contribute to behavioral changes facilitating a healthier lifestyle and sustainable work ability [26, 27].

It is as yet unknown whether these interventions are useful in maintaining work ability and vitality in coach drivers during their peak season. This led to the formulation of

the following research question: Can reductions in coach drivers' work ability and vitality over the coach sector peak season be prevented through the use of self-management interventions targeting work–recovery–rest balance, eating habits, and physical activity at work?

MATERIALS AND METHODS

Participants and Procedure

The study was designed as a randomized controlled trial with equally sized intervention and control arms running in parallel. The follow-up period was seven months (the length of the peak season) and comprised three discrete measurement points. We performed a sample size calculation with a significance level of 0.05 and a power of 0.80. However, the estimated effect size was impossible to predict. When aiming to be able to pick up changes between groups with a medium effect size of 0.5, we estimated needing a group size of 31. Because we did not know what effect size to expect, we chose to include as many drivers as possible for this study to increase the ability of finding differences with a lower effect size. In collaboration with the coach sector organization, we recruited drivers through a variety of strategies—contacting coach companies, visiting events where coach drivers would be present, and placing adverts in job-specific media—over the period from November 2017 to March 2018. After giving informed consent, all drivers who agreed to participate received a baseline questionnaire. No protocol changes were made after the start of the trial.

Baseline questionnaires were administered at the beginning of the peak season (March). The follow-up questionnaires were administered in the middle (July) and at the end of the peak season (October). All questionnaires were digitally administered, meaning that drivers could fill out the questionnaire at the time and location of their choice.

Our research was conducted in accordance with the Declaration of Helsinki [28]. The research proposal was submitted to and approved by the Medical Ethical Committee of the Academic Medical Center, who decreed that a comprehensive evaluation was not required since this study was not subject to the Medical Research Involving Human Subjects Act (W16_153#16.177). This research was registered in the trial register as NTR 7125. The privacy impact assessment was registered as AMC2017-422. For this study, CONSolidated Standards Of Reporting Trials (CONSORT) checklists for reporting e-health interventions and parallel group randomized trials were used as guidelines [29, 30].

Randomisation

All drivers who completed the baseline questionnaire were numbered based on their email addresses. Each number was randomly assigned to either the intervention or control group by an online research randomizer, generating two equally sized lists of numbers (block randomization with blocks of 46 drivers). This was done by an independent researcher (J.S.) who did not possess any personal or other information regarding the drivers nor the allocation details of individual numbers. This researcher reported back the numbers (1–96) as either belonging to the control or intervention group; intervention was subsequently allocated based on these groups.

Measurements

Three questionnaires were developed to study the work-related health of coach drivers. At each of the three time points, work ability, vitality, work-related fatigue, psychosomatic health, sleep complaints, and perceived mental exertion of coach drivers were assessed.

Main Outcome

This trial centers upon the outcome of work ability. Given that the concepts of vitality and work-related fatigue were shown to be closely related to this outcome [18, 19], these were also examined. Work ability was measured using the work ability score (WAS) [31], the first question of the work ability index (WAI). In addition to general work ability, physical and mental work ability were also appraised by drivers, using a scale of 0 to 10. These two questions were based on the second question of the WAI, where 0 represents no work ability at all, and 10 describes the best work ability ever experienced [32]. A higher value represents a greater work ability. This single-item assessment showed sufficient convergent validity with the complete WAI and is suitable for the systematic screening of work ability [33].

Vitality was measured with the vitality subscale of the Short Form-36 (SF-36) [34]. Items from this subscale concern user evaluation of levels of energy and fatigue, for example, “How much of the time during the past four weeks did you have a lot of energy?”. This scale comprises four items and gives a total score between 0 and 100. Those with high scores felt lively and energetic over the past four weeks, while those with low scores felt tired and exhausted. The internal consistency of this scale was 0.82 (α) and the test–retest correlations were 0.76 and 0.63 after two and six months, respectively [34].

Work-related fatigue was measured through the proxy of need for recovery, using the need for recovery scale of the questionnaire on experience and evaluation of work [15]. This scale contains 11 items and, after transformation, gives a score of between 0 and 100, with a higher score indicating a greater recovery requirement. Need for

recovery was shown to be able to predict sickness absence in truck drivers [35]. This scale can be used in both individual and group assessments of need for recovery, with a Cronbach's α value between 0.81 and 0.92 in different subgroups based on education, age, and gender [36].

Secondary Outcomes

In addition to the primary outcomes described above, other factors relevant to work-related health during the coach sector peak season were measured. Psychosomatic health was examined using a questionnaire to measure health complaints (Vragenlijst Onderzoek Ervaren Gezondheid/VOEG) [37, 38]. This 13-item dichotomous questionnaire details a number of common (work-related) health complaints (such as fatigue, headache, and back pain). The total sum value is converted to a 0–100 score, with a higher score representing a greater number of health complaints. These 13 items have a Cronbach's α value of 0.67 [37].

Sleep complaints were recorded using the Groningen sleep quality scale (GSKS) [39]. This scale consists of 14 items with a total score of between 0 and 100. The items on this scale are ranked in order of the severity of the complaints, with a greater number of sleep complaints giving a greater score. The Cronbach's α value of this scale is 0.89 [39].

Perceived mental exertion was measured using the perceived mental exertion scale (in Dutch, SEB) [38, 40, 41]. This scale comprises 19 items in which a driver indicates on a five-point scale which of two answers best describes their situation. From these items, the perceived mental exertion is expressed on a scale between 0 and 100, with a higher score being less favorable. For example, one of the items included in the SEB is “difficulty in planning your own actions vs. working effortlessly”. The Cronbach's α value ranges between 0.95 and 0.97 according to age group and shift work type [41].

Process Measurements

The questionnaire contained questions detailing individual characteristics including age, body mass index (BMI), number of years worked as a coach driver, working hours, sleep hours, work characteristics, regular food and drink intake, and physical activity. This allows for the analysis of driver behavior during the peak season. In addition, changes in behavior can be observed, as well as the points in time that these changes took place. Alongside these characteristics, eight questions were included to examine the influence of work on private life. For example, one of these questions asks “Has your private life been adversely affected by irregular working hours?”.

The follow-up questionnaires distributed to the intervention group also included questions regarding the use of the intervention and its three constituent domains. The

process measurements contained questions about the extent to which specific suggestions may have helped preserve vitality the most, and which suggestions were most commonly used.

Intervention

The drivers assigned to the intervention group received a digital self-management toolbox at the start of the peak season after completing the baseline measurements. This toolbox was an interactive pdf document with suggestions and corresponding assignments based around three domains: work–recovery–rest balance, food and drink intake, and physical activity. Drivers were free to print it and write their data and remarks down on paper or use it digitally on any (mobile) device. Each domain of the toolbox included a general introduction of the topic and its importance. After a general introduction, between 11 and 14 (to coach drivers tailored) suggestions were given for each domain, followed by assignments. The advice provided is based on scientific research and is aimed at influencing behavioral changes in order to improve health parameters, and ranging from easy-to-implement tips to suggestions that required more effort, but which could improve health over a longer period. Examples of suggestions include the ideal duration of a power nap [42], ways by which to increase alertness [43], modifying the eating pattern, advice regarding posture, and suggestions for performing quick on-the-spot workouts. Research showed, for example, that physical activity can maintain work ability and reduce levels of work-related fatigue [44]. Each of the three domains of the toolbox was preceded by suggestions for intrinsic motivation, and drivers were asked to record their motivations for engaging with the toolbox. The suggestions provided in the toolbox were derived through consulting professionals in the relevant scientific fields, such as a dieticians and movement scientists. In the assignments, drivers were required to record their behaviors, set a goal, and indicate which suggestion they intended to use and why they felt that this was going to be successful. In addition, the toolkit sought to include the concept of “peer support”. Drivers were divided into groups of three, all working over the peak season, and were instructed to keep in regular contact through the assignments in order to motivate each other, to encourage engagement with the toolbox, and for evaluative purposes. Peer support and interpersonal contact were previously shown to be of benefit in stimulating behavioral changes in e-health interventions [22]. There was no contact with researchers or other professionals after the distribution of the toolbox to the intervention group, other than through the digital questionnaires to evaluate the intervention.

Statistical Analysis

In order to analyze the differences between the intervention and control groups over the measurement period, analyses of covariance (ANCOVA) were used. Differences in

baseline measurements were corrected for using the baseline values as covariates, with the grouping variable as the fixed factor. The assumptions of the ANCOVA analyses were tested to make sure no abnormalities in the data were observed [45]. For the analysis, an intention-to-treat analysis was used to study the results as measured.

To identify temporal trends in work ability, vitality, and work-related fatigue during the peak season, data from the baseline, intermediate, and final measurements were compared using different paired sample *t*-tests. This allowed the identification of changes within groups that occur between the three time points during the peak season. Differences in recovery opportunities between the three measurements were tested with McNemar tests for binary paired outcomes.

To check for selective dropout, a missing values analysis was performed. To control for missing values, an additional analysis was performed with the last observation carried forward method to correct for missing data in a conservative way, when representing data of the entire population.

Secondary Analyses

A separate secondary analysis was performed to check for the influence of compliance with the intervention on the outcome measures. In the per-protocol analysis, only the participants who complied with the protocol were considered (sufficient use of the toolbox: >10%).

Process Evaluation

In an additional process evaluation, we asked whether or not the drivers had the feeling that the tips from the toolbox helped them to feel more vital. This was done for each individual tip per domain, where drivers could indicate what domain they used the most, and which tips within the domain were most helpful.

RESULTS

A total of 124 of an estimated eligible 6000 drivers working in the private passenger transport sector gave informed consent to participate in this study. The exact number of drivers invited is unknown because the coach sector organization was responsible for inclusion of drivers due to privacy data sharing restrictions. These drivers received the first questionnaire in March 2018. The 96 drivers who completed the baseline questionnaire were randomly assigned to either the intervention or control group, creating two equal groups of 48 drivers. Of the 96 drivers, the average age was 53 ± 10.8 years, and 85% were male. Mean BMI was 28 ± 4.6 , and drivers were working as a coach driver for an average of 16 ± 11.2 years. In total, 28% drove shuttle services

and multi-day trips, 24% drove single-day trips, and 48% drove a combination of both. Population demographics are shown in Table 1.

Table 1: Population demographics (mean and standard deviation) for the intervention and control groups.

Demographic	Intervention Group	Control Group
Number of drivers	48	48
Percentage of male drivers	88	83
Age	55 (9.1)	52 (12.1)
Body mass index (BMI)	28 (4.6)	28 (4.6)
Years working as coach driver	14 (11.1)	17 (11.2)
Percentage not physically active at baseline	31	31

At the second time point, in the middle of the peak season (July 2018), 70 drivers (33 from the intervention arm; 37 from the control arm) filled out the questionnaire. A total of 62 drivers (34 from the intervention arm; 28 from the control arm) completed the final questionnaire at the end of the peak season (October 2018). A flow diagram of driver participation is shown in Figure 1.

Figure 1: Flow diagram of the enrolment, allocation, follow-up, and analysis process.

Differences between Intervention and Control Groups

At baseline, no significant differences between the control and intervention group were observed for any of the variables. Both primary and secondary outcomes of drivers who received the intervention did not differ from the control group over the measurement period (in either the intermediate or final measurements). Access to the toolbox was not associated with any significant change in measures of work ability, vitality, work-related fatigue, psychosomatic health, sleep complaints, or perceived mental exertion, recorded at the middle and end of the peak season when corrected for the baseline score (Table 2). The effect sizes ranged between 0 and 3.5%, demonstrating that use of the toolbox had no effect on the primary and secondary outcomes.

Table 2: Values (mean $\pm$ SD) for the primary and secondary outcome variables for the intervention and control group at the baseline, intermediate, and final measurement, based on the intention-to-treat protocol. The estimated effect size, significance (*p*-value), and *F*-value of the ANCOVA analyses of both analyses are shown in the right columns. Both the intention-to-treat and the per protocol analysis results are presented.

Outcome measure	Baseline		Intermediate		Final		Est. Effect size		Sign.		F	
	Int.	Contr.	Int.	Contr.	Int.	Contr.	ITT	PP	ITT	PP	ITT	PP
Number of drivers	48	48	33	37	28	34						
Work ability score	7.8 (1.4)	7.8 (1.2)	7.7 (1.5)	7.3 (1.6)	7.0 (1.5)	7.5 (1.6)	0.021	0.016	0.261	0.328	1.290	0.972
Physical work ability	7.8 (1.3)	7.8 (1.3)	7.5 (1.9)	7.2 (1.7)	7.3 (1.6)	7.5 (1.4)	0.000	0.000	0.945	0.894	0.005	0.018
Mental work ability	7.9 (1.2)	7.8 (1.4)	7.5 (1.8)	7.4 (1.8)	6.9 (1.8)	7.2 (1.5)	0.009	0.012	0.473	0.393	0.522	0.742
Vitality	63 (15.2)	63 (18.2)	57 (18.4)	57 (18.1)	55 (20.1)	54 (17.8)	0.000	0.010	0.954	0.934	0.003	0.007
Work-related fatigue	33 (30.6)	37 (33.0)	46 (35.6)	48 (37.0)	53 (34.5)	52 (36.6)	0.002	0.010	0.719	0.447	0.131	0.587
Psychosomatic health	25 (19.7)	28 (22.0)	30 (24.6)	31 (24.0)	36 (27.1)	39 (25.4)	0.001	0.001	0.806	0.773	0.806	0.084
Sleep complaints	19 (23.3)	26 (24.6)	23 (21.3)	36 (29.8)	28 (25.8)	33 (29.7)	0.001	0.000	0.788	0.903	0.073	0.015
Perceived mental exertion	25 (19.2)	26 (19.6)	28 (20.9)	32 (22.1)	27 (16.5)	33 (23.5)	0.035	0.031	0.151	0.176	2.121	1.878

Int. = intervention group, Contr. = control group, ITT = intention-to-treat analysis, PP = per protocol analysis. Estimated effect size shown as partial eta squared, Sign. = significance level, F = F-statistic of ANCOVA.

No significant differences between intervention and control groups were found in any of the parameters at any of the three measurement points. Food and drink intake did not change significantly between the intervention and control group. The number of drivers not being physically active increased significantly over the measurement period in both groups, as did the number of hours worked. Neither of these trends, however, differed significantly between groups. The mean number of hours of sleep per night remained unchanged over the measurement periods in both groups. These values are shown in Table 3. In a secondary analysis using the per protocol analyses, no significant differences in the primary and secondary outcomes were found between the intervention and control groups. The numbers of drivers included in the per protocol analysis were 26 and 47 for the intermediate measurement, and 26 and 36 in the final measurement for the intervention and control group, respectively.

Table 3: Values (mean $\pm$ SD or %) for the process measures for the entire group of drivers at the baseline, intermediate, and final measurements.

Process measure	Baseline	Intermediate	Final
Number of drivers	96	70	62
Work hours per week ¹	37 (15.4)	50 (18.4) **	45 (17.6) **
Hours of sleep ²	7.8 (2.2)	7.7 (2.9)	7.8 (2.6)
Not physically active (%) ³	31	42	52 *
Trouble staying alert during evening and night hours (%) ³	29	36	42
Self-assessed as very tired (%) ³	22	41 *	53 **

¹ Mean of last three weeks, ² between last two working days, ³ during the last two weeks. Significant differences compared to baseline measurement are marked with asterisks: * $p < 0.05$, ** $p < 0.01$. For the McNemar test, baseline–intermediate $n = 70$, baseline–final $n = 62$.

Trajectory of Work Ability, Vitality, and Work-Related Fatigue

Because there were no significant changes between the intervention and control group over the measurement period, the combined cohort of coach drivers was evaluated as one group. For this group, a decline in work ability and vitality, and an increase in work-related fatigue were evident. Figure 2 and Figure 3 illustrate these changes in primary outcomes. Also, the secondary outcomes recorded also demonstrate a global decline in health parameters. Psychosomatic health complaints increased from 26 (at baseline measurement) to 38, on a scale of 0 to 100 ($p < 0.01$). Sleep complaints and perceived mental exertion increased from 22 to 31, and from 26 to 31, respectively ($p < 0.05$).

Figure 2: Trajectory of general, physical, and mental work ability during the peak season. Mean and SD are shown with error bars. A score of 0 stands for no work ability at all, and 10 corresponds to the best work ability ever experienced. Values with * or ** showed significant changes compared to baseline measurements: * $p < 0.05$, ** $p < 0.01$.

Figure 3: Trajectory of vitality and work-related fatigue during the peak season. Mean and SD are shown with error bars. A lower score indicates a lower vitality and less work-related fatigue. Values with ** showed significant changes compared to baseline measurements: ** $p < 0.01$.

The work ability score decreased significantly from 7.8 at baseline to 7.3 at the final measurement of the peak season. It is noteworthy that mental work ability declined the most for all work ability parameters recorded, falling from 7.9 at baseline to 7.1 at the final measurement, while physical work ability was reduced from 7.8 to 7.4.

Vitality decreased, and work-related fatigue increased most dramatically over the period preceding the intermediate measurement, with vitality falling from 63 to 55 and fatigue increasing from 35 to 47 between baseline and intermediate measurements. The values of the final measurements were all significantly different from the baseline measurements. This is depicted in Figure 3.

The values in Figure 2 and Figure 3 illustrate that all values changed in the adverse direction over the measurement period. Aside from the primary and secondary outcomes, recovery opportunities also became limited during the peak season. This shows that there is less time for leisure due to increased working hours while maintaining the hours of sleep (Table 3). Driver perception of recovery opportunities is shown in Table 4. Statistical differences were calculated only between drivers that filled out both questionnaires.

Table 4: Number of drivers and percentage of drivers that answered “yes” on the corresponding questions on recovery opportunities.

Recovery opportunity	Baseline	Intermediate	Final
Number of drivers	96	68	62
Could you interrupt your work at times when you felt it necessary?	55	46	50
Could you determine the start and end time of your work yourself?	11	12	10
Could you decide when you took a break?	38	44	45
Could you include a separate day off when you wanted?	60	38 **	44
Have you been recalled from leave/a free day?	13	18 *	16
Were your work and rest times well organized?	91	82	76
Were there opportunities for you to work at hours that fit your private situation?	69	43 **	42 **
Has your private life been adversely affected by irregular working hours?	45	53	61

Significant differences compared to baseline measurement are marked with asterisks: * $p < 0.05$, ** $p < 0.01$. For the McNemar test, baseline–intermediate $n = 68$, baseline–final $n = 62$.

Table 4 illustrates that drivers perceived the number of recovery opportunities to decrease as the peak season progressed. Table 4 presents percentages derived from

the entire driver population participating at that time point. For statistical analyses, only the drivers that completed both questionnaires were included, to allow a true comparison of changes in driver perception (i.e., comparing the same divers at both measurement points) to eliminate bias.

On average, drivers in the intervention group reported that they used $34 \pm 25.5\%$ and $34 \pm 20.0\%$ of the suggestions included in the toolbox during the periods preceding the intermediate and final measurements (Table 5). The process evaluation showed that the domain and corresponding tips on food and drink uptake were used most. The tips that were most helpful for each domain were keeping contact with passengers and colleagues to remain alert on the road in the work–recovery–rest domain, suggestions to have variations in food and drink intake in the food and drink intake domain, and a tip to improve posture during driving and other work activities in the physical activity domain.

Table 5: Number of drivers and usage (mean % $\pm$ SD) of the separate domains in the toolbox in the intervention group in the intermediate and final measurements.

Toolbox domain	Intermediate	Final
Number of drivers	30	28
Work–recovery–rest balance	32 (27.6)	31 (22.3)
Food and drink intake	38 (32.0)	38 (27.8)
Physical activity	31 (24.7)	32 (20.4)

Although the mean usage of the toolbox was not very high, drivers reported an increase in vitality attained by applying the advice given in the toolbox. Most of the drivers did not engage in peer support, but those who did (18%) contacted their peers more than four times per month.

The driver-perceived effectiveness of the toolbox on vitality is presented in Table 6. Despite the low usage, some drivers described feelings of maintained or increased vitality. Only one driver reported a reduction in perceived vitality following the use of the toolbox.

Table 6: Perception of effectiveness (*n* (%)) of the toolbox on vitality according to the drivers' perception in the intermediate and final measurements, based on the following question: Do you have the feeling that you are more vital now than you would have been without the use of the toolbox?

Perceived vitality	Intermediate		Final	
	n	%	n	%
Number of drivers	30		28	
Less vital	0	(0)	1	(4)
Equally vital	17	(55)	15	(54)
More vital	4	(14)	3	(11)
I do not know	9	(31)	9	(32)

DISCUSSION

Key Results

A specifically designed toolbox based on principles of self-management did not lead to maintained levels of work ability, vitality, work-related fatigue, psychosomatic health, sleep complaints, or perceived mental exertion during the peak season. All measured work-related aspects of health showed significant deterioration as the peak season progressed, regardless of whether the driver received the toolbox. The process evaluation revealed that coach drivers used the toolbox to such a limited extent that differences in these parameters between intervention and control groups were unlikely to occur.

Comparison with Literature

Compared to a study conducted during the (August) 1996 coach sector peak season [46], greater values were obtained for work-related fatigue in this study of the 2018 peak season. Work-related fatigue was 33 at the end of the peak season of 1996, measured on the same scale as used in our research, while the score at the start of the peak season of 2018 was 35, increasing to 52 over the measurement period. Psychosomatic health, sleep complaints, and experienced mental stress at the end of the peak season were also higher than the values obtained in 1996 (24, 22, and 27, respectively). It is hard to pinpoint the causes for these differences, but they may be caused by an increased workload and reduced driver capabilities for dealing with this due to more advanced age. The average age of participating drivers increased by nine years from the 1996 study (53 compared to 44 years of age).

With an average WAS of 7.3 at the end of the peak season, the work ability values of this study are considered moderate in the classification of Gould et al. (2008) [47]. A mean WAS score of 7.3 is considerably lower than scores for ambulance workers, an

average working population, a heterogeneous sample of workers, elderly construction workers, or elderly workers, (8.5, 8.1, 7.95, 8.0, and 8.57, respectively). The score is comparable to people returning to work after sick leave (7.4) [32, 33, 48-51]. Vitality scores at the beginning of the peak season are comparable with values found in the literature [52]. These values in the literature are averages at a certain point and do not reflect a period with increased workload.

Strengths and Limitations

A strength of this study was that it adhered strictly to the recommendations for randomized controlled trials (RCTs). For instance, randomization took place after the baseline measurement, eliminating any selection bias [53]. Individual differences in baseline values were controlled for by using the baseline values as covariates to avoid conditional bias. To enhance the quality of the intervention, recommendations for self-management interventions were implemented in a structured way. In self-management interventions, it is important to have a strong theoretical foundation, carefully designed structure, and pathway of action, and to include user reflection on behavior [21]. The toolbox aimed to increase intrinsic motivation and reflection on current and desired driver behavior. Previous research studied measures that can promote health and work ability in truck drivers [54]. The authors considered several criteria to be of importance in this study: a wide range of options, measures that help overcome obstructions, and the inclusion of educational material. The final criterion is that both employer and employee are involved. The toolbox presented in our study met all criteria other than this final one. The reason to not actively involve the employer was that it was the wish of the coach sector organization to implement an intervention that drivers could use on their work ability without the assistance of other parties.

A positive attribute of this study was that it succeeded in conducting an RCT in a population with a high workload, minimal free time, irregular hours, and no fixed workplace location. This study stayed close to practice while following the methodological design. A realistic intervention was implemented in such a way that allowed the effects of this intervention to be monitored over the peak season. High external validity was established because there was no check on compliance, as there would not be any checks after this toolbox is implemented in the coach sector in the future; therefore, the true effects of this intervention were studied. It is important to keep in mind that the intention was for the intervention to work under the conditions of normal practice and not in a restricted controlled situation.

This study aimed to include as many coach drivers working over the peak season as possible. However, the sample size was still low. One reason for this may be the difficulty of reaching these coach drivers. The coach sector association only had

contact details for individual coach companies, and not of individual drivers. The low sample size did not affect the results because no trend was observed in the differences in primary or secondary outcomes between intervention and control groups. The small effect size predicts that even a larger sample size would not lead to significant results. Despite the small sample size, significant trends can be seen when examining the entire cohort of drivers over time. In our measurement process, we chose to include all drivers that filled out questionnaires instead of only the drivers that filled out all questionnaires in order to avoid bias; not doing so would have allowed selective dropout (i.e., sick leave) to change the mean values at baseline. The intention-to-treat analysis showed the population as it was during the peak season.

Generalizability

We would expect the outcomes of this study to apply to the entire population of Dutch coach drivers, as participating drivers were drawn from a number of different companies and locations. The high average age and percentage of males in the cohort are representative of the wider Dutch coach driver population. Selection bias was eliminated through randomization to intervention and control groups. It is plausible that some drivers did not participate in the study because they felt too busy during the peak season, or that only drivers who were busy during the peak season participated because they endorsed the importance of this study.

Unfortunately, we do not have information regarding the reasons for dropout. There was no attrition bias between groups; however, when selective dropout rates were examined, those who dropped out after the baseline measurement possessed more positive baseline values. This would indicate that the worsening in health parameters seen over the study period is slightly exaggerated. However, additional conservative analyses with the 'last observation carried forward'-method still showed significant changes in the negative direction on both the primary and secondary outcomes (see Table A1). These values are without a possible decrease in health parameters in the higher values of the drivers that discontinued after baseline measurement. The values analyzed using the last observation carried forward method are, therefore, an underestimation of the effect of working during the peak season on health parameters. In their potential impact on value differences between groups, we would expect this effect to be minor and balance out between groups because of randomization.

The results of this study can be extrapolated to other countries within the European Union, as the driving and rest times for drivers are controlled by European legislation (EG nr. 561/2006) [55]. We would expect that the working conditions of drivers in other European countries are comparable and drivers experience similar difficulties. In the United States, Canada, and Australia, however, the maximum number of driving

hours per day permitted is higher than in Europe [56-58]. Therefore, it can be hypothesized that, if these driving hours are also irregular, sustainable employment may be harder to achieve in drivers from these countries.

In The Netherlands, the number of rest hours required between journeys was increased over 20 years ago through a collective labor agreement. A study found that drivers who received more rest time between trips had lesser values for work-related fatigue and fewer psychosomatic health complaints [3]. Since this change in legislation, driver workload may have increased, resulting in an increased need for recovery.

In other driving occupations, the same rules apply that govern working hours. However, work ability and work-related fatigue in these professions may benefit from more predictable start and end times, or a more consistent schedule. Coach drivers also have to take a large number of passengers and their wishes into account, which may cause extra stress.

Interpretation

The results of this study clearly demonstrate a significant downward trend in health parameters during the peak season. Values at the start of the peak season were comparable to other working populations as stated above. However, toward the end of the peak season, these values became alarming. The percentage of individuals exhibiting poor work ability at the start of the peak season was 6%. This increased to 11% at the end of the peak season. Although this increase may seem low, this represents nearly a twofold increase of workers at risk of sick leave. Our findings show that 20 to 27% of drivers had a relevant decrease in work ability (general, mental, or physical) over the peak season [49]. Although the mean work ability score was still acceptable, 35% of drivers scored higher than the threshold (over 50) for work-related fatigue at baseline. This percentage increased to 52% at the intermediate measurement and 61% at the final measurement, illustrating that the majority of drivers experience too much work-related fatigue. Since work-related fatigue is related to occupational accidents, these figures underline the necessity for interventions in the coach sector [16, 17, 59]. Vitality decreased over time; however, a cutoff point was not determined. Since significant decreases occurred in vitality between measurements, this should be interpreted as significantly more fatigue corresponding to the baseline scores in Table 2.

Personal demographic characteristics might explain work-related fatigue, as it is known that older workers have a higher need for recovery [13]. Also, BMI can cause difficulties in certain parts of work as a coach driver. Since gender and BMI did not change over the peak season and the analyses were corrected for baseline values, the

changes in the primary and secondary variables were not attributable to these characteristics. In an additional analysis, we found no evidence for influence of BMI, gender, and age on the decrease in work ability, vitality, and increase in work-related fatigue. Future studies may shift focus toward which personal and work aspects cause the work ability and vitality to decrease and the work-related fatigue to increase.

The fact that the work-related health of a large proportion of drivers can worsen significantly over a relatively short period demonstrates that measures to improve sustainable employability are of great importance in this sector.

Implications

Although no significant differences were found between the intervention and control group, our findings show the need for interventions to combat reductions in work ability during the peak season. Such interventions should be aimed at improving sustainable employment in an aging workforce of coach drivers. Given the increased average age in this population (two-thirds are now older than 50 years, half of whom are also older than 60 years), ensuring sustainable employability is of particular importance because the load-bearing capacity decreases on average. This, in combination with the increased work-related fatigue and a higher mental workload experienced by older drivers, has the potential to result in driving errors that may endanger the safety of both passengers and other road users [8, 60, 61].

Previous research advised that monitoring work-related fatigue can help employers and occupational health agencies to develop preventative strategies to increase sustainable employability, given that the need for recovery is an important predictor of future sickness absence [35, 62]. A study in older taxi drivers recommended preventive monitoring and early interventions [63]. This monitoring can be in the form of a preventive medical examination with an associated advice on identified problem areas. Since the reductions in the mean health parameter scores observed in this research were not only attributable to a few low-scoring drivers, interventions should be aimed at the entire coach driver workforce. Future research should aim to find alternate ways to maintain work ability and to improve sustainable employability since the use of a self-management intervention did not yield the desired results. The effectiveness of a more active approach should be tested in the future.

Work ability score, vitality, work-related fatigue, psychosomatic health, sleep complaints, and perceived mental exertion were proven to be useful outcome measures, which can be used in future research because they are able to detect changes over the peak season. Future effectiveness studies for this group should, therefore, include these measurements during the peak season in order to demonstrate the preventive effect of interventions.

Research on the effective underlying mechanisms of interventions is desired in a population that is always on the road in order to improve the preventive effect of interventions. Additional research on the personal characteristics that influence the main outcomes can contribute to improving occupational care and improving sustainable employability. In research in this older population, it is important to address the aspects that may affect the effectiveness of interventions. Therefore, it is suggested to shift the focus of research toward identifying the working component in interventions while still focusing on health improvement.

CONCLUSIONS

A decrease in work ability was not prevented by means of a self-management toolbox in this study. Uptake of the intervention was too low to be able to reliably determine whether such an intervention could lead to an effect between groups. User evaluation showed that the content of the intervention was considered to be positive, but without yielding any significant results. Overall work ability and vitality decreased significantly, and work-related fatigue accumulated as the peak season progressed. Passive intervention with the interactive toolbox was not used enough in this study population and, therefore, the coach sector should explore active interventions to ensure that work ability is maintained in the peak season and that long-term sustainable employability is attained.

Author Contributions: A.v.S., K.N., M.F.-D., and Judith Sluiter (J.S.) (passed away 14 May 2018) were involved in the conceptualization of this study. The methodology was shaped by A.v.S., J.S., M.F.-D., and K.N. Randomization was performed by J.S., and intervention was distributed accordingly by A.v.S. Statistical analyses were executed by A.v.S., and data checks were completed by K.N. A.v.S. wrote the original draft, and K.N. and M.F.-D. reviewed and edited the draft. Project administration was the responsibility of A.S. Funding for this project was acquired by M.F.-D. and J.S.

Funding: This research was funded by the European Social Fund (Directorate General for Employment, Social Affairs, and Inclusion), the Ministry of Social Affairs and Employment in The Netherlands (subsidy granted: ESF 2014-2020 Sustainable employability of regions and sectors, subsidy number: 2016EUSF20164).

Acknowledgments: The authors would like to thank all drivers participating in this research, and the coach sector organization for their help with the recruitment of the participating drivers.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

Table A1: Number of drivers and values (mean $\pm$ SD) for the outcome variables for the entire group of drivers at the baseline, intermediate, and final measurements, with the last observation carried forward and intention-to-treat analysis.

Outcome measure	Baseline	Intermediate	Final
Number of drivers	96	96	96
Work ability score	7.8 (1.3)	7.6 (1.4)	7.5 (1.4) *
Physical work ability	7.8 (1.3)	7.5 (1.7) *	7.5 (1.5) *
Mental work ability	7.9 (1.3)	7.6 (1.6) **	7.3 (1.6) **
Vitality	63 (16.7)	58 (17.1) **	58 (17.4) **
Work-related fatigue	35 (31.9)	44 (34.5) **	45 (34.2) **
Psychosomatic health	26 (20.8)	28 (23.1)	31 (24.6) **
Sleep complaints	22 (24.0)	27 (25.2) *	27 (25.6) *
Perceived mental exertion	26 (19.3)	29 (21.3)	30 (21.4) *

Values with * or ** showed significant changes compared to baseline measurements: * $p < 0.05$, ** $p < 0.01$.

REFERENCES

1. Finnish Centre for Pensions. Retirement ages in 2050 [Webpage]. 2018. Available from: <https://www.etk.fi/en/the-pension-system/international-comparison/retirement-ages/> (Accessed 14th December 2018).
2. Ilmarinen J, Tuomi K. Work ability of aging workers. *Scand J Work Environ Health*. 1992;18 Suppl 2:8-10.
3. Schuring M, Sluiter JK, Frings-Dresen MH. Evaluation of top-down implementation of health regulations in the transport sector in a 5-year period. *Int Arch Occup Environ Health*. 2004;77(1):53-9.
4. Anund A, Fors C, Ihlstrom J, Kecklund G. An on-road study of sleepiness in split shifts among city bus drivers. *Accid Anal Prev*. 2018;114:71-6.
5. Lee S, Kim HR, Byun J, Jang T. Sleepiness while driving and shiftwork patterns among Korean bus drivers. *Ann Occup Environ Med*. 2017;29:48.
6. Evans GW, Carrere S. Traffic congestion, perceived control, and psychophysiological stress among urban bus drivers. *J Appl Psychol*. 1991;76(5):658-63.
7. de Zwart BC, Frings-Dresen MH, van Dijk FJ. Physical workload and the aging worker: a review of the literature. *Int Arch Occup Environ Health*. 1995;68(1):1-12.
8. Cantin V, Lavalliere M, Simoneau M, Teasdale N. Mental workload when driving in a simulator: effects of age and driving complexity. *Accid Anal Prev*. 2009;41(4):763-71.
9. Tse JL, Flin R, Mearns K. Bus driver well-being review: 50 years of research. *Transportation research part F: traffic psychology and behaviour*. 2006;9(2):89-114.
10. Costa G, Sartori S. Ageing, working hours and work ability. *Ergonomics*. 2007;50(11):1914-30.
11. van den Berg TI, Elders LA, de Zwart BC, Burdorf A. The effects of work-related and individual factors on the Work Ability Index: a systematic review. *Occup Environ Med*. 2009;66(4):211-20.
12. Kloimüller I, Karazman R, Geissler H, Karazman-Morawetz I, Haupt H. The relation of age, work ability index and stress-inducing factors among bus drivers. *International Journal of Industrial Ergonomics*. 2000;25(5):497-502.
13. Kiss P, De Meester M, Braeckman L. Differences between younger and older workers in the need for recovery after work. *Int Arch Occup Environ Health*. 2008;81(3):311-20.
14. Ilmarinen J, Rantanen J. Promotion of work ability during ageing. *Am J Ind Med*. 1999;Suppl 1:21-3.
15. Sluiter JK, de Croon EM, Meijman TF, Frings-Dresen MHW. Need for recovery from work related fatigue and its role in the development and prediction of subjective health complaints. *Occup Environ Med*. 2003;60 Suppl 1:i62-70.
16. Duke J, Guest M, Boggess M. Age-related safety in professional heavy vehicle drivers: a literature review. *Accid Anal Prev*. 2010;42(2):364-71.

17. Useche SA, Gomez V, Cendales B, Alonso F. Working Conditions, Job Strain, and Traffic Safety among Three Groups of Public Transport Drivers. *Saf Health Work*. 2018;9(4):454-61.
18. Ahlstrom L, Grimby-Ekman A, Hagberg M, Dellve L. The work ability index and single-item question: associations with sick leave, symptoms, and health—a prospective study of women on long-term sick leave. *Scand J Work Environ Health*. 2010;36(5):404-12.
19. Wentz K, Gyllensten K, Archer T. Recording Recovery Opportunities at Work and Functional Fatigue after Work: Two Instruments Adapted to the Swedish Context. *COJ Nursing & Healthcare*. 2017;1(2):27-36.
20. Sluiter JK, van der Beek AJ, Frings-Dresen MHW. The influence of work characteristics on the need for recovery and experienced health: a study on coach drivers. *Ergonomics*. 1999;42(4):573-83.
21. Murray E. Web-based interventions for behavior change and self-management: potential, pitfalls, and progress. *Med 2.0*. 2012;1(2):e3.
22. Morrison LG, Yardley L, Powell J, Michie S. What design features are used in effective e-health interventions? A review using techniques from critical interpretive synthesis. *Telemedicine and e-Health*. 2012;18(2):137-44.
23. Stratton E, Lampit A, Choi I, Calvo RA, Harvey SB, Glozier N. Effectiveness of eHealth interventions for reducing mental health conditions in employees: A systematic review and meta-analysis. *PLoS One*. 2017;12(12):e0189904.
24. Lehr D, Geraedts A, Asplund RP, Khadjesari Z, Heber E, de Bloom J, et al. Occupational e-mental health: current approaches and promising perspectives for promoting mental health in workers. *Healthy at Work*. 257-281: Springer; 2016.
25. Andersson G, Titov N. Advantages and limitations of Internet-based interventions for common mental disorders. *World Psychiatry*. 2014;13(1):4-11.
26. Free C, Phillips G, Galli L, Watson L, Felix L, Edwards P, et al. The effectiveness of mobile-health technology-based health behaviour change or disease management interventions for health care consumers: a systematic review. *PLoS Med*. 2013;10(1):e1001362.
27. McKay FH, Cheng C, Wright A, Shill J, Stephens H, Uccellini M. Evaluating mobile phone applications for health behaviour change: A systematic review. *J Telemed Telecare*. 2018;24(1):22-30.
28. World Medical Association. Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*. 2013;310(20):2191-4.
29. Eysenbach G. CONSORT-EHEALTH: implementation of a checklist for authors and editors to improve reporting of web-based and mobile randomized controlled trials. *Stud Health Technol Inform*. 2013;192:657-61.
30. Schulz KF, Altman DG, Moher D, Group C. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *PLoS Med*. 2010;7(3):e1000251.
31. Tuomi K, Ilmarinen J, Jahkola A, Katajarinne L, Tulkki A. Work ability index: Institute of Occupational Health Helsinki; 1994.

32. van Schaaik A, Nieuwenhuijsen K, Frings-Dresen MHW, Sluiter JK. Reproducibility of work ability and work functioning instruments. *Occup Med (Lond)*. 2018;68(2):116-9.
33. El Fassi M, Bocquet V, Majery N, Lair ML, Couffignal S, Mairiaux P. Work ability assessment in a worker population: comparison and determinants of Work Ability Index and Work Ability score. *BMC Public Health*. 2013;13:305.
34. Van der Zee K, Sanderman R. RAND-36. Groningen: Northern Centre for Health Care Research, University of Groningen, the Netherlands. 1993;28.
35. de Croon EM, Sluiter JK, Frings-Dresen MH. Need for recovery after work predicts sickness absence: a 2-year prospective cohort study in truck drivers. *J Psychosom Res*. 2003;55(4):331-9.
36. van Veldhoven M, Broersen S. Measurement quality and validity of the "need for recovery scale". *Occup Environ Med*. 2003;60 Suppl 1:i3-9.
37. Jansen M, Sikkel D. Verkorte versie van de VOEG-schaal [Short version of the VOEG-scale]. *Gedrag & samenleving*. 1981;2:78-82.
38. van Veldhoven M, Meijman T. Het meten van psychosociale arbeidsbelasting met een vragenlijst: de vragenlijst beleving en beoordeling van de arbeid (VBBA)[Measuring psychosocial job demands with a questionnaire: the questionnaire on the experience and evaluation of work (QEEW)]. Conditions DfW, editor. Amsterdam: Nederlands Instituut voor Arbeidsomstandigheden; 1994.
39. Meijman T, de Vries-Griever A, De Vries G, R. K. The evaluation of the Groningen sleep quality scale. Groningen: Heymans Bulletin. 1988;88(13-EX).
40. Van Veldhoven M, Prins J, Van der Laken P, Dijkstra L. VBBA 2. 0: Update van de standaard voor vragenlijstonderzoek naar werk, welbevinden en prestaties [VBBA 2. 0: Update of the standard for questionnaire research into work, well-being and performance]. Tilburg: SKB; 2014.
41. Van Veldhoven M, Meijman T, Broersen J, Fortuin R. Handleiding VBBA [Manual VBBA]. Amsterdam: SKB Vragenlijst Services. 2002:2002.
42. Brooks A, Lack L. A brief afternoon nap following nocturnal sleep restriction: which nap duration is most recuperative? *Sleep*. 2006;29(6):831-40.
43. Rosekind MR, Smith RM, Miller DL, Co EL, Gregory KB, Webbon LL, et al. Alertness management: strategic naps in operational settings. *J Sleep Res*. 1995;4(S2):62-6.
44. Lidegaard M, Sogaard K, Krusturup P, Holtermann A, Korshoj M. Effects of 12 months aerobic exercise intervention on work ability, need for recovery, productivity and rating of exertion among cleaners: a worksite RCT. *Int Arch Occup Environ Health*. 2018;91(2):225-35.
45. Van Breukelen GJ. ANCOVA versus change from baseline: more power in randomized studies, more bias in nonrandomized studies [corrected]. *J Clin Epidemiol*. 2006;59(9):920-5.
46. Sluiter JK, Van der Beek AJ, Frings-Dresen MHW. Werkbelasting touringcarchauffeurs (Workload of coach drivers). Amsterdam: Academic Medical Center/Coronel Institute of Occupational Health; 1997.

47. Gould R, Ilmarinen J, Järvisalo J, Koskinen S. Dimensions of work ability. Helsinki: Finnish Centre for Pensions (ETK) The Social Insurance Institution (Kela) National Public Health Institute (KTL) Finnish Institute of Occupational Health (FIOH); 2008.
48. van Schaaik A, Boschman JS, Frings-Dresen MH, Sluiter JK. Appraisal of work ability in relation to job-specific health requirements in ambulance workers. *Int Arch Occup Environ Health*. 2017;90(1):123-31.
49. van Schaaik A, Nieuwenhuijsen K, Frings-Dresen MHW, Sluiter JK. Work ability and work functioning: measuring change in individuals recently returned to work. *Int Arch Occup Environ Health*. 2019;92(3):423-33.
50. De Zwart B, Frings-Dresen M, Van Duivenbooden J. Test–retest reliability of the Work Ability Index questionnaire. *Occupational medicine*. 2002;52(4):177-81.
51. Schouten LS, Bultmann U, Heymans MW, Joling CI, Twisk JW, Roelen CA. Shortened version of the work ability index to identify workers at risk of long-term sickness absence. *Eur J Public Health*. 2016;26(2):301-5.
52. Ware JE, Snow KK, Kosinski M, Gandek B. SF36 Health Survey: Manual and Interpretation Guide. Boston, Massachusetts: The health institute; 1993.
53. Berger VW, Exner DV. Detecting selection bias in randomized clinical trials. *Control Clin Trials*. 1999;20(4):319-27.
54. Staats U, Lohaus D, Christmann A, Woitschek M. Fighting against a shortage of truck drivers in logistics: Measures that employers can take to promote drivers' work ability and health. *Work*. 2017;58(3):383-97.
55. The European parliament and the council of the European Union. Regulation on the harmonisation of certain social legislation relating to road transport 2006. Available from: <https://eur-lex.europa.eu/legal-content/EN/TXT/HTML/?uri=CELEX:32006R0561&from=NL> (Accessed 8th April 2019).
56. Federal Motor Carrier Safety Administration (United States Department of Transportation). Electronic Code of Federal Regulations 49: Transportation PART 395—HOURS OF SERVICE OF DRIVERS 2019. Available from: <https://www.ecfr.gov/cgi-bin/retrieveECFR?gp=1&ty=HTML&h=L&mc=true&=PART&n=pt49.5.395> (Accessed 8th April 2019).
57. Ministry of Transportation and Infrastructure. Commercial Vehicle Drivers Hours of Service Regulations (SOR/2005-313) 2019. 28-11-2009:[Available from: <https://laws-lois.justice.gc.ca/eng/regulations/SOR-2005-313/index.html>] (Accessed 08th April 2019).
58. National Transport Commission. Heavy Vehicle National Law: Heavy vehicle driver fatigue handbook for the Bus and Coach Industry. New South Wales: Roads and Maritime Services; 2008.
59. Swaen G, Van Amelsvoort L, Bultmann U, Kant I. Fatigue as a risk factor for being injured in an occupational accident: results from the Maastricht Cohort Study. *Occupational environmental medicine*. 2003;60(suppl 1):i88-i92.

60. Langford J, Koppel S. Epidemiology of older driver crashes—identifying older driver risk factors and exposure patterns. *Transportation research part F: traffic psychology and behaviour*. 2006;9(5):309-21.
61. Makishita H, Matsunaga K. Differences of drivers' reaction times according to age and mental workload. *Accid Anal Prev*. 2008;40(2):567-75.
62. Janssen N, Kant I, Swaen G, Janssen P, Schröer C. Fatigue as a predictor of sickness absence: results from the Maastricht cohort study on fatigue at work. *Occupational environmental medicine*. 2003;60(suppl 1):i71-i6.
63. Chan ML, Wong Y, Ng R, Koh GC. Medical conditions and driving fitness of older Singaporean taxi drivers. *Occup Med (Lond)*. 2019:211-4.

8

General discussion

OBJECTIVES

In this chapter, the aim and subordinate findings derived from our publications will be addressed. First, the main research findings are summarized followed by a discussion on the interpretation of our findings. Subsequently, methodological considerations are described followed by implications and recommendations for occupational health professionals and researchers, after which the conclusions are summarized.

The aim of this dissertation is to increase knowledge about various aspects of work ability. This aim was divided into three objectives:

1. To gather knowledge about characteristics of work ability measurement in groups of healthy employees and in groups of employees with health complaints and related functional limitations;
2. To gather knowledge about the association between functional limitations and work ability in working populations;
3. To gather knowledge about the course of work ability in work situations characterized by aspects of high workload or in work situations with seasonal changes in workload.

These three objectives are discussed below in light of our findings, current research and practice.

MAIN FINDINGS

To fulfil objective one, we first studied to what extent the Work Ability Score (WAS) is suitable for detecting changes in populations and individuals. In order to use the WAS instrument with individuals, the smallest detectable change is an important parameter that gives information about the smallest change score that the instrument is able to detect. This minimal change was 2 points on a scale of 0-10 (Chapter 2 and 3). The capacity to detect change of the WAS thus proved to be insufficient for use in individuals because the instrument was unable to correctly differentiate between improved, deteriorated or unchanged individuals. The instrument can be used to a certain extent as a monitoring instrument for large (>2) changes or to assist in mapping aspects of work ability in specific populations on a group level.

To fulfil objective two, the association between functional limitations and work ability was studied. In two populations, one with stress complaints the other with upper extremity musculoskeletal (UEMSD) complaints, we found that work ability is influenced by specific factors related to experienced functional limitations. Variance in self-reported work ability in patients with upper extremity complaints can be explained up to 52 % by the degree of difficulties performing daily activities in work, limitations in daily activities as a consequence of health issues, limitations in work and

other daily activities due to physical health issues, and limitations in performing a precision grasp or grip (Chapter 4). Stress had a negative impact on work ability over time in young adults. This negative influence is mediated by several aspects of the work- and private life balance. Stress experienced five years later, job demands affecting private life, feelings of control over private life and feeling well-rested upon waking accounted for 81 % of this relationship (Chapter 5).

To study the third objective, work ability was related to job demands in two populations. First, in ambulance workers, a population characterized by periods of peak workload, self-reported overall work ability was most strongly related to the raised alertness and judgment ability category. Job-specific physical abilities and raised alertness and judgment ability significantly explained variance in physical work ability, while emotional peak load and raised alertness and judgment ability significantly explained variance in mental/emotional work ability (Chapter 6). Second, in coach drivers, a population characterized by seasonal changes in workload, work ability and vitality decreased significantly during peak season, while work-related fatigue increased. The negative impacts of increased workload could not be prevented by a self-management intervention toolbox (Chapter 7).

INTERPRETATION OF FINDINGS

When it comes to the measurement of work ability in individuals, information on measurement properties of the WAS was lacking. In our studies we assessed the reproducibility and responsiveness (sensitivity to change) of the WAS (Figure 1).

Figure 1: Assessing clinimetric properties of the WAS.

Our studies show that the WAS instrument is not suited for monitoring individuals. Despite its questionable validation, the WAS has been used in many monitoring studies without being validated for that aim [1-4]. El Fassi et al. (2013) and Mokarami et al. (2017) recommended the use of the WAS for individual monitoring based merely on its convergent validity with the complete Work Ability Index [5, 6]. However, measurement error in individuals was not taken into account when they issued that conclusion. Contradicting the recommendations of El Fassi and Mokarami, we would advise against using the WAS for individual monitoring based on the results of our studies. Large changes need to occur to conclude that a measurable and relevant change has taken place. Additionally, we found large measurement error and a fair amount of overlap between change scores in the improved, deteriorated and unchanged categories which underlines the lack of ability to adequately detect change in individuals. Using the WAS would result in many misclassifications of individuals as improved, deteriorated or unchanged.

Despite limited usability on the individual level, the WAS is not useless as it can be used in determining important determinants of work ability in groups. We related the WAS to functional limitations and identified pathways in which work ability is affected, which gives the opportunity to direct interventions (Figure 2).

Figure 2: Assessing the relationship between work ability and functional limitations.

Many previous studies have demonstrated the relationship between functional limitations and work ability [7-9]. However, only a few studies go deeper in providing detailed information on the exact work-related functional limitations influencing work ability [10-12]. When the WAS is related to functional limitations as a consequence of health complaints, the aspects that have the strongest impact on work ability can be identified. Boschman et al. (2017) already demonstrated that enquiring about musculoskeletal health and specific experienced limitations is relevant when monitoring work ability to identify possible causes of work ability loss [13]. The relationship between pain and work ability is still not completely clear in each

population, although it seems that more specific information on exact limitations trumps general pain when estimating work ability. Our results show that specific work-related functional limitations can be identified, which in turn can aid improving work or the workplace to avoid decreases in work ability due to functional limitations in UEMSD patients.

One example of more specific enquiry of work ability is that used in our study, in which general pain did not significantly explain variance in work ability in individuals with UEMSD complaints, but specific functional limitations did. Focusing on pain does not help identify aspects that might be changed when aiming to improve work ability. We found that pain was not a predictor of work ability in UEMSD patients when information on limitations was present. This is contradictory to other findings. Lindegård et al. (2014) studied the effect of both musculoskeletal disorders and stress on work ability and found that pain is a risk factor for decreased work ability [14]. Additionally, Oakman et al. (2019) and Kamaleri et al. (2009) found that the number of pain sites was a strong predictor of work ability [15, 16]. In contrast with these findings, de Vries et al. (2013) found that personal and work-related factors are related to work ability in UEMSD patients, rather than pain itself [11]. Our study confirms the results of de Vries and found that although pain was taken into account, this was not a predictor of work ability, whereas functional limitations were. Certain limitations, (e.g. limitations in daily activities or performing a precision grip) are more important for work ability than general pain and can be targeted to improve work ability in UEMSD patients.

Another functional limitation, stress, has been proven to have a negative impact on work ability before [17-21]. Because stress is a prevalent phenomenon during working life, taking stress into account as a potential source of work ability decline is necessary. Although the effect of stress on work ability is widely acknowledged, few studies have been carried out that aim to determine pathways of how stress impacts work ability [22]. Many studies proved a straightforward relationship between stress and work ability [14, 17, 19-21]. Compared with current literature, our study offers a possible approach to inhibit the influence of stress on work ability [19, 21-25]. Based on our study we conclude that the relationship between stress and work ability is complex. For instance, aspects such as job demands affecting private life and feelings of control over private life mediate this stress–work ability relationship in young adults. Boström et al. (2012) previously found an effect of the work-private life balance on work ability [26]. Other authors reported a relation between recovery and feelings of control over private life and work ability [27-32]. However, these aspects were not linked to the stress–work ability relationship before. Our study thus provides novel insight into how stress influences work ability.

We expected that in addition to stress, other factors affecting work ability have their own comprehensive pathways. For example, multiple studies have concluded that work ability is dependent of job demands [33-37]. We were able to identify job demands that strongly relate to work ability, and which can direct actions to avoid declines in the entire workforce over time (Figure 3) [34, 38].

Figure 3: Assessing the relationship between work ability and job demands.

Van den Berg et al. (2009) concluded that most work ability issues arise when either (or both) demands increase or resources diminish [39]. Peak loads (shorts bouts or seasonal increases) put a lot of pressure on high demand populations to meet job demands. To study the relationship between job demands and work ability, we studied high-risk populations because, especially in these populations, the risk of work-related health problems increased [40-42]. Most studies in emergency personnel focus on stressful events and do not take all job demands and health requirements into account [40, 43-47]. In contrast to previous studies on emergency workers we did not find that stressful events were the major influencing factor of work ability. In our study, we included all job and health requirements that are essential to fulfil the job and related these to work ability. We concluded that job demands linked to raised alertness and judgment ability showed the strongest relationship to work ability in ambulance workers. This state of alertness has since been determined as a risk factor for reduced work ability in nurses and was previously related to reduced work ability in young adults [31, 48, 49].

The factors relating strongest to work ability in our studies are important aspects of work ability in their respective populations and might be targeted in prevention and intervention. Aiming for workers to work in good health until their retirement while being able to continually meet job demands is a complex task where prevention and intervention form a major bottleneck (Figure 4).

Figure 4: Work ability in relation to prevention and intervention.

Our intervention in coach drivers during peak season consisted of an interactive toolbox comprising tips and assignments in multiple areas (work-recovery-rest balance, food and drink intake, and physical activity). Despite following the recommendations for developing interventions, the intervention itself did not result in a difference in work ability between intervention and control group [50, 51]. Although well-received, the use of the intervention was too low to expect results. Insufficient adherence to the intervention is an often-reported issue in occupational health interventions [52-56]. Longer follow-up, supervisor involvement, (more frequent) training sessions, intervention during working hours, including sessions to increase awareness of the urgency of the situation or involving third parties, have been suggested to improve adherence [52-55, 57-59]. A different kind of care opposed to self-management interventions can be considered to enhance the effect of care. Blended care is an upcoming promising alternative where e-health is combined with supervision [60]. This type of care might be especially suited for coach drivers that travel frequently.

METHODOLOGICAL CONSIDERATIONS

The measurement instrument

In the studies in this dissertation, we chose the WAS as measurement instrument based on its convergent validity with the complete WAI. Because the WAI is complex and long, the WAS was suggested as suitable alternative for the WAI [5, 6, 61].

In all chapters of this dissertation, the WAS is used without the proposed four categories (0-5, poor; 6-7, moderate; 8-9, good; 10, excellent) which are distribution based [17]. An advantage of this continuous method, is that the continuous variable holds more information, particularly in subgroups where distribution can vary from the population wherein the proposed score-groups were determined [62]. Especially in determining change and cut-off values, a continuous variable can increase the precision of the instrument [63]. Improvement in measurement precision will result in a more solid basis for model development and decision making. In our study we found that a change of 2 points is a measurable and relevant change. When using the category-based method, a difference of 1 point could mean a change in category, and meanwhile a change of 2 points could keep someone in the same category. Classifying change would therefore be inconsistent over categories. Determining the reliability of the WAI or WAS as an ordinal variable is less appropriate because it highly depends on where the individual stands in that categorization [64]. Therefore, using the WAS as a continuous variable is favorable.

To be able to use WAS in research and practice, the clinimetric properties of the WAS were studied in two different populations. The first study was aimed at determining the measurement error in this instrument, in a population that had not changed. This measurement characteristic needs to be studied in a stable population [65]. In our study, the timeframe between the two measurements was 7 days. This is rather short when compared with other studies assessing clinimetric properties that used a larger time interval of 7-15 days, 4 weeks or even 12 weeks between the two measurement points [64, 66, 67]. However, these studies did not always ensure that the work ability of the group was stable. Yang et al. (2013) argue that even a 12-week period is too short to make work ability change. In contrast, we argue that in a period of 7 days, a change in work ability is possible given the fact that 13 % of participants indicated on the selection question that their work ability had changed [68]. Work ability assesses the ability to meet job demands at the current time. Therefore, keeping the time between measurements short and adding a selection question assured us that we measured work ability in stable individuals as is intended in reproducibility measurements [69]. Therefore, we argue that a period of 7 days is justified in our studies.

An often heard critique on the WAS and the complete WAI is that both instruments can only be used for signaling problems and not for establishing the cause of the problem [61]. Relating the WAS to functional limitations and job demands can counter the critique to some extent by providing information on aspects that cause decreases in work ability in specific populations. In this way, we showed that the WAS can be used in populations to identify problem areas.

The populations

Our research focused mainly on populations with specific functional limitations or job demands. The results are applicable to populations with the same functional limitations and job demands and therefore the generalizability of these results to the general working population is limited. Work ability often varies widely between individuals. Therefore, the generalizability of work ability research often remains limited to its application on the group level, as the findings relate to group averages while work ability itself remains personal [17, 39, 42, 70]. Although the generalizability of exact numbers remains limited, pathways in which work ability is affected are relevant in groups and individuals. For in-depth analysis of work ability, a personal approach is desired.

To draw conclusions in research, a sufficient sample size is needed to reach the desired statistical power and significance. Our mediation analyses took place in a longitudinal cohort of young adults. An invitation went out to 20,000 individuals, equally divided between men and women. Because of the origin of this cohort, the number of participants was quite large, which supplied us with sufficient statistical power to substantiate our results, even in subgroups [71].

In contrast, the cohort of coach drivers investigated was small due to recruitment difficulties. Although this study was performed at the request of the coach driver sector organization, this organization was not in the possession of contact details of coach drivers. Therefore, coach companies had to be contacted and inclusion was dependent on their managers to forward the opportunity to participate in our study. Therefore, we visited places where coach drivers get together to invite them to participate. Increasing sample size by rewarding participants with incentives was used to stimulate participation [72]. Although the number of participants was not as high as we would have wanted, we do not expect that a larger study population or statistical alterations would have yielded different results. Our effect size was too small to lead to any clinically relevant changes in work ability. To find a statistical difference, we would need an enormous study population based on the found effect sizes and the desired power level [73].

The analyses

Many different types of statistical analyses were applied in the chapters of this dissertation. One of the frequently used methods was multiple linear regression, which was used to study to what extent the variance in work ability could be explained by functional limitations or job demands.

Multiple linear regression is an appropriate method for analyzing the relationship between job demands or functional limitations and work ability [74]. Using this

method, significant contributors to explained variance in work ability were determined. An alternative to linear regression would be structural equation modelling [75, 76]. Structural equation modelling is perfectly suited to map the complete picture of work ability as a causal system (complex and multifaceted) and all its determinants. This method is able to correct for measurement errors and different types of dependent variables that affect other dependent variables [77]. However, structural equation modelling is most often used for latent variables and multiple dependent variables, which adds error variance to the equation. An advantage of linear regression is that it adjusts for other variables in the model, avoiding collinearity, meaning that linear regression is suited when there is a single dependent variable and a set of predictors as independent variables [78]. In our study of UEMSD patients, we measured exact variables for which multiple linear regression is a better match. In our study on ambulance workers, we conceptualized job demands in overarching categories explained by several variables. In hindsight, structural equation modelling would have been more appropriate.

For the analysis of mediators in the relationship between stress and work ability, we considered structural equation modelling. However, because of the nature of our variables, we chose to model mediation with a mediation analysis as proposed by Hayes [79], which is well suited for research into pathways related to work ability [80]. In mediation analyses, causal pathways are of great importance to prevent reversed causality [81]. It is suggested that mediation in cross-sectional data leaves the possibility for reversed causation [82]. Therefore, we substantiated our theoretical framework and introduced a time-component to support our models to guarantee causation in our mediation analysis.

The intervention

In our intervention study, we aimed at maintaining or even improving work ability in coach drivers during the peak season. To achieve this goal, we tailored the intervention based on pre-conducted interviews. Additionally, the intervention had a theoretical foundation based on scientific literature, acts in multiple areas and the separate items of the intervention were individually evaluated [50, 51, 83]. Another strength of our toolbox was that the real-life effect of the intervention could be measured because no extra proceedings were added during peak season. The implementation of this intervention was as it would be in practice and not a protocolled research setting where the effect might be artificially boosted due to supervision by researchers. Measuring the real-life effects was explicitly the desire of the coach driver sector organization as they did not want to engage in activities or supervision. An associated drawback was that there was no control over the implementation and adherence to the intervention. This realistic approach and

evaluation of the components in the intervention showed that the intervention in this way was not feasible during peak season. No significant differences were found between the intervention and control group in several outcomes.

Because of our process analysis, we know that the content of the toolbox was well-received, which indicates that a different type of implementation might have produced the desired results. Possible options for alternative implementation is supported intervention, which is known to improve the effectiveness of interventions [84]. Other options that might improve effectiveness are blended care as discussed above, which also might improve participation and coherence, or gamification, especially in unsupervised interventions [85, 86].

IMPLICATIONS AND RECOMMENDATIONS

In the previous chapters we aimed to map the relationship between work ability and functional limitations and work ability and job demands in the bigger picture. We showed that the WAS can be useful in work ability research. Although we selected specific populations, the results of our study can help in determining relationships and pathways between these aspects. Our studies provide starting points for further research and the development of interventions.

Implications for occupational health professionals

Occupational health professionals can use the findings derived from our studies to improve work ability in occupational or patient groups, or prevent work ability from deteriorating by advising how to adjust to certain job demands or by introducing tools to ease the job and decrease the workload e.g. a tool to help perform a precision grip.

The present findings suggest that the WAS is not suited for individual monitoring. However, an insufficient WAS can be reason to further study an individual's work ability to check if it has decreased or that this insufficient score is due to measurement error. Further investigation of insufficient WAS scores is required because the WAS instrument itself does not identify in what area the problem occurs [61]. For use in individuals, subsequent testing is required to identify the cause of the problems. An instrument that might be used for additional monitoring is the novel Work Ability Personal Radar (WA-PR) [87]. This WA-PR is based on the dimensions of the work ability house and has the advantage that it detects which aspects of work ability are affected, so that interventions can be aimed at these specific domains. This instrument divided the work ability house into six dimensions, and went further in-depth within each domain, to be able to identify which domain is the problem area of insufficient work ability. The advantage of this instrument is that domains can be identified, which can be addressed for immediate action [37]. Application of this

instrument is promising; however, this instrument is not yet validated for the ability to detect change in individuals.

A downside of the WAS is that only large changes in work ability can be detected on an individual level. We established certain patterns in work ability over a short time-period in workers returning to work after sick leave. In these individuals with rapidly changing work ability levels, the WAS can only be used for a first global assessment of large changes. Changes of 2 or more points on the WAS scale can be seen as a measurable, relevant change in an individual. If this change is negative, occupational physicians could be asked to intervene.

The WAS can be used to identify aspects of work ability in groups of workers. Chapter 4, 5, 6 and 7, showed that several functional limitations and job demands were predictive of work ability. This shows that the WAS can aid occupational health professionals in linking functional limitations or job demands to work ability when both are assessed. Both functional limitations or job demands, and work ability can be assessed in preventive medical examination (PME) in workers. In PME the health of a workers is assessed in relation to job demands. Because the relationship between job demands and work ability is strong, the ability to meet job demands is seen as prominent contributor to variance in work ability. This indicates that PME (with respect to job demands) in combination with the WAS can contribute towards the identification of problem areas in specific populations. Hence, occupational health professionals might be able to gather more information on aspects that cause decreases in work ability which can be targeted with interventions, e.g. tools that help meet job demands, education, multidisciplinary workplace programs or workplace adaptation [88-90].

Identifying problem areas that are most strongly related to work ability is especially important in high-demand occupations as ambulance workers as people's lives depend on it. Therefore, occupational physicians need to know exactly how job demands relate to work ability, and when not being able to meet such demands results in insufficient ability to work in that occupation at that particular time. Solely utilizing the WAS as instrument proved insufficient as fewer than 1 % percent of ambulance workers appraised their own work ability as insufficient, while 68 % of ambulance workers scored insufficient on one or more job requirements. Following up on the identified aspects that affect work ability most strongly is desired in the intervention and treatment of insufficient work ability [91, 92].

It is stated before that work ability declines with increasing age [93, 94]. However, it is expected that job demands affect work ability at every single age; however, aging workers experience fewer resources (e.g. health, lower recovery) to meet these demands, whereas younger workers have the resources to cope with these demands

[95]. The majority of young workers with the same job demands will eventually experience the same problems when they grow older when no interventions take place [96, 97]. Helping young workers cope with job demands and maintain the balance between demands and resources can pay off later in life. It has been suggested that starting work ability promotion early in the career is the best way to improve sustainable work ability [17].

When focusing on younger workers, it is important to keep in mind that workers of different ages run into different problems over the course of their working life [17]. Job demands and resources change over time, which means that prevention needs to be focused on various matters throughout one's working life. Young workers are in a different phase of life and have a different lifestyle than older workers. Therefore, interventions need to be tailored to the needs of the worker at that time in their work lifecycle. For example, in a young population with stress complaints, we would recommend that attention goes to the work/private life balance to reduce the effect of stress on work ability. A direct measure that can be taken by companies is to counteract the negative influence of stress on work ability by minimizing the spillover of job demands into private life. These stressors need to be discussed between employer and employee to avoid decreases in work ability early on in one's working life.

Overall, mapping the strength of the relationships between work ability and functional limitations and job demands allows for better understanding of how (job demands may be adjusted and) a loss in work ability can be prevented. Possibly in the future, an employer can design age-adequate jobs, or even preferably, work ability-adequate jobs when taking the aspects that affect work ability into account [98]. Knowing the importance of job demands and functional abilities for work ability can be valuable information for occupational health professionals. However, occupational health professionals need to keep in mind that although certain functional limitations and job demands affect work ability in populations, work ability is personal, and the personal character of work ability needs to be taken into account.

Implications for research

The WAS is eminently suited for research into aspects of work ability. Because it is a single-item question, it can easily be added to questionnaires and other work ability assessment tools. The WAS can be related to contents of e.g. physical examinations or the ability to meet certain job demands to determine what aspects are most important for work ability in various populations.

The results of our clinimetric studies can help researchers put measured work ability scores in the perspective of change. The goals to aim for in improving work ability with

interventions is to achieve a measurable relevant change of minimally 2 points on the WAS. This relates to individual work ability scores. Group scores can significantly change; however, it can still be the case that no single individual has improved relevantly on his/her own work ability. Therefore, not only population scores matter, researchers should also take individual scores into account when studying whether the work ability of a population has changed.

Our studies show that population characteristics exist that influence work ability in a certain way. These characteristics can be job demands, but also functional limitations that certain groups have in common. We suggest that various aspects that might affect work ability are mapped to assemble a more complete overview of how work ability is affected in various populations. When more of these aspects are mapped, an all-encompassing model of work ability can be modelled including latent variables and pathways between several dependent variables. This model can be developed and tested with structural equation modelling as mentioned above [76].

Interventions are historically aimed at two target groups, namely unhealthy individuals that need health improvement, and healthy individuals at risk that need preventive measures [99]. However, because work ability is known to decline with age, essentially all workers are at risk of a decreased work ability over time. Consequently, we suggest that interventions should be aimed at various groups of workers during the entire working life span. First, it is necessary to identify the needs of employees and how work ability is affected at each stage of working life [17, 31]. This can be achieved by determining which characteristics are most strongly linked to work ability during certain life phases. We concluded that work ability in young adults is related to aspects of the work ability house model; more specifically, the balance between work and private life. Mapping the changing needs of workers during working life can improve the fit and effectiveness of interventions to the worker.

In many studies targeting work ability, interventions are at most slightly effective [100, 101]. Because it is challenging to develop effective interventions, we recommend that first, aspects of work ability are understood in greater detail to ensure that targeted recommendations on possible interventions can be made. Second, other types of interventions than unsupported self-management should be explored to find a format capable of achieving sufficient adherence. Third, it is suggested that interventions need to be carried out longer to achieve an effect [102]. Possibly changing behavior and creating awareness early on in working life can result in lasting effects on the preservation of work ability.

CONCLUSIONS

In this dissertation we aimed to fulfil three objectives by answering three subsequent research questions. The questions were aimed at the clinimetric properties of the Work Ability Score (WAS), and it's relation to functional limitations and job demands. The following questions were composed and subsequently answered:

- I. What are the reproducibility and responsiveness of the work ability score of workers who return to work after sick leave?
- II. How are functional limitations related to work ability in working populations with specific health complaints?
- III. What is the course of work ability in occupations characterized by specific health requirements and peak loads, and to what extent can this be influenced?

Reproducibility of the WAS was found to be moderately reliable. Additionally it was shown that the responsiveness of the instrument was insufficient. Change scores of at least 2 points on the 0–10 scale have to occur to demonstrate change. In addition, the instrument was inadequate in correctly classifying individuals as improved, deteriorated or unchanged. Therefore, we concluded that the WAS is not suitable for individual monitoring of work ability.

The WAS can be used to relate functional limitations and job demands to work ability in groups. This enables identifying aspects that are most important in work ability. Exploring these aspects of work ability in populations with functional limitations and high job demands contributes to the knowledge of how work ability is constructed in these populations. Consequently, these specific functional limitations or job demands most strongly related to work ability can be identified and then targeted with preventive measures. In UEMSD patients, specific functional limitations were established which can help occupational health professionals to intervene and possibly improve work ability. In young adults, the balance between work and private life was prominent in the stress-work ability relationship. Discussing this issue can be a strategy to avoid stress that negatively affects work ability.

Job demands with the strongest relationship to work ability were identified in jobs characterized by peak loads as ambulance workers. These demands are the most crucial to be met in order to avoid losses in work ability. Occupational health professionals should use this information to direct attention to these demands when they are insufficiently met.

A declining trend was seen in the course of work ability and the vitality of coach drivers, when job demands increased during peak season. However, distributing a self-management intervention without any further support was unsuccessful in

influencing this course and maintaining work ability in coach drivers during this period. Suggestions were proposed to improve interventions with the aim of maintaining or improving work ability.

In work ability research and the implementation of preventive measures, attention needs to be placed on sustainable work ability. Therefore, knowledge of work and health aspects and their relationship to work ability needs to be mapped to form a more complete picture, which occupational health professionals can use in their assessment and improvement of work ability. However, although groups can show similar patterns in work ability and associated determinants, work ability is subject to individual variation.

REFERENCES

1. Ahlstrom L, Grimby-Ekman A, Hagberg M, Dellve L. The work ability index and single-item question: associations with sick leave, symptoms, and health--a prospective study of women on long-term sick leave. *Scand J Work Environ Health*. 2010;36(5):404-12.
2. Jaaskelainen A, Kausto J, Seitsamo J, Ojajarvi A, Nygard CH, Arjas E, et al. Work ability index and perceived work ability as predictors of disability pension: a prospective study among Finnish municipal employees. *Scand J Work Environ Health*. 2016;42(6):490-9.
3. Mache S, Danzer G, Klapp BF, Groneberg DA. Surgeons' work ability and performance in surgical care: relations between organisational predictors, work engagement and work ability. *Langenbecks Arch Surg*. 2013;398(2):317-25.
4. von Bonsdorff ME, Kokko K, Seitsamo J, von Bonsdorff MB, Nygard CH, Ilmarinen J, et al. Work strain in midlife and 28-year work ability trajectories. *Scand J Work Environ Health*. 2011;37(6):455-63.
5. El Fassi M, Bocquet V, Majery N, Lair ML, Couffignal S, Mairiaux P. Work ability assessment in a worker population: comparison and determinants of Work Ability Index and Work Ability score. *BMC Public Health*. 2013;13:305.
6. Mokarami H, Mortazavi SB, Asgari A, Choobineh A. Work Ability Score (WAS) as a suitable instrument to assess work ability among Iranian workers. *Health Scope*. 2017;6(1).
7. Lindstrom I, Pallasaho P, Luukkonen R, Suojalehto H, Karjalainen J, Lauerma A, et al. Reduced work ability in middle-aged men with asthma from youth--a 20-year follow-up. *Respir Med*. 2011;105(6):950-5.
8. Hedenrød T, Love J, Staland-Nyman C, Hensing G. Frequent headache and work ability: a population-based study in Sweden. *J Occup Environ Med*. 2014;56(5):472-6.
9. Krstovic-Spremo V, Racic M, Joksimovic BN, Joksimovic VR. The effects of diabetes mellitus and hypertension on work productivity. *Acta Med Acad*. 2014;43(2):122-33.
10. Gouttebarger V, Wind H, Kuijter PP, Sluiter JK, Frings-Dresen MH. How to assess physical work-ability with Functional Capacity Evaluation methods in a more specific and efficient way? *Work*. 2010;37(1):111-5.
11. de Vries HJ, Reneman MF, Groothoff JW, Geertzen JH, Brouwer S. Self-reported work ability and work performance in workers with chronic nonspecific musculoskeletal pain. *J Occup Rehabil*. 2013;23(1):1-10.
12. Alavinia SM, de Boer AG, van Duivenbooden JC, Frings-Dresen MH, Burdorf A. Determinants of work ability and its predictive value for disability. *Occup Med (Lond)*. 2009;59(1):32-7.
13. Boschman JS, Noor A, Lundstrom R, Nilsson T, Sluiter JK, Hagberg M. Relationships between work-related factors and musculoskeletal health with current and future work ability among male workers. *Int Arch Occup Environ Health*. 2017;90(6):517-26.
14. Lindegård A, Larsman P, Hadzibajramovic E, Ahlberg G, Jr. The influence of perceived stress and musculoskeletal pain on work performance and work ability in Swedish health care workers. *Int Arch Occup Environ Health*. 2014;87(4):373-9.

15. Oakman J, Neupane S, Prakash KC, Nygard CH. What Are the Key Workplace Influences on Pathways of Work Ability? A Six-Year Follow Up. *Int J Environ Res Public Health*. 2019;16(13).
16. Kamaleri Y, Natvig B, Ihlebaek CM, Bruusgaard D. Does the number of musculoskeletal pain sites predict work disability? A 14-year prospective study. *Eur J Pain*. 2009;13(4):426-30.
17. Gould R, Ilmarinen J, Järvisalo J, Koskinen S. Dimensions of work ability. Helsinki: Finnish Centre for Pensions (ETK) The Social Insurance Institution (Kela) National Public Health Institute (KTL) Finnish Institute of Occupational Health (FIOH); 2008.
18. Goedhard RG, Goedhard WJA, editors. Work ability and perceived work stress. International Congress Series; 2005: Elsevier.
19. Gharibi V, Mokarami H, Taban A, Yazdani Aval M, Samimi K, Salesi M. Effects of Work-Related Stress on Work Ability Index among Iranian Workers. *Saf Health Work*. 2016;7(1):43-8.
20. Nevanpera N, Seitsamo J, Ala-Mursula L, Remes J, Hopsu L, Auvinen J, et al. Perceived Work Ability in the Light of Long-Term and Stress-Related Unhealthy Behaviors-a Prospective Cohort Study. *Int J Behav Med*. 2016;23(2):179-89.
21. Habibi E, Dehghan H, Safari S, Mahaki B, Hassanzadeh A. Effects of work-related stress on work ability index among refinery workers. *J Educ Health Promot*. 2014;3:18.
22. Yang T, Liu T, Lei R, Deng J, Xu G. Effect of Stress on the Work Ability of Aging American Workers: Mediating Effects of Health. *Int J Environ Res Public Health*. 2019;16(13).
23. Bernburg M, Vitzthum K, Groneberg DA, Mache S. Physicians' occupational stress, depressive symptoms and work ability in relation to their working environment: a cross-sectional study of differences among medical residents with various specialties working in German hospitals. *BMJ Open*. 2016;6(6):e011369.
24. Marklund S, Mienna CS, Wahlstrom J, Englund E, Wiesinger B. Work ability and productivity among dentists: associations with musculoskeletal pain, stress, and sleep. *Int Arch Occup Environ Health*. 2020;93(2):271-8.
25. Yong M, Nasterlack M, Pluto RP, Lang S, Oberlinner C. Occupational stress perception and its potential impact on work ability. *Work*. 2013;46(3):347-54.
26. Boström M, Sluiter JK, Hagberg M. Changes in work situation and work ability in young female and male workers. A prospective cohort study. *BMC Public Health*. 2012;12:694.
27. Camerino D, Conway PM, Sartori S, Campanini P, Estry-Behar M, van der Heijden BI, et al. Factors affecting work ability in day and shift-working nurses. *Chronobiol Int*. 2008;25(2):425-42.
28. Lian Y, Xiao J, Liu Y, Ning L, Guan S, Ge H, et al. Associations between insomnia, sleep duration and poor work ability. *J Psychosom Res*. 2015;78(1):45-51.
29. Yong M, Fischer D, Germann C, Lang S, Vetter C, Oberlinner C. Are chronotype, social jetlag and sleep duration associated with health measured by Work Ability Index? *Chronobiol Int*. 2016;33(6):721-9.

30. Lallukka T, Sivertsen B, Kronholm E, Bin YS, Overland S, Glozier N. Association of sleep duration and sleep quality with the physical, social, and emotional functioning among Australian adults. *Sleep Health*. 2018;4(2):194-200.
31. Boström M, Holmgren K, Sluiter JK, Hagberg M, Grimby-Ekman A. Experiences of work ability in young workers: an exploratory interview study. *Int Arch Occup Environ Health*. 2016;89(4):629-40.
32. Eriksson J, Dellve L, Eklof M, Hagberg M. Early inequalities in excellent health and performance among young adult women and men in Sweden. *Gend Med*. 2007;4(2):170-82.
33. McGonagle AK, Fisher GG, Barnes-Farrell JL, Grosch JW. Individual and work factors related to perceived work ability and labor force outcomes. *J Appl Psychol*. 2015;100(2):376-98.
34. Noone JH, Bohle P, Mackey M. Matching Work Capacity and Job Demands: Toward an Enhanced Measure of Work Ability. *J Occup Environ Med*. 2015;57(12):1360-4.
35. Ruitenburg MM, Frings-Dresen MH, Sluiter JK. Physical job demands and related health complaints among surgeons. *Int Arch Occup Environ Health*. 2013;86(3):271-9.
36. van der Doef MP, Schelvis RMC. Relations Between Psychosocial Job Characteristics and Work Ability in Employees with Chronic Headaches. *J Occup Rehabil*. 2019;29(1):119-27.
37. Ilmarinen J. From Work Ability Research to Implementation. *Int J Environ Res Public Health*. 2019;16(16).
38. Nabe-Nielsen K, Thielen K, Nygaard E, Thorsen SV, Diderichsen F. Demand-specific work ability, poor health and working conditions in middle-aged full-time employees. *Appl Ergon*. 2014;45(4):1174-80.
39. van den Berg TI, Elders LA, de Zwart BC, Burdorf A. The effects of work-related and individual factors on the Work Ability Index: a systematic review. *Occup Environ Med*. 2009;66(4):211-20.
40. Sluiter JK. High-demand jobs: age-related diversity in work ability? *Appl Ergon*. 2006;37(4):429-40.
41. Tuomi K, Huuhtanen P, Nykyri E, Ilmarinen J. Promotion of work ability, the quality of work and retirement. *Occup Med (Lond)*. 2001;51(5):318-24.
42. Tuomi K, Eskelinen L, Toikkanen J, Jarvinen E, Ilmarinen J, Klockars M. Work load and individual factors affecting work ability among aging municipal employees. *Scand J Work Environ Health*. 1991;17 Suppl 1:128-34.
43. Young KM, Cooper CL. Occupational stress in the ambulance service: a diagnostic study. *Health Manpow Manage*. 1997;23(4-5):140-7.
44. Oldenburg M, Wilken D, Wegner R, Poschadel B, Baur X. Job-related stress and work ability of dispatchers in a metropolitan fire department. *Journal of Occupational medicine and Toxicology*. 2014;9(1):31.
45. Klasan A, Madzarac G, Milosevic M, Mustajbegovic J, Keleuva S. Predictors of lower work ability among emergency medicine employees: the Croatian experience. *Emerg Med J*. 2013;30(4):275-9.

46. Stanley IH, Boffa JW, Smith LJ, Tran JK, Schmidt NB, Joiner TE, et al. Occupational stress and suicidality among firefighters: Examining the buffering role of distress tolerance. *Psychiatry Res.* 2018;266:90-6.
47. Kim MJ, Jeong Y, Choi YS, Seo AR, Ha Y, Seo M, et al. The Association of the Exposure to Work-Related Traumatic Events and Work Limitations among Firefighters: A Cross-Sectional Study. *Int J Environ Res Public Health.* 2019;16(5).
48. da Silva TPD, Araujo WN, Stival MM, Toledo AM, Burke TN, Carregaro RL. Musculoskeletal discomfort, work ability and fatigue in nursing professionals working in a hospital environment. *Rev Esc Enferm USP.* 2018;52:e03332.
49. Kaliyaperumal D, Elango Y, Alagesan M, Santhanakrishnan I. Effects of Sleep Deprivation on the Cognitive Performance of Nurses Working in Shift. *J Clin Diagn Res.* 2017;11(8):CC01-CC3.
50. Webb TL, Joseph J, Yardley L, Michie S. Using the internet to promote health behavior change: a systematic review and meta-analysis of the impact of theoretical basis, use of behavior change techniques, and mode of delivery on efficacy. *J Med Internet Res.* 2010;12(1):e4.
51. Nielsen KT, la Cour K, Christensen JR, Pilegaard MS, von Bulow C, Brandt AA, et al. Lessons learned about occupation-focused and occupation-based interventions: A synthesis using group concept mapping methodology. *Scand J Occup Ther.* 2019:1-12.
52. Oude Hengel KM, Blatter BM, Joling CI, van der Beek AJ, Bongers PM. Effectiveness of an intervention at construction worksites on work engagement, social support, physical workload, and need for recovery: results from a cluster randomized controlled trial. *BMC Public Health.* 2012;12:1008.
53. van Dongen JM, van Berkel J, Boot CR, Bosmans JE, Proper KI, Bongers PM, et al. Long-Term Cost-Effectiveness and Return-on-Investment of a Mindfulness-Based Worksite Intervention: Results of a Randomized Controlled Trial. *J Occup Environ Med.* 2016;58(6):550-60.
54. Jorgensen MB, Faber A, Hansen JV, Holtermann A, Sogaard K. Effects on musculoskeletal pain, work ability and sickness absence in a 1-year randomised controlled trial among cleaners. *BMC Public Health.* 2011;11:840.
55. Notenbomer A, Roelen C, Groothoff J, van Rhenen W, Bultmann U. Effect of an eHealth Intervention to Reduce Sickness Absence Frequency Among Employees With Frequent Sickness Absence: Randomized Controlled Trial. *J Med Internet Res.* 2018;20(10):e10821.
56. Maes L, Van Cauwenberghe E, Van Lippevelde W, Spittaels H, De Pauw E, Oppert JM, et al. Effectiveness of workplace interventions in Europe promoting healthy eating: a systematic review. *Eur J Public Health.* 2012;22(5):677-83.
57. Wang Y, Xue H, Huang Y, Huang L, Zhang D. A Systematic Review of Application and Effectiveness of mHealth Interventions for Obesity and Diabetes Treatment and Self-Management. *Adv Nutr.* 2017;8(3):449-62.
58. Miles C, Arden-Close E, Thomas M, Bruton A, Yardley L, Hankins M, et al. Barriers and facilitators of effective self-management in asthma: systematic review and thematic

- synthesis of patient and healthcare professional views. *NPJ Prim Care Respir Med*. 2017;27(1):57.
59. Chinn D. Critical health literacy: a review and critical analysis. *Soc Sci Med*. 2011;73(1):60-7.
 60. Wentzel J, van der Vaart R, Bohlmeijer ET, van Gemert-Pijnen JE. Mixing Online and Face-to-Face Therapy: How to Benefit From Blended Care in Mental Health Care. *JMIR Ment Health*. 2016;3(1):e9.
 61. Ebener M, Hasselhorn HM. Validation of Short Measures of Work Ability for Research and Employee Surveys. *Int J Environ Res Public Health*. 2019;16(18).
 62. Cadiz DM, Brady G, Rineer JR, Truxillo DM. A review and synthesis of the work ability literature. *Work, Aging and Retirement*. 2019;5(1):114-38.
 63. Wagner RF, Beiden SV, Metz CE. Continuous versus categorical data for ROC analysis: some quantitative considerations. *Acad Radiol*. 2001;8(4):328-34.
 64. De Zwart B, Frings-Dresen M, Van Duivenbooden J. Test-retest reliability of the Work Ability Index questionnaire. *Occupational medicine*. 2002;52(4):177-81.
 65. de Vet HC, Terwee CB, Knol DL, Bouter LM. When to use agreement versus reliability measures. *J Clin Epidemiol*. 2006;59(10):1033-9.
 66. Silva Junior SH, Vasconcelos AG, Griep RH, Rotenberg L. [Test-retest reliability of the Work Ability Index (WAI) in nursing workers]. *Rev Bras Epidemiol*. 2013;16(1):202-9.
 67. Yang DJ, Kang D, Kim YK, Kim YH, Yang YA, Cha SM, et al. Reliability of self-administered Work Ability Index questionnaire among Korean workers. *Ergonomics*. 2013;56(11):1652-7.
 68. van Schaaik A, Nieuwenhuijsen K, Frings-Dresen MHW, Sluiter JK. Reproducibility of work ability and work functioning instruments. *Occup Med (Lond)*. 2018;68(2):116-9.
 69. Terwee CB, Bot SD, de Boer MR, van der Windt DA, Knol DL, Dekker J, et al. Quality criteria were proposed for measurement properties of health status questionnaires. *J Clin Epidemiol*. 2007;60(1):34-42.
 70. Fischer FM, Martinez MC. Individual features, working conditions and work injuries are associated with work ability among nursing professionals. *Work*. 2013;45(4):509-17.
 71. Pan H, Liu S, Miao D, Yuan Y. Sample size determination for mediation analysis of longitudinal data. *BMC Med Res Methodol*. 2018;18(1):32.
 72. Smith MG, Witte M, Rocha S, Basner M. Effectiveness of incentives and follow-up on increasing survey response rates and participation in field studies. *BMC Med Res Methodol*. 2019;19(1):230.
 73. Hemming K, Girling AJ, Sitch AJ, Marsh J, Lilford RJ. Sample size calculations for cluster randomised controlled trials with a fixed number of clusters. *BMC Med Res Methodol*. 2011;11:102.
 74. Marill KA. Advanced statistics: linear regression, part II: multiple linear regression. *Academic emergency medicine*. 2004;11(1):94-102.

75. Gefen D, Straub D, Boudreau M-C. Structural equation modeling and regression: Guidelines for research practice. *Communications of the association for information systems*. 2000;4(1):7.
76. Ilmarinen J, Tuomi K, Seitsamo J, editors. *New dimensions of work ability*. International congress series; 2005: Elsevier.
77. Kline RB. *Principles and practice of structural equation modeling*. New York City: Guilford publications; 2015.
78. Nimon KF, Oswald FL. Understanding the results of multiple linear regression: Beyond standardized regression coefficients. *Organizational Research Methods*. 2013;16(4):650-74.
79. Hayes AF. *Introduction to mediation, moderation, and conditional process analysis: A regression-based approach*. New York, USA: Guilford Publications; 2017.
80. Hayes AF. Beyond Baron and Kenny: Statistical mediation analysis in the new millennium. *Communication monographs*. 2009;76(4):408-20.
81. Oude Groeniger J, Burdorf A. Advancing mediation analysis in occupational health research. *Scand J Work Environ Health*. 2020;46(2):113-6.
82. VanderWeele T. *Explanation in causal inference: methods for mediation and interaction*: Oxford University Press; 2015.
83. Bonell C, Fletcher A, Morton M, Lorenc T, Moore L. Realist randomised controlled trials: a new approach to evaluating complex public health interventions. *Soc Sci Med*. 2012;75(12):2299-306.
84. Baumeister H, Reichler L, Munzinger M, Lin J. The impact of guidance on Internet-based mental health interventions—A systematic review. *internet Interventions*. 2014;1(4):205-15.
85. Looyestyn J, Kernot J, Boshoff K, Ryan J, Edney S, Maher C. Does gamification increase engagement with online programs? A systematic review. *PLoS One*. 2017;12(3):e0173403.
86. Birk MV, Atkins C, Bowey JT, Mandryk RL, editors. *Fostering intrinsic motivation through avatar identification in digital games*. *Proceedings of the 2016 CHI conference on human factors in computing systems*; 2016.
87. Ilmarinen V, Ilmarinen J, Huuhtanen P, Louhevaara V, Nasman O. Examining the factorial structure, measurement invariance and convergent and discriminant validity of a novel self-report measure of work ability: work ability--personal radar. *Ergonomics*. 2015;58(8):1445-60.
88. Kettunen O, Vuorimaa T, Vasankari T. 12-mo intervention of physical exercise improved work ability, especially in subjects with low baseline work ability. *Int J Environ Res Public Health*. 2014;11(4):3859-69.
89. de Boer AG, Burdorf A, van Duivenbooden C, Frings-Dresen MH. The effect of individual counselling and education on work ability and disability pension: a prospective intervention study in the construction industry. *Occup Environ Med*. 2007;64(12):792-7.

90. Cloostermans L, Bekkers MB, Uiters E, Proper KI. The effectiveness of interventions for ageing workers on (early) retirement, work ability and productivity: a systematic review. *Int Arch Occup Environ Health*. 2015;88(5):521-32.
91. Lele DV. Occupational Health Surveillance. *Indian J Occup Environ Med*. 2018;22(3):117-20.
92. Porru S, Placidi D, Carta A, Alessio L. Prevention of injuries at work: the role of the occupational physician. *Int Arch Occup Environ Health*. 2006;79(3):177-92.
93. Ilmarinen J, Tuomi K. Work ability of aging workers. *Scand J Work Environ Health*. 1992;18 Suppl 2:8-10.
94. Costa G, Sartori S. Ageing, working hours and work ability. *Ergonomics*. 2007;50(11):1914-30.
95. Converso D, Sottimano I, Guidetti G, Loera B, Cortini M, Viotti S. Aging and Work Ability: The Moderating Role of Job and Personal Resources. *Front Psychol*. 2017;8:2262.
96. Kiss P, De Meester M, Braeckman L. Differences between younger and older workers in the need for recovery after work. *Int Arch Occup Environ Health*. 2008;81(3):311-20.
97. Ilmarinen J, Rantanen J. Promotion of work ability during ageing. *Am J Ind Med*. 1999;Suppl 1:21-3.
98. Feissel A, Peter R, Swart E, March S. Developing an Extended Model of the Relation between Work Motivation and Health as Affected by the Work Ability as Part of a Corporate Age Management Approach. *Int J Environ Res Public Health*. 2018;15(4).
99. Murray E. Web-based interventions for behavior change and self-management: potential, pitfalls, and progress. *Med 2 O*. 2012;1(2):e3.
100. Oakman J, Neupane S, Proper KI, Kinsman N, Nygard CH. Workplace interventions to improve work ability: A systematic review and meta-analysis of their effectiveness. *Scand J Work Environ Health*. 2018;44(2):134-46.
101. Rongen A, Robroek SJW, van Lenthe FJ, Burdorf A. Workplace health promotion: a meta-analysis of effectiveness. *Am J Prev Med*. 2013;44(4):406-15.
102. Lidegaard M, Sogaard K, Krusturup P, Holtermann A, Korshoj M. Effects of 12 months aerobic exercise intervention on work ability, need for recovery, productivity and rating of exertion among cleaners: a worksite RCT. *Int Arch Occup Environ Health*. 2018;91(2):225-35.

9

Summary

SUMMARY

Work ability related to job demands and functional limitations in various occupational populations

Work ability is of growing importance in life. As the retirement age rises, so does the expected duration of working life. Three decades ago, Dr. Juhani Ilmarinen introduced the concept of work ability because of the challenge of an aging workforce. Ilmarinen aimed to improve and measure the ability to meet job demands satisfactorily until later age without harming an individual's health. Work ability was conceptualized as a house model, in which both health and work aspects play an important role. Among others, functional limitations due to health complaints, and the characteristics of job demands, affect the ability to meet these job demands. Work ability depends on the balance between the demands put on the individual and the personal resources that he or she has to meet those demands.

The concept of work ability can be measured by the Work Ability Score (WAS), which is a self-assessment instrument of work ability. This scale assesses an individual's ability to fulfil job demands in his or her current job position and consists of a single item question comprising an 11-point scale. Zero corresponds with no work ability at all, and ten corresponds with the best work ability ever experienced by the individual.

The focus in work ability research lies on improving work ability to maintain working in good health until later age and maintaining good health after retirement. Measuring work ability can give insight into aspects associated with changing work ability. Two significant aspects of the work ability house model are health and job demands. In order to direct interventions and improve occupational healthcare, knowledge of these aspects is essential. We therefore need research to show which functional limitations or which job demands have the strongest impact on work ability. Once these aspects are identified, targeted intervention and occupational care is possible. To study these aspects of work ability in specific populations, populations with Upper Extremity Musculoskeletal Complaints (UEMSD) and stress were studied. The relationship between job demands and work ability were furthermore studied in populations characterized by high demands; in this case, ambulance workers and coach drivers. The work of ambulance workers is characterized by high peak loads, whereas coach drivers experience a seasonal workload increase during peak season.

The goal of this dissertation was to advance information on the concept of work ability and hence, three objectives were constructed to address scientific gaps on the topic:

1. To gather knowledge about characteristics of work ability measurement in groups of healthy employees and in groups of employees with health complaints and related functional limitations;

2. To gather knowledge about the association between functional limitations and work ability in working populations;
3. To gather knowledge about the course of work ability in work situations characterized by aspects of high workload or in work situations with seasonal changes in workload.

Based on the three objectives, three study questions were formulated and subsequently answered:

1. *What are the reproducibility and responsiveness of the work ability score of workers who return to work after sick leave?*

Prior to relating work ability to functional limitations and job demands, we studied the clinimetric properties of the WAS to identify whether this instrument could be used to monitor work ability in individuals. To fulfil the overarching first objective, the reproducibility and responsiveness to relevant changes of the work ability score were determined to identify the smallest detectable change in individuals.

In the first part of this dissertation, the work ability of a stable working population was analyzed. The WAS showed moderate reliability on repeated measures which was demonstrated by an ICC of 0.52. Additionally, the smallest detectable change (SDC) was determined. The SDC of work ability related to two points on the 0–10 scale, indicating that a score of two points entails a measurable change. A change of one consequently falls within the measurement error of repeated measurements and can therefore not be determined as significant change in an individual. Therefore only large changes of two or more points can be detected.

In elaboration of these measurement properties, additional clinimetrics of both instruments were assessed. In the third chapter, the clinical relevance of change in work ability and work functioning was assessed. To identify changes in an individual's work ability, measurement properties of the WAS were determined. In a population returning to work after sick leave, the SDCs of the work ability questions was 2 points. The minimal important change was determined to be 1 for work ability improvements or deterioration. Because the minimal important change was smaller than the smallest detectable change, we concluded that a change of two points also meant that a relevant change had occurred in an individual. This led to the conclusion that the WAS had limited responsiveness because the ability of the WAS to detect change was insufficient.

Because the minimal important change was smaller than the measurement error, the SDC needs to be used as cut-off value to determine change. Using the cut-off value of two points will result in many missed, or wrongly identified cases. However, this cut-off value is the most justified, resulting in the most correctly identified cases.

The second study question was:

II. How are functional limitations related to work ability in working populations with specific health complaints?

Although the WAS has limited use in individuals, it can be used to increase knowledge on aspects of work ability in groups of employees. Studying the relationship between functional limitations and work ability can shine a light on how work ability is affected and might be improved. With this purpose, two populations with specific functional limitations as a result of health complaints were studied: a population of workers with UEMSD complaints, and a young working population with stress complaints.

In workers with UEMSD (Chapter 4), functional limitations explained 52 % of the variance in work ability. The functional limitations that explained a significant amount of variance were the degree of difficulties performing daily activities at work, limitations in daily activities as a consequence of health issues, limitations at work and other daily activities due to physical health issues, and the ability to perform a precision grip. This shows that work ability is a concept determined to a large extent by functional limitations in UEMSD patients.

Most work ability research is performed in aging populations. While work ability declines with age, work ability in young workers can also be affected. Stress is a common phenomenon experienced by young workers. Starting a career with a demanding social environment while developing a family life can cause stress to accumulate. In fact, in young workers, stress had a negative impact on work ability five years down the road (Chapter 5). A large part of this relationship was mediated by several other factors. Stress that occurred five years later, job demands affecting private life, feelings of control over private life, and feeling well-rested upon waking accounted for 81% of the relationship between stress and work ability five years later. With these mediators in our model, the direct effect was no longer significant indicating that these mediators are important in preventing stress from affecting work ability, as well as preventing declines in actual work ability. Physical activity in leisure time was not associated with the mediation between stress and work ability.

In the third research question, job demands were related to work ability.

III. What is the course of work ability in occupations characterized by specific health requirements and peak loads, and to what extent can this be influenced?

Just as functional limitations, job demands constitute an important determinant of work ability. The more demanding those demands are, the more they affect work ability. If job demands increase, the resources to cope with these demands need to follow suit to avoid the loss of work ability. With our third objective, we aimed to

identify the relationship between the ability to meet specific job demands and work ability, and how the course of work ability is influenced by this balance.

In Chapter 6, the job demands of ambulance workers were studied. This occupation is characterized by high peak loads as high physical and mental demands are placed on workers. Self-reported work ability assessed by the WAS was related to the ability to meet the job demands to identify which job demands have the largest influence on work ability. For general work ability, the outcomes in alertness and judgment ability, and emotional peak load contributed most. Job-specific physical abilities and raised alertness and judgment ability aspects significantly explained variance in physical work ability, while variance in mental/emotional work ability was significantly explained by emotional peak load and raised alertness and judgment ability. All significant categories together only explained a variance of 17 % which indicated that other aspects are of larger importance for work ability in ambulance workers than those studied.

Additional value is attributed to workers health surveillance because merely 1 % of ambulance workers appraised their work ability as insufficient, while in actuality 68 % scored insufficient on one or more of the specific demands required to satisfactorily meet their job standards.

In the case of coach drivers, the peak spring and summer seasons places an increasing workload on them. An increasing workload and decreasing ability to recover from it threatens work ability. Additionally, the percentage of drivers who reported being very tired increased from 22 % at the beginning of peak season to 53 % at the end of it, demonstrating this situation puts drivers at risk of failing to meet their job demands satisfactorily until pensionable age. In a randomized controlled trial, the use of a tailored self-management intervention during the peak season is evaluated (Chapter 7). During the peak season, the work ability and vitality decreased significantly while work-related fatigue accumulated in both the intervention and control group. The trajectory of work ability showed a significant declining trend, indicating an imbalance between job demands and the resources needed to meet these demands

Although the separate components of the intervention were well-received, the uptake was too low to expect differences between the two groups. The low adherence indicated that this type of intervention is not suitable for coach drivers during peak season and that alternative implementation of the intervention should be sought.

In conclusion, the work ability score is not suited for individual assessment of work ability due to its difficulties detecting change in individuals. However, when related to functional limitations or job demands in groups of workers it can be useful. Linking functional limitation and job demands with work ability contributes to knowledge of specific working populations. This knowledge can help improve occupational health

care, direct preventive measures and help develop interventions in order to assist workers in maintaining their ability to meet their job demands satisfactorily while preserving their health.

10

Samenvatting

SAMENVATTING

Werkvermogen gerelateerd aan functie-eisen en functionele beperkingen in verschillende beroepspopulaties

Het evenwicht tussen de eisen die het werk aan een individu stelt en de middelen die de werkende ter beschikking heeft om de werkzaamheden uit te voeren wordt het werkvermogen genoemd. Werkvermogen wordt steeds belangrijker in het huidige leven. De pensioenleeftijd stijgt en daarmee wordt ook de verwachte arbeidsduur langer. Er komt daarom meer aandacht voor het in goede gezondheid kunnen werken tot de pensioengerechtigde leeftijd en behoud van een goede gezondheid na pensionering.

Het begrip werkvermogen is door prof. dr. Ilmarinen gedefinieerd als 'het vermogen van werkenden om op dit moment en in de nabije toekomst hun werk naar tevredenheid uit te voeren met inachtneming van de werkeisen, hun gezondheid en mentale capaciteit'.

Het werkvermogen van een individu is een complex concept waarop veel factoren van invloed zijn. Zowel het type werk dat iemand doet als specifieke gezondheidsklachten kunnen het werkvermogen van werkenden beïnvloeden. In groepen mensen met bijvoorbeeld klachten aan hand, arm, nek of schouders (RSI-klachten) of met stress kan het werkvermogen negatief worden beïnvloed. Daarnaast kan hoge werkdruk invloed hebben op het werkvermogen. Dit kan het geval zijn voor beroepen als ambulancemedewerker of touringcarchauffeur.

Prof. dr. Ilmarinen visualiseert het concept werkvermogen in de vorm van een huis, waarin vele aspecten samenkomen. De basis wordt gevormd door een goede fysieke en mentale gezondheid. Deze fundering heeft gevolgen voor de andere verdiepingen. De eerste verdieping bevat de vaardigheden en kennis die van belang zijn bij het werk en bouwt voort op de fundering van het huis, de gezondheid. De vaardigheden en kennis kunnen constant veranderen, omdat individuen bijleren en vaardigheden ontwikkelen. Op de volgende verdieping bevinden zich de normen en waarden, niet alleen van de werkende zelf, maar ook van het bedrijf waarvoor de werkende werkt. Wanneer de competenties voldoende zijn, komen de houding en motivatie om te werken in het spel. De eerste drie verdiepingen van het huis vertegenwoordigen de mogelijkheden en middelen van werkenden, en de bovenste verdieping bestaat uit het werk zelf en de bijbehorende werkeisen. Hieronder vallen onder andere de werkomgeving, de werkinhoud en de werkorganisatie. Alle verdiepingen samen dragen bij aan een goed werkvermogen. Het werkvermogen kan daarnaast zowel positief als negatief worden beïnvloed door zaken van buiten het huis zoals familie, directe sociale en operationele omgeving en de samenleving. Alle verdiepingen en de

omgeving van het huis moeten met elkaar in balans zijn om voldoende niveau van werkvermogen te houden. Twee van de belangrijkste aspecten van het huis zijn de gezondheids- en werkaspecten.

Het werkvermogen hangt dus sterk af van de balans tussen de functie-eisen en de mogelijkheden en middelen (zoals gezondheid) die een werkende ter beschikking heeft om aan deze eisen te voldoen. Als de gezondheid afneemt, neemt het vermogen om aan de functie-eisen te voldoen en daarmee het werkvermogen ook af. Het evenwicht tussen de eisen en mogelijkheden en middelen om daarmee om te gaan is daarom belangrijk om een goed werkvermogen te behouden gedurende het werkende leven.

Werkvermogen kan worden gemeten aan de hand van de werkvermogenscore (Work Ability Score, vanaf nu de WAS genoemd). Met behulp van de WAS kan een werkende zelf het huidige werkvermogen aangeven ten opzichte van het beste werkvermogen dat hij of zij ooit in het werkende leven heeft ervaren. Op een schaal van 0–10 komt ‘helemaal geen werkvermogen’ overeen met nul, en het beste werkvermogen dat het individu ooit heeft ervaren overeen met tien.

Het accent bij onderzoek naar werkvermogen ligt op het verbeteren van het werkvermogen om ervoor te zorgen dat dit tot op latere leeftijd behouden kan worden. Het meten van het werkvermogen kan inzicht geven in aspecten die verband houden met een verminderd werkvermogen. Om interventies in te zetten en de kwaliteit van bedrijfsgezondheidszorg te verbeteren, is daarnaast kennis nodig over de aspecten die het nauwst verband houden met werkvermogen. Door te onderzoeken welke functionele beperkingen en werkeisen het sterkst samenhangen met werkvermogen wordt informatie verzameld over welke aspecten belangrijk zijn voor het kunnen uitvoeren van werk. Functionele beperkingen als gevolg van gezondheidsproblemen (zoals RSI of stressklachten) of het niet kunnen voldoen aan functie-eisen kunnen invloed hebben op het werkvermogen. De aspecten die werkvermogen negatief beïnvloeden kunnen worden opgespoord met als doel deze te verbeteren. Aspecten die de grootste invloed hebben op werkvermogen zijn belangrijk voor het verbeteren of behouden van werkvermogen. Het doel van dit proefschrift is om informatie te verzamelen over het concept werkvermogen. Daartoe zijn drie doelstellingen opgesteld om wetenschappelijke hiaten in de kennis op dit gebied op te vullen.

Deze doelstellingen zijn als volgt omschreven:

1. Kennis vermeerderen over het verloop van werkvermogen in groepen gezonde werkenden en bij groepen werkenden met gezondheidsklachten en gerelateerde functionele beperkingen;

2. Kennis vermeerderen over het verband tussen functionele beperkingen en werkvermogen in werkende populaties;
3. Kennis vermeerderen over het verloop van het werkvermogen in werksituaties die worden gekenmerkt door hoge werkbelasting en in werksituaties met seizoensgebonden veranderingen in werkdruk.

Gekoppeld aan elke doelstelling is een onderzoeksvraag geformuleerd. De vragen en de antwoorden hierop worden hieronder per vraag besproken:

1. *Hoe betrouwbaar zijn herhaalde metingen en welk verschil in WAS is in de praktijk een relevant verschil?*

Om het verloop van werkvermogen te kunnen bestuderen, hebben we de eigenschappen van de WAS onderzocht in hoofdstuk 2 en 3. Zo konden we bepalen of dit meetinstrument geschikt is om het werkvermogen bij individuen in de tijd te monitoren. Om aan doelstelling 1 te voldoen, zijn de betrouwbaarheid en het vermogen van de WAS om veranderingen in de tijd te registreren bepaald. Om dit te kunnen doen is de kleinst meetbare verandering (KMV) bij individuen voor dit instrument bepaald. Het verschil op een schaal van 0–10 dat daadwerkelijk een verschil in de praktijk betekende is vastgesteld.

Uit ons onderzoek bleek dat de WAS een matige betrouwbaarheid bij herhaalde metingen heeft en niet kan worden gebruikt om een verandering bij een individu te meten. Pas wanneer er grote veranderingen van meer dan twee punten op de schaal van het meetinstrument optreden kan dit worden gezien als een relevante verandering. Omdat niet iedereen die twee punten op de schaal veranderde, zelf ook vond dat zijn/haar werkvermogen veranderd was, kan dus niet met zekerheid onderscheid worden gemaakt tussen een verbeterd, een verslechterd en een onveranderd persoon, als deze grenswaarde wordt gebruikt. Dit resulteerde in veel foutief geclassificeerde individuen. Hieruit wordt de conclusie getrokken dat de WAS ongeschikt is voor het monitoren van individuele personen.

De conclusie van hoofdstuk 2 en 3 is dat de WAS niet geschikt is voor individuele monitoring van werkvermogen. Hoewel de WAS voor individuen beperkt bruikbaar is –doordat deze alleen geschikt is om grote veranderingen op te merken– kan de WAS wel worden gebruikt om de kennis te vergroten over aspecten van het werkvermogen bij groepen werkenden. Het gebruik van de WAS kan nuttig zijn om te bepalen welke gezondheids- en werkaspecten het sterkst zijn gerelateerd aan het werkvermogen van groepen met functionele beperkingen door gezondheidsklachten zoals RSI-klachten of stress. Door de tweede onderzoeksvraag te beantwoorden wordt inzicht verkregen in de mogelijke relatie tussen functionele beperkingen en het ervaren werkvermogen:

II. Hoe zijn functionele beperkingen gerelateerd aan het werkvermogen in werkende populaties met specifieke gezondheidsklachten?

In het 'huis van werkvermogen' van prof. dr. Ilmarinen vormt gezondheid de basis van het werkvermogen. Voldoende gezondheid is een van de middelen die nodig zijn om werk uit te voeren en aan de werkeisen te voldoen. Dit leidt tot de aanname dat functionele beperkingen het werkvermogen nadelig kunnen beïnvloeden. Het bestuderen van de relatie tussen functionele beperkingen en werkvermogen kan licht werpen op de wijze waarop het werkvermogen wordt beïnvloed en kan worden verbeterd. Zo kunnen bepaalde klachten ertoe leiden dat een werkende niet meer kan voldoen aan de werkeisen waardoor het werkvermogen daalt. Dit kan op zijn beurt weer leiden tot een afname van het aantal gewerkte uren.

De functionele beperkingen verklaarden 52 % van de verschillen in werkvermogen tussen werkenden met RSI-klachten. Dat wil zeggen dat de verschillen in werkvermogen voor ongeveer de helft worden bepaald door het wel of niet kunnen uitvoeren van bepaalde werkgerelateerde handelingen. De functionele beperkingen die een aanzienlijk verschil verklaarden, waren de mate van problemen bij het uitvoeren van dagelijkse activiteiten op het werk, beperkingen in dagelijkse activiteiten als gevolg van gezondheidsproblemen, beperkingen op het werk en van andere dagelijkse activiteiten als gevolg van lichamelijke gezondheidsproblemen, en het vermogen om een precisiegrip uit te voeren. Wanneer deze werkgerelateerde handelingen niet konden worden uitgevoerd, had dat de grootste invloed op werkvermogen. Dit toont aan dat werkvermogen een concept is dat nauw verband houdt met functionele beperkingen bij RSI-patiënten.

Omdat bekend is dat het werkvermogen afneemt naarmate een werkende ouder wordt, wordt het meeste onderzoek naar werkvermogen uitgevoerd onder oudere bevolkingsgroepen. Maar ondanks het feit dat werkvermogen afneemt met de leeftijd, kan het werkvermogen ook bij jonge werkenden zijn aangetast. Stress komt al bij jonge werkenden voor. Het starten van een carrière in een veeleisende sociale omgeving en daarnaast een gezinsleven ontwikkelen kan ertoe leiden dat stress zich opstapelt. De resultaten van het onderzoek in hoofdstuk 5 laten zien dat stress bij jongeren een negatieve invloed heeft op het werkvermogen van jonge werkenden vijf jaar later. Een groot deel (81 %) van deze relatie kan verklaard worden door verschillende andere factoren. De factoren die deze relatie sterk beïnvloedden waren de mate van stress die vijf jaar later aanwezig was, werkeisen die het privéleven negatief beïnvloedden, gevoelens van controle over het privéleven en het gevoel goed uitgerust wakker te worden. Aandacht voor deze factoren is belangrijk om te voorkomen dat door stress het werkvermogen negatief wordt beïnvloed.

De derde onderzoeksvraag betreft de relatie tussen werkeisen en het ervaren werkvermogen. Een verhoging van werkdruk, zonder vergroting van middelen om hiermee om te gaan kan het werkvermogen negatief beïnvloeden. De derde onderzoeksvraag is als volgt geformuleerd:

III. Hoe verloopt het werkvermogen in beroepen die worden gekenmerkt door specifieke gezondheidseisen en piekbelastingen, en in hoeverre kan dit worden beïnvloed?

Net als functionele beperkingen zijn functie-eisen belangrijk voor het niveau van het werkvermogen. Hoe veeleisender het werk is, hoe meer deze beperkingen het werkvermogen negatief kunnen beïnvloeden. Als de werkdruk wordt verhoogd, dienen ook de middelen om aan deze eisen te voldoen te worden vergroot, om verlies van werkvermogen te voorkomen.

Het werk van ambulancepersoneel wordt gekenmerkt door hoge piekbelastingen. Aan deze werkenden worden hoge fysieke en mentale eisen gesteld. Tijdens de periodieke gezondheidscontrole van werkenden wordt getest of werkenden aan deze werkeisen kunnen voldoen. In hoofdstuk 6 is onderzocht in welke mate zelf-gerapporteerd werkvermogen is gerelateerd aan het kunnen voldoen aan de functie-eisen van het beroep ambulancemedewerker. Hierbij zijn de functie-eisen die de grootste invloed hebben op het werkvermogen geïdentificeerd. Daarbij is onderscheid gemaakt tussen drie typen werkvermogen die met de WAS zijn gemeten: 1) het algemene werkvermogen, 2) het fysieke werkvermogen en 3) het mentale werkvermogen. Door de werkeisen te relateren aan het werkvermogen kan worden beoordeeld welke werkeisen het belangrijkst zijn om aan te voldoen om voldoende werkvermogen te hebben.

Voor algemeen werkvermogen waren de uitkomsten met betrekking tot alertheid en beoordelingsvermogen met daarnaast emotionele piekbelasting het belangrijkst. Werkspecifieke fysieke vaardigheden, verhoogde alertheid en beoordelingsvermogen droegen het sterkst bij aan fysiek werkvermogen, terwijl mentaal/emotioneel werkvermogen het sterkst gerelateerd was aan emotionele piekbelasting en daarnaast verhoogde alertheid en beoordelingsvermogen. De belangrijkste categorieën samen voorspelden 17 % van de verschillen in werkvermogen, wat betekent dat er meer aspecten van belang zijn voor het werkvermogen van ambulancepersoneel.

Het met de WAS beoordelen van het werkvermogen in ambulancemedewerkers voldoet niet omdat slechts 1 % van het ambulancepersoneel zelf het werkvermogen op de WAS als onvoldoende beoordeelde, terwijl 68 % onvoldoende scoorde op een of meer van de specifieke functie-eisen die nodig zijn om naar tevredenheid te kunnen

functioneren. De periodieke gezondheidscontrole heeft een toegevoegde waarde, omdat het kunnen uitvoeren van de functie-eisen niet goed kan worden ingeschat door ambulancemedewerkers zelf.

Voor touringcarchauffeurs zorgt het hoogseizoen (voorjaar en zomer) voor een hogere werkdruk. Deze hogere werkdruk en verminderde mogelijkheden om van deze werkdruk te herstellen, bedreigen het werkvermogen en de vitaliteit. Daarnaast is het moeilijk om er een gezonde levensstijl op na te houden. In een onderzoek waarbij twee groepen met elkaar zijn vergeleken, is het gebruik van op maat gemaakte digitale zelfhulptips met bijbehorende opdrachten tijdens het hoogseizoen geëvalueerd (hoofdstuk 7). Voor dit onderzoek zijn de chauffeurs willekeurig in twee groepen verdeeld waarbij één groep de digitale zelfhulptips en opdrachten ontving (interventiegroep,) en de andere groep niet (controlegroep).

Tijdens het hoogseizoen namen het werkvermogen en de vitaliteit aanzienlijk af, terwijl werkgerelateerde vermoeidheid zich opstapelden in zowel de groep die de digitale zelfhulp tips kreeg, als de controlegroep. Het werkvermogen daalde tijdens het hoogseizoen aanzienlijk en dit duidt op een disbalans tussen de werkeisen en de middelen om aan deze eisen te voldoen.

Ter illustratie: het percentage touringcarchauffeurs dat zich erg vermoeid voelde steeg van 22 % aan het begin van het hoogseizoen naar 53 % aan het einde van het hoogseizoen. Hoewel de tips goed werden ontvangen, was het gebruik ervan te gering om verschillen in werkvermogen en vitaliteit tussen de twee groepen te kunnen verwachten. Het geringe gebruik van de digitale zelfhulptips duidt erop dat dit type interventie niet geschikt is voor touringcarchauffeurs tijdens het hoogseizoen en dat er gezocht moet worden naar een andere manier voor het geven van tips dan digitale zelfhulp met opdrachten. Een voorbeeld kan zijn om de tips aan te bieden door middel van face-to-face voorlichting en gedurende het hoogseizoen begeleiding te geven.

De conclusie van de onderzoeken in dit proefschrift is dat de werkvermogenscore (WAS) niet geschikt is voor individuele beoordeling van het werkvermogen, vanwege het onvermogen om veranderingen bij individuele personen te detecteren. Echter, wanneer de score wordt gerelateerd aan functionele beperkingen of functie-eisen in groepen werkenden kan de WAS wel van nut zijn. In dat geval draagt deze bij aan kennis over de manier waarop het werkvermogen wordt beïnvloed in specifieke beroepspopulaties en kan het ondersteuning bieden bij het opzetten en in de praktijk brengen van preventieve maatregelen in de bedrijfsgezondheidszorg. Daarnaast kan de opgedane kennis helpen bij de ontwikkeling van interventies om werkenden te helpen hun vermogen te behouden om naar tevredenheid, en met behoud van hun gezondheid aan hun werkeisen te voldoen.

11

Appendices

- I. List of abbreviations
- II. Curriculum Vitae
- III. List of publications
- IV. Co-authors and their affiliation
- V. Portfolio
- VI. Acknowledgments/Dankwoord

LIST OF ABBREVIATIONS

AMC	Academic Medical Center
ANCOVA	ANalyses of COVariance
AUC	Area Under the Curve
BMI	Body Mass Index
BSI	Brief Symptoms Inventory
CI	Confidence Interval
CONSORT	CONsolidated Standards Of Reporting Trials
COSMIN	Consensus-based Standards for the selection of health Measurement INstruments
CPR	CardioPulmonary Resuscitation
DASH	Disability of the Arm, Shoulder and Hand
DF	Degrees of Freedom
ESS	Epworth Sleepiness Scale
GRC	Global Ratings of Change
ICC	Intraclass Correlation Coefficient
ISCO	International Standard Classification of Occupations
ITT	Intention-To-Treat
MCAR	Missing Completely at Random
MIC	Minimal Important Change
NFR	Need For Recovery
NS	Not Significant
PME	Preventive Medical Examination
PPA	Per-Protocol-Analysis
RCT	Randomized controlled Trial
ROC	Receiver Operating Characteristic
RSI	Repetitive Strain Injury
RTW	Return To Work
SD	Standard Deviation
SDC	Smallest Detectable Change
SEB	Schaal Ervaren Belasting
SEM	Standard Error of Measurement
SF-36	Short-Form-36
UEMSD	Upper Extremity MusculoSkeletal Disorders
VIF	Variance Inflation Factor

VOEG	Vragenlijst Onderzoek Ervaren Gezondheid
WA	Work Ability
WA-PR	Work Ability-Personal Radar
WAI	Work Ability Index
WAS	Work Ability Score
WAYA	Work Ability in Young Adults
WF	Work Functioning
WHS	Workers Health Surveillance

CURRICULUM VITAE

Art van Schaaijk is geboren op 7 april 1986 in 's-Hertogenbosch. In 2005 voltooide hij het Gymnasium aan het R.K. Gymnasium Beekvliet, te Sint-Michielsgestel. Na dit middelbaar onderwijs volgde de reis naar Maastricht om daar de studie Algemene Gezondheidswetenschappen te volgen aan de Universiteit Maastricht. Direct koos hij als major de specialisatie Bewegingswetenschappen, met Bioregulatie & Gezondheid als minor.

Gedurende deze studie had Art verschillende nevenfuncties waaronder voetbaltrainer en fitnessinstructeur bij de sportschool van de Universiteit Maastricht. Daarnaast heeft hij verschillende bestuursfuncties bekleed waaronder die van voorzitter van de studentenvoetbalvereniging van Maastricht.

Na het afronden van de bachelor bewegingswetenschappen is hij meteen gestart met de master Biology of human performance and health. Tijdens zijn master heeft hij onderzoek verricht in de jeugdopleiding van PSV waarbij de fysieke en fysiologische belasting werd vergeleken tussen verschillende trainingsvormen en wedstrijden. Na het afronden van deze master heeft hij nog anderhalf jaar meerdere functies bekleed bij de sportschool van de Universiteit Maastricht, waarna de kans zich voordeed om zich verder te ontwikkelen in het onderzoek door middel van het doen van promotieonderzoek bij het Coronel Instituut voor Arbeid en Gezondheid in het AMC, nu Amsterdam UMC aan de Universiteit van Amsterdam.

Dit promotieonderzoek werd uitgevoerd onder begeleiding van Prof. dr. Judith Sluiter, MBA en Prof. dr. Monique Frings-Dresen, waarna later dr. Karen Nieuwenhuijsen aan het begeleidingsteam werd toegevoegd. In de periode van juli 2015–juni 2020 heeft hij onderzoek verricht naar werkvermogen in verschillende populaties. Tijdens deze periode heeft hij verschillende onderzoeks- en analysetechnieken toegepast bij onderzoek in diverse populaties. Tijdens zijn promotietraject heeft hij zijn onderzoeksresultaten gepresenteerd op meerdere nationale en internationale congressen. Daarbij is ook een gezamenlijk project ontstaan met een onderzoeksgroep uit Zweden, waar tijdens een werkbezoek aan een wetenschappelijk onderzoek is gewerkt. Daarnaast nam hij deel aan cursussen om zichzelf verder te ontwikkelen en specialiseren in het onderzoek en was hij betrokken bij het opstarten van een junior council van het Amsterdam Public Health research institute. Tijdens deze periode is hij daarnaast betrokken geweest bij het onderwijzen van geneeskundestudenten op het gebied van academische vorming en professionele ontwikkeling. Op dit moment is hij werkzaam in de functie van post-doc waarin het zijn taak is om subsidies te verwerven en onderzoek voort te zetten.

LIST OF PUBLICATIONS

Articles included in this dissertation

van Schaaik, A., Boschman, J. S., Frings-Dresen, M. H. W., & Sluiter, J. K. (2017). Appraisal of work ability in relation to job-specific health requirements in ambulance workers. *International archives of occupational and environmental health*, 90(1), 123-131.

van Schaaik, A., Nieuwenhuijsen, K., Frings-Dresen, M. H. W., & Sluiter, J. K. (2018). Reproducibility of work ability and work functioning instruments. *Occupational Medicine*, 68(2), 116-119.

van Schaaik, A., Nieuwenhuijsen, K., Frings-Dresen, M. H. W., & Sluiter, J. K. (2019). Work ability and work functioning: measuring change in individuals recently returned to work. *International archives of occupational and environmental health*, 92(3), 423-433.

van Schaaik, A., Nieuwenhuijsen, K., & Frings-Dresen, M. (2019). Work Ability and Vitality in Coach Drivers: An RCT to Study the Effectiveness of a Self-Management Intervention during the Peak Season. *International Journal of Environmental Research and Public Health*, 16(12), 2214.

van Schaaik, A., Noor Baloch, A., Thomée, S., Frings-Dresen, M., Hagberg, M., & Nieuwenhuijsen, K. (2020). Mediating Factors for the Relationship between Stress and Work Ability over Time in Young Adults. *International Journal of Environmental Research and Public Health*, 17(7), 2530.

van Schaaik A, Nieuwenhuijsen K, Frings-Dresen MHW. (2020). Work ability and percentage of hours worked related to limitations in patients with upper extremity musculoskeletal disorders: a cross-sectional cohort study. *BMC Musculoskeletal Disorders*, 21(1):389.

Articles not included in this dissertation

Schaaik A van, Nieuwenhuijsen K, Frings-Dresen MHW. *Vitaler het touringcar hoogseizoen door: een evaluatie van interventies*. Coronel Instituut voor Arbeid en Gezondheid, Academisch Medisch Centrum, Universiteit van Amsterdam, Amsterdam Public Health research institute, 2018:124 p. (Coronel rapportnummer; 18-03).

Schaaik A van, Nieuwenhuijsen K, Frings-Dresen MHW. *Vitaler het touringcar hoogseizoen door: Toolbox*. Coronel Instituut voor Arbeid en Gezondheid, Academisch Medisch Centrum, Universiteit van Amsterdam, Amsterdam Public Health research institute, 2018:24 p. (Behorend bij Coronel rapportnummer; 18-03).

Schaaik A van, Nieuwenhuijsen K, Frings- Dresen MHW. *Vitaler het touringcar hoogseizoen door: Achtergronddocument Toolbox*. Coronel Instituut voor Arbeid en Gezondheid, Academisch Medisch Centrum, Universiteit van Amsterdam, Amsterdam Public Health research institute, 2018:30 p. (Behorend bij Coronel rapportnummer; 18-03).

CO-AUTHORS AND THEIR AFFILIATION

Julitta Boschman¹

Contributed to Chapter 6.

Monique Frings-Dresen¹

Contributed to all chapters in this dissertation.

Mats Hagberg²

Contributed to Chapter 5.

Karen Nieuwenhuijsen¹

Contributed to Chapter 1, 2, 3, 4, 5, 7 and 8 in this dissertation

Adnan Noor Baloch³

Contributed to Chapter 5.

Judith Sluiter¹

Contributed to Chapter 2, 3 and 6.

Sara Thomee⁴

Contributed to Chapter 5.

All authors contributed by reading and revising the respective manuscripts. Conceptualising the study, and methodology was a joint task of all co-authors. Data checks and checks of the statistical analyses were performed by Julitta Boschmann and Karen Nieuwenhuijsen for the studies that they contributed to. Judith Sluiter and Monique Frings-Dresen shared ultimate responsibility as PI's and signed off on projects.

¹ Amsterdam UMC, University of Amsterdam, Coronel Institute of Occupational Health, Amsterdam Public Health research institute, Meibergdreef 9, P.O. Box 22660, 1100 DE Amsterdam, The Netherlands

² Occupational and Environmental Medicine, School of Public Health and Community Medicine, Sahlgrenska Academy and University Hospital, University of Gothenburg, P.O. Box 414, 405 30 Gothenburg, Sweden

³ Biostatistics, School of Public Health and Community Medicine, Institute of Medicine, University of Gothenburg, P.O. Box 463, 405 30 Gothenburg, Sweden

⁴ Department of Psychology, University of Gothenburg, P.O. Box 500, 405 30 Gothenburg, Sweden

PHD PORTFOLIO

Name PhD candidate: Art van Schaaijk
 Name PhD supervisor: Prof. dr. J.K. Sluiter, MBA (until May 2018) and
 Prof. dr. M.H.W. Frings-Dresen
 PhD period: July 2015 – June 2020

1. PhD training	Year	Workload (ECTS)
General courses		
The AMC World of Science	2015	0.7
Endnote course	2015	0.1
Scientific writing in English for publication	2015	1.5
e-BROK (Basic course on Regulations and Organisation for clinical investigators)	2016	1.5
Didactical skills	2018	0.4
e-BROk re-registration	2020	0.5
Specific courses		
Practical biostatistics	2015	1.4
Course Clinical Epidemiology 1: Randomized Clinical Trials	2017	0.6
Advanced topics in biostatistics	2018	2.1
Entrepreneurship in health and life sciences	2020	1.5
Seminars, workshops and master classes		
<i>APROVE workshops</i>		
Pimp my EXCEL	2015	0.1
Pimp my tax return	2016	0.1
Pimp my thesis	2019	0.1
<i>Other</i>		
Seminar Informed Consent, Amsterdam	2016	0.1
Masterclass prof. John Ioannidis; Higher scientific standards	2016	0.2
(Inter)national conferences		
- Poster presentations		
Amsterdam Public Health Annual meeting (4)	2016-2019	1
Occupational Health days, Papendal (3)	2017-2019	0.8
International Congress on Occupational Health, Dublin	2018	1.2
Science Exchange Day, Amsterdam	2019	1.3

- Oral presentations		
Research meeting Coronel Institute (7)	2016-2020	1.4
Occupational Health days, Papendal	2016	0.2
CaRe days, Maastricht	2017	1
Science day, Ministry of social affairs and Employment	2017	0.5
PREMUS, Bologna	2019	1
- Attended conferences without presentation		
Heijermanslezing, Amsterdam (8)	2015-2019	0.8
International Symposium on Occupational Health Care, Amsterdam	2016	0.3
Work Disability Preventions & Integration conference, Amsterdam	2016	1.4
Societal Participation and Health meetings (3)	2017-2019	0.3
Amsterdam Public Health Spring meeting (3)	2017-2019	0.6
Science day, Ministry of social affairs and Employment (2)	2018-2019	0.5
CaRe days, Eindhoven	2019	1
Societal Participation and Health junior meeting	2019	0.1
Day of sport research, Amsterdam	2019	0.3
- Other courses/meetings		
Get together conference (Europe's biggest conference for movement professionals), Amsterdam (2)	2016-2018	0.8
PhD candidate committee Research Institute APH (4)	2017-2020	4
Coronel Institute: Commission Annual report	2017-2020	1.1
Organising junior committee APH Annual day (2)	2019-2020	0.3
2. Teaching		
Lecturing		
Academic education; Academic skills (5)	2016-2020	4
Designing a scientific poster	2019	0.1
Physician and top sports	2019-2020	0.2
Other		
Supervising junior internships	2017	0.2
Prevention day for coach drivers(2)	2018	0.4
Supervising elective education	2019	0.4
Total (28 hrs = 1 ECTS)		36.1

ACKNOWLEDGMENTS/DANKWOORD

Bij het afsluiten van deze wetenschappelijke periode zou ik graag even de aandacht willen vestigen op iedereen die dit mogelijk heeft gemaakt. Reden dus om mijn dank te tonen aan allen die me in deze periode op welke mogelijke manier dan ook geholpen hebben. Allereerst wil ik iedereen bedanken die heeft deelgenomen aan een van mijn onderzoeken. Misschien zullen velen dit nooit lezen maar toch, zonder jullie was het voltooien van dit proefschrift nooit gelukt.

Ondanks dat er soms frustrerende periodes tussen zaten, is de moed me nooit in de schoenen gezakt. Leerzame en leuke momenten voerden gelukkig altijd de boventoon. In het begin werd ik in het diepe gegooid, maar tegelijkertijd toch scherp in de gaten gehouden. Vervolgens heb ik mijn eigen weg gezocht en merkte ik dat ik vrijer werd gelaten. In een periode van vijf jaar is het begrijpelijk dat leuke momenten afgewisseld worden met momenten waarop het soms wat moeizamer gaat. Altijd waren er op die momenten verschillende personen op wie ik kon rekenen en die me er doorheen hebben gesleept, die ik nu wil bedanken.

Allereerst wil ik mijn dank uitspreken aan mijn promotor en co-promotor. Monique, Jij hebt me van het begin tot het eind begeleid. Vanaf mijn sollicitatie tot aan hier. Vanwege je pensioen zou je een stap terug zetten en zou je Judith het grootste deel voor haar rekening laten nemen. Helaas liep dat anders. Je stapte direct terug in de ring en bleef zo je vrije ochtenden opofferen om mij te begeleiden. Ik heb de laatste tijd steeds vaker van je gehoord: 'Zoals dit nu gaat heb ik in al die tijd nog nooit meegemaakt', wanneer bijvoorbeeld het ging over de twee weken die ik in Zweden verbleef, en terugkwam met een bijna volledig manuscript, of wanneer er een aparte procedure doorlopen moest worden. Soms zorgde je ook voor wat frustratie. Bijvoorbeeld wanneer je tijdens het weekoverleg bij een bijna volledig manuscript bedacht, dat we er beter twee studies van konden maken. Gelukkig kon ik deze frustraties altijd in de avond botvieren op je, wanneer je naar mijn trainingen kwam in het USC. Voor straf kreeg je dan twee dagen spierpijn.

Karen, vanaf het tweede project ben je verbonden geweest aan mijn onderzoeken. In het begin verzorgde je de data-checks, en daarna werd je meer betrokken bij het schrijfproces. Vanaf het moment dat bekend werd dat Judith ziek was, is besloten je definitief toe te voegen aan het begeleidingsteam. Ontzettend nauwkeurig en precies ging je met alles te werk. Dat paste wel goed bij mijn instelling. Dankzij deze alertheid hield je mij ook altijd scherp. Je vertelde me exact waar mensen iets verkeerd konden begrijpen en spoorde me aan om zinsconstructies anders te verwoorden. Daarnaast was het fijn om korte lijntjes met je te hebben als coördinator van het onderwijs, zo waren problemen altijd snel verholpen. Toch heb ik nog wat voor jullie verborgen

kunnen houden. Bij het schrijven van publicaties wilde ik altijd graag het onderwerp van mijn promotietraject direct naar voren laten komen. Volgens mij is dat aardig gelukt als je kijkt naar ieder eerste woord van de introducties.

Graag bedank ik ook alle leden van de leescommissie; Prof dr. Nynke van Dijk, Prof dr. Raoul Engelbert, Prof dr. Anton Kunst, Prof dr. Ute Bültmann en Prof dr. Willem van Rhenen, voor het beoordelen van mijn dissertatie, en de bereidheid om te opponeren tijdens de openbare verdediging hiervan.

Mijn allereerste stappen als PhD student maakte ik aan de zijde van Julitta. Jij had het hele traject net afgerond en hebt me laten zien wat de valkuilen zijn, en waar ik extra op moest letten. Fijn dat je me meteen op sleeptouw nam.

Tijdens mijn periode bij het Coronel Instituut heb ik met vele collega's de kamer gedeeld. Door tijdelijke contracten en het promoveren van mijn kamergenoten heb ik er veel versleten: Martijn, Inge, Maria, Marjolein en Bedrah, bedankt voor de gezellige, leerzame en vaak korte tijd. Gelukkig was er wel altijd tijd voor een praatje over wat dan ook. Een langere periode heb ik de kamer gedeeld met Maryam en Michiel. Maryam, je schreef het zelf al: Als complete tegenpolen vonden we elkaar in liefde voor muziek, sport en Michiko. Of deze laatste de afgelopen tijd overleefd heeft weet ik niet, maar ik heb in ieder geval goed aan mijn groene vingers kunnen werken. Dank voor je altijd vrolijke en enthousiaste kijk op alles. Daarnaast heb ik het laatste deel van mijn promotietraject bij Michiel op de kamer doorgebracht. Doordat we samen in de afrondende fase zaten van ons proefschrift, hebben we veel aan elkaar gehad door ervaringen uit te wisselen en procedures door te nemen. Doordat dat jij vaak de late avonden werkte en ik de vroege ochtenden, was er in ieder geval altijd bedrijvigheid in onze kamer. Dank voor de momenten samen waarop we elkaar tips konden geven, ondanks dat onze onderzoeken anders in elkaar staken. Bedankt dat jij nu ook naast me staat.

Ook wil ik de rest van mijn (oud)collega's van het Coronel Instituut bedanken. Na vergaderingen en op de gang, er kon altijd snel wat besproken worden. Daarnaast vonden we ons in groter gezamenlijk doel: om de (bedrijfs)gezondheidszorg en arbeidsomstandigheden te (blijven) verbeteren. Dankzij alle gesprekken over wat dan ook bleef ik scherp omdat er altijd een inzicht vanuit een andere hoek was die interessant bleek. Ook veel dank naar de collega's van het secretariaat, Bibi en Els. De ondersteuning die ik daar heb ervaren heeft me zeer geholpen.

Daarnaast wil ik nog in het bijzonder noemen de collega's met wie ik veel leuke congressen heb meegemaakt. Fietsen door Dublin, een voetbalwedstrijd van donkeys bezoeken, wandelingen over ontzettend veel trappen naar een plek met prachtig uitzicht en de vele etentjes waren allemaal hoogtepunten van de afgelopen 5 jaar. Paul, Karen, Maryam, Michiel, Anne-Claire, Marloes, Henk, Carlien, en de andere

collega's die ik ook elders heb vermeld, dankzij deze congressen zijn we als collega's dichter naar elkaar gegroeid en hebben we elkaar beter leren kennen. Dankjulliewel voor deze gezellige en leerzame perioden. Daarnaast komt er in ieder geval nog een congres aan waar ik met veel plezier naar uitkijk.

Dear Mats, Adnan and Sara, thank you for welcoming me in your home, Sahlgrenska universitetssjukhuset Arbets- Och Miljömedicin. The first morning you gave me the keys to the building, 'to work evenings and weekends'. In those two weeks I learned a new statistical method and produced an almost completed manuscript. Thank you for supervising and teaching me during those two weeks.

Naast mijn werk bij het Coronel Instituut werkte ik ook twee avonden bij het USC. Een perfecte afwisseling van het soort werk. In de sportschool kon ik me fysiek lekker uitleven waar ik juist in het AMC me mentaal kon pijnigen. Mijn collega's van het USC, in het bijzonder Koen en Emile, vroegen bijna elke week: 'Hoe gaat het met je promotietraject?'. Waarop mijn standaardantwoord vaak 'te langzaam' was.

Daarnaast heeft de tijd in Maastricht mij gevormd tot wie ik nu ben. Ik heb dankzij Nucleus Durus vrienden voor het leven gemaakt die ik nu nog vaak zie. Dankzij jullie ben ik wie ik nu ben. In deze periode hebben we elkaar goed leren kennen, en weten we wat we aan elkaar hebben. Altijd is er altijd een helpende hand wanneer het nodig was. Verhuizen, bruiloft, kinderen en natuurlijk de etentjes en borrelavonden. Rolf, Roeland, Jelle, Joris, Jasper, Jasper, Joep, Jeroen, Vincent, Dirk, Jakko en jullie betere helften natuurlijk, heel erg bedankt voor alles. Ook kapitein Jurgen, bedankt voor onze vele bbq boottochtjes over de Amsterdamse grachten. Inmiddels ken ik Amsterdam beter vanaf het water dan vanaf het land. In het bijzonder wil ik Job nog bedanken die me altijd aanspoorde om (door) te studeren. Het heeft geholpen.

Daar waar ik nu sta komt met name door mijn familie, in het bijzonder mijn ouders. Ze stonden, en staan nog steeds altijd voor me klaar. Ik weet dat jullie soms hebben getwijfeld aan jullie opvoedkundige kwaliteiten, maar dat is nergens voor nodig. Jullie hebben het uitstekend gedaan, en wel 4 keer. Heel erg bedankt voor alle liefde die jullie mij gegeven hebben, ik had het niet anders gewild.

Iets korter in mijn leven, maar niet minder belangrijk zijn mijn broertje en zusjes. Koen, Mijke en Tinka, ik besef dat jullie het niet altijd makkelijk hebben gehad. Ik was altijd 'ver' van huis, eerst Maastricht, daarna Amsterdam en nu Utrecht. Daarnaast heb ik voor mezelf altijd de lat zo hoog mogelijk gelegd, me niet realiserend dat jullie dat vaak ook als jullie lat konden ervaren. Ik ben ontzettend trots op hoe jullie met elke situatie omgaan, en wat jullie al hebben bereikt. Hopelijk blijft onze band net zo goed als deze is gebleven ondanks de afstand, en ik zal ook altijd voor jullie klaar blijven staan. Daarnaast brengt Femm heel veel vreugde in onze levens. Femm, ik kijk altijd zo uit naar trips naar Brabant en het vrolijke (en nu ook steeds vaker ondeugende)

gezicht van jou. Ik kijk er naar uit dat wanneer je ouder bent, je kunt komen logeren en spelen, en we samen de wereld gaan veroveren.

Een van de belangrijkste personen tijdens de periode van mijn PhD is Judith. Judith, dankjewel voor alles. Altijd was je er voor me. Of het nu 7 uur in de ochtend was wanneer ik op onze afdeling aankwam, of wanneer ik er een keer tot laat was, jij was er altijd. Altijd was er de mogelijkheid om aan te kloppen bij je kamer, waar letterlijk de deur altijd open stond. Je hebt me zo ongelooflijk veel geleerd. Zelfs toen duidelijk was dat je ziek was en niet lang meer te leven had, ben je zo vaak mogelijk naar het AMC gekomen, en heb je me, toen het echt niet anders meer kon, vanuit huis tot het einde begeleid. Al voor deze periode gaf je me zelfvertrouwen en het gevoel van zelfstandigheid. Vlak voordat je kwam te overlijden ben je nog naar mijn kamer toe gekomen. Om te zeggen dat je vertrouwen in me had, en zag dat het allemaal wel ging lukken, ook zonder jou. Met daarbij de toevoeging dat je van boven mee zou blijven kijken. Deze is ook voor jou.

Last but not least, Sophie, dankjewel voor alles wat je voor me gedaan hebt en voor me doet. Ik voel me gelukkig met en door jou. Sorry voor alle keren dat ik hard aan het werk was en geen tijd voor je had, ik maak het snel goed. Je hebt het voor me over om, ondanks je hekel aan vliegen, me op te komen zoeken in Zweden. Dat doet me heel veel. Als ik zie hoe je met iedere situatie omgaat, denk ik dat ik nog heel veel van je kan leren. Gelukkig hebben we nog een lange tijd om naar uit te kijken. Ik kijk al vooruit naar alle leuke dingen en avonturen die we samen gaan beleven. We zijn een goed team.

LIST OF HOUSES:

Front cover	The Netherlands, Amsterdam, view from Amstel river
Chapter 1	Sweden, Southern archipelago of Gothenburg
Chapter 2	Ireland, Cobh
Chapter 3	Ireland, Killarney, Ross Castle
Chapter 4	Ireland, Kerry, Ardtully Castle
Chapter 5	Sweden, Southern archipelago of Gothenburg
Chapter 6	Belgium, Bruges, St Bonifacius bridge
Chapter 7	Italy, Bologna, view from Asinelli tower
Chapter 8	The Netherlands, Utrecht, Kasteel de Haar
Chapter 9	Malta, Popeye village
Chapter 10	The Netherlands, Oud-Zuilen, Slot Zuylen
Chapter 11	Ireland, Cork, Sheep's head
Back cover	The Netherlands, Utrecht, Kasteel de Haar